



REVIEW

Exploring long-term effects of biochar on mitigating methane emissions from paddy soil: a review

Qiong Nan¹ · Liqing Xin¹ · Yong Qin¹ · Muhammad Waqas² · Weixiang Wu¹

Received: 27 November 2020 / Accepted: 30 March 2021 / Published online: 5 May 2021
© The Author(s) 2021

Abstract

Biochar has been reported to mitigate short-term methane (CH₄) emissions from paddy soil. Currently, CH₄ mitigation by biochar has primarily focused on the abundance and variations of methanogens and methanotrophs, and changes in their activities during methane production and consumption. However, long-term effects of biochar on methane mitigation from paddy soil remain controversial. This review overviewed the existing mechanisms for CH₄ mitigation as a result of biochar application. In addition, the two existing opinions on the long-term CH₄ mitigation effect upon biochar application were highlighted. Combining the already explored mechanisms of fresh biochar on CH₄ mitigation from paddy soil and a novel discovery, the potential mechanisms of biochar on long-term methane emission response were proposed. This review also revealed the uncertain responses of biochar on long-term CH₄ mitigation. Therefore, to achieve carbon neutral goal, it is important to further explore the mechanisms of long-term CH₄ mitigation under biochar application.

Keywords Biochar · Methane mitigation · Methanogens · Methanotrophs

1 Introduction

Global warming (GW) is the rise in the average earth surface temperature as a result of increase in the concentration of greenhouse gases (GHGs) including methane (CH₄), nitrous oxide (N₂O), water vapor, ozone (O₃), chlorofluorocarbons (CFCs) and carbon dioxide (CO₂) (Al-Ghussain 2019). Basically, it is the greenhouse effect that made the earth a suitable place to live. Without GHGs, there would be no life on earth because the surface temperature would be too low to live on (Anderson et al. 2016). However, the exponential increase in the concentration of GHGs in the atmosphere results in the catastrophic phenomenon, which is termed as GW. Among the GHGs, CH₄ is one of the most widespread greenhouse gases emitted from wetlands, paddy fields, coal mines, ruminants and anthropogenic activities

including livestock raising and leakage from natural gas systems (Waqas et al. 2020). Methane was expected to contribute GW around 18% over next 50 years after 1999 (Milich 1999). Whereas, GW contribution of CH₄ emissions over 27% worldwide in 2015 (Ma et al. 2019). In the recent years, it has been noticed that CH₄ emission from wetlands has significantly increased to about 164 Tg year⁻¹ (Bridgham 2013; Singh 2017). The total occupied earth's surface by wetlands accounts to about 3.8%, with the total global CH₄ emissions of 20–40% (Tiwari et al. 2020). Among the wetlands, paddy fields are considered as a significant contributor of CH₄ emission (Xiao et al. 2018). In the flooded paddy soils, CH₄ is produced by a group of bacteria termed as methanogens. The flooded paddy soils, such as rice fields, restrict the oxygen (O₂) supply to the soil and produce anaerobic conditions that result in the fermentation of soil organic matter and release CH₄ to the atmosphere. The release of CH₄ from deeper layer of flooded soil to the atmosphere is carried out by ebullition, diffusion, and through aerenchyma conduits of paddy plant (Singh 2017). The recent report by Food and Agriculture Organization Corporate Statistical Database (FAOSTAT) demonstrated that among the agricultural soil all around the world, the rice cultivation contributed up to 20% of the global CH₄ emissions by the end of year of 2018

✉ Yong Qin
qinyong@zju.edu.cn

¹ Institute of Environmental Science and Technology, College of Environment and Resource Science, Zhejiang University, Hangzhou 310029, People's Republic of China

² Department of Botanical and Environmental Science, Kohat University of Science and Technology, Kohat 26000, Khyber Pakhtunkhwa, Pakistan

(FAOSTAT 2020). Hence, it is imperative to reduce methane emissions from paddy fields.

Biochar has been well reported as a promising material for mitigating GHGs especially for CH₄ emissions from paddy soil (Awad et al. 2018; Han et al. 2016; Wu et al. 2019a). Biochar is a black-colored carbon (C) rich material produced as a result of thermal decomposition of biomass under O₂ deficient conditions (Mia et al. 2017a; Waqas et al. 2018). The potential of biochar has been extensively explored due to its benefits in mitigation of CH₄ emissions (Nan et al. 2020b; Pratiwi and Shinogi 2016; Yoo et al. 2016). Ji et al. (2020) reported significant reduction of CH₄ emissions using biochar during an incubation experiment conducted for 77 days. In addition, the low rate (2.8 t ha⁻¹) application of biochar on paddy fields has also been reported to reduce CH₄ emissions by 41% for the short-term application (Nan et al. 2020a, b). Furthermore, a meta-analysis on biochar also revealed that the soil with application of various types of biochar significantly reduce CH₄ emissions (Awad et al. 2018). These results suggest that the ecological benefit of biochar application on CH₄ emission has been widely demonstrated.

The potential of biochar to mitigate CH₄ emission from paddy soil has been well reported in the pot experiments and short-term field studies. In contrast, the observations from long-term field effects, especially after years of biochar aging, are seldom reported. Hence, most of the mechanisms

for CH₄ mitigation are limited to the application of fresh biochar. The mechanisms developed by fresh biochar on mitigation of CH₄ emissions from paddy soil mainly focus on the impacts on soil physicochemical structure and microbial dynamics. The reductions in CH₄ emission as a result of biochar addition are mainly attributed to lowering the ratio of methanogens/methanotrophs (*mcrA/pmoA*) (Table 1). In addition to the substantial reduction in the ratio of methanogens/methanotrophs, the other specified mechanisms are either significantly decreased activities of methanogens (Dong et al. 2013; Han et al. 2016) or substantially increased activity changes of methanotrophs (Han et al. 2016; Nan et al. 2020a; Wu et al. 2019a; Yang et al. 2013). On the other side, few studies also demonstrated the potential of biochar to mitigate CH₄ emission after years of aging process; however, their results are in controversy with similar studies (Nan et al. 2020a; Wang et al. 2019c; Wu et al. 2019a) and the exact mechanisms are still blurred. Due to lacking the long-term continuous CH₄ emission observation under biochar application in paddy fields, it is difficult to evaluate CH₄ mitigating effect of biochar amendment scientifically and realistically over decades scale. Hence, the efficacy and the mechanism of biochar in mitigating CH₄ emissions after years of aging in paddy fields remain to be explored.

The present review offers a comprehensive overview to fully understand the long-term effects of biochar amendment and the potential mechanisms in reducing CH₄ emissions

Table 1 Mechanism of biochar on methane mitigation from paddy soil

Feedstocks	Temperature (°C)	Experiment type	Experiment duration (year)	Effective-ness duration (year)	Mechanism	References
Wheat straw	450	Incubation	–		Decreased methanogens and increased methanotrophs	He et al. (2020)
Rice straw	500	Incubation	–		Decreased methanogens	Ji et al. (2020)
Wheat straw	~500	Incubation	–		Increased methanotrophs	Wu et al. (2019b)
Rice straw	~500	Incubation	–		Decreased methanogens	(Cai et al. 2018)
Rice straw	500	Pot	1		Decreased methanogens and increased methanotrophs	Han et al. (2016)
Rape straw	500	Pot	2	2	Decreased methanogens	Qi et al. (2021)
Rice straw	–	–	–		Increased methanotrophic activity	Singh (2017)
Tobacco straw	450	Field	1		Decreased methanogens and increased methanotrophs	Huang et al. (2019)
Wheat straw	500	Field	3	2	Improved soil aeration and Eh induced enhanced methanotrophic activity	Chen et al. (2018)
Rice straw	500	Field	4	2	Increased methanotrophs	Nan et al. (2020a)
Rice straw	500	Field	4	4	Increases in soil dissolved organic carbon, NH ₄ ⁺ -N, and porosity induced larger increase of methanotrophs than methanogens in the first year suppressed the abundance of methanogens for the rest three years	Wang (2019a)

from paddy soil. Emphasis was given on the current explored mechanism of fresh biochar application on CH₄ mitigation. Furthermore, the existing results of biochar on CH₄ emission after long-term application were explored. Finally, the potential impacts of biochar aging on CH₄ emission and underlying mechanism were briefly discussed.

2 Effects of biochar application on methanogenic activity

2.1 Inhibition of methanogenic activity through decreasing dissolved organic carbon

Most of the published research studies suggested that biochar applications into paddy soil inhibit the numbers and activities of methanogens by decreasing the soil dissolved organic carbon (DOC). Soil DOCs are important substrates for methanogens and thus for CH₄ production (Conrad 2007). The high porous biochar material is composed of a large portion of recalcitrant (aromatic) C and small fraction of labile C (Zimmerman 2010). Yu et al. (2012) reported that when biochar was applied into paddy field, the soil DOC was decreased, which accounted to the adsorption via biochar pore. Similarly, Zheng et al. (2016) demonstrated that contents of DOC were significantly decreased by 52% and 71% under biochar amendment at 20 and 40 t ha⁻¹, respectively. In other study, Liu et al. (2011) conducted an experiment to explore the bamboo chip and rice straw biochar (BC and SC) amendment on CH₄ mitigation. Their result depicted that both BC and SC biochars significantly reduced CH₄ emission during incubation period by decreasing methanogenic activity derived by methanogens substrates. In addition, Han et al. (2016) also observed decreased methanogenic activity and soil DOC content when biochar was amended into paddy soil. Therefore, reducing soil DOC content as a result of biochar amendment is an important mechanism on inhibition methanogenic activity.

2.2 Inhibition of methanogenic activity through increased oxygen input

An important mechanism to inhibit the activities of methanogens in paddy soil is the high porosity of biochar that increases the O₂ flux due to its high porosity and hence increases the toxicity to methanogens. Furthermore, being a nutrient-rich source (Chen et al. 2020; He et al. 2019), biochar application into paddy soil promotes rice roots growth (Xiang et al. 2017) and thus increases O₂ secretion (Dong et al. 2013; Ma et al. 2010; Zhao et al. 2014). Consequently, the activities of methanogens and methanogenic activity are inhibited. Kim et al. (2017) examined the effect of biochar on rice yield and CH₄ emission and reported that biochar

amendment increased rice yield and reduced CH₄ emission by inhibiting methanogenesis through enhancing the soil aeration and O₂ availability.

Correspondingly, biochar amendment into paddy also decreases soil bulk density and increases soil aeration and thus inhibits methanogenic activity. The responsible factor for reducing the soil bulk density is the high porosity of biochar (Waqas et al. 2018). Carvalho et al. (2014) illustrated that even 3 years application of 0.5 and 1.6% eucalyptus timber biochar into sandy loam significantly increased soil total porosity. The pot experiment of Peake et al. (2014) demonstrated that 2.5% biochar application decreased bulk density by circa 4.2–19.2%. Devereux et al. (2012) demonstrated amelioration in bulk density with biochar application rate of 5% (w/w), as Githinji (2014) observed in sandy loam and Mukherjee et al. (2014) observed in an artificially degraded soil. In addition, they also stated that methanogenic surviving cells dropped exponentially upon exposure to the increasing O₂ level in the soil as a result of biochar addition (Kiener and Leisinger 1983). The concept figure of biochar inhibition on methanogenic activity is shown in Fig. 1a.

2.3 Promotion of methanogenic activity

Few studies on soil biochar application also showed promotion of methanogenic activity. The supporting concept behind promotion of methanogenic activity is that biochar amendment increases soil NH₄⁺-N (Wang et al. 2019b). Hence, the ammonia-based fertilizers promote the growth of both methanogens and methanotrophs (Wang et al. 2019b). Further, biochar application may increase rhizo deposit and plant litter as substrates for methanogens due to increased rice biomass (Banger et al. 2012). Moreover, biochar amendment also increases soil pH and is capable of ameliorating the acidic nature of paddy soil (Zhang et al. 2018). The increased soil pH is beneficial to both methanogens and methanotrophs (Le Mer and Roger 2001). However, methanotroph is more sensitive to soil pH and is promoted to a larger extent compared to methanogens (Hanson and Hanson 1996; Jeffery et al. 2016), which leads to a lower ratio of methanogens/methanotrophs. Therefore, the lower ratio of methanogens/methanotrophs will lead to the reduction in CH₄ emission. The similar case has been reported from the results of Wang et al. (2019b), who observed that both the *mcrA* and *pmoA* copy numbers were higher than the control for the first year of biochar application, while with time, the ratio of *mcrA/pmoA* got lower in biochar treatments than the control.

Similarly, the other methanogenic activity promotion occurred when biochar under low pyrolysis temperature (300 °C) was applied into paddy soil. In this case, biochar application into paddy would stimulate CH₄ emission rather




Fig. 1 Biochar effect on methanogenic inhibition (a) and methanotrophic (b) activity in the paddy soil

than mitigate. Cai et al. (2018) demonstrated that biochar produced at 300 °C significantly increased paddy CH₄ emissions in comparison to the control treatment. Conversely, the CH₄ emissions were significantly reduced when biochar was produced at 500 and 700 °C, respectively. The responsible mechanism of promoting methanogenic activity under low pyrolysis temperature is the aromatic structure that is not formed well under low temperature (< 500 °C, with the high production of fulvic acids and humic acids at lower temperature (300 °C)) (Mia et al. 2017a). Therefore, biochar produced at 300 °C is largely applied to increase soil DOC and C source for methanogens (Ji et al. 2020). Singla and Inubushi (2014) reported that application of biochar produced at 300 °C significantly increased CH₄ emissions due to an increase in the soil DOC and significant growth of methanogens. In contrast, the biochar produced at elevated temperature has more stable aromatic structures (Guo and Chen 2014) and enrichment porosity (Ippolito et al. 2020), hence will not increase soil DOC due to its ability to absorb soil DOC. The comprehensive mechanism of methanogenic activity promotion as a result of biochar addition is shown in the given Fig. 1b.

3 Impact of biochar application on methanotrophic activity

3.1 Biochar impact on promoting the aerobic methane oxidation activity

Promotion of methanotrophic activity under biochar amendment has been widely recognized. This promotion mainly contributes to physical (porosity and high surface

area) and chemical (alkaline nature, functional groups, nutrients supply) characteristics of biochar. The porous structure of biochar has been well reported to adsorb CH₄ and provide habitat for methanotrophs (Ji et al. 2020) and ameliorate soil aeration, thus stimulating methanotrophic activity. The alkaline nature of biochar increases soil pH to an optimum range for methanotrophs (Wang et al. 2019c). Likewise, biochar amendment as nutrients supplement also benefits the plants by increasing the rice yield increase and root growth, thus bringing more O₂ through rhizosphere (Dong et al. 2013; Watanabe et al. 1997). The figure of biochar promotion on methanotrophic activity is shown in Fig. 2.




Fig. 2 Biochar effect on methanotrophic activity in the paddy soil

3.1.1 Biochar porous structure on methane mitigation responses

The high porous characteristic of biochar contributes to CH₄ adsorption and soil aeration amelioration. Rising pyrolysis temperature is accompanied with the solid matrix shrinking, various organic compounds volatilizing, and transformation of relatively large pores to smaller ones (de Melo Carvalho et al. 2014; Weber and Quicker 2018). Thus, the porosity and overall specific surface area (SSA) of biochar increases with increasing pyrolysis temperature. High porosity and large SSA of biochar exert a function of CH₄ adsorption and also provide habitat for methanotrophs (Liu et al. 2011). Further, biochar adsorption could also be benefit for nutrients retention and thus promote O₂ delivery from rice rhizosphere aerenchyma tissues. In addition, high porosity of biochar also increases the exposure of methanogens to O₂ through ameliorating soil aeration and reducing bulk density as discussed earlier (de Melo Carvalho et al. 2014; Mukherjee et al. 2014; Peake et al. 2014). The research findings of Conrad and Rothfuss (1991) on biochar application in the paddy field illustrated that 80% of produced CH₄ was consumed by methanotrophs through diffusion. Therefore, biochar porosity plays pivotal roles in enhancing aerobic methanotrophic activity in the paddy soil.

3.1.2 Liming effect of biochar on methane mitigation responses

The increase of soil pH as a result of biochar addition is another important mechanism for the promotion of methanotrophic activity. The alkaline nature of biochar is mainly due to the ash content, solid phase hydroxide and carbonate phases of biochar (Ippolito et al. 2020). In the study of Si et al. (2018), biochar when applied at the rate of 2.5 t ha⁻¹ significantly increased soil pH due to liming effect. Similarly, a meta-analysis also reported that biochar application increased 11.5–11.9% of soil pH (Awad et al. 2018). Correspondingly, the optimum pH for methanotrophs is 5.0–6.5 and methanotrophs are very sensitive to the pH changes (Jeffery et al. 2016). The pH of the paddy soil is usually around 5; thus, biochar incorporation into paddy soil enhances the methanotrophic activity due to liming effect. Weber and Quicker (2018) observed a large decrease on methanotrophic when soil pH decreases from 6.3 to 5.6. Generally, at higher pyrolysis temperature, the biochar produced is higher alkaline and hence its application significantly promotes the activity of methanotrophic in acid paddy soil (Wang et al. 2014). In addition, other cooperation of liming effect of biochar with methanotrophs is lowering Al³⁺ availability, thus providing protection to methanotrophs because lower pH Al³⁺ is highly toxic to methanotrophs (Tamai et al. 2007). Therefore, maintaining the liming effect duration of biochar

can prolong the promotion of biochar on methane oxidation activity (Jeffery et al. 2016).

3.2 Promotion of anaerobic methane oxidation activity

Recent studies explored the positive effects of biochar on anaerobic CH₄ oxidation activity induced by the electronic accepting capacities (EAC) of biochar. The surface of biochar contains various oxygen functional groups like carboxyl, carbonyl, quinone phenolic hydroxyl groups (Wu et al. 2016). The O-containing functional groups are the main entities responsible for biochar redox potential (Klupfel et al. 2014). Specifically, carbonyl and quinone are capable of accepting electron and determine the EAC properties of biochar (Zhang et al. 2019b). Thus, theoretically, carbonyl and quinone moieties can act as an electronic acceptor and consume CH₄ in the paddy. Zhang et al. (2019a) reported the anaerobic CH₄ oxidation due to the presence of quinone structure (C=O) using biochar. CH₄ anaerobic oxidation can account for 50% of the total CH₄ consumption in wetland environments (Segarra et al. 2015). Hence, biochar might mitigate CH₄ emission partly through anaerobic CH₄ consumption due to its EAC in paddy soil. Additionally, the aromatic structure of biochar can also act as a good electron acceptor due to the conjugated π -electron systems and thus may contribute to CH₄ consumption. However, the condensed aromatic structure of biochar usually formed as a result of higher pyrolysis temperature (above 700 °C), which is not usually used in paddy soil considering the agricultural ecosystem benefits (McBeath et al. 2011).

4 Long-term effect of biochar on methane mitigation in paddy soil

4.1 Short-term effectiveness of biochar mitigation responses from paddy soil

Two cases showed a short-term methane mitigation effectiveness with biochar amendment. Liu et al. (2019) conducted a six-year field observation from 2010 to 2015 and the result showed that biochar at 20 and 40 t ha⁻¹ application rate reduced CH₄ emission in the first year only. Likewise, a four-year field study conducted by Nan et al. (2020a) also observed the significant CH₄ mitigation responses only for two years after the biochar application at 22.5 t ha⁻¹ to the paddy soil. They also reported that no significant differences were shown in CH₄ mitigation for the following two years. The vanished CH₄ mitigation effect in the subsequent two years was ascribed to the lowered impact of biochar on methanotrophic activity. Spokas (2013) also reported that biochar aging decreased the ability to promote methanotroph

activity. These studies suggested that biochar provided limited and temporary benefits on methane mitigation over a long term. To maximize the long-term CH₄ mitigation effect of biochar from paddies, the mechanism underlying the mitigation effect of aged biochar needs to be further explored.

4.2 Long-term effectiveness of biochar mitigation responses on paddy soil

Few studies reach the consensus that biochar application can mitigate CH₄ emission for a prolonged period in paddy soil. A detailed prolonged (> 4 years) research of biochar application on various soils has been conducted, which found biochar application is positive to CH₄ mitigation into paddy soil. Several trials were established in 2012 at various locations i.e., Huizhou (Qin et al. 2016), Changsha (Wang et al. 2018), Nanjing (Wu et al. 2019b), respectively. From the 4-year CH₄ mitigation effectiveness of biochar application, Wang et al. (2018; 2019c) revealed the dynamic mechanism behind this response. They reported that biochar stimulated both the methanogens and methanotrophs abundance, but with a higher promotion on methanotrophs, which caused the CH₄ reduction in the first year. However, for the subsequent three years, biochar showed an inhibition on methanogens abundance probably attributing to increased soil aeration and no effect on methanotrophs, thus mitigating CH₄ emission. These findings were in agreement with those of Wang and Qin et al. (2016), who also suggested that, three out of four years, biochar application at 5 and 20 t ha⁻¹ significantly mitigated CH₄ emission. The related mechanism was roughly ascribed to promotion of methanotrophs and the improved soil aeration, bulk density and pH. In addition, in a 6-year field experiment, Wu et al. (2019b) concluded that biochar application at 20 and 40 t ha⁻¹ reduced CH₄ emission by 11.2% and 17.5% on average 6 years, respectively. However, in this study, none cumulative methane difference analysis was conducted on single experimental year. Therefore, the 6-year long-term effectiveness of biochar on methane mitigation is doubtful.

Even there exists field experiment which demonstrated 4 years of effectiveness in CH₄ mitigation with biochar application, high rate (> 10 t ha⁻¹) (Awad et al. 2018) biochar application only in the first amendment year may not prolong methane mitigation effect in paddy soil. First, CH₄ mitigation only 1 or 2 years of effectiveness of biochar application on CH₄ mitigation was also reported. Hence, long-term (> 4 years) observation of biochar amendment on CH₄ mitigation should be widely conducted. Theoretically, in terms of high-rate biochar application in single year strategy, it is hard to prolong CH₄ emission reduction effectively. First, biochar particle size must be smaller and smaller under agricultural activities and aging process (Martin et al. 2012; Mia et al. 2017a). In addition, biochar liming effect would

decrease gradually, which would lose the ability to ameliorate soil pH, especially for acid paddy soil. Furthermore, biochar degradation becomes faster in paddy soil under the plant growth ambient, which means that recalcitrant carbon of biochar would change to liable carbon more quickly and with increment in amount. A nine-year study on biochar physicochemical changes and transformation from paddy soil conducted by Yi et al. (2020) demonstrated that aromatic carbon (recalcitrant carbon) of biochar decreased by 5.0% in bamboo biochar and 8.7% in rice straw biochar, respectively. Smaller biochar particle size and the gradually lost liming effect would weaken the promotion of biochar on methanotrophic activity, while the accelerated biochar degradation would enhance methanogenic activity. Hence, it is probably that, after years of aging process, biochar will lose the ability to mitigate CH₄ emission in paddy soil. However, the biochemical processes in soil and the interaction between biochar and soil biochemical factors are such complex. The actual long-term CH₄ mitigation effect exerted by biochar amendment should be further explored.

4.3 Potential impact of aged biochar on methanotrophic and methanogenic activities

Though the few studies reported short-term effectiveness of biochar on CH₄ mitigation, the mechanism behind this phenomenon is still unclear. The lower potential of biochar for CH₄ emission may also be accompanied with aging process. In paddy soil, the force of agricultural activities and crop growth accelerate physicochemical changes, plowing and tillage would expose biochar to air for oxidation and experience physical breakdown (Mia 2017a; Yi et al. 2020). Biochar would also be oxidized by O₂ secreted from crop roots (Joseph et al. 2010). Accordingly, the exploration of the mechanism should first focus on changes in biochar attributes over time in field.

4.3.1 Changes in the porosity and its impact on methanotrophic activity

The blockage and breaking of pores on biochar surface may contribute to lowering the promotion of methanotrophic activity. As a result of continuous farming practices, such as rice growing, the biochar particles become smaller and the pore structure may be blocked or in-filling with SOM (Martin et al. 2012). As a result, the smaller size and blocked pore of biochar would probably reduce the improvement of soil aeration. This would weaken the positive promotion of biochar on methanotrophic activity. However, there are also some studies which reported that with the process of biochar aging, the porosity and SSA increased significantly. Through a 5-year field experiment, Dong et al. (2017)

found decreased average diameter of biochar pores through increasing new small pores and 98% to 114.3% increase in SSA. The increased porosity and SSA would probably increase CH₄ adsorption and thus adhere more substrates to methanotrophs. Further, the increased porosity would increase nitrogen retention ability and benefit the methanotrophs. Wang et al. (2020) also illustrated that the porosity and SSA increases lead to the maximum adsorption of NH₄⁺-N. However, porous structure influences on methanotrophic activity with aging process need a comprehensive assessment.

4.3.2 Reduction in biochar liming effect and its impact on methanotrophic activity

The gradual reduction in the liming effect of biochar may contribute to lowering the activity of methanotrophs. As discussed, the alkaline nature of biochar comes from ash content composed of oxide/carbonate minerals of various elements, such as phosphorus, potassium and calcium (Ippolito et al. 2020). The ash content also acts as a nutrient supplement and is easily absorbed by the plants when dissolved in water. During rice tillering and jointing growth stage (flood period), the ash content of biochar dissolves and leaches, leading to the gradual disappearance of the liming effect (Chang et al. 2019; Lou et al. 2012). The same has been observed by a four-year study of Wang et al. (2019c) who reported that soil pH was constantly decreasing in the subsequent years. In addition, Nan et al. (2020a) also conducted a four-year field trial and observed the weak liming effect that probably lost the effect of ameliorating acid environment for methanotrophs and methanotrophic activities.

4.3.3 Biochar electronic transfer ability and its impacts on methanogenic and methanotrophic activities

Changes in oxygen functional groups during aging process affect the electronic transfer ability of biochar and thus exert difference on methanogenic and methanotrophic activities. Biochar experiences accelerated aging process in paddy than that without agricultural activities and introduces more oxygen functional groups with aging process. Various studies reported an increase of oxygen functional groups like hydroxyl, carbonyl, carboxyl, ketone and phenol groups in natural aged biochar (Cheng et al. 2008; Mia et al. 2017b; Qian and Chen 2014). It has been reported that after four months of soil incubation, biochar quinone content decreased (Mukome et al. 2014). Quinones and carbonyl mainly contribute to EAC, and phenolic OH is the main electron donating ability (EDC) source. Thus, changes in quinones and carbonyl functional group content inevitably affect methanotrophic processes (Klupfel et al. 2014; Zhang et al. 2019a). Even no studies illustrated methanogenic

activity through biochar EDC in paddy system, plenty of studies have demonstrated the promotion effect through EDC of biochar in anaerobic digestion trail (Shao et al. 2019; Shen et al. 2017). The oxygen functional groups induced EDC increase may act on methanogenic activity promotion. Thus, calculation of EAC and EDC variation may help a lot on long-term methane mitigation responses of biochar revealing.

5 Prospects for future works

5.1 Necessity of long-term field experiments

As discussed, recent studies mainly focus on short-term CH₄ mitigation effectiveness of biochar application and lack the long-term field experiment observation on CH₄ mitigation effect. Currently, very few long-term (> 4 years) studies exist with controversy results of the CH₄ mitigation. To achieve the C peak in 2030 and C neutral in 2060, exploring biochar strategies that prolong CH₄ mitigation effectiveness is of great importance. Whereas, the current studies failed to underpin a pleasant strategy that guarantees stable and long-term CH₄ mitigation effect. Consequently, to provide theory basis and help with drafting the long-term stable effectiveness of biochar application on CH₄ mitigation strategy, long-term and *in-situ* field experiments focus on CH₄ mitigation from paddy should be conducted widely.

5.2 Key factors of biochar influencing methane mitigation

To figure out the exact mechanism of biochar aging process on long-term CH₄ mitigation effect, the key factors of biochar attributes on CH₄ mitigation should be specified. Currently, the mechanisms about biochar effect on CH₄ mitigation usually focus on the comprehensive responses of methanogenic and methanotrophic activities after biochar amendment. Researches can attribute CH₄ mitigation effect to methanogenic activity inhibition or methanotrophic activity promotion. The reason of biochar on methanogenic and methanotrophic activities was developed mainly on soil properties change (soil pH, DOC, bulk density and soil aeration etc.). However, detailed studies on biochar physical (porosity, surface area), chemical properties (oxygen functional groups, electronic transfer ability) and their significance on CH₄ mitigation are lacking. Combined with biochar physiochemical attributes changes during biochar aging process, the mechanism behind the long-term CH₄ mitigation effect under biochar amendment would be easier to illustrate. The specification of biochar attributes on CH₄ mitigation will also help to develop novel biochar materials

and strategy management for enhancing long-term CH₄ mitigation effect.

Acknowledgements This research was supported by National Natural Science Foundation of China [Grant numbers 42077032 and 41571241] and the National Key Technology Research and Development Program of the Ministry of Science and Technology of China [Grant number 2015BAC02B01]. We thank the editors and reviewers for their constructive comments.

Declarations

Conflict of interest The authors declare no conflicts of interest.

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

References

- Al-Ghussain L (2019) Global warming: review on driving forces and mitigation. *Environ Prog Sustain Energy* 38:13–21
- Anderson TR, Hawkins E, Jones PD (2016) CO₂, the greenhouse effect and global warming: from the pioneering work of Arrhenius and Callendar to today's earth system models. *Endeavour* 40:178–187
- Awad YM, Wang J, Igalavithana AD, Tsang DCW, Kim K-H, Lee SS et al (2018) Chapter One—biochar effects on rice paddy: meta-analysis. In: Sparks DL (ed) *Advances in agronomy*. Academic Press, pp 1–32
- Banger K, Tian H, Lu C (2012) Do nitrogen fertilizers stimulate or inhibit methane emissions from rice fields? *Glob Change Biol* 18:3259–3267
- Bridgman SDC-Q, Hinsby K, Jason K (2013) Methane emissions from wetlands: biogeochemical, microbial, and modeling perspectives from local to global scales. *Glob Change Biol* 19:1325–1346
- Cai F, Feng Z, Zhu L (2018) Effects of biochar on CH₄ emission with straw application on paddy soil. *J Soils Sediments* 18:599–609
- Chang R, Sohi SP, Jing F, Liu Y, Chen J (2019) A comparative study on biochar properties and Cd adsorption behavior under effects of ageing processes of leaching, acidification and oxidation. *Environ Pollut* 254:113123
- Chen D, Wang C, Shen J, Li Y, Wu J (2018) Response of CH₄ emissions to straw and biochar applications in double-rice cropping systems: insights from observations and modeling. *Environ Pollut* 235:95–103
- Chen L, Liu M, Ali A, Zhou Q, Zhan S, Chen Y et al (2020) Effects of biochar on paddy soil fertility under different water management modes. *J Soil Sci Plant Nutr* 20:1810–1818
- Cheng C-H, Lehmann J, Engelhard MH (2008) Natural oxidation of black carbon in soils: changes in molecular form and surface charge along a climosequence. *Geochim Cosmochim Acta* 72:1598–1610
- Conrad R (2007) Microbial ecology of methanogens and methanotrophs. *Advances in agronomy*, vol 96. Academic Press, pp 1–63
- Conrad R, Rothfuss F (1991) Methane oxidation in the soil surface layer of a flooded rice field and the effect of ammonium. *Biol Fertil Soils* 12:28–32
- de Melo Carvalho MT, de Holanda A, Madari BE, Bastiaans L, van Oort PAJ, Heinemann AB et al (2014) Biochar increases plant-available water in a sandy loam soil under an aerobic rice crop system. *Solid Earth* 5:939–952
- Devereux RC, Sturrock CJ, Mooney SJ (2012) The effects of biochar on soil physical properties and winter wheat growth. *Earth Environ Sci Trans R Soc Edinb* 103:13–18
- Dong D, Yang M, Wang C, Wang H, Li Y, Luo J et al (2013) Responses of methane emissions and rice yield to applications of biochar and straw in a paddy field. *J Soils Sediments* 13:1450–1460
- Dong X, Li G, Lin Q, Zhao X (2017) Quantity and quality changes of biochar aged for 5 years in soil under field conditions. *CATENA* 159:136–143
- FAOSTAT (2020) Food and agriculture organization of the united nations (FAO)
- Githinji L (2014) Effect of biochar application rate on soil physical and hydraulic properties of a sandy loam. *Arch Agronomy Soil Sci* 60:457–470
- Guo J, Chen B (2014) Insights on the molecular mechanism for the recalcitrance of biochars: interactive effects of carbon and silicon components. *Environ Sci Technol* 48:9103–9112
- Han X, Sun X, Wang C, Wu M, Dong D, Zhong T et al (2016) Mitigating methane emission from paddy soil with rice-straw biochar amendment under projected climate change. *Sci Rep* 6:24731
- Hanson RS, Hanson TE (1996) Methanotrophic bacteria. *Microbiol Rev* 60:439
- He L, Shan J, Zhao X (2019) Variable responses of nitrification and denitrification in a paddy soil to long-term biochar amendment and short-term biochar addition. *Chemosphere* 234:558–567
- He T, Yuan J, Luo J, Lindsey S, Xiang J, Lin Y et al (2020) Combined application of biochar with urease and nitrification inhibitors have synergistic effects on mitigating CH₄ emissions in rice field: a three-year study. *Sci Total Environ* 743:140500
- Huang Y, Wang C, Lin C, Zhang Y, Chen X, Tang L et al (2019) Methane and nitrous oxide flux after biochar application in subtropical acidic paddy soils under tobacco-rice rotation. *Sci Rep* 9:17277
- Ippolito JA, Cui L, Kammann C, Wrage-Mönnig N, Estavillo JM, Fuertes-Mendizabal T et al (2020) Feedstock choice, pyrolysis temperature and type influence biochar characteristics: a comprehensive meta-data analysis review. *Biochar* 2:421–438
- Jeffery S, Verheijen FGA, Kammann C, Abalos D (2016) Biochar effects on methane emissions from soils: a meta-analysis. *Soil Biol Biochem* 101:251–258
- Ji M, Zhou L, Zhang S, Luo G, Sang W (2020) Effects of biochar on methane emission from paddy soil: focusing on DOM and microbial communities. *Sci Total Environ* 743:140725
- Joseph SD, Camps-Arbestain M, Lin Y, Munroe P, Chia CH, Hook J et al (2010) An investigation into the reactions of biochar in soil. *Soil Res* 48:501–515
- Kiener A, Leisinger T (1983) Oxygen sensitivity of methanogenic bacteria. *Syst Appl Microbiol* 4:305–312
- Kim J, Yoo G, Kim D, Ding W, Kang H (2017) Combined application of biochar and slow-release fertilizer reduces methane emission but enhances rice yield by different mechanisms. *Appl Soil Ecol* 117:57–62
- Klupfel L, Keiluweit M, Kleber M, Sander M (2014) Redox properties of plant biomass-derived black carbon (biochar). *Environ Sci Technol* 48:5601–5611
- Le Mer J, Roger P (2001) Production, oxidation, emission and consumption of methane by soils: a review. *Eur J Soil Biol* 37:25–50

- Liu Y, Yang M, Wu Y, Wang H, Chen Y, Wu W (2011) Reducing CH₄ and CO₂ emissions from waterlogged paddy soil with biochar. *J Soils Sediments* 11:930–939
- Liu X, Zhou J, Chi Z, Zheng J, Li L, Zhang X et al (2019) Biochar provided limited benefits for rice yield and greenhouse gas mitigation 6 years following an amendment in a fertile rice paddy. *CATENA* 179:20–28
- Lou L, Luo L, Cheng G, Wei Y, Mei R, Xun B et al (2012) The sorption of pentachlorophenol by aged sediment supplemented with black carbon produced from rice straw and fly ash. *Biores Technol* 112:61–66
- Ma KE, Qiu Q, Lu Y (2010) Microbial mechanism for rice variety control on methane emission from rice field soil. *Glob Change Biol* 16:3085–3095
- Ma J, Chen W, Niu X, Fan Y (2019) The relationship between phosphine, methane, and ozone over paddy field in Guangzhou. *China Glob Ecol Conserv* 17:e00581
- Martin SM, Kookana RS, Van Zwieten L, Krull E (2012) Marked changes in herbicide sorption–desorption upon ageing of biochars in soil. *J Hazard Mater* 231:70–78
- McBeath AV, Smernik RJ, Schneider MPW, Schmidt MWI, Plant EL (2011) Determination of the aromaticity and the degree of aromatic condensation of a thermosequence of wood charcoal using NMR. *Org Geochem* 42:1194–1202
- Mia SD, Feike A, Singh B (2017b) Aging induced changes in biochar's functionality and adsorption behavior for phosphate and ammonium. *Environ Sci Technol* 51:8359–8367
- Mia S, Dijkstra FA, Singh B (2017a) Long-term aging of biochar, pp 1–51
- Milich L (1999) The role of methane in global warming: where might mitigation strategies be focused? *Global Environ Change* 9(3):179–201
- Mukherjee A, Lal R, Zimmerman AR (2014) Effects of biochar and other amendments on the physical properties and greenhouse gas emissions of an artificially degraded soil. *Sci Total Environ* 487:26–36
- Mukome FND, Kilcoyne ALD, Parikh SJ (2014) Alteration of biochar carbon chemistry during soil incubations: SR-FTIR and NEXAFS investigation. *Soil Sci Soc Am J* 78:1632–1640
- Nan Q, Wang C, Wang H, Yi Q, Wu W (2020a) Mitigating methane emission via annual biochar amendment pyrolyzed with rice straw from the same paddy field. *Sci Total Environ* 746:141351–141351
- Nan Q, Wang C, Yi Q, Zhang L, Ping F, Thies JE et al (2020b) Biochar amendment pyrolysed with rice straw increases rice production and mitigates methane emission over successive three years. *Waste Manag* 118:1–8
- Peake LR, Reid BJ, Tang X (2014) Quantifying the influence of biochar on the physical and hydrological properties of dissimilar soils. *Geoderma* 235:182–190
- Pratiwi EPA, Shinogi Y (2016) Rice husk biochar application to paddy soil and its effects on soil physical properties, plant growth, and methane emission. *Paddy Water Environ* 14:521–532
- Qi L, Ma Z, Chang SX, Zhou P, Huang R, Wang Y et al (2021) Biochar decreases methanogenic archaea abundance and methane emissions in a flooded paddy soil. *Sci Total Environ* 752:141958
- Qian L, Chen B (2014) Interactions of aluminum with biochars and oxidized biochars: implications for the biochar aging process. *J Agric Food Chem* 62:373–380
- Qin X, Ye Li, Wang H, Liu C, Li J, Wan Y et al (2016) Long-term effect of biochar application on yield-scaled greenhouse gas emissions in a rice paddy cropping system: a four-year case study in south China. *Sci Total Environ* 569:1390–1401
- Segarra FS, Samarkin V, Yoshinaga MY, Hinrichs KU, Joye SB (2015) High rates of anaerobic methane oxidation in freshwater wetlands reduce potential atmospheric methane emissions. *Nat Commun* 6:7477
- Shao L, Li S, Cai J, He P, Lü F (2019) Ability of biochar to facilitate anaerobic digestion is restricted to stressed surroundings. *J Clean Prod* 238:117959
- Shen Y, Yu Y, Zhang Y (2017) Role of redox-active biochar with distinctive electrochemical properties to promote methane production in anaerobic digestion of waste activated sludge. *J Clean Prod* 278:123212
- Si L, Xie Y, Ma Q (2018) The short-term effects of rice straw biochar, nitrogen and phosphorus fertilizer on rice yield and soil properties in a cold waterlogged paddy field. *Sustainability* 10:537
- Singh CT, Boudh S, Singh JS (2017) Biochar application in management of paddy crop production and methane mitigation. *Manag Environ Pollut* 2:123–145
- Singla A, Inubushi K (2014) Effect of biochar on CH₄ and N₂O emission from soils vegetated with paddy. *Paddy Water Environ* 12:239–243
- Spokas KA (2013) Impact of biochar field aging on laboratory greenhouse gas production potentials. *Glob Change Biol Bioenergy* 5:165–176
- Tamai N, Takenaka C, Ishizuka S (2007) Water-soluble Al inhibits methane oxidation at atmospheric concentration levels in Japanese forest soil. *Soil Biol Biochem* 39:1730–1736
- Tiwari S, Singh C, Singh JS (2020) Wetlands: a major natural source responsible for methane emission. A trajectory towards a sustainable environment, *Restoration of Wetland Ecosystem*, pp 59–74
- Wang C, Liu J, Shen J, Chen D, Li Y, Jiang B et al (2018) Effects of biochar amendment on net greenhouse gas emissions and soil fertility in a double rice cropping system: a 4-year field experiment. *Agr Ecosyst Environ* 262:83–96
- Wang XL, Chunxing A, Li Z (2019) Effect of pyrolysis temperature on characteristics, chemical speciation and risk evaluation of heavy metals in biochar derived from textile dyeing sludge. *Ecotoxicol Environ Saf* 168:45–52
- Wang C, Shen J, Liu J (2019) Microbial mechanisms in the reduction of CH₄ emission from double rice cropping system amended by biochar: a four-year study. *Soil Biol Biochem* 2:251–263
- Wang C, Shen J, Liu J, Qin H, Yuan Q, Fan F et al (2019) Microbial mechanisms in the reduction of CH₄ emission from double rice cropping system amended by biochar: a four-year study. *Soil Biol Biochem* 135:251–263
- Wang Z, Li J, Zhang G, Zhi Y, Yang D, Lai X et al (2020) Characterization of acid-aged biochar and its ammonium adsorption in an aqueous solution. *Materials (Basel)* 13:2270
- Waqas M, Nizami AS, Aburiazza AS, Barakat MA, Ismail IMI, Rashid MI (2018) Optimization of food waste compost with the use of biochar. *J Environ Manag* 216:70–81
- Waqas M, Asam Z, Rehan M, Anwar MN, Khattak RA, Ismail IMI et al (2020) Development of biomass-derived biochar for agronomic and environmental remediation applications. *Biomass Convers Biorefin*. <https://doi.org/10.1007/s13399-020-00936-2>
- Watanabe I, Hashimoto T, Shimoyama A (1997) Methane-oxidizing activities and methanotrophic populations associated with wetland rice plants. *Biol Fertil Soils* 24:261–265
- Weber K, Quicker P (2018) Properties of biochar. *Fuel* 217:240–261
- Wu M, Han X, Zhong T, Yuan M, Wu W (2016) Soil organic carbon content affects the stability of biochar in paddy soil. *Agr Ecosyst Environ* 223:59–66
- Wu Z, Song Y, Shen H, Jiang X, Li B, Xiong Z (2019a) Biochar can mitigate methane emissions by improving methanotrophs for prolonged period in fertilized paddy soils. *Environ Pollut* 253:1038–1046
- Wu Z, Zhang X, Dong Y, Li B, Xiong Z (2019b) Biochar amendment reduced greenhouse gas intensities in the rice-wheat rotation system: 6-year field observation and meta-analysis. *Agric For Meteorol* 278:107625

- Xiang Y, Deng Q, Duan H, Guo Y (2017) Effects of biochar application on root traits: a meta-analysis. *GCB Bioenergy* 9:1563–1572
- Xiao Y, Yang S, Xu J, Ding J, Sun X, Jiang Z (2018) Effect of biochar amendment on methane emissions from paddy field under water-saving irrigation. *Sustainability* 10:1371
- Yang M, Liu Y, Sun X, Dong D, Wu W (2013) Biochar improves methane oxidation activity in rice paddy soil. *Trans Chin Soc Agric Eng* 29:145–151
- Yi Q, Liang B, Nan Q, Wang H, Zhang W, Wu W (2020) Temporal physicochemical changes and transformation of biochar in a rice paddy: insights from a 9-year field experiment. *Sci Total Environ* 721:137670
- Yoo G, Kim YJ, Lee YO, Ding W (2016) Investigation of greenhouse gas emissions from the soil amended with rice straw biochar. *KSCE J Civ Eng* 20:2197–2207
- Yu L, Tang J, Zhang R, Wu Q, Gong M (2012) Effects of biochar application on soil methane emission at different soil moisture levels. *Biol Fertil Soils* 49:119–128
- Zhang Y, Chen H, Ji G (2018) Effect of rice-straw biochar application on rice (*Oryza sativa*) root growth and nitrogen utilization in acidified paddy soil. *Int J Agric Biol* 20:2529–2536
- Zhang X, Xia J, Pu J, Cai C, Tyson GW, Yuan Z et al (2019a) Biochar-mediated anaerobic oxidation of methane. *Environ Sci Technol* 53:6660–6668
- Zhang Y, Xu X, Zhang P, Ling Z, Qiu H, Cao X (2019b) Pyrolysis-temperature depended quinone and carbonyl groups as the electron accepting sites in barley grass derived biochar. *Chemosphere* 232:273–280
- Zhao X, Wang J, Wang S, Xing G (2014) Successive straw biochar application as a strategy to sequester carbon and improve fertility: a pot experiment with two rice/wheat rotations in paddy soil. *Plant Soil* 378:279–294
- Zheng JF, Chen JH, Pan GX, Liu XY, Zhang XH, Li LQ, Zheng JW (2016) Biochar decreased microbial metabolic quotient and shifted community composition 4 years after a single incorporation in a slightly acid rice paddy from southwest China. *Sci Total Environ* 571:206–217
- Zimmerman AR (2010) Abiotic and microbial oxidation of laboratory-produced black carbon (biochar). *Environ Sci Technol* 44:1295–1301