

Review Article

Tumor exosomes: a double-edged sword in cancer therapy

Wei SUN¹, Ju-dong LUO¹, Hua JIANG^{1, *}, Dayue Darrel DUAN^{2, *}

¹Department of Oncology, Changzhou Second People's Hospital, Changzhou 213003, China; ²Laboratory of Cardiovascular Phenomics, Department of Pharmacology, University of Nevada School of Medicine, Reno, NV 89557, USA

Abstract

Tumor cells produce and secrete more nucleic acids, proteins and lipids than normal cells. These molecules are transported in the blood or around the cells in membrane-encapsulated exosomes. Tumor-derived or tumor-associated exosomes (usually 30–100 nm in diameter) contain abundant biological contents resembling those of the parent cells along with signaling messengers for intercellular communication involved in the pathogenesis, development, progression, and metastasis of cancer. As these exosomes can be detected and isolated from various body fluids, they have become attractive new biomarkers for the diagnosis and prognosis of cancer. Furthermore, tumor exosomes have also attracted increasing attention due to their potential as novel therapeutic strategies for the treatment of cancers. On the one hand, the lipid bilayer membrane-encapsulated vesicles are promising carriers of drugs and other therapeutic materials targeting specific cancer cells. On the other hand, tumor exosomes are important mediators for modulation of the microenvironment that orchestrates events critical to the growth and metastasis of cancer cells as well as chemoresistance. Here, we summarize the advances in our understanding of tumor-associated or tumor-derived exosomes in recent years, and discuss their roles in cancer development, progression, invasion, and metastasis of cancers and, more importantly, their potential in strategies for precision therapy of various cancers as well as important caveats.

Keywords: exosomes; tumor; metastasis; chemotherapy; precision medicine

Acta Pharmacologica Sinica (2018) 39: 534–541; doi: 10.1038/aps.2018.17; published online 15 Mar 2018

Introduction

Tumor cells, similar to normal cells, are factories that constantly produce and export various biological molecules. These molecules are released from the cells and transported in the blood or around the cells in unique nano-sized lipid bilayer membrane vesicles^[1]. A group of these membrane vesicles with a diameter of approximately 40 nanometer (nm) were initially identified from two types of cancer cells with electron microscopy by Trams *et al* in 1981^[2]. Later, Johnstone *et al* reported vesicle externalization from multivesicular bodies using immunoelectron microscopy during reticulocyte maturation, and then, they defined the exosomes with a diameter of 30–100 nm as microvesicles^[3]. At this very early stage, exosomes were erroneously identified as cellular garbage until the emergence of dendritic cell (DC)-derived exosomes that contributed to the inhibition of cancer cell growth by increasing the proliferation of cytotoxic T lymphocytes^[4]. Indeed,

these pioneer studies promoted the application of exosomes not only in cancer research but also in the general field of cell biology, physiology, and pharmacology. In 2013, the Nobel Prize in Physiology or Medicine was awarded to three scientists "for their discoveries of machinery regulating vesicle traffic, a major transport system in our cells"^[5]. Randy SCHEKMAN discovered a set of genes that were required for vesicle traffic^[6]. James ROTHMAN identified protein machinery that allows vesicles to fuse with their targets to permit vesicle transfer^[7]. Thomas SÜDHOF revealed how signals instruct the cell to organize microvesicle exosomes and deliver them to the right place at the right time^[8]. Indeed, later studies consistently found that disturbances in the exquisitely precise control system for the transport and delivery of microvesicles/exosomes have deleterious effects and contribute to many pathological conditions, such as immunological disorders^[9, 10] and cancers^[11, 12]. More details about the general physiology, pathology, functionality, and roles in cancer metastasis and therapy of these molecules can be found in several recent excellent review articles^[13–17]. To date, tumor-associated or tumor-derived exosomes have been shown to be extensively involved in the intercellular substance exchange and signal

*To whom correspondence should be addressed.

E-mail dduan@medicine.nevada.edu (Dayue Darrel DUAN);

lygcancer@126.com (Hua JIANG)

Received 2017-06-29 Accepted 2018-02-24

communication between the tumor and the host and play crucial roles in the pathogenesis, development, invasion, and metastasis of cancer cells; more importantly, their potential as novel therapeutic strategies for precision therapy of various cancers has been demonstrated. In this review, we summarize the advances in our understanding of tumor exosomes in recent years and discuss their roles in the development, invasion, and metastasis of cancer cells, and more importantly, the potential and limitations for the development of tumor exosome-based therapeutic strategies for cancer treatment.

Characteristics of tumor-derived and tumor-associated exosomes

Tumor cells continuously secrete membrane vesicles to the extracellular environment. These tumor-derived exosomes represent novel vehicles for cell-cell communication networks and new components of a supportive microenvironment for localization, proliferation, and survival of tumor cells^[2, 18-21]. Trams *et al* first used the term “exosomes” to describe membrane-enclosed structures of variable sizes budding from the cell membranes of C-6 glioma or N-18 neuroblastoma cells^[2]. Exosomes are now often used to describe small extracellular membrane vesicles with a diameter of 30–150 nm generated via an endocytic pathway. Exosomes released by malignant tumor cells contain specific repertoires of molecules, including proteins, lipids, DNA molecules, miRNAs, mRNAs and non-coding RNAs, which are important for cancer cell communication with the environment. In the ExoCarta database, 286 studies focused on exosomes were found together with the contents in the exosomes, which included 41860 proteins, 7540 RNA molecules and 1116 lipids^[22-24]. Specific sorting mechanisms have been proposed for controlling the sorting and loading of selected molecules into exosomes^[25].

In addition to cancer cells, multiple other cell types, including adult stem cells, stromal cells, and cancer stem cells, are known to communicate with each other through their exosomes within the tumor microenvironment^[26-32]. Accumulating evidence has demonstrated that these tumor-associated exosomes mediate the interactions among different types of cells within the tumor microenvironment, thereby modulating tumor development, progression, and metastasis^[29-32]. These studies have also provided new insight into how these cells interact with each other to promote resistance to anti-cancer drug therapy and contribute to tumor recurrence through exosome signaling^[33-37]. While the detailed cellular and molecular mechanisms for the interaction between cancer cells and their surrounding cells, and thereby tumor progression and metastasis, remain to be further elucidated, current evidence has demonstrated that extracellular microvesicles, especially exosomes, are transferred among cancer cells and their surrounding cancer stem cells, adult stem cells and stromal cells, providing another layer of complexity in cancer cell biology associated with the tumor microenvironment and tumor progression.

Although there are differences in the chemical composition and biological functions of exosomes in various tissues and

cells, the process of exosome formation is similar. Initially, primary endosomes are formed after entrapment of the cytoplasmic membrane, and then, the endosomes may fuse with the lysosomes, which trigger the formation of vesicles in the lumen after releasing the cellular contents into the endosomes. Then, these primary endosomes are shifted to secondary endosomes and are called multivesicular bodies^[38]. Next, the multivesicular bodies fuse with a specific part of the cellular membrane. The multivesicular bodies fuse with the lysosomes, followed by fusing with the cellular membrane and secretion of vesicles through exocytosis via regulation of the GTPase and NsF coupling protein receptor. These substances were reported to be detected in blood, lymph fluid, urine, saliva, breast milk and ascites^[39].

Upon uptake of exosomes by the target cells, the exosome content may affect the biological behaviors and induce phenotypic changes in target recipient cells by several mechanisms. Exosome miRNAs, for example, can silence mRNA targets and influence cellular functions^[40-42]. Exosomes released by acute myeloid leukemia cells affected the proliferation and migration of bone marrow stromal cells^[43], and multiple myeloma exosomes enhanced angiogenesis^[44].

Tumor exosomes in invasion and metastasis of cancer cells

Not all cancer cells, or at least cancer cells in the initial early stages, show an invasive capacity. Invasion gradually progresses during the pathogenesis and development of most cancer cells. The biological and genetic alterations of the daughter cells of cancer cells may be transferred through exosomes after multiple rounds of division and proliferation, which then result in the sharing of biological features among cancer cells. Metastasis is a complex multistep process that involves cancer cell invasion, survival in blood vessels, and attachment to and colonization of distant host organs. A variety of molecular and functional mediators in cell interactions are involved in cancer metastasis. Tumor exosomes are now known to influence almost every step of the signaling cascade of cancer cell metastasis and thus can be targeted as a therapeutic approach. Al-Nedawi *et al* studied the effects of intercellular transfer of the oncogenic receptor EGFRvIII by microvesicles derived from glioma cells with EGFR mutations^[45]. Abundant expression of anti-apoptotic genes was observed in the recipient cells, together with increased non-adherent growth of cancer cells and invasion. Similarly, transfer of DKO-1 (mutant KRAS allele only) to DKs-8 (wild-type KRAS allele only) resulted in enhanced three-dimensional growth of the wild-type KRAS-expressing non-transformed cells as exosomes from mutant KRAS cells contained many tumor-promoting proteins, including KRAS, EGFR, SRC family kinases and integrins^[46]. These findings indicated that exosomes may serve as an important signal transfer vehicle and a vector for the delivery of genetic information^[47]. Le *et al* indicated that in murine cancer and human xenograft models, extracellular vesicles from highly metastatic breast cancer cells promoted metastasis of otherwise weakly metastatic cells either nearby or at distant

sites and induced these cells to colonize distant tissues in a miR-200-dependent manner^[48]. These results demonstrated that uptake of extracellular vesicles contributed to the transfer of metastatic capability in breast cancer cells.

It has been well established that organs of future cancer metastasis are not passive receivers of circulating cancer cells. The tumor microenvironment is closely related to cancer metastasis. Before metastasis, cancer cells may modulate the microenvironment of the target organs through exosomes to facilitate the metastasis and growth of cancer cells^[49]. The metastasis of cancer cells to the specific target organ is highly reliant on the expression of integrins contained in the tumor exosomes^[50]. Upon entering the recipient cells, exosomes may promote conditions that are suitable for the microenvironment formation before metastasis by activating Src phosphorylation and increasing S100 expression. Zhang *et al* reported that exosomal microRNA could induce microenvironmental PTEN loss in brain tissues, which then resulted in astrocytoma metastasis^[51].

Increasing evidence indicates that cancer cell-derived exosomes cause complex effects on stromal cells, vascular endothelial cells (VECs), and fibroblasts, which then promote the pathogenesis and development of cancer. Alterations in vascular permeability are indicators of microenvironmental formation before metastasis^[52]. Exosomes with abundant miR-105 from highly metastatic breast cancer cells could inhibit ZO-1 protein expression in endothelial cells, which then altered the vascular permeability and increased the susceptibility to cancer. This mechanism was attributed to the hepatic metastasis of breast cancer^[53]. Pancreatic cancer exosomes containing integrin $\beta 5$ and αv could be specifically recognized by Kupffer cells in the hepatic tissues. After phagocytosis of Kupffer cells, exosome-released macrophage migration inhibitory factor (MIF) upregulated the secretion of TGF β and contributed to the generation of fibronectin by Kupffer cells in the liver. This fibrotic microenvironment could enhance the recruitment of bone marrow-derived macrophages, which finally induced the formation of the microenvironment before hepatic metastasis, as well as the possibility of liver metastasis. Otherwise, strategies that inhibit MIF expression could attenuate the microenvironment formation in the liver before metastasis and thus limit hepatic metastasis of cancer cells^[54, 55].

In 2016, Greening *et al* reported that exosomes from malignant mesothelioma were involved in the reconstitution and regeneration of blood vessels by enhancing the migration of VECs and fibroblasts^[56]. Similarly, Cui *et al* reported that exosomes from pulmonary adenocarcinoma may be involved in the angiogenesis of cancer cells by upregulating Ephrin $\alpha 3$ expression in stromal cells in a miR-210-dependent manner^[57]. Moreover, proteins from tumor-derived exosomes, such as Rab3D^[58], TGF- $\beta 1$ ^[59], and LMP1^[60], were reported to promote epithelial-mesenchymal transition (EMT), which then increased the oncogenicity and invasion of cancer cells. In addition to the effects on proliferation of endothelial cells and mesenchymal remodeling, tumor-derived exosomes can also directly regulate the distal lymph node microenvironment,

which provided suitable conditions for lymph node metastasis and melanoma growth^[61]. Furthermore, exosomes from stromal cells were reported to affect the biological behaviors of cancer cells. For example, Luga *et al* revealed that upon trafficking of CD81-positive fibroblast exosomes in breast cancer cells, the fibroblast WNT-11 autocrine protein contributed to the invasion and metastasis of cancer by activating the Wnt-PCP signaling pathway^[62].

Exosomes involved in the degradation of extracellular matrix (ECM)

ECM is an important tissue barrier for cancer metastasis. Alterations and remodeling of ECM affect the pace of cancer cell invasion. Exosomes from human prostate cancer have varying microRNA contents, such as miR-100-5p, miR-21-5p, and miR-139-5p, which could trigger the expression of matrix metalloproteinases (MMPs), such as MMP2, MMP9 and MMP13. These exosomes are involved in the degradation of ECM and release of growth factors, which finally trigger the invasion and metastasis of cancer cells^[63]. Additionally, tumor exosomes can mediate the direct transmission of MMP13 to recipient cells, which then lead to the degradation of ECM and metastasis of cancer cells^[64]. Moreover, McCready *et al* showed that invasive carcinoma exosome-released Hsp90 protein could promote cellular migration by activating fibrinogenase^[65].

Potential of exosomes in cancer therapy

To date, cancer therapy has primarily relied on surgery, radiotherapy, chemotherapy, immunotherapy, and targeted therapy. However, the prognosis remains poor due to the recurrence and metastasis of cancer. Based on the roles of exosomes in the invasion and metastasis of cancer cells, tumor exosomes may be promising new therapeutic candidates for the treatment of cancer and may improve the prognosis and outcomes of cancer therapy.

Tumor-derived exosomes or tumor-related exosomes are considered to be closely related to the pathogenesis and microenvironmental formation of cancer, as the number of exosomes in cancer cells is higher than that of the normal cells^[66]. Therefore, inhibition of the synthesis, release and absolute circulating level of exosomes may serve as an effective treatment regimen for cancer. The formation and secretion of exosomes may be modulated through several approaches. For example, sphingomyelinase 2 could facilitate the synthesis of tumor-derived exosomes. Intravenous injection of an inhibitor of sphingomyelinase 2, GW4869, effectively inhibited the secretion of exosomes in tumor-bearing mice and decreased the metastatic rate of cancer cells^[67]. Chalmin *et al* reported that 5-*N,N*-dimethylamiloride could decrease the secretion of exosomes from cancer cells by blocking the H⁺/Na⁺ and Na⁺/Ca²⁺ channels, which subsequently delayed the growth of cancer cells^[68]. In contrast, these phenomena were not observed in PC3 cells^[69], which indicated that these inhibitory effects might be selective to specific cell types. A few conserved protein families are closely related to the secretion of exosomes,

among which Rab27a and Rab27b have been regarded as important mediators of these biological processes^[70]. Bobrie *et al* showed that silencing of Rab27a and Rab27b by RNA interference may alter the cancer microenvironment and delay the growth and metastasis of cancer cells^[71]. Nevertheless, the release of exosomes was similar in CD9- and MFG8-positive cells, which indicates that the release of exosomes was partially independent of Rab27a. Similarly, not all regulatory factors could effectively interfere with the release of exosomes. Although the expression of integrin facilitates the migration of cancer cells to specific organs, Mason *et al* reported confounding results of a phase III trial using cilengitide, an inhibitor of integrin $\alpha\beta3$ and $\alpha\beta5$, for treating glioblastoma^[72]. Interestingly, in a recent study, utilization of a hemofiltration system (*eg*, the Aethlon ADAPT™ system) to capture the exosomes in the circulating system contributed to the decrease of exosome secretion in the circulation, which provided a potential strategy to inhibit the growth and metastasis of cancer cells^[73].

The unique lipid bilayer structure of the exosome protects it from degradation and RNase damage. Compared with conventional liposome preparations, exosomes showed comparatively lower toxicity and satisfactory tissue tolerance^[74]. Moreover, the integrin expression spectrum contained in these molecules is abundant, which can mediate the migration of exosomes into target organs^[49]. These advantages have attracted increasing interest in developing exosomes as an excellent vector *in vivo* for the effective delivery of agents for cancer treatment. Exosomes can be used as specific vectors for the intracellular transmission of selective DNAs, miRNAs, non-coding RNAs, and proteins^[75]. For example, an exosome-based delivery platform for adriamycin^[76] or paclitaxel^[77] was developed for targeted cancer therapy and shown to have minimal immunogenicity and toxicity. Compared with conventional drug delivery systems, the exosome-based adriamycin delivery system showed higher drug availability with lower toxicity, especially toxicity to the heart. A previous study showed that the let-7 miRNA could inhibit the PI3K signaling pathway in cancer cells, which shares the same target PI3K signaling pathway as that for gefitinib for lung cancer^[78]. The overlap of targets for let-7 miRNA and gefitinib may induce sensitization of these two agents. Given that anti-miR-135 contributed to the sensitivity of lung cancer cells to taxanes, we speculated that delivery of specific miRNAs into specific cells using exosomes, which then increase the sensitivity of cancer cells to chemotherapy, could be a promising approach^[79]. According to previous results, the resistance of pulmonary adenocarcinoma to paclitaxel and cisplatin was related to the loss of PTEN on chromosome^[24]. Therefore, delivery of miR-181a through exosomes into pulmonary adenocarcinoma may increase the sensitivity of the cancer cells to paclitaxel and platinum-based chemotherapy^[80].

Another advantage for the use of exosomes as a therapeutic vector is their small size. Exosomes can easily pass through various biological barriers, including the blood-brain barrier^[81].

Extensive studies have been carried out to investigate the

roles of exosomes in immunological reactions after Zitvogel *et al* reported that exosomes are involved in immunological reactions in 1998^[4]. Exosomes are now considered a new hot topic in research on vaccine development in cancer treatment. In a previous study, it was reported that after stimulation of cancer-related antigens, DCs could produce exosomes containing specific cancer antigens^[82]. The tumor-derived exosomes and DC-derived exosomes could migrate into specific lymph nodes and then activate CD4⁺ and CD8⁺ T cells to trigger anti-tumor immunological reactions. To date, there are indeed some promising reports on the development of cancer vaccines for treating lung cancer. For example, in a clinical trial, TG4010 could extend the survival duration of patients with non-small cell lung cancer (NSCLC) at stages IIIb and IV^[83]. Meanwhile, the EGF vaccination was reported to be effective for the treatment of NSCLC patients^[84]. Additionally, in a recent phase II clinical trial focused on the treatment of NSCLC patients, generation of exosomes by DCs in the presence of tumor-related antigens could extend the progression-free survival of advanced cancer patients with low expression of NKp30^[85].

Tumor exosomes were reported to facilitate the apoptosis of cancer cells. Cancer cells could deliver survivin into the lung cancer cells through exosomes, which then facilitated the apoptosis of recipient cells by inhibiting cell growth. Survivin-D53A, a dominant-negative mutant survivin, could promote the apoptosis of pulmonary adenocarcinoma^[86]. Some components in vegetal cell-derived exosomes also showed anti-cancer effects. For example, the combination of curcumin and triptolide could increase the apoptosis of ovarian cancer cells^[87].

Caveats and limitations of exosomes for precision therapy of various cancers

Currently, clinical application of exosomes in cancer therapy is still not possible due to the following major caveats and limitations: i) specific markers for the identification of extracellular vesicles are still lacking; ii) there is currently no standard for the isolation and purification of specific tumor exosomes^[88]; iii) more effective methods are needed to obtain homogenous exosomes; iv) regulatory control of exosome contents and secretion is still poorly understood. Some exosomes could also promote growth and invasion as well as metastasis of cancer cells. For example, several studies confirmed that exosomes could induce complete metastasis of cancer cells^[50, 54]. Hoshino *et al* reported that unconventional secretion of exosome vesicles from multivesicular endosomes (MVE) occurs across a broad set of systems and is increased in cancer, where it promotes aggressive behavior. Analysis of cancer cells identified specialized invasive actin structures called invadopodia as specific and critical docking and secretion sites for CD63- and Rab27a-positive MVE^[50]. Inhibition of invadopodia formation reduced exosome secretion into conditioned media. Addition of purified exosomes or inhibition of exosome biogenesis or secretion significantly affected multiple invadopodia life cycle steps, including invadopodia formation, stabilization, and exocytosis of proteinases. Therefore, exosome cargoes may

play a key role in promoting invasive activity and providing in situ signaling feedback. Exosome secretion also controlled cellular invasion through a 3-dimensional matrix. The synergistic interaction between exosome secretion and invadopodia biogenesis strongly suggests that regulation of tumor exosome secretion may play a fundamental role in promoting cancer cell invasiveness^[50]. Pancreatic ductal adenocarcinomas (PDACs) are highly metastatic and have a poor prognosis due to delayed detection. PDAC-derived exosomes were reported to induce liver pre-metastatic niche formation in naive mice and consequently increase liver metastatic burden. Uptake of PDAC-derived exosomes by Kupffer cells caused TGF β secretion and upregulated fibronectin production by hepatic stellate cells. This fibrotic microenvironment enhanced recruitment of bone marrow-derived macrophages. MIF was substantially higher in PDAC-derived exosomes from stage I PDAC patients who later developed liver metastasis than exosomes from patients whose pancreatic tumors did not progress. When MIF expression in the PDAC-derived exosomes was blocked, liver pre-metastatic niche formation and metastasis were also prevented. These findings suggest that MIF in tumor exosomes may prime the liver for metastasis and may be a prognostic marker for the development of PDAC liver metastasis^[54]. Leal *et al* reported that tumor-derived exosomes and neutrophils may act cooperatively in cancer-associated thrombosis because the thrombosis in the cancer-bearing mice was faster than that of the normal controls^[89].

Understanding the underlying and complex mechanisms of exosome-mediated immunosuppression is important for preventing the immune escape of tumor cells and for identifying novel treatments for cancer. DCs and lymphocytes in the tumor microenvironment may modulate immune responses; however, tumor-derived exosomes may mediate the death receptor pathway to induce CD8⁺ T lymphocyte apoptosis. In addition, these molecules could mediate T lymphocyte imbalance by regulating the proliferation of T lymphocytes and inhibiting the proliferation of effector T lymphocytes, which then inhibited the immune function of the tumor microenvironment^[90, 91]. As a means of cellular communication, tumor-derived exosomes can inhibit the immune function of the receptor cells or trigger the distribution of immunologic receptors and ligands of the receptor cells with a similar pattern to that of oncocytes. Thereafter, they may interfere with immune therapies via sequestration of therapeutic antibodies or elimination of vaccine-induced or adoptively transferred immune effector cells^[92]. In a recent study, tumor-derived exosomes were shown to activate the transfer of epidermal growth factor receptor to macrophages in the hosts, inhibit innate immunity and induce immunocompromise^[93].

Therefore, in contrast to exosomes from other sources used to deliver therapeutic agents, tumor exosomes may be a double-edged sword when used as a therapeutic tool for cancer treatment. Full elucidation of the formation, secretion, and networking function of tumor exosomes is urgently needed for the realization of this attractive and promising strategy for cancer therapy. More extensive clinical or animal studies with

large sample sizes are needed in the future to investigate the features and safety of tumor exosomes serving as either a target or a vector for genes and/or drugs.

Conclusion and perspectives

In the future, we will need to broaden our understanding of the biological features and biochemical characteristics as well as the functional roles of tumor exosomes in the pathogenesis and development of cancer. Only when the specificity, efficacy, functionality, toxicity, and safety of exosomes have been fully elucidated can tumor exosomes serve as promising candidates for anti-cancer therapy and be developed as new strategies for precision therapy of various cancers.

Acknowledgements

This work was supported by the National Natural Science Foundation of China (81773224), the Scientific Program of Changzhou (CE20175025) and the Scientific and Technology Program of Nanjing Medical University (2015NJMU172).

References

- 1 Kreger BT, Dougherty AL, Greene KS, Cerione RA, Antonyak MA. Microvesicle cargo and function changes upon induction of cellular transformation. *J Biol Chem* 2016; 291: 19774–85.
- 2 Trams EG, Lauter CJ, Salem N Jr, Heine U. Exfoliation of membrane ecto-enzymes in the form of micro-vesicles. *Biochim Biophys Acta* 1981; 645: 63–70.
- 3 Johnstone RM, Adam M, Hammond JR, Orr L, Turbide C. Vesicle formation during reticulocyte maturation. Association of plasma membrane activities with released vesicles (exosomes). *J Biol Chem* 1987; 262: 9412–20.
- 4 Zitvogel L, Regnault A, Lozier A, Wolfers J, Flament C, Tenza D, *et al*. Eradication of established murine tumors using a novel cell-free vaccine: dendritic cell-derived exosomes. *Nat Med* 1998; 4: 594–600.
- 5 Pfeffer SR. A prize for membrane magic. *Cell* 2013; 155: 1203–6.
- 6 Kaiser CA, Schekman R. Distinct sets of SEC genes govern transport vesicle formation and fusion early in the secretory pathway. *Cell* 1990; 61: 723–33.
- 7 Balch WE, Dunphy WG, Braell WA, Rothman JE. Reconstitution of the transport of protein between successive compartments of the Golgi measured by the coupled incorporation of N-acetylglucosamine. *Cell* 1984; 39: 405–16.
- 8 Hata Y, Slaughter CA, Sudhof TC. Synaptic vesicle fusion complex contains unc-18 homologue bound to syntaxin. *Nature* 1993; 366: 347–51.
- 9 Chaput N, Thery C. Exosomes: immune properties and potential clinical implementations. *Semin Immunopathol* 2011; 33: 419–40.
- 10 Harada T, Yamamoto H, Kishida S, Kishida M, Awada C, Takao T, *et al*. Wnt5b-associated exosomes promote cancer cell migration and proliferation. *Cancer Sci* 2017; 108: 42–52.
- 11 Hendrix A, Hume AN. Exosome signaling in mammary gland development and cancer. *Int J Dev Biol* 2011; 55: 879–87.
- 12 Hewson C, Morris KV. Form and function of exosome-associated long non-coding RNAs in cancer. *Curr Top Microbiol Immunol* 2016; 394: 41–56.
- 13 Gilligan KE, Dwyer RM. Engineering exosomes for cancer therapy. *Int J Mol Sci* 2017; 18. pii: E1122.
- 14 Javeed N, Mukhopadhyay D. Exosomes and their role in the micro-

- macro-environment: a comprehensive review. *J Biomed Res* 2017; 31: 386–94.
- 15 Steinbichler TB, Dudas J, Riechelmann H, Skvortsova II. The role of exosomes in cancer metastasis. *Semin Cancer Biol* 2017; 44: 170–81.
- 16 Wang J, Sun X, Zhao J, Yang Y, Cai X, Xu J, *et al*. Exosomes: a novel strategy for treatment and prevention of diseases. *Front Pharmacol* 2017; 8: 300.
- 17 Weidle UH, Birzele F, Kollmorgen G, Ruger R. The multiple roles of exosomes in metastasis. *Cancer Genomics Proteomics* 2017; 14: 1–15.
- 18 Paggetti J, Haderk F, Seiffert M, Janji B, Distler U, Ammerlaan W, *et al*. Exosomes released by chronic lymphocytic leukemia cells induce the transition of stromal cells into cancer-associated fibroblasts. *Blood* 2015; 126: 1106–17.
- 19 Palazzolo G, Albanese NN, DI CG, Gygax D, Vittorelli ML, Pucci-Minafra I. Proteomic analysis of exosome-like vesicles derived from breast cancer cells. *Anticancer Res* 2012; 32: 847–60.
- 20 Pan J, Ding M, Xu K, Yang C, Mao LJ. Exosomes in diagnosis and therapy of prostate cancer. *Oncotarget* 2017; 8: 97693–700.
- 21 Panagiotara A, Markou A, Lianidou ES, Patrinos GP, Katsila T. Exosomes: a cancer theranostics road map. *Public Health Genomics* 2017; 20: 116–25.
- 22 Gangoda L, Liem M, Ang CS, Keerthikumar S, Adda CG, Parker BS, *et al*. Proteomic profiling of exosomes secreted by breast cancer cells with varying metastatic potential. *Proteomics* 2017; 17. doi: 10.1002/pmic.201600370.
- 23 Keerthikumar S, Gangoda L, Liem M, Fonseka P, Atukorala I, Ozcitti C, *et al*. Proteogenomic analysis reveals exosomes are more oncogenic than ectosomes. *Oncotarget* 2015; 6: 15375–96.
- 24 Keerthikumar S, Chisanga D, Ariyaratne D, Al SH, Anand S, Zhao K, *et al*. ExoCarta: a web-based compendium of exosomal cargo. *J Mol Biol* 2016; 428: 688–92.
- 25 Villarroya-Beltri C, Baixauli F, Gutierrez-Vazquez C, Sanchez-Madrid F, Mittelbrunn M. Sorting it out: regulation of exosome loading. *Semin Cancer Biol* 2014; 28: 3–13.
- 26 Cho JA, Park H, Lim EH, Lee KW. Exosomes from breast cancer cells can convert adipose tissue-derived mesenchymal stem cells into myofibroblast-like cells. *Int J Oncol* 2012; 40: 130–8.
- 27 Milane L, Singh A, Mattheolabakis G, Suresh M, Amiji MM. Exosome mediated communication within the tumor microenvironment. *J Control Release* 2015; 219: 278–94.
- 28 Nabet BY, Qiu Y, Shabason JE, Wu TJ, Yoon T, Kim BC, *et al*. Exosome RNA unshielding couples stromal activation to pattern recognition receptor signaling in cancer. *Cell* 2017; 170: 352–66.
- 29 Naito Y, Yoshioka Y, Yamamoto Y, Ochiya T. How cancer cells dictate their microenvironment: present roles of extracellular vesicles. *Cell Mol Life Sci* 2017; 74: 697–713.
- 30 Whiteside TL. Exosome and mesenchymal stem cell cross-talk in the tumor microenvironment. *Semin Immunol* 2017. pii: S1044-5323(17)30010-6.
- 31 Bhome R, Goh RW, Bullock MD, Pillar N, Thirdborough SM, Mellone M, *et al*. Exosomal microRNAs derived from colorectal cancer-associated fibroblasts: role in driving cancer progression. *Aging (Albany NY)* 2017; 9: 2666–94.
- 32 Devhare PB, Ray RB. Extracellular vesicles: Novel mediator for cell to cell communications in liver pathogenesis. *Mol Aspects Med* 2017. pii: S0098-2997(17)30089-4.
- 33 Shenoy GN, Loyall JL, Maguire O, Iyer V, Kelleher RJ, Minderman H, *et al*. Exosomes associated with human ovarian tumors harbor a reversible checkpoint of T cell responses. *Cancer Immunol Res* 2018; 6: 236–47.
- 34 Wu K, Xing F, Wu SY, Watabe K. Extracellular vesicles as emerging targets in cancer: recent development from bench to bedside. *Biochim Biophys Acta* 2017; 1868: 538–63.
- 35 Wang J, Yeung BZ, Cui M, Peer CJ, Lu Z, Figg WD, *et al*. Exosome is a mechanism of intercellular drug transfer: application of quantitative pharmacology. *J Control Release* 2017; 268: 147–58.
- 36 Sueta A, Yamamoto Y, Tomiguchi M, Takeshita T, Yamamoto-Ibusuki M, Iwase H. Differential expression of exosomal miRNAs between breast cancer patients with and without recurrence. *Oncotarget* 2017; 8: 69934–44.
- 37 Patel GK, Khan MA, Bhardwaj A, Srivastava SK, Zubair H, Patton MC, *et al*. Exosomes confer chemoresistance to pancreatic cancer cells by promoting ROS detoxification and miR-155-mediated suppression of key gemcitabine-metabolising enzyme, DCK. *Br J Cancer* 2017; 116: 609–19.
- 38 Hanson PI, Cashikar A. Multivesicular body morphogenesis. *Annu Rev Cell Dev Biol* 2012; 28: 337–62.
- 39 Vlassov AV, Magdaleno S, Setterquist R, Conrad R. Exosomes: current knowledge of their composition, biological functions, and diagnostic and therapeutic potentials. *Biochim Biophys Acta* 2012; 1820: 940–8.
- 40 Romano G, Kwong LN. miRNAs, melanoma and microenvironment: an intricate network. *Int J Mol Sci* 2017; 18. pii: E2354.
- 41 Takano Y, Masuda T, Iinuma H, Yamaguchi R, Sato K, Tobo T, *et al*. Circulating exosomal microRNA-203 is associated with metastasis possibly via inducing tumor-associated macrophages in colorectal cancer. *Oncotarget* 2017; 8: 78598–613.
- 42 Fan Q, Yang L, Zhang X, Peng X, Wei S, Su D, *et al*. The emerging role of exosome-derived non-coding RNAs in cancer biology. *Cancer Lett* 2018; 414: 107–15.
- 43 Umez T, Tadokoro H, Azuma K, Yoshizawa S, Ohyashiki K, Ohyashiki JH. Exosomal miR-135b shed from hypoxic multiple myeloma cells enhances angiogenesis by targeting factor-inhibiting HIF-1. *Blood* 2014; 124: 3748–57.
- 44 Umez T, Imanishi S, Azuma K, Kobayashi C, Yoshizawa S, Ohyashiki K, *et al*. Replenishing exosomes from older bone marrow stromal cells with miR-340 inhibits myeloma-related angiogenesis. *Blood Adv* 2017; 1: 812–23.
- 45 Al-Nedawi K, Meehan B, Micallef J, Lhotak V, May L, Guha A, *et al*. Intercellular transfer of the oncogenic receptor EGFRvIII by microvesicles derived from tumour cells. *Nat Cell Biol* 2008; 10: 619–24.
- 46 Demory Beckler M, Higginbotham JN, Franklin JL, Ham AJ, Halvey PJ, Imasuen IE, *et al*. Proteomic analysis of exosomes from mutant KRAS colon cancer cells identifies intercellular transfer of mutant KRAS. *Mol Cell Proteomics* 2013; 12: 343–55.
- 47 Gezer U, Ozgur E, Cetinkaya M, Isin M, Dalay N. Long non-coding RNAs with low expression levels in cells are enriched in secreted exosomes. *Cell Biol Int* 2014; 38: 1076–9.
- 48 Le MT, Hamar P, Guo C, Basar E, Perdigo-Henriques R, Balaj L, *et al*. miR-200-containing extracellular vesicles promote breast cancer cell metastasis. *J Clin Invest* 2014; 124: 5109–28.
- 49 Hood JL, San RS, Wickline SA. Exosomes released by melanoma cells prepare sentinel lymph nodes for tumor metastasis. *Cancer Res* 2011; 71: 3792–801.
- 50 Hoshino A, Costa-Silva B, Shen TL, Rodrigues G, Hashimoto A, Tesic Mark M, *et al*. Tumour exosome integrins determine organotropic metastasis. *Nature* 2015; 527: 329–35.

- 51 Zhang L, Zhang S, Yao J, Lowery FJ, Zhang Q, Huang WC, *et al*. Microenvironment-induced PTEN loss by exosomal microRNA primes brain metastasis outgrowth. *Nature* 2015; 527: 100–4.
- 52 Psaila B, Lyden D. The metastatic niche: adapting the foreign soil. *Nat Rev Cancer* 2009; 9: 285–93.
- 53 Zhou W, Fong MY, Min Y, Somlo G, Liu L, Palomares MR, *et al*. Cancer-secreted miR-105 destroys vascular endothelial barriers to promote metastasis. *Cancer Cell* 2014; 25: 501–15.
- 54 Costa-Silva B, Aiello NM, Ocean AJ, Singh S, Zhang H, Thakur BK, *et al*. Pancreatic cancer exosomes initiate pre-metastatic niche formation in the liver. *Nat Cell Biol* 2015; 17: 816–26.
- 55 Tomasetti M, Lee W, Santarelli L, Neuzil J. Exosome-derived microRNAs in cancer metabolism: possible implications in cancer diagnostics and therapy. *Exp Mol Med* 2017; 49: e285.
- 56 Greening DW, Ji H, Chen M, Robinson BW, Dick IM, Creaney J, *et al*. Secreted primary human malignant mesothelioma exosome signature reflects oncogenic cargo. *Sci Rep* 2016; 6: 32643.
- 57 Cui H, Seubert B, Stahl E, Dietz H, Reuning U, Moreno-Leon L, *et al*. Tissue inhibitor of metalloproteinases-1 induces a pro-tumourigenic increase of miR-210 in lung adenocarcinoma cells and their exosomes. *Oncogene* 2015; 34: 3640–50.
- 58 Yang J, Liu W, Lu X, Fu Y, Li L, Luo Y. High expression of small GTPase Rab3D promotes cancer progression and metastasis. *Oncotarget* 2015; 6: 11125–38.
- 59 Ko H, Jeon H, Lee D, Choi HK, Kang KS, Choi KC. Sanguin H6 suppresses TGF-beta induction of the epithelial-mesenchymal transition and inhibits migration and invasion in A549 lung cancer. *Bioorg Med Chem Lett* 2015; 25: 5508–13.
- 60 Hood JL, Pan H, Lanza GM, Wickline SA; Consortium for Translational Research in Advanced Imaging and Nanomedicine (C-TRAIN). Paracrine induction of endothelium by tumor exosomes. *Lab Invest* 2009; 89: 1317–28.
- 61 Aga M, Bentz GL, Raffa S, Torrisi MR, Kondo S, Wakisaka N, *et al*. Exosomal HIF1alpha supports invasive potential of nasopharyngeal carcinoma-associated LMP1-positive exosomes. *Oncogene* 2014; 33: 4613–22.
- 62 Luga V, Zhang L, Vitoria-Petit AM, Ogunjimi AA, Inanlou MR, Chiu E, *et al*. Exosomes mediate stromal mobilization of autocrine Wnt-PCP signaling in breast cancer cell migration. *Cell* 2012; 151: 1542–56.
- 63 Sanchez CA, Andahur EI, Valenzuela R, Castellon EA, Fulla JA, Ramos CG, *et al*. Exosomes from bulk and stem cells from human prostate cancer have a differential microRNA content that contributes cooperatively over local and pre-metastatic niche. *Oncotarget* 2016; 7: 3993–4008.
- 64 You Y, Shan Y, Chen J, Yue H, You B, Shi S, *et al*. Matrix metalloproteinase 13-containing exosomes promote nasopharyngeal carcinoma metastasis. *Cancer Sci* 2015; 106: 1669–77.
- 65 McCreedy J, Sims JD, Chan D, Jay DG. Secretion of extracellular hsp90alpha via exosomes increases cancer cell motility: a role for plasminogen activation. *BMC Cancer* 2010; 10: 294.
- 66 Jenjaroenpun P, Kremenska Y, Nair VM, Kremenskoy M, Joseph B, Kurochkin IV. Characterization of RNA in exosomes secreted by human breast cancer cell lines using next-generation sequencing. *PeerJ* 2013; 1: e201.
- 67 Fabbri M, Paone A, Calore F, Galli R, Gaudio E, Santhanam R, *et al*. MicroRNAs bind to Toll-like receptors to induce prometastatic inflammatory response. *Proc Natl Acad Sci U S A* 2012; 109: E2110–6.
- 68 Chalmin F, Ladoire S, Mignot G, Vincent J, Bruchard M, Remy-Martin JP, *et al*. Membrane-associated Hsp72 from tumor-derived exosomes mediates STAT3-dependent immunosuppressive function of mouse and human myeloid-derived suppressor cells. *J Clin Invest* 2010; 120: 457–71.
- 69 Phuyal S, Hessvik NP, Skotland T, Sandvig K, Llorente A. Regulation of exosome release by glycosphingolipids and flotillins. *FEBS J* 2014; 281: 2214–27.
- 70 Hendrix A, De Wever O. Rab27 GTPases distribute extracellular nanomaps for invasive growth and metastasis: implications for prognosis and treatment. *Int J Mol Sci* 2013; 14: 9883–92.
- 71 Bobrie A, Krumeich S, Reyat F, Recchi C, Moita LF, Seabra MC, *et al*. Rab27a supports exosome-dependent and -independent mechanisms that modify the tumor microenvironment and can promote tumor progression. *Cancer Res* 2012; 72: 4920–30.
- 72 Mason WP. End of the road: confounding results of the CORE trial terminate the arduous journey of cilengitide for glioblastoma. *Neuro Oncol* 2015; 17: 634–5.
- 73 Marleau AM, Chen CS, Joyce JA, Tullis RH. Exosome removal as a therapeutic adjuvant in cancer. *J Transl Med* 2012; 10: 134.
- 74 Lai RC, Yeo RW, Tan KH, Lim SK. Exosomes for drug delivery - a novel application for the mesenchymal stem cell. *Biotechnol Adv* 2013; 31: 543–51.
- 75 Thakur BK, Zhang H, Becker A, Matei I, Huang Y, Costa-Silva B, *et al*. Double-stranded DNA in exosomes: a novel biomarker in cancer detection. *Cell Res* 2014; 24: 766–9.
- 76 Tian Y, Li S, Song J, Ji T, Zhu M, Anderson GJ, *et al*. A doxorubicin delivery platform using engineered natural membrane vesicle exosomes for targeted tumor therapy. *Biomaterials* 2014; 35: 2383–90.
- 77 Yang T, Martin P, Fogarty B, Brown A, Schurman K, Phipps R, *et al*. Exosome delivered anticancer drugs across the blood-brain barrier for brain cancer therapy in *Danio rerio*. *Pharm Res* 2015; 32: 2003–14.
- 78 Sebio A, Pare L, Paez D, Salazar J, Gonzalez A, Sala N, *et al*. The LCS6 polymorphism in the binding site of let-7 microRNA to the KRAS 3'-untranslated region: its role in the efficacy of anti-EGFR-based therapy in metastatic colorectal cancer patients. *Pharmacogenet Genomics* 2013; 23: 142–7.
- 79 Holleman A, Chung I, Olsen RR, Kwak B, Mizokami A, Saijo N, *et al*. miR-135a contributes to paclitaxel resistance in tumor cells both *in vitro* and *in vivo*. *Oncogene* 2011; 30: 4386–98.
- 80 Li H, Zhang P, Sun X, Sun Y, Shi C, Liu H, *et al*. MicroRNA-181a regulates epithelial-mesenchymal transition by targeting PTEN in drug-resistant lung adenocarcinoma cells. *Int J Oncol* 2015; 47: 1379–92.
- 81 Alvarez-Erviti L, Seow Y, Yin H, Betts C, Lakhai S, Wood MJ. Delivery of siRNA to the mouse brain by systemic injection of targeted exosomes. *Nat Biotechnol* 2011; 29: 341–5.
- 82 Romagnoli GG, Zelante BB, Toniolo PA, Migliori IK, Barbuto JA. Dendritic cell-derived exosomes may be a tool for cancer immunotherapy by converting tumor cells into immunogenic targets. *Front Immunol* 2014; 5: 692.
- 83 Rotonda C, Anota A, Mercier M, Bastien B, Lacoste G, Limacher JM, *et al*. Impact of TG4010 vaccine on health-related quality of life in advanced non-small-cell lung cancer: results of a Phase IIB clinical trial. *PLoS One* 2015; 10: e0132568.
- 84 Crombet Ramos T, Rodriguez PC, Neningen Vinageras E, Garcia Verdecia B, Lage Davila A. CIMAvax EGF (EGF-P64K) vaccine for the treatment of non-small-cell lung cancer. *Expert Rev Vaccines* 2015; 14: 1303–11.
- 85 Besse B, Charrier M, Lapierre V, Dansin E, Lantz O, Planchard D, *et al*. Dendritic cell-derived exosomes as maintenance immunotherapy after first line chemotherapy in NSCLC. *Oncoimmunology* 2016; 5: e1071008.
- 86 Yu M, Peng X, Lu Y, Huang M. Inhibition function of dominant-negative

- mutant gene survivin-D53A to SPC-A1 lung adenocarcinoma xenograft in nude mice models. *Sheng Wu Yi Xue Gong Cheng Xue Za Zhi* 2015; 32: 624–8.
- 87 Cai YY, Lin WP, Li AP, Xu JY. Combined effects of curcumin and triptolide on an ovarian cancer cell line. *Asian Pac J Cancer Prev* 2013; 14: 4267–71.
- 88 Kowal J, Arras G, Colombo M, Jouve M, Morath JP, Primdal-Bengtson B, *et al*. Proteomic comparison defines novel markers to characterize heterogeneous populations of extracellular vesicle subtypes. *Proc Natl Acad Sci U S A* 2016; 113: E968–77.
- 89 Leal AC, Mizurini DM, Gomes T, Rochael NC, Saraiva EM, Dias MS, *et al*. Tumor-derived exosomes induce the formation of neutrophil extracellular traps: implications for the establishment of cancer-associated thrombosis. *Sci Rep* 2017; 7: 6438.
- 90 Whiteside TL. The effect of tumor-derived exosomes on immune regulation and cancer immunotherapy. *Future Oncol* 2017; 28: 2583–92.
- 91 Gao L, Wang L, Dai T, Jin K, Zhang Z, Wang S, *et al*. Tumor-derived exosomes antagonize innate antiviral immunity. *Nat Immunol* 2018. doi: 10.1038/s41590-017-0043-5.
- 92 Peng P, Yan Y, Keng S. Exosomes in the ascites of ovarian cancer patients: origin and effects on anti-tumor immunity. *Oncol Rep* 2011; 25: 749–62.
- 93 Miller IV, Grunewald TG. Tumour-derived exosomes: tiny envelopes for big stories. *Biol Cell* 2015; 107: 287–305.