

ORIGINAL ARTICLE

Contribution of breast milk and formula to arsenic exposure during the first year of life in a US prospective cohort

Courtney C. Carignan^{1,2,3}, Margaret R. Karagas^{1,4}, Tracy Punshon^{1,2}, Diane Gilbert-Diamond^{1,4} and Kathryn L. Cottingham^{1,2}

Arsenic is a carcinogen that can also affect the cardiac, respiratory, neurological and immune systems. Children have higher dietary arsenic exposure than adults owing to their more restricted diets and greater intake per unit body mass. We evaluated the potential contributions of breast milk and formula to arsenic exposure throughout the first year of life for 356 infants in the prospective New Hampshire Birth Cohort Study (NHBCS) using infant diets reported by telephone at 4, 8 and 12 months of age; measured household water arsenic concentrations; and literature data. Based on our central-tendency models, population-wide geometric mean (GM) estimated arsenic exposures in the NHBCS were relatively low, decreasing from 0.1 $\mu\text{g}/\text{kg}/\text{day}$ at 4 months of age to 0.07 $\mu\text{g}/\text{kg}/\text{day}$ at 12 months of age. At all three time points, exclusively formula-fed infants had GM arsenic exposures ~ 8 times higher than exclusively breastfed infants owing to arsenic in both tap water and formula powder. Estimated maximum exposures reached 9 $\mu\text{g}/\text{kg}/\text{day}$ among exclusively formula-fed infants in households with high tap water arsenic (80 $\mu\text{g}/\text{l}$). Overall, modeled arsenic exposures via breast milk and formula were low throughout the first year of life, unless formula was prepared with arsenic-contaminated tap water.

Journal of Exposure Science and Environmental Epidemiology (2016) **26**, 452–457; doi:10.1038/jes.2015.69; published online 4 November 2015

Keywords: breastfeeding; child; cohort studies; diet; environmental exposure; infant formula

INTRODUCTION

Early life is a period of heightened vulnerability to arsenic exposure.¹ Arsenic is a known human carcinogen that can also adversely affect the neurological, respiratory, cardiovascular, immunological and endocrine systems.² Most studies have investigated populations with chronic exposure to high concentrations of arsenic via drinking water, such as those in Bangladesh, Chile and Taiwan, where concentrations of arsenic in drinking water can be substantially $> 50 \mu\text{g}/\text{l}$.^{3,4} In such populations, early-life exposure has been associated with increased fetal mortality,^{3,4} decreased birth weight⁴ and diminished cognitive function,^{5,6} although results are not always consistent across studies.⁷ Moreover, effects of chronic early-life exposure can manifest in adulthood with increased occurrences and/or severity of lung disease, cardiovascular disease and cancer.^{1,2,8} Although less is known about the short- and long-term consequences of exposure below the current US EPA and World Health Organization maximum contaminant level (MCL) of 10 $\mu\text{g}/\text{l}$,⁹ *in utero* low-dose exposure has been associated with increased infant respiratory infections and the severity of these infections.¹⁰

There are two primary pathways for arsenic exposure in the US population: water and food. Arsenic is a naturally occurring element in aquifer bedrock and, as a result, is found in well water throughout the world.^{11–15} Drinking water primarily contains As^{III} and As^{V} , inorganic forms of arsenic (iAs) with known toxicity.¹⁶ Foods can contain iAs; organic metabolites of iAs such as dimethylarsinic acid (DMA) and monomethylarsonic acid (MMA); and arsenobetaine, arsenosugars and arsenolipids.^{17,18} Arsenobetaine is considered non-toxic and passes through the body

unmetabolized.¹⁹ Evidence from *in vitro* studies suggests that trivalent forms of MMA and DMA may have toxic and potentially carcinogenic properties,^{20–23} whereas the effects of the pentavalent forms of these compounds are less certain,²⁴ especially when consumed in food.²⁵

Dietary exposure to arsenic is expected to be about three times higher for infants and young children than for adults,²⁶ in part because their intake per unit body mass is higher²⁷ and their dietary diversity is lower than adults.²⁶ For example, newborn infants subsist on a diet of breast milk and/or formula. Powdered infant formula contains some arsenic,^{28–31} including in the inorganic form,^{31,32} and can contribute to exposure, especially when reconstituted with arsenic-contaminated water.³³ In contrast, we recently reported low levels of arsenic in breast milk from New Hampshire mothers with low exposure to arsenic in drinking water (median = 0.26 $\mu\text{g}/\text{l}$).³³ Other studies have found that arsenic in breast milk is low regardless of arsenic exposure via drinking water.^{34–39} The dominant species of arsenic in breast milk is currently unclear but may depend on the source or magnitude of exposure. In a Bangladeshi population exposed to high arsenic in drinking water (median = 78 $\mu\text{g}/\text{l}$),⁴⁰ breast milk samples ($n = 79$) contained predominantly iAs,³⁷ whereas only organic arsenic was detectable in breast milk from a small sample of Swedish mothers ($n = 3$) who had no known source of arsenic in their drinking water.³⁴

Here we quantify the potential arsenic exposure via breast milk and formula for individual infants in the New Hampshire Birth Cohort Study (NHBCS) over their first year of life. Our goals are to use exposure modeling to (1) provide insight into the relative contribution of breast milk vs formula to exposure, assuming use

¹Children's Environmental Health and Disease Prevention Research Center at Dartmouth, Hanover, NH, USA; ²Department of Biological Sciences, Dartmouth College, Hanover, NH, USA; ³Department of Environmental Health, Harvard T.H. Chan School of Public Health, Boston, MA, USA and ⁴Department of Epidemiology, Geisel School of Medicine at Dartmouth, Hanover, NH, USA. Correspondence: Professor Kathryn L. Cottingham, Department of Biological Sciences, Dartmouth College, 78 College St, Hanover, NH 03755, USA. Tel.: +603 646 0216. Fax: +603 646 1347.

E-mail: Kathryn.L.Cottingham@dartmouth.edu

Received 22 December 2014; revised 25 September 2015; accepted 25 September 2015; published online 4 November 2015

of home tap water to reconstitute infant formula, and (2) explore how population-wide exposure varies as breastfeeding prevalence decreases and formula use increases during the period from birth to 1 year of age. Given that breast milk is expected to be lower in arsenic than formula, we hypothesized that arsenic exposure would increase during infancy owing to the increased prevalence of formula feeding.⁴¹

SUBJECTS AND METHODS

The New Hampshire Birth Cohort Study

In January 2009, we began recruiting pregnant women ranging in age from 18 to 45 years who were receiving prenatal care at study clinics in New Hampshire, USA, as described previously.^{10,42} Enrollment criteria included a singleton pregnancy; the use of a private, unregulated well in the home since their last menstrual period; and plans to stay in the current residence through delivery. This study was reviewed and approved by the Committee for the Protection of Human Subjects (CPHS) at Dartmouth College, Hanover, NH, and all participants in the study provided informed consent in accordance with CPHS guidelines. Household tap water arsenic was measured in samples returned by subjects after enrollment.^{10,33,42}

Maternal Questionnaire

Women who agreed to participate were asked to complete a prenatal medical history and lifestyle questionnaire that included questions about sociodemographic factors, health history, personal habits, home water source and home water consumption.

Infant Feeding

At 4, 8 and 12 months postpartum, participants were contacted for a brief telephone interview that included questions about feeding practices on an average day, such as frequency of breastfeeding, amount of formula consumed and type of water used to reconstitute powdered formula (e.g., home tap vs bottled). Based on these responses, we assigned infants to one of the three feeding categories for their liquid diet: Breastfed (fed only breast milk), Formula-fed (fed only formula), or Mixed (fed both breast milk and formula). We assumed that all formula was prepared using home tap water and that infants in the Mixed feeding type received exactly half breast milk and half formula. This was done in the absence of more detailed data on infant feeding practices and local data on bottled water arsenic and to better reflect potential upper-bound exposures among New Hampshire infants.

Arsenic Exposure Estimates via Breast Milk and Formula for NHBCS Infants

We calculated potential arsenic exposure for each NHBCS infant at 4, 8 and 12 months of age based on (1) current feeding type (Breastfed, Mixed or Formula-fed, as described above); (2) the measured concentration of arsenic in household tap water (Supplementary Table S1) for Mixed and Formula-fed infants; and (3) literature-based values for age-specific breast milk ingestion rate (IR_{BW}), reported in the US EPA Child Specific Exposure Factors Handbook⁴³ Table 15-4) and the arsenic concentration in breast milk³³ or formula powder.³¹ The feeding type of each individual was allowed to vary at each time point (4, 8 and 12 months) depending on parental reports.

For each infant at each time point, we used two different models to estimate the distribution of potential arsenic exposure among NHBCS infants owing to ingestion of breast milk and/or formula—a central tendency model and an upper-bound model—based on individual-level concentrations of arsenic in household tap water and reported feeding type at 4, 8 and 12 months of age. Exposures were estimated using the equations in Supplementary Appendix S1 and the following numeric values:

- (1) The central tendency model multiplied the median concentration of arsenic in breast milk ($0.31 \mu\text{g/l}$)³³ and/or reconstituted formula ($1.1 \mu\text{g/l}$ from powder³¹ plus the measured household tap water arsenic)³³ by the mean IR_{BW} ⁴³ ($0.112 \text{ liter/kg/day}$ at 4 months, $0.075 \text{ liter/kg/day}$ at 8 months and $0.047 \text{ liter/kg/day}$ at 12 months).
- (2) The upper-bound model multiplied the maximum concentration of arsenic in breast milk ($0.62 \mu\text{g/l}$)³³ and/or reconstituted formula ($1.8 \mu\text{g/l}$ from powder³¹ plus the measured household tap water arsenic)³³ by the upper percentile IR_{BW} ⁴³ (defined as the mean plus 2 SDs: 0.148 liter/

kg/day at 4 months, $0.125 \text{ liter/kg/day}$ at 8 months and $0.101 \text{ liter/kg/day}$ at 12 months). This model is likely best for estimating short-term upper-bound exposures, as data for upper-bound IR_{BW} were collected over short periods (i.e., 3–10 days).

In making these calculations, we assumed that arsenic concentrations did not vary across formula brands based on published reports of comparable arsenic concentrations across brands and products;³¹ moreover, the median concentration used ($1.1 \mu\text{g/l}$) was similar to concentrations reported by the FDA³² ($1 \mu\text{g/l}$ of total arsenic when converted from $\mu\text{g/g}$ of formula powder assuming arsenic-free water). Similarly, median and maximum concentrations of arsenic in breast milk were applied from our previous study of NHBCS mothers ($n=9$) whose home tap water arsenic concentrations ranged from <0.01 to $8.9 \mu\text{g/l}$, a concentration range that represents approximately 90% of our study population.³³ Other studies around the world also report low concentrations of arsenic in breast milk relative to drinking water.^{34–39}

To test the effect of assuming that milk ingestion rates were consistent across feeding types, we performed two sensitivity analyses. First, we calculated the IR_{BW} for a subset of Formula-fed NHBCS infants for whom data on both daily formula consumption rates and body weight were available ($n=34, 48$ and 58 for 4, 8 and 12 months, respectively), and compared these values to the modeled inputs. We then estimated exposure for the Formula-fed and Mixed-fed infants using summary statistics for this NHBCS-specific formula IR_{BW} ,⁴³ instead of the breast milk IR_{BW} . In addition, we estimated exposure using the tap water IR_{BW} determined by the US EPA,⁴³ which also includes water intake other than formula (i.e., mixing with cereal).

RESULTS

NHBCS Characteristics

We selected 356 infants from the NHBCS with complete records of feeding type at the 4, 8 and 12 month time points (Table 1). In this subset, mean (SD) maternal age was 31.5 (4.6) years at the time of delivery. Most of the mothers were college graduates (39%) or had attended some postgraduate schooling (31%), and the majority reported being married (85%). Slightly more than half of the infants were female (55%) and all were white (100%). Less than half of the infants attended daycare at 4, 8 and 12 months (33, 38 and 41%, respectively). Demographics of the larger cohort are similar to this subset (data not shown).

Tap water arsenic within the study subset was generally low: 86% of the families had household tap water arsenic concentrations $<10 \mu\text{g/l}$, the current US EPA MCL (Table 1). However, the maximum household tap water arsenic for the NHBCS reached levels eight times the MCL ($79.7 \mu\text{g/l}$), with some differences among feeding types and time points (Supplementary Table S1).

Infant Feeding Patterns in the NHBCS

The predominant feeding type shifted from Breastfed to Formula-fed over the first year of life (Figure 1; Supplementary Tables S2 and S3). At 4 months of age, 45% of mothers reported that their infants were Breastfed, 23% were Mixed-fed and 32% were Formula-fed. By 12 months of age, the percentage of Breastfed infants had decreased to 21% and the percentage of Formula-fed infants had doubled to 66%.

Among Formula-fed infants, formula ingestion rates reported by parents decreased as infants got older (Supplementary Table S2). The average reported volume of formula consumed by Formula-fed infants decreased from 839 ml/day at 4 months to 710 ml/day at 12 months of age. Estimates of IR_{BW} for the subset of Formula-fed NHBCS infants were similar to the values recommended by the US EPA Child Specific Exposure Factors Handbook for exclusively breastfed babies (Table 15-4)⁴³ at 4 months ($n=34$; mean = $0.13 \text{ liter/kg/day}$; mean+2 SD = $0.20 \text{ liter/kg/day}$), 8 months ($n=48$; mean = $0.08 \text{ liter/kg/day}$; mean+2 SD = $0.15 \text{ liter/kg/day}$) and 12 months of age ($n=58$; mean = $0.06 \text{ liter/kg/day}$; mean+2 SD = $0.14 \text{ liter/kg/day}$) (Supplementary Table S4). For infants receiving both breast milk and formula, the average reported volume of

Table 1. Selected characteristics of mothers and infants in the New Hampshire Birth Cohort sub-study reported here ($n = 356$).

Population characteristic	Mean (range) or n (%) ^a
Maternal variables	
Maternal age, years	31.5 (18.6–44.5)
< 20	4 (1%)
20–29	104 (29%)
30–35	176 (49%)
> 35	72 (20%)
Maternal education	
< 11th grade	3 (1%)
High school graduate/GED	29 (8%)
Junior college, some college, technical school	70 (20%)
College graduate	135 (39%)
Postgraduate schooling	106 (31%)
Relationship status	
Single	37 (11%)
Married	292 (85%)
Separated or divorced	14 (4%)
Infant variables	
Infant sex	
Male	161 (45%)
Female	195 (55%)
Infant race	
White	337 (100%)
Other	0 (0%)
Attended day care	
4 months of age	118 (33%)
8 months of age	135 (38%)
12 months of age	147 (41%)
Household tap water (As)	0.48 ^b (< 0.01–79.7)
< 1 $\mu\text{g/l}$	205 (58%)
1–10 $\mu\text{g/l}$	101 (28%)
> 10 $\mu\text{g/l}$	50 (14%)

^aSum of subjects for the following variables are less than the total sample size (n missing): maternal education (13), relationship status (13), and infant race (19). ^bMedian concentration and range is reported, as data are log-normally distributed.

formula decreased from 337 ml/day and 6 breast-feedings/day at 4 months to 277 ml/day and 4 breast-feedings per day at 12 months of age (Supplementary Tables S2 and S3). Breastfed infants had an average of 7 feedings/day at 4 months; by 12 months this decreased to 4 feedings/day (Supplementary Table S3).

Our models assumed all formula was made with tap water in order to estimate potential exposure among users of private wells. Many families reported using home tap water to reconstitute formula powder, and the use of home tap water increased slightly over the first year of life. Among Formula-fed infants, the percentage of mothers who reported always using tap water to prepare formula increased from 57% at 4 months to 65% at 12 months. However, there were some differences in tap water arsenic concentrations by use of tap water to prepare formula: a higher geometric mean (GM) tap water arsenic concentration was found among those who reported never using tap water to prepare formula (0.88 $\mu\text{g/l}$) compared with than those who reported using tap water to mix formula most of the time at 8 months of age (0.45 $\mu\text{g/l}$; $P = 0.04$) and 12 months of age (1.11 vs 0.36 $\mu\text{g/l}$; $P = 0.002$), but not at 4 months of age (0.71 vs 0.54 $\mu\text{g/l}$, $P = 0.42$).

Arsenic Exposure Estimates for NHBCS Infants via Breast Milk and Formula Made with Home Tap Water

Results from both the central tendency and upper-bound models have distributions owing to the use of individual-level data on feeding type (Breastfed, Mixed or Formula-fed) and the concentration of arsenic in household tap water at 4, 8 and 12 months. Based on the central tendency model, GM estimated arsenic exposure




Figure 1. Proportion of NHBCS infants fed formula and breast milk at 4, 8 and 12 months of age.

across the NHBCS population was relatively low, ranging from 0.1 $\mu\text{g/kg/day}$ at 4 months of age to 0.07 $\mu\text{g/kg/day}$ at 12 months of age (Table 2). However, GM estimated exposure from this model was approximately eight times higher for Formula-fed infants than for Breastfed infants at all time points (Table 2, Supplementary Figure S1), owing to the presence of low concentrations of arsenic in both the formula powder and household tap water (Table 3). GM exposure among Formula-fed, Mixed and Breastfed infants was 55, 67 and 58% lower at 12 months compared with 4 months, respectively. Including all feeding types, population-wide GM exposure decreased approximately 27% from 4 to 12 months owing to decreasing IR_{BW} (Figure 2a).

Based on the upper-bound model, population-wide GM estimated arsenic exposures were slightly higher across time points, ranging from 0.22 $\mu\text{g/kg/day}$ at 4 months of age to 0.24 $\mu\text{g/kg/day}$ at 12 months of age (Table 2). GM estimated exposures were 5–6 times higher for Formula-fed infants than for Breastfed infants (Table 2, Supplementary Figure S1), and exposures decreased from 4 to 12 months in all feeding types. Unlike the central tendency model, population-wide GM exposure from the upper-bound model increased 8% from 4 to 12 months of age in conjunction with increased prevalence of formula consumption at the population level (Figure 2b).

Variability in estimated exposure among Formula-fed infants was due to the concentration of arsenic in household tap water. The most exposed Formula-fed infant from our central tendency model was estimated to be exposed to 73 times more arsenic per kg body mass per day than the least exposed Formula-fed infant (Supplementary Table S5), and >300 times more arsenic than Breastfed infants (Table 2, Supplementary Table S5). At low concentrations of tap water arsenic, approximately half of exposure among Formula-fed infants was attributable to the formula powder itself (Table 3).

Using the central tendency model but excluding the 14% of infants whose tap water exceeded the current US EPA MCL of 10 $\mu\text{g/l}$, GM exposure among Formula-fed infants was ~30% lower than for the full subset (0.19 $\mu\text{g/kg/day}$ at 4 months and 0.09 $\mu\text{g/kg/day}$ at 12 months of age; Supplementary Table S6) and 6–9 times higher for Formula-fed infants compared with Breastfed infants (Table 2, Supplementary Table S7).

DISCUSSION

Estimated GM arsenic exposures in the NHBCS were generally low regardless of feeding type and exposure model. However, the estimated upper-bound exposures for infants fed exclusively formula reflected the elevated arsenic concentrations in tap water, which reached 79.7 $\mu\text{g/l}$. The 8 \times increase in arsenic exposure among Formula-fed compared with Breastfed infants is consistent with our previous finding for 6-week-old NHBCS infants,³³ with

Table 2. Estimated exposure to arsenic ($\mu\text{g}/\text{kg}/\text{day}$) during the first year of life.

	<i>All</i>		<i>Breastfed</i>		<i>Mixed</i>		<i>Formula fed</i>	
	% (n)	GM (95% CI)	% (n)	Value ^a	% (n)	GM (95% CI)	% (n)	GM (95% CI)
<i>(A) Central tendency model</i>								
4 months	100 (356)	0.10 (0.09, 0.12)	45 (159)	0.03	24 (85)	0.20 (0.16, 0.25)	31 (112)	0.28 (0.23, 0.34)
8 months	100 (356)	0.09 (0.08, 0.10)	33 (119)	0.02	16 (58)	0.13 (0.10, 0.17)	50 (179)	0.20 (0.17, 0.24)
12 months	100 (356)	0.07 (0.07, 0.08)	21 (73)	0.01	13 (48)	0.07 (0.05, 0.09)	66 (235)	0.13 (0.11, 0.15)
<i>(B) Upper-bound model</i>								
4 months	100 (356)	0.22 (0.20, 0.24)	45 (159)	0.09	24 (85)	0.37 (0.30, 0.44)	31 (112)	0.51 (0.43, 0.61)
8 months	100 (356)	0.24 (0.21, 0.26)	33 (119)	0.08	16 (58)	0.30 (0.24, 0.38)	50 (179)	0.45 (0.40, 0.52)
12 months	100 (356)	0.24 (0.21, 0.26)	21 (73)	0.06	13 (48)	0.21 (0.17, 0.25)	66 (235)	0.37 (0.33, 0.41)

Abbreviations: CI, confidence interval; GM, geometric mean. These estimates were calculated using central tendency or upper-bound inputs for the body weight-adjusted ingestion rate, the concentration of arsenic in infant formula powder and the concentration of arsenic in breast milk. Variability among infants is due to individual-level data on the concentration of arsenic in tap water (for Formula-fed and Mixed feeding infants) and change in feeding mode over time.

^aThere is no distribution for breastfed infants because all infants had the same input values for body weight-adjusted ingestion rate and breast milk arsenic concentration.

Table 3. Estimated exposure to arsenic ($\mu\text{g}/\text{kg}/\text{day}$) from formula powder compared with mixing water among exclusively Formula-fed infants.

	<i>Formula powder</i>		<i>Mixing water</i>		<i>Total arsenic</i>		<i>% from powder</i>	
	GM	Max	GM	Max	GM	Max	GM	Max
<i>(A) Central tendency model</i>								
4 months	0.12	0.16	8.92	0.28	9.05	44%	1%	
8 months	0.08	0.12	5.98	0.20	6.06	41%	1%	
12 months	0.05	0.07	3.74	0.13	3.80	41%	1%	
<i>(B) Upper-bound model</i>								
4 months	0.27	0.24	11.8	0.51	12.1	52%	2%	
8 months	0.23	0.23	9.96	0.45	10.2	50%	2%	
12 months	0.18	0.19	8.05	0.37	8.23	49%	2%	

Abbreviations: GM, geometric mean; Max, maximum value across individuals.

approximately half of exposure attributable to the powdered component of formula when the arsenic concentration in mixing water was relatively low ($\sim 1 \mu\text{g}/\text{l}$). Importantly, GM exposures were similar even when the analysis was restricted to infants whose tap water arsenic was below the current MCL of $10 \mu\text{g}/\text{l}$, suggesting that this finding is generalizable to the US infant population consuming formula reconstituted with water from regulated sources. The reduction in exposure within each feeding type from 4 to 12 months reflects the decrease in liquid consumption per unit body weight over the first year of life, owing both to growth and the introduction of solid foods. However, this decrease was attenuated at the population level, when all feeding types were included, owing to the 35% increase in the percentage of Formula-fed infants in the population from 4 to 12 months.

Although our models are for total arsenic and did not explicitly consider arsenic speciation, the majority of arsenic in tap water is iAs. Thus, we expect that much of the exposure to arsenic via formula could be in the more toxic, inorganic form, especially for infants receiving formula made with water high in arsenic. Estimates of iAs in formula powder range from at least 50%, an assumption applied by the FDA,³² to nearly 100% inorganic.³¹ In contrast, the dominant form of arsenic in breast milk is currently unclear: we were unable to measure the percentage of iAs in breast milk owing to the low total arsenic concentrations,³³ and previous studies have reported contradictory findings.^{34,37}

The presence of iAs in reconstituted infant formula is of particular concern because, at 4 months of age, we estimate that

16% of our study population (29% when restricted to Mixed- and Formula-fed infants) exceeded the reference dose (RfD) for chronic oral ingestion of arsenic calculated by the US EPA⁴⁴— $0.3 \mu\text{g}/\text{kg}/\text{day}$ —using the central tendency model. Although a useful benchmark, it must be acknowledged that the EPA RfD is based on a no-effect level calculated for adult chronic lifetime exposure to arsenic in drinking water and does not indicate potential cancer risks. The US EPA is in the process of re-evaluating the arsenic RfD based on additional data that allow a more comprehensive assessment of potential non-cancer risks of arsenic for vulnerable early-life stages as well as for lifetime exposure.⁴⁵

Our exposure models included a number of assumptions that were applied in the absence of subject-specific body weights and precise measurements of breast milk consumption and were selected with the aim of estimating potential exposure for the population. These assumptions lead to expected underestimates and overestimates of true exposure, which we evaluate below using sensitivity analysis and/or qualitative discussion:

- (1) Our models assumed that all formula was prepared using home tap water, whereas 30% of mothers in our study population reported using primarily bottled water to prepare formula. Because bottled water is regulated to have arsenic concentrations below the current MCL,⁴⁶ this assumption would lead to an expected overestimate in our upper percentile estimates from either model. However, it is expected to have minimal impact on our GM estimates because 86% of our study population had tap water concentrations below the MCL and our findings were robust to the inclusion of NHBCS infants with home tap water arsenic concentrations above the MCL (Supplementary Tables S6 and S7). Also, this assumption is expected to improve generalizability of our findings to the general population of New Hampshire infants from homes with residential wells, as our study population was provided with their tap water results and thus may have been less likely to mix formula using tap water containing elevated concentrations of arsenic than the typical New Hampshire family.
- (2) Our models assumed that Mixed-fed infants received 50% breast milk, which would lead to an expected overestimate of exposure in this feeding type at 4 months, but an underestimate at 12 months, owing to a switch from predominantly breast milk at 4 months of age to predominantly formula at 12 months of age (Supplementary Tables S2 and S3).
- (3) Our models applied the IR_{BW} for breast milk⁴³ to all feeding types and generally assumed no exposure to tap water other than via formula. These assumptions may underestimate exposure as the IR_{BW} —and thus arsenic exposure—may be




Figure 2. Arsenic exposure ($\mu\text{g}/\text{kg}/\text{day}$) from formula and breast milk among NHBCS infants ($n = 356$) at 4, 8 and 12 months of age using the (a) central tendency model and (b) upper-bound model, which use central tendency or upper-bound inputs, respectively, for the body weight-adjusted ingestion rate, infant formula powder and breast milk. Variability between infants is due to individual-level data on feeding type and measured concentrations of arsenic in home tap water. Each symbol indicates one infant, the boxes denote the first and third quartile (Q1 and Q3) with the median as a line in the middle, and the whiskers denote points within 1.5 times the interquartile range of Q1 and Q3. There is no variability in Breastfed infants because all were modeled with the same inputs.

higher for formula, as confirmed by our sensitivity analysis using the calculated IR_{BW} for the subset of Formula-fed NHBCS infants that had data available for both ingestion rate and body weight. In models using this NHBCS-derived IR_{BW} , central tendency estimated exposures were 11–35% higher than estimates from the primary models (Supplementary Table S6).

- (4) The IR_{BW} applied in the upper-bound model (mean+2 SD) for all feeding types may lead to an expected overestimate as this rate is based on a 24-h time frame, and extreme values in the short term will not be maintained.
- (5) The maximum concentration of arsenic in breast milk may be underestimated as it is based on a small sample and the maximum concentration of arsenic in home tap water in that sample ($8.9 \mu\text{g}/\text{l}$) was lower than in our study population ($79.7 \mu\text{g}/\text{l}$). However, populations with high exposures, such as those in Bangladesh and Chile, have also reported low levels in breast milk,^{34–39} suggesting that the impact of this underestimate on exposures from our upper-bound model would be relatively small.

Finally, the concentrations of arsenic in breast milk and formula powder used in our model were taken from previous studies in the absence of individual-level data. However, our confidence in the appropriateness of these data for our study population is high as they were measured for the NHBCS population and values are comparable to those from previous studies: total arsenic concentrations in formula were similar to those reported by the US FDA³² and concentrations in breast milk were similar to reports from other countries.^{34–39}

In conclusion, modeled arsenic exposures via breast milk and formula were relatively low throughout the first year of life, unless formula was prepared with arsenic-contaminated tap water. We expect that the transition to solid foods, which typically occurs beginning at ~ 6 months, presents an additional source of exposure to arsenic for the typical NHBCS infant, as many foods commonly fed to infants during weaning, such as infant rice cereal, contain elevated concentrations of arsenic.^{31,47,48} Thus

future work should seek to quantify daily arsenic intake during this critical period of development.

CONFLICT OF INTEREST

The authors declare no conflict of interest.

ACKNOWLEDGEMENTS

We thank the participants and staff members of the New Hampshire Birth Cohort Study and members of the Cottingham Laboratory for their assistance. Archana Ramanujan and David Fried conducted the medical record reviews for infant body weights. Patricia Fabian and three anonymous reviewers provided helpful comments on a draft manuscript. This publication was supported in part by grants P01 ES022832 and P20 ES018175 from the National Institute of Environmental Health Sciences (NIEHS) and RD83459901 and RD83544201 from the Environmental Protection Agency (EPA). Contents are solely the responsibility of the grantee and do not necessarily represent the official views of the US EPA. Further, the US EPA does not endorse the purchase of any commercial products or services mentioned in the publication.

REFERENCES

- 1 Farzan SF, Karagas MR, Chen Y. In utero and early life arsenic exposure in relation to long-term health and disease. *Toxicol Appl Pharmacol* 2013; **272**: 384–390.
- 2 Naujokas MF, Anderson B, Ahsan H, Vasken Aposhian H, Graziano JH, Thompson C *et al*. The broad scope of health effects from chronic arsenic exposure: update on a worldwide public health problem. *Environ Health Perspect* 2013; **121**: 8.
- 3 Rahman A, Persson LA, Nermell B, El Arifeen S, Ekstrom EC, Smith AH *et al*. Arsenic exposure and risk of spontaneous abortion, stillbirth, and infant mortality. *Epidemiology* 2010; **21**: 797–804.
- 4 Vahter M. Health effects of early life exposure to arsenic. *Basic Clin Pharmacol Toxicol* 2008; **102**: 204–211.
- 5 Roy A, Kordas K, Lopez P, Rosado JL, Cebrian ME, Vargas GG *et al*. Association between arsenic exposure and behavior among first-graders from Torreon, Mexico. *Environ Res* 2011; **111**: 670–676.
- 6 Hamadani JD, Tofail F, Nermell B, Gardner R, Shiraji S, Bottai M *et al*. Critical windows of exposure for arsenic-associated impairment of cognitive function in pre-school girls and boys: a population-based cohort study. *Int J Epidemiol* 2011; **40**: 1593–1604.

- 7 Kwok RK, Kaufmann RB, Jakariya M. Arsenic in drinking-water and reproductive health outcomes: a study of participants in the Bangladesh integrated nutrition programme. *J Health Popul Nutr* 2006; **24**: 190–205.
- 8 Steinmaus C, Ferreccio C, Acevedo J, Yuan Y, Liaw J, Duran V *et al*. Increased lung and bladder cancer incidence in adults after in utero and early-life arsenic exposure. *Cancer Epidemiol Biomarkers Prev* 2014; **23**: 1529–1538.
- 9 Bodwell JE, Kingsley LA, Hamilton JW. Arsenic at very low concentrations alters glucocorticoid receptor (GR)-mediated gene activation but not GR mediated gene repression; complex dose response effects are closely correlated with levels of activated GR and require a functional GR DNA binding domain. *Chem Res Toxicol* 2004; **17**: 1064–1076.
- 10 Farzan SF, Korrick S, Li Z, Enelow R, Gandolfi AJ, Madan J *et al*. In utero arsenic exposure and infant infection in a United States cohort: a prospective study. *Environ Res* 2013; **126**: 24–30.
- 11 Argos M, Kalra T, Rathouz PJ, Chen Y, Pierce B, Parvez F *et al*. Arsenic exposure from drinking water, and all-cause and chronic-disease mortalities in Bangladesh (HEALS): a prospective cohort study. *Lancet* 2010; **376**: 252–258.
- 12 Chen CJ, Wang CJ. Ecological correlation between arsenic level in well water and age-adjusted mortality from malignant neoplasms. *Cancer Res* 1990; **50**: 5470–5474.
- 13 Samanta G, Sharma R, Roychowdhury T, Chakraborti D. Arsenic and other elements in hair, nails, and skin-scales of arsenic victims in West Bengal, India. *Sci Total Environ* 2004; **326**: 33–47.
- 14 Cubadda F, Aureli F, D'Amato M, Raggi A, Turco AC, Mantovani A. Speciated urinary arsenic as a biomarker of dietary exposure to inorganic arsenic in residents living in high-arsenic areas in Latium, Italy. *Pure Appl Chem* 2012; **84**: 203–214.
- 15 Karagas MR, Stukel TA, Tosteson TD. Assessment of cancer risk and environmental levels of arsenic in New Hampshire. *Int J Hyg Environ Health* 2002; **205**: 85–94.
- 16 International Agency for Research on Cancer (IARC). *Some Drinking-Water Disinfectants and Contaminants, Including Arsenic*. World Health Organization: Lyon, France, 2004.
- 17 Edmonds JS, Francesconi KA. Arsenic in seafoods - human health-aspects and regulations. *Mar Pollut Bull* 1993; **26**: 665–674.
- 18 Meharg A, Sun G, Williams PN, Adomako EE, Deacon C, Zhu Y-G *et al*. Inorganic arsenic levels in baby rice are of concern. *Environ Pollut* 2008; **152**: 746–749.
- 19 Tseng CH. A review on environmental factors regulating arsenic methylation in humans. *Toxicol Appl Pharmacol* 2009; **235**: 338–350.
- 20 Gosse JA, Taylor VF, Jackson BP, Hamilton JW, Bodwell JE. Monomethylated trivalent arsenic species disrupt steroid receptor interactions with their DNA response elements at non-cytotoxic cellular concentrations. *J Applied Toxicol* 2014; **34**: 498–505.
- 21 Medeiros M, Zheng X, Novak P, Wnek SM, Chyan V, Escudero-Lourdes C *et al*. Global gene expression changes in human urothelial cells exposed to low-level monomethylarsonous acid. *Toxicology* 2012; **291**: 102–112.
- 22 Styblo M, Del Razo LM, Vega L, Germolec DR, LeCluyse EL, Hamilton GA *et al*. Comparative toxicity of trivalent and pentavalent inorganic and methylated arsenicals in rat and human cells. *Arch Toxicol* 2000; **74**: 289–299.
- 23 Petrick JS, Ayala-Fierro F, Cullen WR, Carter DE, Vasken Aposhian H. Monomethylarsonous acid (MMA(III)) is more toxic than arsenite in Chang human hepatocytes. *Toxicol Appl Pharmacol* 2000; **163**: 203–207.
- 24 Cohen SM, Arnold LL, Eldan M, Lewis AS, Beck BD. Methylated arsenicals: the implications of metabolism and carcinogenicity studies in rodents to human risk assessment. *Crit Rev Toxicol* 2006; **36**: 99–133.
- 25 Cohen SM, Arnold LL, Beck BD, Lewis AS, Eldan M. Evaluation of the carcinogenicity of inorganic arsenic. *Crit Rev Toxicol* 2013; **43**: 711–752.
- 26 European Food Safety Authority (EFSA). EFSA Panel on Contaminants in the Food Chain (CONTAM): Scientific Opinion on Arsenic in Food. *EFSA J* 2009; **7**: 1351.
- 27 Tsuji JS, Yost LJ, Barraj LM, Scrafford CG, Mink PJ. Use of background inorganic arsenic exposures to provide perspective on risk assessment results. *Regul Toxicol Pharmacol* 2007; **48**: 59–68.
- 28 Hernandez-Martinez R, Navarro-Blasco I. Survey of total mercury and arsenic content in infant cereals marketed in Spain and estimated dietary intake. *Food Control* 2013; **30**: 423–432.
- 29 Ljung K, Palm B, Grandér M, Vahter M. High concentrations of essential and toxic elements in infant formula and infant foods - a matter of concern. *Food Chem* 2011; **127**: 943–951.
- 30 Sorbo A, Turco AC, Di Gregorio M, Ciaralli L. Development and validation of an analytical method for the determination of arsenic, cadmium and lead content in powdered infant formula by means of quadrupole inductively coupled plasma mass spectrometry. *Food Control* 2014; **44**: 159–165.
- 31 Jackson BP, Taylor VF, Punshon T, Cottingham KL. Arsenic concentration and speciation in infant formulas and first foods. *Pure Appl Chem* 2012; **84**: 215–223.
- 32 U.S. Food and Drug Administration. Analytical Results from Inorganic Arsenic in Rice and Rice Products Sampling. Arsenic in Rice and Rice Products. 2013; <http://www.fda.gov/downloads/Food/FoodbornellnessContaminants/Metals/UCM352467.pdf>. (accessed 4 September 2015).
- 33 Carignan CC, Cottingham KL, Jackson BP, Farzan SF, Gandolfi AJ, Punshon T *et al*. Estimated exposure to arsenic in breastfed and formula-fed infants in a United States Cohort. *Environ Health Perspect* 2015; **123**: 500–506.
- 34 Björklund KL, Vahter M, Palm B, Grandér M, Lignell S, Berglund M. Metals and trace element concentrations in breast milk of first time healthy mothers: a biological monitoring study. *Environ Health* 2012; **11**: 92.
- 35 Samanta G, Das D, Mandal BK, Chowdhury TR, Chakraborti D, Pal A *et al*. Arsenic in the breast milk of lactating women in arsenic-affected areas of West Bengal, India and its effect on infants. *J Environ Sci Health A Tox Hazard Subst Environ Eng* 2007; **42**: 1815–1825.
- 36 Concha G, Vogler G, Nermell B, Vahter M. Low-level arsenic excretion in breast milk of native Andean women exposed to high levels of arsenic in the drinking water. *Int Archiv Occup Environ Health* 1998; **71**: 42–46.
- 37 Fängström B, Moore S, Nermell B, Kuenstl L, Goessler W, Grandér M *et al*. Breast-feeding protects against arsenic exposure in Bangladeshi infants. *Environ Health Perspect* 2008; **116**: 963–969.
- 38 Sternowsky HJ, Moser B, Szadkowsky D. Arsenic in breast milk during the first 3 months of lactation. *Int J Hyg Environ Health* 2002; **205**: 405–409.
- 39 Islam MR, Attia J, Alauddin M, McEvoy M, McElduff P, Slater C *et al*. Availability of arsenic in human milk in women and its correlation with arsenic in urine of breastfed children living in arsenic contaminated areas in Bangladesh. *Environ Health* 2014; **13**: 101.
- 40 Vahter ME, Li L, Nermell B, Rahman A, El Arifeen S, Rahman M *et al*. Arsenic exposure in pregnancy: a population-based study in Matlab, Bangladesh. *J Health Popul Nutr* 2006; **24**: 236–245.
- 41 Centers for Disease Control and Prevention (CDC). Breastfeeding Report Card, United States, 2013. Centers for Disease Control and Prevention, National Center for Chronic Disease Prevention and Health Promotion, Division of Nutrition, Physical Activity and Obesity. <http://www.cdc.gov/breastfeeding/pdf/2013breastfeedingreportcard.pdf>.
- 42 Gilbert-Diamond D, Cottingham KL, Gruber JF, Punshon T, Sayarath V, Gandolfi AJ *et al*. Rice consumption contributes to arsenic exposure in US women. *Proc Natl Acad Sci USA* 2011; **108**: 20656–20660.
- 43 U.S. Environmental Protection Agency (US EPA). *Child-Specific Exposure Factors Handbook (Final Report)*. U.S. Environmental Protection Agency: Washington, DC, USA, 2008. <http://cfpub.epa.gov/ncea/cfm/recordisplay.cfm?deid=1992432008>.
- 44 U.S. Environmental Protection Agency. Chronic Health Hazard Assessments for Noncarcinogenic Effects, Reference Dose for Chronic Oral Exposure (RfD). Integrated Risk Information System (IRIS) on inorganic arsenic. Verification date February 3, 1994. <http://www.epa.gov/iris/subst/0278.htm>.
- 45 National Research Council. *Critical Aspects of EPA's IRIS Assessment of Inorganic Arsenic: Interim Report*. National Academies Press: Washington, DC, USA, 2013. http://www.nap.edu/catalog.php?record_id=18594.
- 46 Josyula AB, McClellan H, Hysong TA, Kurzius-Spencer M, Poplin GS, Sturup S *et al*. Reduction in urinary arsenic with bottled-water intervention. *J Health Popul Nutr* 2006; **24**: 298–304.
- 47 Meharg AA, Sun GX, Williams PN, Adomako E, Deacon C, Zhu YG *et al*. Inorganic arsenic levels in baby rice are of concern. *Environ Pollut* 2008; **152**: 746–749.
- 48 Lai PY, Cottingham KL, Steinmaus C, Karagas MR, Miller MD. Arsenic and rice: translating research to address health care providers' needs. *J Pediatr* 2015; **167**: 797–803.

Supplementary Information accompanies the paper on the Journal of Exposure Science and Environmental Epidemiology website (<http://www.nature.com/jes>)