

REVIEW

Markers of microglia in post-mortem brain samples from patients with Alzheimer's disease: a systematic review

KE Hopperton¹, D Mohammad^{1,2}, MO Trépanier¹, V Giuliano¹ and RP Bazinet¹

Neuroinflammation is proposed as one of the mechanisms by which Alzheimer's disease pathology, including amyloid- β plaques, leads to neuronal death and dysfunction. Increases in the expression of markers of microglia, the main neuroimmune cell, are widely reported in brains from patients with Alzheimer's disease, but the literature has not yet been systematically reviewed to determine whether this is a consistent pathological feature. A systematic search was conducted in Medline, Embase and PsychINFO for articles published up to 23 February 2017. Papers were included if they quantitatively compared microglia markers in post-mortem brain samples from patients with Alzheimer's disease and aged controls without neurological disease. A total of 113 relevant articles were identified. Consistent increases in markers related to activation, such as major histocompatibility complex II (36/43 studies) and cluster of differentiation 68 (17/21 studies), were identified relative to nonneurological aged controls, whereas other common markers that stain both resting and activated microglia, such as ionized calcium-binding adaptor molecule 1 (10/20 studies) and cluster of differentiation 11b (2/5 studies), were not consistently elevated. Studies of ionized calcium-binding adaptor molecule 1 that used cell counts almost uniformly identified no difference relative to control, indicating that increases in activation occurred without an expansion of the total number of microglia. White matter and cerebellum appeared to be more resistant to these increases than other brain regions. Nine studies were identified that included high pathology controls, patients who remained free of dementia despite Alzheimer's disease pathology. The majority (5/9) of these studies reported higher levels of microglial markers in Alzheimer's disease relative to controls, suggesting that these increases are not solely a consequence of Alzheimer's disease pathology. These results show that increased markers of microglia are a consistent feature of Alzheimer's disease, though this seems to be driven primarily by increases in activation-associated markers, as opposed to markers of all microglia.

Molecular Psychiatry (2018) **23**, 177–198; doi:10.1038/mp.2017.246; published online 12 December 2017

INTRODUCTION

Elevations in neuroinflammatory markers are widely reported in Alzheimer's disease (AD), both in animal models^{1–3} and human subjects.^{4–7} This has contributed to the development of the neuroinflammatory hypothesis of AD that suggests that aberrant activation of immune cells may drive neuronal death and dysfunction in AD.⁸ This is supported by genome-wide association studies that have identified polymorphisms in inflammation-associated genes as risk factors for the development of AD.^{9–11}

Microglia are the resident immune cells of the brain, and are thought to be the main cells responsible for initiating the immune response to AD pathology. Several of the inflammation-associated genetic risk factors for AD, including human leukocyte antigen (HLA)-DRB1/B5,⁹ cluster of differentiation (CD) 33,¹² triggering receptor expressed on myeloid cell-2 (TREM2)¹⁰ and phospholipase C γ 2,¹⁰ are highly expressed in microglia where they are involved in cell function and activation. This suggests that aberrant microglial activation is a causal factor in the development of AD, as opposed to a consequence of AD pathology. Although it is commonly accepted that there are increased microglia markers in the brains of patients with AD relative to controls, no one has yet systematically synthesized the literature to see whether this is

supported by the totality of the evidence. Here, we describe the results of a systematic review examining microglia in post-mortem human brain samples from patients with AD and healthy controls. We find that some markers associated with cell activation, such as major histocompatibility complex- II (MHCII) and CD68, are consistently increased in the AD brain, but that studies using other common microglial markers that stain both resting and activated cells, such as ionized calcium-binding adaptor molecule-1 (Iba1) and CD11b, are heterogeneous and do not demonstrate a consistent elevation. We further identify brain regions, such as the white matter and the cerebellum, that appear to be more resistant to inflammation in AD.

METHODS

The systematic search was conducted in Medline, Embase and PsychINFO covering articles published up to 23 February 2017. The search protocol was developed based on Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) and World Health Organization (WHO) Review Protocol Template Guidelines where applicable for a systematic review of descriptive (non-interventional) data, and is provided in Appendix 1 of the Supplementary Materials section. The search queried the

¹Department of Nutritional Sciences, Faculty of Medicine, University of Toronto, Toronto, ON, Canada and ²Department of Pharmacology and Toxicology, University of Toronto, Toronto, ON, Canada. Correspondence: Dr RP Bazinet, Department of Nutritional Sciences, Faculty of Medicine, University of Toronto, FitzGerald Building, 150 College Street, Room 306, Toronto, ON M5S 3E2, Canada.

E-mail: richard.bazinet@utoronto.ca

Received 2 June 2017; revised 15 August 2017; accepted 14 September 2017; published online 12 December 2017




Figure 1. Flow diagram of systematic search.

following terms with numerous synonyms and related words as both MeSH/Emtree terms (where applicable) and as keywords (for title, abstract and keyword searches): Alzheimer's Disease AND brain AND inflammation. The additional term 'AND post-mortem' with its synonyms and related words was included for the Embase and PsychINFO searches. A full list of terms used for each search can be found in Appendix 2.

Studies were included if they used human brain samples from patients with AD, included a measure of inflammation, were conducted post-mortem and included a comparison to a control group without AD or a confounding neurological disease. This review initially set out to include all inflammatory markers, including for astrocytes, complement, cytokines, lipid mediators and other immune cells, but the title and abstract screening returned 757 eligible papers, making it unfeasible to summarize all the evidence in a single paper. It was decided to proceed with a review of microglia, as these are the main immune effector cells in the brain, and are the source of many of the other inflammatory mediators measured in other studies. Microglia terms in the initial search include the MeSH term: neuroglia and neurogenic inflammation, the Emtree terms: neurogenic inflammation, glia and leukocyte antigen, and keywords: microglia, HLA, MHC, CD11b, CD68, Iba1, OX-42 and CD45 along with their synonyms and alternate spellings (see Appendix 2 of the Supplementary Materials for the full list of terms). In addition, any other papers returned in the full search that used other markers identified by the study authors as being specific to microglia or their activation were included. Studies were excluded if they were not in English or not published in full in a peer-reviewed journal. Papers that measured markers associated with an M1 or M2 phenotype, such as interleukin (IL)-1 β , tumour necrosis factor- α , IL-4 or IL-10 but that did not localize these markers specifically to microglia, were not included.

Data from included studies were extracted by at least two independent reviewers (KEH and DM, VG or MT), with a third reviewer employed in the case of a conflict. Extracted data included origin of brain samples, number of subjects, sex, age, apolipoprotein E (APOE) genotype, histological confirmation of AD status, Braak stage, control history of neurological or psychiatric

disease, post-mortem interval (length of time between death and retrieval of brain), brain regions examined, medications at the time of death, anti-inflammatory drug use at death, technique for measuring microglia, marker of microglia used and the results of the comparison between the AD subjects and healthy controls. Where available, information on relationship between labelled microglia and amyloid- β plaques or neurofibrillary tangles was also extracted. The terms 'higher' or \uparrow and 'lower' or \downarrow are used in the text and tables referring to significantly higher or lower levels of the microglial marker used in the study relative to the nonneurological aged control group. As the type of outcome reporting was extremely heterogeneous, results were reported as higher, lower or unchanged for AD relative to control as identified by the study authors. Meta-analysis or other summary statistics were not used because of the large variability in assessment techniques and brain regions examined between studies. Data were listed as 'Not Reported' if the relevant information could not be found in the article, or in a previous article specifically referred to by the authors in the Methods section. In a few instances where information in the article was unclear, the corresponding authors were contacted to provide clarification or additional detail.

RESULTS

A total of 22 229 articles were screened, of which 757 met inclusion for full text review for inflammatory markers, including 224 that examined microglia (Figure 1). A total of 113 papers that quantified microglia and compared the results statistically between control and AD groups were fully extracted and are presented in the tables below (see Figure 2 for a visual summary of the results). Fifty-three papers that made a nonquantitative comparison between AD and control, often as a qualitative assessment of the intensity of immunohistochemistry staining, were not fully extracted but are summarized under the 'Non-quantitative comparisons' heading.

The microglial markers analysed in this review all serve distinct functions within the cell, and as such, the interpretation of an up-




Figure 2. Summary results of systematic search. Size of circle is proportional to the number of identified studies, whereas the colour and position on the graph illustrates the percentage of studies that identified an increase in AD in at least one brain region. AD, Alzheimer's disease; CD, cluster of differentiation; Iba1, ionized calcium-binding adaptor molecule 1; IL, interleukin; MHC, major histocompatibility complex; RCA-1, Ricinus communis agglutinin-1; TSPO, translocator protein; TREM2, triggering receptor expressed on myeloid cells 2.

or down-regulation of expression will vary depending on the marker used. The functions of all major markers included in this review, their association with polarization and their expression on other cell types besides microglia is shown in Supplementary Table 1. M1 and M2 polarization is increasingly regarded as an oversimplification of many diverse functions of activated microglia; however, it is used here to provide a reference for whether a marker is associated with the pro-inflammatory or phagocytic microglial phenotype. As shown in the table, it is important to note that all the major markers in this review are also expressed on other cell types, particularly on macrophages, and hence it is possible that other infiltrating or perivascular immune cells may have contributed to the results.

Major histocompatibility complex II

MHCII is expressed on the surface of antigen-presenting cells and is responsible for antigen recognition and the activation of the adaptive immune system. Within the brain, MHCII is primarily expressed on microglia, where it is generally considered a marker of activated cells, though it may have weaker expression in resting cells.¹⁴ Forty-three papers were identified that quantitatively compared markers of MHCII between AD and control in post-mortem human brains (Table 1). The majority (41/43) used immunohistochemistry for HLA-DR or for multiple isoforms of HLA (HLA-DR-DQ-DP) quantified by cell counts, scoring or staining area, whereas other techniques included gene expression by quantitative PCR (qPCR) or protein quantification by western blot.

Thirty-six papers reported higher MHCII in AD relative to control brains in at least one of the measured brain regions,^{7,15-49} whereas 7 identified no difference between AD and control,⁵⁰⁻⁵⁶ and 5 identified no difference in at least one brain region.^{21,32,34,41,44} Increased MHCII staining, counts or expression was noted in: the entorhinal cortex,^{7,18,20,35,43,47} frontal and temporal gyri,^{7,19,24,32,33,39,45,56} hippocampus,^{20,25,27,30,32-38,40,42-44,47} and frontal,^{18,21,25,40,41,44,46,48,49,52} temporal^{16-18,28,30,35,43,44,48,52} and occipital cortices.^{25,35,41,43,44} HLA-DR was found to increase with AD plaque stage and clinical dementia rating in the entorhinal cortex, hippocampus and occipital and temporal cortices.^{43,47} HLA-DR-stained microglia were reported to take on an activated morphology in AD,^{19,23,45,48,49} indicating that AD pathology may stimulate the activation of microglia and upregulation of MHCII. In contrast, seven papers identified no difference in MHCII between AD and control in several regions including the hippocampus,^{21,55} frontal, temporal and parietal grey matter,⁵² the

temporal polar association cortex⁵⁴ and the subventricular zone of the lateral ventricle.⁵³

Of the five studies examining HLA-DR immunoreactivity in the cerebellum, four found no differences between AD and control,^{25,34,44,51} suggesting that regional differences may explain some of the null findings. Overmyer *et al.*⁵² noted that women had higher HLA-DR reactivity than men in AD, whereas the reverse was true in the controls, suggesting that sex differences in the AD and control groups could also contribute to between-study variability. HLA-DR reactivity increased in both AD and control patients >75 of age in one study,⁵² but decreased in AD patients >80 of age in another,²⁸ making the effect of age on HLA-DR expression unclear. APOE genotype can also impact the results, with the ε4 risk allele increasing HLA-DR-positive cells or area in the frontal and temporal cortices⁴⁸ and middle frontal gyrus.⁵⁶

Ionized calcium-binding adaptor molecule 1

Iba1 is a cytoplasmic protein expressed in monocyte lineage cells and in the brain and is primarily restricted to microglia.⁵⁷ Although its expression is thought to increase with microglial activation,⁵⁸ where it may be involved in membrane ruffling and phagocytosis,⁵⁹ it is considered a marker of all microglia, rather than an activated subset.⁶⁰ Twenty papers quantitatively compared Iba1 between AD and control post-mortem human brains (Table 2). Ten studies identified increases in Iba1 cell counts, staining intensity or expression in AD compared with control samples in at least one brain region,^{20,60-68} whereas 10 studies identified no difference from controls^{32,54,69-76} and 5 reported lower Iba1 in AD than controls^{32,54,56,75,76} in at least one of the regions measured.

The results of studies measuring Iba1 are relatively heterogeneous. Higher Iba1 was noted in the AD frontal cortex in four studies,^{20,63,64,68} but was the same as control in three others^{32,70,74} and lower in the white matter of the frontal lobe in one study.³² Iba1-positive cell density⁷² and gene expression⁷⁴ was the same in AD and control in the temporal cortex in two studies, but greater by western blot or immunocytochemistry in three other studies.^{20,61,66} Similarly, although Iba1 was higher in AD than control in the hippocampus in three studies,^{20,62,67} it was unchanged in five studies^{32,69,71,73,75} and reduced in one study.⁷⁶ Iba1 was higher in the entorhinal cortex^{20,62} and inferior parietal cortex⁶⁵ in AD compared with control, but this is based on a limited number of studies. All but 1 of the 10 studies that identified increases in Iba1 in AD used expression-based assays.

Table 1. Major histocompatibility complex II (MHCII)

First Author	Brain bank	N	Sex	Age	AD genetic risk factors	AD histologically confirmed	Braak stage	C history of neurological or psychiatric disease	PMI (h)	Brain region	Technique	Marker	Results (AD vs C unless otherwise specified)
Akiyama ¹⁵	NR	AD: 9 C: 6	AD: 77 C: 69	NR	NR	NR	NR	No neurological disease	All within 2–12 h	Temporal lobe	IHC	HLA-DR	↑
Carpenter ¹⁹	NR	AD: 5 C: 5	AD 4/1 C: 5/0	AD: 75.4 C: 73.6	NR	Khachaturian	NR	No history of neurological or systemic diseases affecting the brain	AD: 3.6 C: 2.4	Grey matter of the middle temporal gyrus	IHC	HLA-DR (LN3)	↑ Density of cells per mm, staining area and percent of area. Most cells of resting morphology in Cs vs activated in AD (not compared statistically)
Dal Bianco ²⁰	NR	AD: 9 C: 15	AD: 0/9 C: 13/2	AD: 81 C: 70	NR	Braak, CERAD	AD: IV: 2, V: 4, VI: 3	No neurological disease or brain lesions	NR	Cortical areas of the temporal lobe, including entorhinal cortex, hippocampus and temporal cortex	Immunocytochemistry	MHCII	↑ MHCII
Desai ²¹	Religious Orders Study	AD:8 C: 7	AD: 3/5 C: 4/3	AD: 85.2 C: 79.5	NR	NIA-Reagan criteria	AD: III–VI	NR	AD: 5.4 C: 15.5	Hippocampus, midfrontal cortex, locus ceruleus, substantia nigra pars compacta	IHC	HLA-DR-DQ- DP (Cr3/43)	↑ In the midfrontal cortex, and locus ceruleus ↔ In density of HLA-DR microglia in hippocampus
Dhawan ²²	University of Washington ADRC	NR	NR	NR	NR	NR	NR	NR	NR	Temporal lobe	IHC	HLA-DR	↑ HLA-DR+ microglia
Mattiace ⁴⁹	NR	AD: 15 C: 14	AD: 2/13 C: 3/11	AD: 76.6 C: 67.8 (as young as 6 months)	NR	Khachaturian	NR	Depressive psychosis, manic depressive psychosis (n = 2), Parkinson's disease (n = 1)	AD: 6.4 C: 6.3	Midfrontal cortex (BA9)	IHC	HLA-DR, double staining with Leu-M5 (CD11c)	↑ Activated microglia (morphology) ↑ Proportion of HLA-1DR+ microglia in grey matter ↔ In white matter Results not compared statistically
Egensperger ⁴⁸	Institute of Neuropathology of the University of Munich	AD: 20 C: 5	AD: 5/15 C: NR	AD: 75.9 C: 71.6	APOE: AD: 3/3: 7 3/4: 10 4/4: 3 C: NR	Braak, CERAD	NR	No neurological or neuropathological disorder	NR	Frontal and temporal cortex	IHC	HLA-DR-DQ-DP (CR3/43)	↑ Counts and area —Plaque-associated microglia demonstrate activated morphology— Microglia number correlates with neuritic plaques and NFT
Flanary ²⁴	BSHRI	AD: 4 HPC: 3 C: 4	AD: 1/3 HPC: 2/1 C: 3/1	AD: 82.0 HPC: 87.7 C: 81.5	NR	Yes	NR	NR	AD: 2.4 HPC: 3.1 C: 2.6	Superior frontal and temporal gyri	IHC	HLA-DR	↑ Dystrophic microglia in AD vs C, HPC vs C and AD+HPC vs C ↑ HPC vs AD (not clear if statistically significant)
Giulian ²⁵	NR	AD: 6 C: 5	NR	NR	NR	CERAD	NR	No neuropathological disorder	NR	Cerebellum, hippocampus, frontal, occipital, parietal cortices and neocortical white matter	IHC, confocal microscopy	HLA-DR	↑ Hippocampus, frontal, occipital and parietal cortices ↔ Cerebellum, white matter
Gouw ²⁶	NBB and Vrije University Medical Centre	AD: 11 C: 7	AD: 3/8 C: 3/4	AD: 82.6 C: 78.3	NR	Braak, CERAD	AD: V C: I	All had white matter hyperintensities (small vessel disease), other neurological diseases excluded	AD: 6.1 C: < 24 h	Normal white matter and white matter hyperintensities in frontal, parietal and prefrontal lobes	IHC	HLA-DR	↑ Overall than C ↑ Higher in white matter hyperintensities than normal white matter

Table 1. (Continued)

First Author	Brain bank	N	Sex	Age	AD genetic risk factors	AD histologically confirmed	Braak stage	C history of neurological or psychiatric disease	PMI (h)	Brain region	Technique	Marker	Results (AD vs C unless otherwise specified)
Halliday ¹⁶	Dementia clinics at the Repatriation General Hospital Concord and Lidcombe Hospital in Sydney, Australia	AD: 12 C: 10	NR	AD: 79 C: 77	APOE: AD: 3/4: 1 C: 3/4: 1	CERAD, NIA-Reagan Criteria	NR	NR	All < 45 h, mean 19	Anterior cingulate, hippocampal, inferior temporal, parahippocampal and superior frontal regions	IHC	HLA-DR	↑ In AD vs C ↔ AD patients taking NSAIDs and AD patients not taking NSAIDs
Hensley ²⁷	NR	AD: 3 C: 3	Whole sample: AD: 9/13 C: 4/3	Whole sample: AD: 78.1 C: 79.7	NR	Khachaturian, Mirra	NR	No history of neurological and/or psychiatric disorders	Whole sample: AD: 4.6 C: 4.4	Cerebellum, hippocampus, inferior parietal lobule	IHC	HLA-DR	↑ In all regions
Hoozemans ¹⁷	NBB	Braak stage 0: 5, I-II: 16, III-IV: 10, V-VI: 9	Braak stage 0: 3/2, I-II: 6/10, III-IV: 3/6	Braak stage 0: 62, I-II: 83, III-IV: 89, V-VI: 76	NR	Braak	0-VI (not divided into C and AD)	NR	Braak stage 0: 8, I-II: 7.5, III-IV: 6.5, V-VI: 5	Temporal cortex	IHC	HLA-DR-DQ-DP (CR3/43)	↑ With increasing Braak NFT or plaque stage ($P < 0.05$ for trend), significant for NFT group V-VI vs O ↑, ↑ In AD patients >80 years compared with those > 80 years ↑
Hoozemans ²⁸	Netherlands Brain Bank	AD: 19 C: 19	AD: 3/16 C: 8/11	AD: 83.5 C: 76.8	APOE4: AD: 12 C: 8	Braak	AD: avg IV C: avg I	NR	AD: 5.1 C: 8.6	Midtemporal cortex	IHC	HLA	↑
Imamura ²⁹	NR	AD: 6 C: 6	AD: 2/4 C: 2/4	AD: 65.4 C: 62.8	NR	Yes	NR	NR	NR	Temporal lobe	IHC	HLA-DR	↑
Itagaki ³⁰	NR	AD: 10 C: 5	NR	NR	NR	Yes	NR	No neurological complications	All within 2-12	Mixed: hippocampus and temporal cortex	IHC: semiquantitative scoring of staining	HLA-DR, LCA	↑
Jantarotnotai ³¹	Kinsmen Laboratory Brain Bank at the University of British Columbia	AD severe: 9 AD mild: 6 C: 9	NR	AD severe: 74.2 AD mild: 77.7 C: 83	NR	Braak, NIA-Reagan criteria	NR	No neurological disorders	NR	Medial temporal cortex	IHC	HLA-DR	↑
Kellner ¹⁸	NR	AD: 48 C: 48	AD: 19/29 C: 24/24	AD: 80.3 C: 77.5	NR	Braak, CERAD	AD: II-VI (38 > 4) C: I-III (45 = 0)	NR	NR	Entorhinal, frontal cortex, temporal cortex	IHC	HLA-DR	↑
Korvatska ³²	University of Washington Neuropathology Core Brain Bank	AD (normal TREM2): 6 AD (with TREM2 R47H variant): 7 C: 3	NR	Whole sample: 84.9	NR	CERAD	AD: III: 1, V: 4, VI: 1, AD R47H: V: 6, VI: 1, C: 0: 1, I: 1, III: 1	NR, One C CERAD score A and one B	Whole sample: 4.5	Frontal lobe: grey and white matter Hippocampus: CA1, hilus, parahippocampal gyrus and white matter	IHC	MHCII	↑ Staining in hippocampus CA1, hilus, parahippocampal gyrus and white matter ↑ Activated counts in hippocampus white matter
Lopes ³³	BSHRI	AD: 7 young C: 3 Aged C: 7 HPC: 7	AD: 4/3 young C: 2/3 Aged C: 6/1 HPC: 5/2	AD: 80.3 young C: 36.3 aged C: 80.0 HPC: 83.4	NR	Yes	NR	NR	AD: 2.3 young C: 2.8 aged C: 2.5 HPC: 2.90	Amygdala, hippocampus, superior frontal gyrus, superior, middle, and inferior temporal gyri	IHC and morphometric analyses	HLA-DR	Microglia counts HLA-DR: ↑ vs all other groups Dystrophic microglia HLA-DR: ↑ vs C, ↔ vs HPC
Lue ⁷	NR	AD: 6 HPC: 6 C: 6	AD: 3/3 HPC: 5/1 C: 2/4	AD: 81 HPC: 78 C: 77	NR	Markesbery	NR	NR, Cs had minimal AD pathology or sufficient plaques or tangles to qualify for AD diagnosis	AD: 3.2 HPC: 3.2 C: 1.9	Entorhinal cortex, superior frontal gyrus	IHC	HLA-DR (LN3)	↑ AD > HPC > C
Lue ³⁴	BSHRI	AD: 11 C: 10	AD: 5/6 C: 4/6	AD: 80.8 C: 80.5	APOE: AD: 3/4: 4 4/4: 4 C: 3/4: 3 4/4: 1	Braak, CERAD	AD: IV-VI C: I-III	NR	AD: 2.6 C: 2.3	Hippocampus, cerebellum, superior frontal gyrus	IHC	HLA-DR (LN3)	↑ In the hippocampus, parahippocampal gyrus and superior frontal gyrus ↔ Cerebellum
Matsuo ³⁵	NR	AD: 8 C: 5	NR	NR	NR	Yes	NR	Neurologically normal	All 2-24	Angular, entorhinal, hippocampus, occipitotemporal cortices	IHC	HLA-DR	↑ HLA-DR (more intense staining in more severe AD cases)

Table 1. (Continued)

First Author	Brain bank	N	Sex	Age	AD genetic risk factors	AD histologically confirmed	Braak stage	C history of neurological or psychiatric disease	PMI (h)	Brain region	Technique	Marker	Results (AD vs C unless otherwise specified)
McGeer ³⁸	Pathology Department of the University of British Columbia	NR	NR	NR	NR	NR	NR	NR	NR	Hippocampus	PCR	HLA-DR	↑
McGeer ³⁶	Autopsy Service of the University of British Columbia	AD: 9 C: 7	AD: 5/4 C: NR	AD: 77.2 C: 73.4	NR	NR	NR	No neurological disorders	All > 3 days, most > 10 h	Hippocampus, substantia nigra	IHC	HLA-DR	↑
McGeer ³⁷	NR	AD: 6 C: 5	NR	AD: 78, C: 73	NR	NR	NR	No neurological disease	All within 2–12 h	Hippocampus	IHC	HLA-DR	↑
Minett ⁵⁶	Medical Research Council Cognitive Function and Ageing Study—six centres in UK	AD: 83 C: 130	AD: 30/53 C: 64/66	AD: 89 C: 84	NR	CERAD	NR	NR	NR	Middle frontal gyrus	IHC	HLA-DR	↔
Narayan ³⁹	Neurological Foundation of New Zealand Human Brain Bank (Centre for Brain Research, University of Auckland)	AD: 14 C: 17	AD: 6/8 C: 10/7	AD: 74.1 C: 58.9	NR	Braak or CERAD	NR	Neurologically normal, four have high plaque load	AD: 16.7 C: 16.7	Inferior temporal gyrus	IHC	HLA-DP-DQ-DR	↑ HLA-DP, DQ, DR-positive cells correlate with Iba1-positive cells in C but not AD
Overmyer ⁵²	Kuopio University Hospital	AD: 73 C: 22	AD: 12/61, C: 12/10	AD: 84 C: 78	APOE4 carriers: AD: 31 C: 7	CERAD Patients with possible AD and vascular dementia included	NR	NR—55% demonstrated plaque and tangles, 32% enough for diagnosis of possible AD	All within 48 h	Grey and white matter of frontal, temporal and parietal cortices	IHC	HLA-DR	↔ With dementia diagnosis (trend) ↑ With CERAD in grey matter ↔ White matter (counts and area) ↔ With plaque burden but ↑ correlation with NFT
Parachikova ⁴⁰	Institute for Brain Aging and Dementia Tissue Repository, and the BSHRI	AD: 10 HPC: 10 C: 4	AD: 6/4 HPC: 4/6 C: 3/1	AD: 85.3 HPC: 86.6 C: 76.3	NR	Braak	AD: IV–V HPC: 1–V	NR	AD: 2.6 HPC: 2.8 C: 3.0	Hippocampus and prefrontal cortex (gene chip only)	Gene chip, western, IHC	GeneChip: MHCII western: HLA-DR-DQ-DP IHC: CD4/43 HLA-DR	GeneChip and western: ↑ vs HPC +C (pooled) IHC: ↑ MHCII (not quantified)
Pugliese ⁵³	Neurological Tissue Bank (Serveis Científic-Tècnics), Universitat de Barcelona	AD: 7 C: 3	AD: 2/5 C: 1/2	AD: 84.0 C: 63.3	NR	Braak, Newell Criteria	AD: II: 3, V: 1, VI: 3	NR	AD: 8.8 C: 5.1	Subventricular zone of the lateral ventricle	IHC	HLA-DR	↔ Number of microglia ↑ Activated microglia
Rezaie ⁴¹	MRC London Neurodegenerative Diseases Brain Bank	AD: 10 C: 10	AD: 4/6 C: 7/3	AD: 79.3 C: 70.2	Not reported	CERAD	NR	No history of neurological disease or neuropathology	AD: 20.9, C: 43.2	Frontal blocks included agranular-intermediate frontal cortex (BA 6/8), cingulate cortex (BA 24/32) Occipital blocks included the calcarine sulcus (BA 17) and striate cortex	IHC	HLA-DR-DP-DQ (CR3/43)	↑ In frontal white matter, occipital white matter, plaque associated frontal grey matter, plaque associated occipital grey matter ↔ In MHCII in frontal grey matter, or occipital grey matter
Serrano-Pozo ⁵⁴	Massachusetts ADRC Brain Bank	AD: 40 C: 32	AD: 14/26 C: 13/19	AD: 77.3 C: 81.3	APOE4: AD: 21/40 C: 5/27	NIA—Reagan criteria	NR	No clinical history of neurological disorders and did not meet the pathological criteria for any neurodegenerative disease	AD: 14.1 C: 17.9	Temporal polar association cortex (BA38)	IHC, stereology	HLA-DR-DQ-DP	↔ Total microglia ↓ Iba1+/MHC2- microglia ↑ Iba1-/MHC2+ microglia ↔ Iba1+/MHC2+ microglia ↑ Iba1+/MHC2+ microglia in APOE4 + ↔ with microglia by genotype

Table 1. (Continued)

First Author	Brain bank	N	Sex	Age	AD genetic risk factors	AD histologically confirmed	Braak stage	C history of neurological or psychiatric disease	PMI (h)	Brain region	Technique	Marker	Results (AD vs C unless otherwise specified)
Shepherd ⁴²	Collected brains from a regional brain donor program for neurodegenerative diseases in 1993	AD:10 C: 11	NR	AD:76 C: 71	NR	CERAD	AD: V or VI	No history of neurological disease or neuropathology	AD: 16 C: 21	Cortex and hippocampus (grey and white matter)	IHC	HLA-DR	↑ White and grey matter
Szpak ²³	NR	AD: 18 (7 had Lewy body variant of AD) C: 6	NR	Whole sample: 63–86 years old	NR	CERAD	NR	No neuropathological abnormality	NR	Cortical layers of limbic, cingulate cortex and temporal association cortex	IHC	CR 3/43 clone HLA-DP-DQ-DR	↑
Thal ⁴³	Pathological Institute of the University of Leipzig and University of Frankfurt	159 participants (68 non-demented, 24 and 19 in GDS scores 6 and 7)	NR	Ages 46–93 (most between 71 and 90)	NR	Braak	Whole sample: 0: 23, I: 23, II: 42, III: 36, IV: 16, V: 13, VI: 6	No confounding neurological diagnosis	All within 12–72	Entorhinal cortex, hippocampus (CA1, CA4), occipital region (BA 17), temporal cortex	IHC	HLA-DR	↑
Valente ⁵⁵	Hospital Clinic-University of Barcelona	AD: 7 AD with diabetes: 7 C: 6	AD: 2/5 AD with diabetes: 5/2 C:3/3	AD: 83.9 AD with Diabetes: 73.0 C: 70.0	NR	Braak	AD: VI AD with diabetes: VI	NR	AD: 8.9 AD with diabetes: 11.5 C: 9.6	Hippocampus	IHC	HLA	↔ AD vs C ↑ AD +diabetes vs C
Van Everbroeck ⁴⁴	NR	AD: 10 C: 10	NR	NR	NR	Braak	AD: at least III–IV C: 0, Av or A 1	Some had protein deposition and some had core containing plaques (numbers not given)	NR	Cerebellum, hippocampus (CA1, CA4, subiculum), frontal, temporal and occipital neocortices	IHC	HLA-DR	↑ In grey matter and hippocampus ↔ In white matter and cerebellum
Vehmas ⁴⁵	Johns Hopkins University ADRC and Baltimore Longitudinal Study of Aging NBB	AD: 9 HPC: 15 C: 11	AD: 3/6 HPC: 10/5 C: 11/0	AD: 83.2 HPC: 86.3 C: 81.7	NR	Braak, CERAD	AD: II–V HPC: I–IV C: 0–III	NR, free of plaque	NR	Mixed: middle frontal gyrus, middle and superior temporal gyrus	IHC	HLA-DR	↑ Than C ↔ high pathology C
Verwer ⁵⁰	NR	AD: 14 C: 7	AD: 4/10 C: 2/5	AD: 83.9 C: 79.0	NR	Braak	AD: IV: 3 V: 8 VI: 2 NA: 1 C: 0: 3 I: 1 II: 1 III: 1 NA: 1	No neurological causes of death	AD: 4.6 C: 4.6	Neocortex	IHC	HLA-DR-DQ-DP	↔ <i>P</i> = 0.08
Wilcock ⁴⁶	Irvine ADRC, the Maryland Developmental Disorders Brain Bank and the University of Kentucky Alzheimer's Disease Center.	AD: 9 C: 9	IHC: AD: 6/2 C: 3/6 qPCR +western: AD: 6/4 C: 12/4	IHC: AD: 81.3, C: 81.6 qPCR +western: AD: 80 C: 81.6	NR	NR	NR	NR	IHC: AD: 5.5, C: 3.3 qPCR +western: AD: 6.8 C: 3.3	Frontal cortex	IHC for HLA-DR staining, RT-qPCR and western for expression of M2 and M1 markers	HLA-DR M1 markers: IL-1B, IL-6, IL-12, TNF-α M2a markers: CH13L1, IL1Ra, IL-10, MRc1, M2b markers: CD86, FCGR1B M2c markers: TGFB	↑ HLA-DR in AD vs C ↑ HLA-DR in AD vs. AD+DS (in grey and white matter) —Pattern of increases in both M1 and M2 markers: IL6, IL-12, IL-10, CH13L1, TGFB1 in AD vs C
Wojtera ⁵¹	NR	AD: 4 C: 2	NR	NR	NR	NIA-Reagan criteria	NR	NR	NR	Mixed: cerebellum, cerebral cortex	IHC	HLA-DR	↔, ↔ in HLA-DR/CD68 ratio between AD and C (activation)
Xiang ⁴⁷	ADRC of the Mount Sinai School of Medicine	AD: 26 with clinical dementia rating 0.5–5, 6 rated 5	AD: 7/19 C: 0/5	AD: 88.7 C: 83.2	NR	CERAD	NR	NR	AD: 4.2 C: 4.2	Entorhinal cortex and dorsal hippocampus (CA1 pyramidal cell layer, DG granule cell layer and	IHC	HLA-DR	Entorhinal cortex: ↑ In grey and white matter at CDR 5, in grey matter only at CDR 2 ↔ For CDR

Table 1. (Continued)	First Author	Brain bank	N	Sex	Age	AD genetic risk factors	AD histologically confirmed	Braak stage	C history of neurological or psychiatric disease	PMI (h)	Brain region	Technique	Marker	Results (AD vs C unless otherwise specified)
			(very severe) C: 5								upper molecular layer)			scores of 0.5 to 1 vs 0 Hippocampus: ↑ in all regions for CDR > 2 vs 0; for CA1 pyramidal layer and upper molecular layer for CDR 1 and for the upper molecular layer only for CDR 0.5—HLA-DR score correlates with plaque and tangle scores in various regions

Abbreviations: AD, Alzheimer's disease; ADRC, Alzheimer's Disease Research Center; APOE, apolipoprotein E; Avg, average; BSHRI, Banner Sun Health Research Institute; BA, Brodmann area; C, Control; CA, Cornu ammonis; CD, cluster of differentiation; CDR, Clinical Dementia Rating score; CERAD, Consortium to Establish a Registry for Alzheimer's Disease; CH3L, chitinase 3-like; DG, dentate gyrus; GDS, Global Deterioration Scores; HLA, human leukocyte antigen; HPC, high pathology control; Iba1, ionized calcium-binding adapter molecule 1; IHC, immunohistochemistry; IL, interleukin; MHC, major histocompatibility complex; MRC, mannose receptor; MRC, Medical Research Council; NA, not available; NBB, Netherlands Brain Bank; NFT, neurofibrillary tangle; NIA, National Institute on Aging; NR, not reported; NSAID, nonsteroidal anti-inflammatory drug; TNF- α , tumour necrosis factor- α ; TREM2, triggering receptor expressed on myeloid cells 2. Results are expressed relative to control unless specified otherwise. Where there are both young and older controls, values are reported for the older (age-matched controls).

The remaining positive study quantified plaque-associated microglia.²⁰ In contrast, 6 of the 10 null studies used cell counting, 2 of which used stereology, the gold standard for quantifying cells without bias. Expression-based assays like qPCR, western blot, or intensity of immunohistochemistry staining indicate the amount of Iba1 in a sample. An increase could reflect a change in cell numbers, cell size or function, as Iba1 expression is thought to increase with microglial activation. In contrast, the studies using cell counting for Iba1, which is expressed by all microglia,^{58,77} assess the absolute number of microglia in the samples. The discrepancy between the studies using expression-based assays and those that used cell counting suggests that Iba1 expression, and thus microglial activation, increases in AD without affecting the absolute number of microglia.

Cluster of differentiation 68

CD68 is a common marker for macrophage lineage cells, primarily localized to microglia within the brain parenchyma, and perivascular macrophages in the cerebral blood vessels and, occasionally, parenchyma.⁷⁸ Although there is some CD68 expression on resting microglia,¹⁴ it labels the lysosome and is therefore commonly considered a marker of activated phagocytic microglia.⁶⁰ Twenty-one studies were identified that compared CD68 between AD and control post-mortem brain samples (Table 3). Twenty of these studies used immunohistochemistry to visualize CD68-positive cells and measured cell counts or staining area, whereas one study measured CD68 gene expression by qPCR. Seventeen identified an increase in CD68 expression, staining or positive cell counts in AD relative to control samples in at least one region,^{17,18,20,28,41,56,69,76,78–86} whereas four found no difference between AD and control,^{51,87–89} and four reported no difference in at least one of the brain regions measured.^{20,41,69,80}

CD68-positive cell counts, staining area or gene expression were measured in the hippocampus in eight studies: six identified higher levels in AD^{20,69,76,78,79,86} and two identified no differences.^{69,87} CD68 was higher in AD than control in the frontal cortex in three studies,^{18,41,82} though for white matter only in one of the studies.⁴¹ Elevations in CD68 were also reported in the temporal cortex,^{20,28} the olfactory bulb,⁸⁰ the calcarine cortex,^{69,86} the superior temporal sulcus,⁸³ the orbitofrontal cortex,⁸⁶ parahippocampal gyrus⁷⁶ and temporal association isocortex.⁸⁴ No difference between AD and control was reported in the caudate nucleus,⁸⁸ combined cerebellum and cerebral cortex,⁵¹ medio-dorsal nucleus of the thalamus⁸⁸ and the middle frontal gyrus.⁵⁶ CD68 immunoreactivity appears to increase with age in control subjects, but decreases with age in patients with AD.²⁸ It also increases with APOE ϵ 4 genotype.⁵⁶ Characteristics of AD and control groups could therefore contribute to between-study variability. On balance, CD68 appears to be increased in the brains of patients with AD, though there is some variation between studies and brain regions.

Cluster of differentiation 11b

CD11b forms part of complement receptor 3 that aids in the recognition and phagocytosis of antigens, including amyloid- β .⁹⁰ Like Iba1, CD11b is expressed by both resting and activated microglia, though it too is inducible with activation.¹⁴ Five studies were identified that compared CD11b between AD and control post-mortem human brain samples (Supplementary Materials Table 2). Two papers reported an increase in CD11b,^{15,38} whereas three reported no differences between AD and control.^{74,76,91}

Like Iba1, the results of studies measuring CD11b in AD and control post-mortem human brain samples are heterogeneous. Of two studies in the hippocampus, one identified an increase in CD11b gene expression,³⁸ whereas the other identified no difference in expression between AD and controls.⁷⁶ Similarly, Akiyama *et al.* reported an increase in CD11b-positive cells in the

Table 2. Ionized calcium-binding adapter molecule 1 (Iba1)

First author	Brain bank	N	Sex	Age	AD genetic risk factors	AD histologically confirmed and criteria	Braak stage	C history of neurological or psychiatric disease	PMI (h)	Brain region	Technique	Marker	Direction of results
Bachstetter ⁶⁹	University of Kentucky Alzheimer's Disease Center	AD: 7 C: 9	AD: 4/3 C: 6/3	AD: 77 C: 86	APOE: AD: NA: 2, 4/4: 1, 3/4: 2, C: 3/4: 1	CERAD	AD: ~VI C: ~II	NR	AD: 4.2 C: 2.4	Hippocampus: CA1, CA2/3, CA4, DG, subiculum and adjacent white matter Morphology assessed in the CA1 only	IHC	Iba1	↔ Iba1 staining or cell counts in any hippocampal area ↔ in CA1 Iba1+ microglial morphology
Griciuc ⁷⁰	Massachusetts ADRC	AD: 25 C: 15	AD: 7/18 C: 6/9	AD: 79.2 C: 79.9	APOE carrier: AD: 18 (8 homozygous) C: 5 (0 homozygous) AD: all APOE3/3 C: APOE2/3: 2 APOE3/3: 1 NR: 1	NIA-Reagan Institute Criteria	NR	NR	AD: 17 C: 29	Frontal cortex	IHC (stereology), western	Iba1	↔ Iba1+ microglia (data not shown) ↔ Iba1 protein (nonsignificant increase) ↔
Magistri ⁷¹	BSHRI	AD: 4 C: 4	AD: 1/4 C: 2/2	AD: 83.75 (not exact, one age just listed >90) C: 83.5		NIA-Reagan criteria	AD: V: 1 VI: 3 C: I: 1 II: 3	NR	AD: 2.5 C: 2.5	Hippocampus	RNA seq (gene expression)	Iba1	↔
Nielsen ⁶²	NBB	AD: 4 C: 5	AD: 1/3 C: 1/4	AD: 76.3 C: 77.6	NR	Yes	AD: III: 1, IV: 2, V: 1 C: 0: 1, I: 3 II: 1, AD: IV: 2, V: 4, VI: 3	NR	AD: 6.3 C: 6.1	Entorhinal cortex, hippocampus	IHC	Iba1	↑
Dal Bianco ²⁰	NR	AD: 9 C: 15	AD: 0/9 C: 13/2	AD: 81 C: 70	NR	Braak, CERAD		No neurological disease or brain lesions	NR	Cortical areas of the temporal lobe, including entorhinal cortex, hippocampus and temporal cortex Cingulate cortex, inferior temporal cortex	Immunocytochemistry	AIF-1	↑ AIF-1 near plaque only
Davies ⁷²	New South Wales Brain Bank	AD: 7 C: 5	AD: 3/4 C: 2/3	AD: 83.6 C: 83.0	NR	Braak, CERAD, National Institute on Aging-Alzheimer's Association Guidelines for Neuropathological Assessment of AD Braak, CERAD	AD: V: 3 VI: 4 C: 0: 4 I: 1	No co-existing pathology	AD: 12.3 C: 15.4		IHC	Iba1	↔ Cell density ↑ Microglia with dystrophic morphology, activated morphology (lower ramification)
Satoh ⁶⁸	NR	AD: 7 C: 14	AD: 5/5 C: 6/5	AD: 70 C: 75	NR	Braak, CERAD	AD: VI: 10	4 Died of nonneurological causes, 3 with Parkinson's, 4 ALS No neuropathology	NR	Frontal cortex	IHC, qPCR	Iba1	PCR: ↑ IHC: ↔
Tang ⁶⁵	University of Kentucky Alzheimer's Disease Center Autopsy Program	AD: 10 C: 10	AD: 5/5 C: 6/4	AD: 82.1 C: 82.7	NR	NIA-Reagan criteria	AD: VI: 6 C: I-III, (avg 1.6)		AD: 26.2 C: 6.2	Inferior parietal cortex	Western	Iba1	↑
Ekonomou ⁷³	United Kingdom MRC Cognitive Function and Ageing Study	AD: 13 C: 15	14/28 (both AD and C)	Whole sample: 84.8	NR	Braak	Whole sample: 0-II: 12, III-IV: 11, V-VI: 5	No neurological disease	All within 17.5–25.0	Hippocampus DG	IHC	Iba1	↔ Between AD and C ↑ in Braak stage 3–4 than 0–2 or 5–6
Korvatska ³²	University of Washington Neuropathology Core Brain Bank	AD (normal TREM2): 6 AD (with TREM2 R47H variant): 7 C: 3	NR	Whole sample: 84.9	NR	CERAD	AD: III: 1, V: 4, VI: 1 AD R47H: V: 6, VI: 1 C: 0: 1, I: 1, III: 1	NR, one C CERAD score A and one B	All avg 4.5	Frontal lobe: grey and white matter Hippocampus: CA1, hilus, parahippocampal gyrus and white matter	IHC	Iba1	↓ Staining in hippocampus and frontal lobe white matter ↔ Staining in hilus, CA1, parahippocampal gyrus or frontal lobe grey matter ↔ Counts in frontal lobe grey matter or activated counts in hippocampus white matter in AD, though ↓ in R47H

Table 2. (Continued)

First author	Brain bank	N	Sex	Age	AD genetic risk factors	AD histologically confirmed and criteria	Braak stage	C history of neurological or psychiatric disease	PMI (h)	Brain region	Technique	Marker	Direction of results
Lastres-Becker ⁶⁷	Banco de Tejidos de la Fundación CIEN	AD: 4 C: 4	NR	AD: 73–90 C: 78–90	NR	Braak	AD: II–IV	No neuropsychiatric disease or neuropathology	All within 5 h	Hippocampus	IHC, immunoblot	Iba1	↑
Lee ⁷⁴	OPTIMA and Newcastle Brain Tissue Resource (NBTR)	AD: 12 C: 11	AD: 7/5 C: 6/5	AD: 73.1 C: 81.1	NR	Braak, CERAD	AD: V–VI C: I–II	NR	AD: 61.2 C: 41.5	Prefrontal (BA9) and temporal (BA22) cortices	PCR	Iba1	↔
Lue ⁶¹	BSHRI	AD: 11 C: 11 HPC: 11	AD: 6/5 C: 7/4 HPC: 3/8	AD: 82.4 C: 85.4 HPC: 86.5	APOE 4 Carriers: AD: 5/6 C: 1/10 HPC: 2/9	Braak, CERAD	AD: Avg 5.2 C: Avg 2.8 HPC: Avg 2.9 AD: Avg 4.8 C: Avg 1.4 NR	NR	NR	Middle temporal cortices	Western	Iba1	↑ Than C and HPC
Marlatt ⁷⁵	Netherlands Brain Bank	AD: 8 C: 8	AD:4/4 C: 4/4	AD: 81 C: 80	NR	Braak	AD: Avg 4.8 C: Avg 1.4 NR	NR	All within 5–7 NR	Hippocampus (CA1/2, CA3, DG/SCZ, Hilus) Middle frontal gyrus (BA9)	IHC	Iba1	↔ In cell number or in morphology
Minett ⁵⁶	Medical Research Council Cognitive Function and Ageing Study—six centres in UK	AD: 83 C: 130	AD: 64/53 C: 51/66	AD: 89 C: 84	NR	CERAD	NR	NR	NR	Middle frontal gyrus (BA9)	IHC	Iba1	↓ Iba1 no association with cognition (MMSE), positive association with AD pathology (plaques, tangles) ↑ Iba1 staining density, $P=0.06$ for staining intensity
Rangaraju ⁶³	Emory ADRC Neuropathology Core, Atlanta	AD: 10 C: 10	AD: 6/4 C: 6/4	AD:71.5 C: 71.5	APOE: AD: 8 with APOE4 (3 homozygous) C: 1 APOE4 (0 homozygous) APOE4/4: Braak stage 0–1: 1 2–3: 4 4–5: 5 6: 0	Yes	AD: All VI C: 0	NR	NR	Frontal cortex	IHC	Iba1	↑ Iba1 staining density, $P=0.06$ for staining intensity
Rivera ⁶⁴	KPBBS, University Medical Center	Braak stage 0–1: 12, 2–3: 12 4–5: 12 6: 9	Braak stage: 0–1: 6/6 2–3: 5/7 4–5: 3/9 6: 0/9	Braak stage: 0–1: 74.4 2–3: 81.1 4–5: 82.1 6: 71.8	NR	Braak, NIA-Reagan Criteria	AD: II–VI C: 0–I	NR	All < 16 h	Anterior frontal cortex	PCR	AIF1 IBA1	↑
Sanchez-Mejias ⁷⁶	Tissue bank at Fundación CIEN	Braak stage 0: 8 II: 13, III–IV: 9, V–VI: 17	Braak stage 0: 5/3, II: 7/13, III–IV: 4/5, V–VI: 7/11	Braak stage 0: 19, II: 78, III–IV: 80, V–VI: 79	NR	Braak Braak V–VI clinically classified as AD, Braak II age-matched and used as C	Braak stage 0: 8, II: 13, III–IV: 9, V–VI: 17	NR	Braak stage 0: 8 II: 7, III–IV: 6, V–VI: 8	Hippocampus CA1, CA3, parahippocampal gyrus	IHC, PCR	Iba1	PCR: ↔ Iba1 IHC: ↓ in DG and CA3 ↔ CA1 and parahippocampal gyrus More activated morphology
Serrano-Pozo ⁵⁴	Massachusetts ADRC Brain Bank	AD: 40 C: 32	AD: 14/26 C: 13/19	AD: 81.3 C: 77.6	APOE4: AD: 21/40 C: 5/27	NIA-Reagan criteria	NR	No clinical history of neurological disorders and did not meet the pathological criteria for any neurodegenerative disease	AD: 18.0 C: 14.1	Temporal polar association cortex (BA 38)	IHC, stereology	Iba1	↔ Total microglia ↓ IBA1+/MHC2- microglia ↑ IBA1-/MHC2+ microglia ↔ IBA1+/MHC2+ microglia ↑ IBA1+/MHC2+ microglia in APOE4+ ↔ with microglia by genotype ↑ For CD33 rs3865444 allele A/C genotype ↔ For C/C and A/A genotypes
Walker ⁶⁰	BSHRI	AD: 30 C: 41	Whole sample: AD: 49/48 C: 50/46	Whole sample: AD: 82.2 C: 84.9	APOE4/4 genotypes excluded	NIA-Reagan criteria	NR	NR	Whole sample: AD: 3.6 C: 4	Temporal cortex	Western	Iba1	↑ For CD33 rs3865444 allele A/C genotype ↔ For C/C and A/A genotypes

Abbreviations: AD, Alzheimer's disease; ADRC, Alzheimer's Disease Research Center; ALS, amyotrophic lateral sclerosis; APOE, apolipoprotein E; Avg, average; BSHRI, Banner Sun Health Research Institute; BA, Brodmann area; C, Control; CA, Cornu ammonis; CD, cluster of differentiation; CERAD, Consortium to Establish a Registry for Alzheimer's Disease; CIEN, Centro Investigación Enfermedades Neurológicas; DG, dentate gyrus; HPC, high pathology control; Iba1, ionized calcium-binding adapter molecule 1; IHC, immunohistochemistry; KPBBS, Kathleen Price Bryan Brain Bank; MRC, Medical Research Council; MMSE, mini-mental state examination; NA, not available; NBTR, Newcastle Brain Tissue Resource; NBB, Netherlands Brain Bank; NIA, National Institute on Aging; NR, not reported; OPTIMA, Oxford Project to Investigate Memory and Aging; PMI, post-mortem interval; Seq, sequencing; TREM2, triggering receptor expressed on myeloid cells 2. Results are expressed relative to control unless specified otherwise. Where there are both young and older controls, values are reported for the older (age-matched controls).

Table 3. Cluster of differentiation 68 (CD68)

First author	Brain bank	N	Sex	Age	AD genetic risk factors	AD histologically confirmed and criteria	Braak stage	C history of neurological or psychiatric disease	PMI (h)	Brain region	Technique	Direction of results
Alvarez ⁸⁷	NR	AD: 24 (for cortex and CA1) ADaβ-: 5 Control: 24 (16 for cortex and CA1)	NR	AD: 70–86 ADaβ-: 70-76 Control: 70-86	NR	Braak, CERAD ADaβ- group had no aβ pathology	AD: V–VI	No mental disorder	NR	Cerebellar cortex and hippocampus white matter molecular layer, Purkinje cell layer, granule cell layer, white matter core of the folium, central white matter, layer V of the cortex and CA1	IHC	↔ Between AD, ADaβ- and control
Arnold ⁸⁶	University of Pennsylvania Alzheimer Disease Center Core	AD: 10 C: 14	AD: 5/5 C: 6/8	AD: 81.8 C: 75.3	NR	Khachaturian	NR	No major psychiatric illness, no neuropathologic abnormality except 1 patient with lacunar infarct, one with small temporal contusions	AD: 9.8 C: 11.4	Calcarine cortex (BA17), entorhinal cortex (BA 28) hippocampus CA1, midfrontal cortex (BA9 and 46) orbitofrontal cortex (BA11), subiculum	IHC	↑ In all regions
Arnold ⁷⁹	AD and FTD from: University of Pennsylvania's Alzheimer Disease Center Core	AD: 10 C: 10	AD:5/5 C: 4/6	AD: 81.8 C: 76.1	NR	Yes	NR	No history of major neurological or psychiatric disorder, no neuropathologic abnormalities relevant to mental status	AD: 9.8 C: 11.6	Calcarine cortex, frontal lobe, hippocampus	IHC	↑ (Not compared statistically)
Bachstetter ⁶⁹	University of Kentucky Alzheimer's Disease Center	AD:7 C: 9	AD: 4/3 C: 6/3	AD: 77 C: 86	APOE: AD: NA: 2 4/4: 1 3/4: 2 C: 3/4: 1	CERAD	AD: Median VI C: Median II	NR	AD: 4.2 C: 2.4	Hippocampus: CA1, CA2/3, CA4, DG, subiculum and adjacent white matter. Morphology assessed in the CA1 only	IHC	↑ CD68 staining in subiculum, CA1, DG, and mean of hippocampal regions ↔ in CA2/3, CA4 or white matter ↔ in CD68+ amoeboid in any region except ↑ DG
Dal Bianco ²⁰	NR	AD: 9 C: 15	AD: 0/9 C: 13/2	AD: 81 C: 70	NR	Braak, CERAD	AD: IV: 2, V: 4, VI: 3	No neurological disease or brain lesions	NR	Cortical areas of the temporal lobe, including entorhinal cortex, hippocampus and temporal cortex	Immunocytochemistry	↑ CD68 near plaque only
DeLuca ⁸⁰	Oxford Brain Bank	AD: 4 C: 8	AD: 3/1 C: 5/3	AD: 76.3 C: 63.0	NR	NR	AD: V or VI	No neurological disease	NR	Olfactory bulb/tract	IHC	↑ In parenchyma and meninges
Doorn ⁸¹	NBB or Department of Pathology, Vrije Universiteit, University Medical Center in Amsterdam, The Netherlands	AD: 8 C: 11	AD: 3/5 C: 5/6	AD: 74.5 C: 84	NR	Braak	NFT AD: IV–VI C: 0–III Amyloid: AD: C: 7 B: 1 C: 0:4 A: 3 B: 3 C: 1	Without neurological or psychiatric diseases	AD: 6.2 C: 5.9	Olfactory bulb	IHC	↔ Perivascular ↑ Amoeboid microglia ↔ Ramified

Table 3. (Continued)

First author	Brain bank	N	Sex	Age	AD genetic risk factors	AD histologically confirmed and criteria	Braak stage	C history of neurological or psychiatric disease	PMI (h)	Brain region	Technique	Direction of results
Falke ⁸⁸	University of Pennsylvania ARDC	AD: 12 C: 11	AD: 2/10 C: 7/4	AD: 79.4 C: 77.6	NR	NR	NR	No neuropsychiatric disease—3 control subjects had abnormality at autopsy (haemorrhagic microinfarct, bilateral contusion, adenocarcinoma metastasis). 1 AD subject had microinfarct, all had a β plaques and NFT	AD: 10.9 C: 12.4	Caudate nucleus (6 AD, 7 Control), mediodorsal nucleus of the thalamus (12 AD, 10 Control)	IHC	\leftrightarrow In caudate nucleus \leftrightarrow In MEDIODORSAL nucleus of the thalamus ($P=0.06$)
Fiala ⁷⁸	UCLA ADRC Brain Bank	AD: 8 C: 5	NR	AD: 77.6 C: 74.6	NR	Yes, one patient with vascular dementia not excluded	NR	No neuropathological findings	All 5–6 h	Mix of areas (different cases): hippocampus, frontal lobe (mix of left and right), superior temporal lobe	IHC	\uparrow CD68 staining in AD than C
Hoozemans ¹⁷	NBB	Braak stage 0: 5, I–II: 16, III–IV: 10, V–VI: 9	Braak stage 0: 3/2, I–II: 6/10, III–IV: 0/10, V–VI: 3/6	Braak stage 0: 62, I–II: 83, III–IV: 89, V–VI: 76	NR	Braak	Subjects vary from 0–VI (not divided into C and AD)	NR	Braak stage 0: 8, I–II: 7.5, III–IV: 6.5, V–VI: 5	Temporal cortex	IHC	\uparrow With increasing Braak NFT or plaque stage ($P < 0.05$ for trend), significant for NFT group V–VI vs 0
Hoozemans ²⁸	Netherlands Brain Bank	AD: 19 C: 19	AD: 3/16 C: 8/11	AD: 83.5, C: 76.8	APOE4: AD: 12 C: 8	Braak	AD: avg IV C: avg I	NR	AD: 5.1 C: 8.6	Midtemporal cortex	IHC	$\uparrow \uparrow$ In AD patients < 80 years compared with those > 80 years
Hultman ⁸²	KPBBB, Duke University, North Carolina	AD: 36 C: 22	AD: 13/23 C: 10/12	AD: 76.9 C: 79.1	APOE4 Carriers: AD: 14 C: 0	CERAD, NIA-Reagan criteria	AD: III: 11, IV: 3, V: 13, VI: 9 C: I: 18, II: 3, III: 1	NR, some cases and Cs had mild to severe atherosclerosis	AD: 9.2 C: 7.7	Frontal cortex—perivascular	IHC	\uparrow
Kellner ¹⁸	NR	AD: 48 C: 48	AD: 19/29 C: 24/24	AD: 80.3 C: 77.5	NR	Braak, CERAD	AD: II–VI (38 $>$ 4) C: I–III (45 = 0) NR	NR	NR	Entorhinal, frontal cortex, temporal cortex	IHC	\uparrow
Lue ⁸⁹	BSHRI	AD: 16 C: 21	NR	NR	NR	NR	NR	NR	All avg 3.1	Mixed: corpus callosum, superior and middle frontal gyri of the right hemisphere Middle frontal gyrus (BA9)	IHC	\leftrightarrow
Minett ⁵⁶	Medical Research Council Cognitive Function and Ageing Study—six centres in UK	AD: 83 C: 130	AD: 64/53 C: 51/66	AD: 89 C: 84	NR	CERAD	NR	NR	NR	Middle frontal gyrus (BA9)	IHC	\uparrow Negative correlation with cognition (MMSE), positively with AD pathology (plaques, tangles)

Table 3. (Continued)

First author	Brain bank	N	Sex	Age	AD genetic risk factors	AD histologically confirmed and criteria	Braak stage	C history of neurological or psychiatric disease	PMI (h)	Brain region	Technique	Direction of results
Perez-Nievas ⁸³	Massachusetts General Hospital, Mayo Clinic and University of Pittsburgh ADRC Brain Banks	AD:15 LPC: 15 IPC: 12 HPC: 8	NR	AD: 87.2 LPC: 84.4 IPC: 89.8 HPC: 88.4	NR	Braak, CERAD	NR	NR	NR	Superior temporal sulcus	IHC, stereology	↑ In AD vs LPC, IPC and HPC ↔ in IPC or HPC vs C
Rezaie ⁴¹	MRC London Neurodegenerative Diseases Brain Bank	AD: 10 C: 10	AD: 4/6 C: 7/3	AD: 79.3 C: 70.2	NR	CERAD	NR	No history of neurological disease or neuropathology	AD: 20.9, C: 43.2	Frontal blocks included agranular-intermediate frontal cortex (BA 6/8), cingulate cortex (BA 24/32) Occipital blocks included the calcarine sulcus (BA 17) and striate cortex)	IHC	↑ In frontal white matter, occipital white matter, plaque associated frontal grey matter, plaque associated occipital grey matter ↔ Frontal grey matter, or occipital grey matter
Sanchez-Mejias ⁷⁶	Tissue bank at Fundación CIEN	Braak stage 0: 8, II: 13, III–IV: 9, V–VI: 17	Braak stage 0: 5/3, II: 7/13, III–IV: 4/5, V–VI: 7/11	Braak stage 0: 19, II: 78, III–IV: 80, V–VI: 79	NR	Braak Braak V–VI clinically classified as AD, Braak II age-matched and used as C	Braak stage 0: 8, II: 13, III–IV: 9, V–VI: 17	NR	Braak stage 0: 8, II: 7, III–IV: 6, V–VI: 8	Hippocampus CA1, CA3, parahippocampal gyrus	PCR	↑ With increasing Braak stage Braak stage V–VI had clinical AD and were compared with stage II Cs
Serrano-Pozo ⁸⁴	Massachusetts ADRC Brain Bank	AD: 91 C: 15	AD: 33/58 C: 5/10	AD: 79.0 C: 79.9	APOE E4 carriers AD: 59/32 C: 4/11	NIA-Reagan Criteria	NR	NR, 10/15 had some plaque burden	AD: 13.9 C: 22.3	Temporal association isocortex (BA 38)	IHC, stereology	↑ With increasing disease stage and NFT, no correlation with amyloid burden
Van Everbroeck ⁸⁵	NR	AD: 21 C: 40	NR	NR	NR	Yes	NR	NR, 14 cases/Cs suffered from inflammatory conditions	NR	Mixed: Cerebellum (when available), frontal, occipital and temporal cortices	IHC	↑
Wojtera ⁵¹	NR	AD: 4 C: 2	AD: 4 C: 2	NR	NR	NIA-Reagan Criteria	NR	NR	NR	Mixed: cerebellum, cerebral cortex	IHC	↔ Microglia number ↔ cortex and cerebellum ↔ in HLA-DR/CD68 ratio between AD and control (activation)

Abbreviations: AD, Alzheimer's disease; ADRC, Alzheimer's Disease Research Center; APOE, apolipoprotein E; Avg, average; BSHRI, Banner Sun Health Research Institute; BA, Brodmann area; C, Control; CA, Cornu ammonis; CD, cluster of differentiation; CERAD, Consortium to Establish a Registry for Alzheimer's Disease; CIEN, Centro Investigación Enfermedades Neurológicas; DG, dentate gyrus; FTD, frontotemporal dementia; HLA, human leukocyte antigen; HPC, high pathology control; IHC, immunohistochemistry; IPC, intermediate pathology control; KPBBB, Kathleen Price Bryan Brain Bank; LPC, low pathology control; MRC, Medical Research Council; NA, not available; NBB, Netherlands Brain Bank; NFT, neurofibrillary tangle; NIA, National Institute on Aging; NR, not reported; PMI, post-mortem interval; UCLA, University of California Los Angeles. Results are expressed relative to control unless specified otherwise. Where there are both young and older controls, values are reported for the older (age-matched controls).

temporal lobe of AD cases,¹⁵ whereas Lee *et al.*⁷⁴ found no difference in gene expression relative to controls in the same region. Two studies that compared CD11b gene expression in the prefrontal cortex identified no difference between AD and control.⁹¹ Interestingly, two of the studies that identified increases in CD11b also measured MHCII, and both reported greater increases in MHCII than CD11b in the AD brain.^{15,38} Based on these studies, CD11b does not appear to be consistently increased in the AD brain.

Cluster of differentiation 45

CD45 is a cell surface antigen that is expressed on most hematopoietic lineage cells, where it is involved in the regulation of numerous processes, including cell division and differentiation. CD45 is expressed by both resting and activated microglia, but it does not appear to be inducible with activation in humans.⁹² Six papers were identified that compared CD45 in AD and control human brains (Supplementary Materials Table 3). Two used immunohistochemistry with cell counting, three measured gene expression and one used a combination of western analysis and immunohistochemistry with cell counting. Four studies reported higher CD45 in AD in at least one brain region,^{15,76,93,94} whereas two reported no difference relative to control^{71,95} and one reported no difference in at least one region.⁹⁴

All three of the studies that used immunohistochemistry with cell counting reported an increase in CD45-positive cells in AD patients in the temporal lobe,¹⁵ midfrontal cortex,⁹³ frontal cortex⁹⁴ and hippocampus molecular and pyramidal layers,⁹⁴ respectively. In contrast, the polymorphous layer of the hippocampus did not demonstrate increased CD45-positive cells.⁹⁴ Studies using gene expression were more heterogeneous: Sanchez-Mejias *et al.*⁷⁶ detected higher CD45 gene expression in the hippocampus and parahippocampal gyrus in AD, whereas Magistri *et al.*⁷¹ identified no such increase in the hippocampus relative to controls. No difference in expression was also reported in the frontal cortex.⁹⁵ The evidence suggests that there are more CD45-positive cells in the brains of patients with AD, but that this is not necessarily accompanied by increased CD45 gene expression.

Ferritin

Iron is stored in the brain as heavy (H) or light (L) subunits. Neurons and other cells primarily contain H ferritin, which is the less reactive form, whereas glial cells contain the L subunit, which can be used to generate free radicals as part of the inflammatory response.^{96,97} Although not specific to microglial cells, ferritin immunohistochemistry combined with morphological identification can be used to visualize these cells in the central nervous system, and increases in ferritin levels with inflammation are thought to be caused by increases in microglial number and activation. Seven papers measured ferritin in post-mortem human brain samples from patients with AD and controls (Supplementary Materials Table 4). Five of these papers identified greater levels of ferritin, ferritin-positive microglia or microglia-associated plaques in AD in at least one of the brain regions measured,^{96,98–101} whereas two found no difference in cell counts relative to control,^{33,51} and one identified no change in ferritin-associated plaques in one region and a decrease in another.¹⁰⁰

Three papers that measured ferritin-positive cell counts or protein levels in the hippocampus identified higher levels in AD,^{96,99,101} whereas one that combined counts from the hippocampus with the amygdala, superior frontal gyrus and the superior, middle and inferior temporal gyri found no significant difference relative to control, though a nonsignificant increase was noted.³³ Higher ferritin-positive microglia was also reported in AD relative to control in the amygdala, entorhinal cortex, and frontal, occipital, parietal and temporal neocortices.⁹⁹ No difference

between AD and control was reported in one study that counted microglia over 12 slices of combined cerebellum and cerebral cortex.⁵¹

Cluster of differentiation 33

CD33 is a myeloid cell surface marker that is involved in the regulation of the innate immune system. In microglia, CD33 seems to regulate phagocytosis of amyloid- β , with a reduction in CD33 associated with increased phagocytosis.⁷⁰ Two polymorphisms in the CD33 gene have recently been identified that modulate AD risk,^{102,103} with the protective allele associated with reduced CD33^{12,66,70} and increased phagocytosis.¹² Four papers were identified that used qPCR, western blot or immunohistochemistry to compare levels of CD33 in the brains of patients with AD to controls, three of which reported elevations in AD, whereas one reported no change (Supplementary Materials Table 5).

Higher expression of CD33 genes or of CD33 immunolabelled cells was noted in three studies in the superior and middle temporal gyri, frontal cortex and temporal cortex of patients with AD relative to controls.^{66,70,104} In contrast, Sanchez-Mejias *et al.*⁷⁶ reported no difference in CD33 gene expression between patients with AD and controls in the hippocampus cornu ammonis 1 (CA1), CA3 and parahippocampal gyrus.⁷⁶ CD33 expression was shown to correlate with pan-microglial markers Iba1 or CD11b, supporting its microglial localization.^{66,70,104} As CD33 increases when adjusting for Iba1 in most of studies, it is likely a marker of microglial function or activity, and not of the number of microglia. Though few studies are currently available, the evidence indicates that CD33 is likely increased in the brains of patients with AD.

Triggering receptor expressed on myeloid cells 2

TREM2 is a regulatory protein under the control of γ -secretase that controls toll-like receptor 4 signalling. In microglia, TREM2 seems to be important for microglial activation and phagocytosis of apoptotic neurons.¹⁰⁵ Polymorphisms in the R47H allele of the TREM2 gene have recently been identified as a strong genetic risk factor for the development of AD,¹³ though the actual impact of these polymorphisms on microglial function remain unclear. Four papers were identified that compared levels of TREM2 in the brains of patients with AD and controls (Supplementary Materials Table 6). Two identified an increase of TREM2 in AD,^{61,106} whereas one reported no difference relative to control⁷⁶ and one reported opposing results for qPCR and western analysis.¹⁰⁷

Both papers that examined the temporal cortex identified an increase in TREM2, either for the protein,^{61,106} staining intensity by immunohistochemistry⁶¹ or gene expression.¹⁰⁶ Roussos *et al.*¹⁰⁷ examined the impact of R47H genotype on TREM2 mRNA and protein in the superior temporal gyrus. They also reported higher TREM2 gene expression in AD R47H carriers relative to controls and no difference in TREM2 between AD noncarriers and control. Their protein measurements, however, indicated decreased TREM2 in AD R47H carriers relative to controls with no difference between AD noncarriers and control. The authors speculate that the discrepancy between gene expression and protein quantity may be explained by increased immature TREM2 and increased degradation. In contrast, Sanchez-Mejias *et al.*⁷⁶ identified no difference in TREM2 gene expression relative to control in the hippocampus CA1, CA3 and the parahippocampal gyrus. As with CD33, TREM2 expression is increased even when adjusting for Iba1,¹⁰⁶ and hence it can be considered a marker of microglial function as opposed to number.

Cluster of differentiation 11c

CD11c is a transmembrane protein expressed on the surface of various immune cells, including microglia, macrophages and neutrophils. Four studies were identified that compared CD11c in

AD and control post-mortem brains, all of which used immunohistochemistry and reported higher staining scores or cell counts in AD in at least one brain region^{15,108–110} (Supplementary Materials Table 7).

Three studies examined CD11c in the hippocampus. Two reported higher levels of staining quantified by a semiquantitative scoring system in AD in the CA1 and subiculum;^{108,110} however, Paulus *et al.*¹⁰⁹ quantified CD11c by unbiased cell counts and identified no difference in CA1 or the granular layer of the dentate gyrus relative to control. Elevated CD11c was also reported in the temporal lobe,¹⁵ frontal cortex,¹⁰⁹ entorhinal cortex¹¹⁰ and superior temporal gyrus.¹⁰⁸ No difference between AD and control was detected in the frontal white matter.¹⁰⁹ Based on the limited number of studies, it appears that CD11c is increased in brain of patients with AD relative to controls, but this may be influenced by differences in methodology between the studies.

IL-1 α -expressing microglia

IL-1 α is a proinflammatory cytokine that plays a central role in the induction of an immune response. Although IL-1 α is not a microglia marker, three papers measuring it were included in this review because they used immunohistochemistry and cell counting to examine IL-1 α -positive microglia as an indication of activation (Supplementary Materials Table 8). All three papers are from the same research group and reported higher IL-1 α -positive microglia in AD in at least one of the measured brain areas, but two reported no difference relative to control in one region.

Elevated IL-1 α -positive microglia were reported in the parahippocampal cortex,¹¹¹ the hippocampus¹¹² and the frontal,¹¹² occipital¹¹² and temporal lobes.^{112,113} Within the temporal cortex, layers III–VI were found to be enriched in IL-1 α -expressing microglia in AD, whereas layers I–II were no different from controls.¹¹³ No difference between AD and control brains was detected in the cerebellum.¹¹² Based on the available evidence, IL-1 α -expressing microglia seem to be increased in the brains of patients with AD.

Ricinus communis agglutinin-1

Microglia appear to be the only resident brain cells that express the lectin Ricinus communis agglutinin-1 (RCA-1).¹¹⁴ Two papers were identified that used RCA-1 immunohistochemistry to compare microglia counts between AD and control post-mortem human brain samples (Supplementary Materials Table 9). Both reported significantly more RCA-1-labelled microglia in AD in the inferomedial temporal lobe¹¹⁵ and the association cortex, periallocortex/allocortex and primary cortex, respectively.¹¹⁶

Translocator protein

Translocator protein (TSPO), also known as the peripheral benzodiazepine receptor, is a mitochondrial protein expressed on activated microglia in the brain. It is the ligand for (11)C-PK11195 that is commonly used in positron emission tomography imaging of activated microglia *in vivo*. Two papers returned in the systematic search compared *in situ* PK11195 binding or expression of TSPO in control and AD post-mortem human brain samples (Supplementary Materials Table 10). Kravitz *et al.*¹¹⁷ identified higher [3H]-PK11195 binding in the entorhinal cortex, the subiculum, the striatum and various areas of the hippocampus. In contrast, Marutle *et al.*¹¹⁸ identified no difference in PK11195 binding in the hippocampus between control and AD brains, though they reported higher levels in the frontal cortex. Kravitz *et al.*¹¹⁷ also measured TSPO mRNA and identified higher expression in the hippocampus in AD relative to controls, but no difference in the striatum. Because these two studies have somewhat opposing results for the brain region they have in common, the hippocampus, it is unclear whether TSPO is

increased in post-mortem brain samples from patients with AD. PK11195 has a lower affinity for TSPO and a lower signal-to-noise ratio than newer ligands which may explain some of the discrepancy. A recent review of positron emission tomography imaging of microglia in AD *in vivo* found that of the five papers using the PK11195 ligand, three identified increases in AD relative to controls and two identified no differences, whereas for the second-generation radioligands, 5/6 reported increases in AD in various brain regions.¹¹⁹

Cluster of differentiation 163

CD163 is a scavenger receptor expressed on monocyte/macrophage lineage cells. Two papers identified higher CD163 in AD than control post-mortem human brains (Supplementary Materials Table 11). Dal Bianco *et al.*²⁰ identified a greater number of CD163-positive cells in the cortical areas of the temporal lobe, including the hippocampus and entorhinal and temporal cortices. Like Dal Bianco *et al.*²⁰ Pey *et al.*¹²⁰ also found higher levels of CD163 staining in the hippocampus, as well as the frontal and occipital cortices. Based on this small number of studies, CD163 seems to be upregulated in AD relative to control post-mortem brains.

Microglia identified by morphology

Two studies used nonspecific cell stains to visualize microglia that were identified based on morphology (Supplementary Materials Table 12). Shefer¹²¹ identified both a greater absolute number of glia and a greater number of microglia per volume in the subiculum of the archicortex in the hippocampal fissure. This combination of relative and absolute counts provides evidence that the apparent increase in microglia was not just a function of tissue shrinkage. In contrast, Pelvig *et al.*¹²² also identified microglia based on morphology and quantified them using stereology, but found no difference in the total number of cells in the neocortex.

Other

Seventeen papers returned in the systematic search used markers other than those discussed in previous sections (Supplementary Materials Table 13). Some of these, such as Ox-42 and GLUT-5, are known microglial markers, whereas others were identified by the study authors as being generalizable either to microglia or their activation. Most (16/17) reported changes in markers consistent with increased numbers or activation of microglia in AD.

High-throughput techniques: microarray and proteomics

Microarrays and proteomic studies that specifically discussed microglia or their markers were included in the interest of comprehensiveness, but they are presented separately (Supplementary Materials Table 14) as there is significant risk that those presented here do not represent the balance of the literature. Any microarray would include some microglial markers, such as HLA-DR, but studies that did not identify significant differences between AD and control for these markers may be less likely to have mentioned them by name in the title, abstract or article keywords, and would therefore have been missed by the systematic search. Six studies used high-throughput techniques, either microarray^{11,123–126} or proteomics,¹²⁷ to examine genes, proteins or patterns of gene expression associated with microglia in AD and control in various brain regions, including the entorhinal cortex, hippocampus, postcentral gyrus, superior frontal gyrus, prefrontal cortex, cerebellum, dorsolateral prefrontal cortex, visual cortex and precuneus. All high-throughput studies reported increases in AD relative to control in some of the microglial markers measured.

Nonquantitative comparisons

Fifty-three papers that would otherwise have met the inclusion and exclusion criteria for this review were not included in the full extraction because they were nonquantitative, though results and key methodological details are presented in Supplementary Materials Table 15. Papers were considered nonquantitative if quantitative data for the comparison between AD and control were not presented and/or the methods did not indicate that quantification had occurred. These papers used immunohistochemistry or immunohistochemistry with western blot to measure microglia using various markers, and commented qualitatively on the number of cells, amount of staining/expression or morphology. The most common brain region investigated by the nonquantitative papers was the hippocampus (24/53 studies); however, like in the quantitative papers in the review, many other regions were also investigated, including the entorhinal cortex, temporal cortex, occipital cortex, frontal cortex and anterior cingulate gyrus. Nonquantitative studies used many of the same markers discussed previously, such as HLA, ^{39,128–144} Iba1, ^{77,145} CD68, ^{142,146} ferritin, ^{140,147–150} CD45, ^{142,151–156} RCA-1, ^{138,157} and various other markers or combinations of markers, considered to be related to microglial activation. ^{158–175} All but two of these papers, ^{97,176} measuring CD68 and Iba1, respectively, reported elevations in microglial markers in AD relative to control post-mortem brains. Thus, the null finding rate in the nonquantitative papers is less than a third of the rate of quantitatively assessed markers like MHCII.

DISCUSSION

Most (76/113) of the studies included in this review measured microglia using one of three common markers: MHCII, CD68 and Iba1. Although studies measuring MHCII or CD68 consistently identified increases in AD relative to control in most brain regions studied, 10 of the 20 studies that compared Iba1 identified no difference or a decrease relative to controls. Importantly, 9/10 studies noting an increase in Iba1 in AD relative to controls used expression-based quantification methods (qPCR, western, fluorescence intensity) that indicate only the amount of Iba1 in the sample, whereas most that identified no difference used cell counting, including two studies that used stereological quantification. Iba1 is a pan-microglial marker whose expression increases with microglial activation, ^{58,77} and hence these results indicate that there are increases in the expression of Iba1, but not the absolute number of microglia in AD. This, along with the increases in MHCII and CD68, which are both markers of activated microglia, suggest that the apparent increases in microglial markers in AD are attributable to increases in activation rather than the absolute number of microglia. This is supported by findings with CD11b that, like Iba1, labels resting and activated microglia, that also had mixed results, and by studies using other activation markers, such as ferritin, IL-1 α and CD33, that were consistently increased in AD. However, there is still controversy surrounding what markers are indicative of activation, as well as the type of activation they are associated with, ⁶⁰ and hence more research on the physiological significance of increases in these markers is needed.

The nine studies identified by this review that compared microglial markers between AD and a high pathology control group, the cognitively intact subjects with AD neuropathology, may shed light on whether increased microglial activation is a cause or consequence of the disease. Five of these studies reported higher levels of microglia markers in AD, three using HLA-DR, ^{7,33,40} one using CD68, ⁸³ and one using both TREM2 and Iba1. ⁶¹ In contrast, three studies, using CD36, HLA-DR or proteomics, reported no difference between AD and the high-pathology control group despite increases relative to the regular control group, ^{45,127,177} whereas one reported more HLA-DR-

positive microglia in the high pathology control group than in AD. ²⁴ Though there appears to be increased microglia markers in AD relative to high pathology control subjects, more studies using this reference group are needed to help elucidate the role of microglia in AD pathogenesis. Polymorphisms in genes encoding microglial markers HLA-DR, CD33 and TREM2 have been implicated as risk factors for late-onset AD, ^{9,10,103} supporting the notion that increased microglial activation is a contributor to AD development, and not merely a response to established AD pathology.

Some of the heterogeneity identified between studies may be attributed to differences in the brain regions examined. More than half the studies that used tissue from the cerebellum identified no difference between control and AD across a range of microglial markers. ^{25,34,44,112,178} This lack of reactivity in the cerebellum has been remarked upon previously, leading the cerebellum to be proposed as a reference region for positron emission tomography imaging of TSPO. ¹⁷⁹ Similarly, half the studies examining microglia in the white matter showed either no difference between AD and control ^{25,44,52,69,94,109,139} or higher levels in control. ³² Several studies report higher levels of HLA-DR-expressing microglia in the white matter than the gray matter of nondemented cases, suggesting that microglia in this tissue may be constitutively activated. ^{49,52,143} The lack of consistent differences in the white matter between patients with AD and controls in the studies in our review could therefore indicate that AD pathology does not stimulate further increases in microglial markers on top of their already activated state, whereas elevations in AD can be observed in the grey matter that does not demonstrate this constitutive elevation.

Heterogeneity can also exist within a brain region. For example, Bachstetter *et al.* ⁶⁹ counted CD68-positive microglia in the hippocampus CA1, CA2/3, CA4 and the dentate gyrus, and noted increases in the CA1 and dentate gyrus relative to controls, but no difference in CA2/3 or CA4. Similarly, Masliah *et al.* ⁹⁴ quantified microglia in different layers of the hippocampus and reported higher levels relative to control in the molecular and pyramidal layer, but lower levels in the stratum polymorphous. Thus, studies that examined staining in the whole hippocampus could dilute potential differences between AD and control by examining multiple regions simultaneously, and this may help to explain why nearly a third of studies using hippocampal tissue reported no difference between AD and control. The potential for heterogeneity of results between tissues is particularly important for studies that performed their analyses in a mix of different tissues, rather than examining each region separately. ^{30,89,98,122,180}

Only one of the identified studies reported on the use of anti-inflammatory drugs in their subjects. ¹⁶ Though this study did not identify significant differences in microglia counts in users and nonusers, nonsteroidal anti-inflammatory drug treatment reduces activated glial cells in a mouse model of AD, ¹⁸¹ and the use of nonsteroidal anti-inflammatory drugs is associated with less microglial activation in the brains of elderly patients post-mortem, ¹⁸² suggesting that unreported nonsteroidal anti-inflammatory drug use could be a potential source of confounding. It is therefore feasible that unreported differences in medication use between AD and controls within and between studies contributed to the variability in the results.

The genotypes of the subjects could also offer a source of heterogeneity. Microglial activation appears to be affected by APOE genotype, with carriers of the $\epsilon 4$ risk allele exhibiting greater activation in some ^{48,56} but not all studies. ⁸² Only 29 of the studies included in this review reported the APOE genotype of their subjects. Among those that reported genotype, there was variability, with some studies excluding those with the $\epsilon 4$ allele, ^{68,172} and others including a majority of AD subjects carrying the risk allele. ^{30,62,79}

Differences in control group characteristics could also influence the results. Over half the included studies did not report screening controls for neurological or neuropathological abnormalities. In addition, only 11 studies reported excluding subjects with a history of psychiatric disorders. As various neurological and psychiatric diseases have been associated with neuroinflammation in some post-mortem studies,^{183–185} this lack of screening could contribute heterogeneity to the results.

The severity of AD pathology in AD cases and controls also varies between studies. Braak stage was the most widely reported pathological score, and is presented in the tables where available. Most studies used AD cases with a Braak score of V–VI, but some used AD cases with scores as low as I and II.^{18,45,53,64,67,123} Similarly for controls, Braak stages less than III were most commonly used, but some studies included controls with Braak stages IV or higher.^{123,127} This variation may be important, as some studies identified associations between Braak stage or neurofibrillary tangles and microglia markers, including HLA-DR, Iba1 and CD68.^{17,48,52,64,84,186} Few studies reported on the amount of plaque in the brains included in their cohorts, but differences in plaque load between studies could also influence the results, as microglial markers were found to correlate with plaque loads in several studies.^{47,48,52,56,186} These correlations could be functionally important to the progression of AD, as post-mortem analyses from brains of AD patients administered amyloid- β vaccines demonstrate positive correlations between Iba1 and CD68 and markers of neuronal loss and degeneration that are attenuated following immunization and amyloid- β clearance,¹⁸⁷ when the number of activated CD68-positive but not total Iba1-positive microglia are reduced.¹⁸⁶

Very few studies reported using stereology to complete their cell counts,^{54,70,83,84,122} the gold standard for quantifying cells in a tissue without bias. In our review, Serrano-Pozo *et al.*⁵⁴ reported that the absolute number of microglia was the same in AD and control samples when stereology was used, but higher in AD when nonstereology-based counting was applied because of an apparent concentration of cells caused by cortical atrophy. The use of nonstereological counting could therefore have introduced some bias into the results. Interestingly, the three stereology studies that counted all microglia, identified by either Iba1 or morphology, reported no differences in total microglial counts in AD relative to control samples,^{54,122,70} whereas increases were noted in the number of cells positive for activation-associated markers MHCII, CD68 and CD33.^{54,70,83,84} This supports the suggestion that microglial activation is increased in AD without an absolute increase in the number of microglia.

Technical differences could also contribute to variation between studies. For immunohistochemistry, the most commonly used technique in this review, the method and duration of tissue fixation,^{188,189} the thickness of the brain sections,¹⁸⁹ the use of antigen retrieval,¹⁹⁰ the antibody selected^{77,188} and variation in many steps in the immunohistochemistry protocol itself can all influence the intensity of staining that could affect the ability of studies to pick up differences between AD and control subjects. Differences in the methods used for measuring gene expression or for quantifying protein by western blot or enzyme-linked immunosorbent assay could similarly contribute to variability in the results. This technical information, with the exception of technique used and post-mortem interval, was not extracted in the tables because the wide variability in methods and reporting made it unfeasible, but it is an important potential contributor to study heterogeneity.

Limitations

Though efforts were made to identify every published paper that compared microglia markers between AD and control post-mortem

human brain samples, it is possible that some papers have been missed. The large number of papers returned in the initial search (> 20 000) may have increased the risk of misidentifying a paper. In addition, studies that measured a microglial marker, but did not mention it in the title, abstract or keywords would have been missed by the systematic search. Thus, studies that did not examine microglia as a primary objective are more likely to have been omitted which may introduce some bias into the results of this review.

Microglia interact with astrocytes to initiate and drive inflammation within the brain. Like with microglia, markers of astrocytes are reported to be elevated in patients with AD relative to controls,^{191,192} where they seem to colocalize with amyloid- β plaques.^{193,194} Although our review focused only on microglial markers, it had initially set out to include other inflammatory markers as well, including, astrocytes, cytokines, complement, lipid mediators and other immune cells. Our title and abstract screening uncovered over 700 papers, nearly 297 for astrocytes alone, an unfeasibly large number of papers that led us to focus on microglia for this review. Not including astrocytes and other inflammatory mediators is a limitation of our review because it results in an incomplete picture of neuroinflammation in AD.

Another important limitation of this review is that although it describes the results of the included studies in the context of microglia, it should be noted that nearly all microglia markers are also expressed by other myeloid cells such as infiltrating or perivascular macrophages, neutrophils or T cells (see Supplementary Table 1 for a summary of marker expression in other cell types). Although there are generally few of these cells in the brain parenchyma under normal conditions, there may be increased accumulation and infiltration in AD that could lead to an overestimation of the quantity of microglial markers in diseased relative to control brains.

CONCLUSION

Microglia markers are increased in various brain regions in patients with AD relative to controls. The balance of the evidence suggests that this increase is more attributable to increased activation than to an increase in absolute cell number, but more research measuring microglial proliferation in post-mortem human brain samples would be useful for clarifying this point.

CONFLICT OF INTEREST

RPB has received research grants from Bunge Ltd, Arctic Nutrition, the Dairy Farmers of Canada and Nestle Inc, as well as travel support from Mead Johnson and mass spectrometry equipment and support from Sciex. RPB is on the executive of the International Society for the Study of Fatty acids and Lipids and held a meeting on behalf of Fatty Acids and Cell Signaling, both of which rely on corporate sponsorship. RPB has given expert testimony in relation to supplements and the brain. RPB also provides complimentary fatty acid analysis for farmers, food producers and others involved in the food industry, some of whom provide free food samples. The funding body had no role in any step of this paper, or in the decision to publish results. The remaining authors declare no conflicts of interest.

ACKNOWLEDGMENTS

Dr JoAnne McLaurin and Dr Anthony Domenichiello provided insight and resources respectively that aided in the preparation of this review. Dr John Sievenpiper and the staff at the University of Toronto Gerstein Science Information Centre provided assistance in developing the systematic search protocol. This work was supported by a grant from the Canadian Institutes of Health Research (CIHR) to RPB, a CIHR Vanier Canada Graduate Scholarship to KEH, a CIHR Canadian Graduate Scholarship to VG and a National Sciences and Engineering Research Council of Canada Undergraduate Student Research Award to DM. RPB is a Canada Research Chair in Brain Lipid Metabolism.

AUTHOR CONTRIBUTIONS

KEH designed the search with the assistance of MT, reviewed the articles returned by the search for eligibility with the assistance of DM, reviewed all data extraction and wrote the paper. DM, MT and VG assisted with the full-text assessments, data extraction and provided feedback on the paper. RPB oversaw the project, provided feedback on all steps of the search and data extraction and reviewed the manuscript.

REFERENCES

- 1 Ma K, Mount HT, McLaurin J. Region-specific distribution of beta-amyloid peptide and cytokine expression in TgCRND8 mice. *Neurosci Lett* 2011; **492**: 5–10.
- 2 Souza LC, Filho CB, Goes AT, Fabbro LD, de Gomes MG, Savegnago L et al. Neuroprotective effect of physical exercise in a mouse model of Alzheimer's disease induced by beta-amyloid(1)(-)(4)(0) peptide. *Neurotox Res* 2013; **24**: 148–163.
- 3 Hopperton KE, Trepanier MO, Giuliano V, Bazinet RP. Brain omega-3 polyunsaturated fatty acids modulate microglia cell number and morphology in response to intracerebroventricular amyloid-beta 1-40 in mice. *J Neuroinflammation* 2016; **13**: 257.
- 4 Vanzani MC, Iacono RF, Caccuri RL, Berria MI. Immunohistochemical and morphometric features of astrocyte reactivity vs. plaque location in Alzheimer's disease. *Medicina* 2005; **65**: 213–218.
- 5 Flanders KC, Lippa CF, Smith TW, Pollen DA, Sporn MB. Altered expression of transforming growth factor-beta in Alzheimer's disease. *Neurology* 1995; **45**: 1561–1569.
- 6 Griffin WS, Stanley LC, Ling C, White L, MacLeod V, Perrot LJ et al. Brain interleukin 1 and S-100 immunoreactivity are elevated in Down syndrome and Alzheimer disease. *Proc Natl Acad Sci USA* 1989; **86**: 7611–7615.
- 7 Lue LF, Brachova L, Civin WH, Rogers J. Inflammation, A beta deposition, and neurofibrillary tangle formation as correlates of Alzheimer's disease neurodegeneration. *J Neuropathol Exp Neurol* 1996; **55**: 1083–1088.
- 8 Griffin WS. Inflammation and neurodegenerative diseases. *Am J Clin Nutr* 2006; **83**: 470S–474S.
- 9 Lambert JC, Ibrahim-Verbaas CA, Harold D, Naj AC, Sims R, Bellenguez C et al. Meta-analysis of 74,046 individuals identifies 11 new susceptibility loci for Alzheimer's disease. *Nat Genet* 2013; **45**: 1452–1458.
- 10 Sims R, van der Lee SJ, Naj AC, Bellenguez C, Badarinarayan N, Jakobsdottir J et al. Rare coding variants in PLCG2, ABI3, and TREM2 implicate microglial-mediated innate immunity in Alzheimer's disease. *Nat Genet* 2017; **49**: 1373–1384.
- 11 Zhang B, Gaiteri C, Bodea LG, Wang Z, McElwee J, Podtelezchnikov AA et al. Integrated systems approach identifies genetic nodes and networks in late-onset Alzheimer's disease. *Cell* 2013; **153**: 707–720.
- 12 Bradshaw EM, Chibnik LB, Keenan BT, Ottoboni L, Raj T, Tang A et al. CD33 Alzheimer's disease locus: altered monocyte function and amyloid biology. *Nat Neurosci* 2013; **16**: 848–850.
- 13 Guerreiro R, Wojtas A, Bras J, Carrasquillo M, Rogaeva E, Majounie E et al. TREM2 variants in Alzheimer's disease. *N Engl J Med* 2013; **368**: 117–127.
- 14 Lee YB, Nagai A, Kim SU. Cytokines, chemokines, and cytokine receptors in human microglia. *J Neurosci Res* 2002; **69**: 94–103.
- 15 Akiyama H, McGeer PL. Brain microglia constitutively express beta-2 integrins. *J Neuroimmunol* 1990; **30**: 81–93.
- 16 Halliday GM, Shepherd CE, McCann H, Reid WG, Grayson DA, Broe GA et al. Effect of anti-inflammatory medications on neuropathological findings in Alzheimer disease. *Arch Neurol* 2000; **57**: 831–836.
- 17 Hoozemans JJ, van Haastert ES, Veerhuis R, Arendt T, Scheper W, Eikelenboom P et al. Maximal COX-2 and ppRb expression in neurons occurs during early Braak stages prior to the maximal activation of astrocytes and microglia in Alzheimer's disease. *J Neuroinflamm* 2005; **2**: 27.
- 18 Kellner A, Matschke J, Bernreuther C, Moch H, Ferrer I, Glatzel M. Autoantibodies against beta-amyloid are common in Alzheimer's disease and help control plaque burden. *Ann Neurol* 2009; **65**: 24–31.
- 19 Carpenter AF, Carpenter PW, Markesbery WR. Morphometric analysis of microglia in Alzheimer's disease. *J Neuropathol Exp Neurol* 1993; **52**: 601–608.
- 20 Dal Bianco A, Bradl M, Frischer J, Kutzelnigg A, Jellinger K, Lassmann H. Multiple sclerosis and Alzheimer's disease. *Ann Neurol* 2008; **63**: 174–183.
- 21 Desai BS, Schneider JA, Li JL, Carvey PM, Hendey B. Evidence of angiogenic vessels in Alzheimer's disease. *J Neural Transm* 2009; **116**: 587–597.
- 22 Dhawan G, Floden AM, Combs CK. Amyloid-beta oligomers stimulate microglia through a tyrosine kinase dependent mechanism. *Neurobiol Aging* 2012; **33**: 2247–2261.
- 23 Szpak GM, Lechowicz W, Lewandowska E, Bertrand E, Wierzbica-Bobrowicz T, Gwiazda E et al. Neurons and microglia in central nervous system immune response to degenerative processes. Part 1: Alzheimer's disease and Lewy body variant of Alzheimer's disease. Quantitative study. *Folia Neuropathol* 2001; **39**: 181–192.
- 24 Flanary BE, Sammons NW, Nguyen C, Walker D, Streit WJ. Evidence that aging and amyloid promote microglial cell senescence. *Rejuvenation Res* 2007; **10**: 61–74.
- 25 Giulian D, Haverkamp LJ, Li J, Karshin WL, Yu J, Tom D et al. Senile plaques stimulate microglia to release a neurotoxin found in Alzheimer brain. *Neurochem Int* 1995; **27**: 119–137.
- 26 Gouw AA, Seewann A, Vrenken H, van der Flier WM, Rozemuller JM, Barkhof F et al. Heterogeneity of white matter hyperintensities in Alzheimer's disease: post-mortem quantitative MRI and neuropathology. *Brain* 2008; **131**(Pt 12): 3286–3298.
- 27 Hensley K, Hall N, Subramaniam R, Cole P, Harris M, Aksenov M et al. Brain regional correspondence between Alzheimer's disease histopathology and biomarkers of protein oxidation. *J Neurochem* 1995; **65**: 2146–2156.
- 28 Hoozemans JJ, Rozemuller AJ, van Haastert ES, Eikelenboom P, van Gool WA. Neuroinflammation in Alzheimer's disease wanes with age. *J Neuroinflamm* 2011; **8**: 171.
- 29 Imamura K, Sawada M, Ozaki N, Naito H, Iwata N, Ishihara R et al. Activation mechanism of brain microglia in patients with diffuse neurofibrillary tangles with calcification: a comparison with Alzheimer disease. *Alzheimer Dis Assoc Disord* 2001; **15**: 45–50.
- 30 Itagaki S, McGeer PL, Akiyama H. Presence of T-cytotoxic suppressor and leucocyte common antigen positive cells in Alzheimer's disease brain tissue. *Neurosci Lett* 1988; **91**: 259–264.
- 31 Jantarotnai N, Schwab C, Ryu JK, McGeer PL, McLarnon JG. Converging perturbed microvasculature and microglial clusters characterize Alzheimer disease brain. *Curr Alzheimer Res* 2010; **7**: 625–636.
- 32 Korvatska O, Leverenz JB, Jayadev S, McMillan P, Kurtz I, Guo X et al. R47H variant of TREM2 associated with Alzheimer disease in a large late-onset family: clinical, genetic, and neuropathological study. *JAMA Neurol* 2015; **72**: 920–927.
- 33 Lopes KO, Sparks DL, Streit WJ. Microglial dystrophy in the aged and Alzheimer's disease brain is associated with ferritin immunoreactivity. *Glia* 2008; **56**: 1048–1060.
- 34 Lue LF, Walker DG, Brachova L, Beach TG, Rogers J, Schmidt AM et al. Involvement of microglial receptor for advanced glycation endproducts (RAGE) in Alzheimer's disease: identification of a cellular activation mechanism. *Exp Neurol* 2001; **171**(1): 29–45.
- 35 Matsuo A, Walker DG, Terai K, McGeer PL. Expression of CD43 in human microglia and its downregulation in Alzheimer's disease. *J Neuroimmunol* 1996; **71**: 81–86.
- 36 McGeer PL, Itagaki S, Boyes BE, McGeer EG. Reactive microglia are positive for HLA-DR in the substantia nigra of Parkinson's and Alzheimer's disease brains. *Neurology* 1988; **38**: 1285–1291.
- 37 McGeer PL, Itagaki S, Tago H, McGeer EG. Reactive microglia in patients with senile dementia of the Alzheimer type are positive for the histocompatibility glycoprotein HLA-DR. *Neurosci Lett* 1987; **79**: 195–200.
- 38 McGeer PL, McGeer EG, Yasojima K. Alzheimer disease and neuroinflammation. *J Neural Transm Suppl* 2000; **59**: 53–57.
- 39 Narayan P, Lill C, Faull R, Curtis MA, Dragunow M. Increased acetyl and total histone levels in post-mortem Alzheimer's disease brain. *Neurobiol Dis* 2015; **74**: 281–294.
- 40 Parachikova A, Agadjanyan MG, Cribbs DH, Blurton-Jones M, Perreau V, Rogers J et al. Inflammatory changes parallel the early stages of Alzheimer disease. *Neurobiol Aging* 2007; **28**: 1821–1833.
- 41 Rezaie P, Pontikis CC, Hudson L, Cairns NJ, Lantos PL. Expression of cellular prion protein in the frontal and occipital lobe in Alzheimer's disease, diffuse Lewy body disease, and in normal brain: an immunohistochemical study. *J Histochem Cytochem* 2005; **53**: 929–940.
- 42 Shepherd CE, Thiel E, McCann H, Harding AJ, Halliday GM. Cortical inflammation in Alzheimer disease but not dementia with Lewy bodies. *Arch Neurol* 2000; **57**: 817–822.
- 43 Thal DR, Arendt T, Waldmann G, Holzer M, Zedlick D, Rub U et al. Progression of neurofibrillary changes and PHF-tau in end-stage Alzheimer's disease is different from plaque and cortical microglial pathology. *Neurobiol Aging* 1998; **19**: 517–525.
- 44 Van Everbroeck B, Dobbeleir I, De Waele M, De Leenheer E, Lubke U, Martin JJ et al. Extracellular protein deposition correlates with glial activation and oxidative stress in Creutzfeldt-Jakob and Alzheimer's disease. *Acta Neuropathol* 2004; **108**: 194–200.
- 45 Vehmas AK, Kawas CH, Stewart WF, Troncoso JC. Immune reactive cells in senile plaques and cognitive decline in Alzheimer's disease. *Neurobiol Aging* 2003; **24**: 321–331.
- 46 Wilcock DM, Hurban J, Helman AM, Sudduth TL, McCarty KL, Beckett TL et al. Down syndrome individuals with Alzheimer's disease have a distinct

- neuroinflammatory phenotype compared to sporadic Alzheimer's disease. *Neurobiol Aging* 2015; **36**: 2468–2474.
- 47 Xiang Z, Haroutunian V, Ho L, Purohit D, Pasinetti GM. Microglia activation in the brain as inflammatory biomarker of Alzheimer's disease neuropathology and clinical dementia. *Dis Markers* 2006; **22**: 95–102.
- 48 Egensperger R, Kosel S, von Eitzen U, Graeber MB. Microglial activation in Alzheimer disease: association with APOE genotype. *Brain Pathol* 1998; **8**: 439–447.
- 49 Mattiace LA, Davies P, Dickson DW. Detection of HLA-DR on microglia in the human brain is a function of both clinical and technical factors. *Am J Pathol* 1990; **136**: 1101–1114.
- 50 Verwer RW, Sluiter AA, Balesar RA, Baayen JC, Noske DP, Dirven CM *et al*. Mature astrocytes in the adult human neocortex express the early neuronal marker doublecortin. *Brain* 2007; **130**(Pt 12): 3321–3335.
- 51 Wojtera M, Sobow T, Kloszewska I, Liberski PP, Brown DR, Sikorska B. Expression of immunohistochemical markers on microglia in Creutzfeldt-Jakob disease and Alzheimer's disease: morphometric study and review of the literature. *Folia Neuropathol* 2012; **50**: 74–84.
- 52 Overmyer M, Helisalmi S, Soininen H, Laakso M, Riekinen P Sr., Alafuzoff I. Reactive microglia in aging and dementia: an immunohistochemical study of postmortem human brain tissue. *Acta Neuropathol* 1999; **97**: 383–392.
- 53 Pugliese M, Rodriguez MJ, Gimeno-Bayon J, Pujadas L, Billett EE, Wells C *et al*. Alzheimer's disease modifies progenitor cell expression of monoamine oxidase B in the subventricular zone. *J Neurosci Res* 2010; **88**: 2588–2597.
- 54 Serrano-Pozo A, Gomez-Isla T, Growdon JH, Frosch MP, Hyman BT. A phenotypic change but not proliferation underlies glial responses in Alzheimer disease. *Am J Pathol* 2013; **182**: 2332–2344.
- 55 Valente T, Gella A, Sole M, Durany N, Unzeta M. Immunohistochemical study of semicarbazide-sensitive amine oxidase/vascular adhesion protein-1 in the hippocampal vasculature: pathological synergy of Alzheimer's disease and diabetes mellitus. *J Neurosci Res* 2012; **90**(10): 1989–1996.
- 56 Minnett T, Classey J, Matthews FE, Fahrenhold M, Taga M, Brayne C *et al*. Microglial immunophenotype in dementia with Alzheimer's pathology. *J Neuroinflamm* 2016; **13**(1): 135.
- 57 Imai Y, Ibata I, Ito D, Ohsawa K, Kohsaka S. A novel gene iba1 in the major histocompatibility complex class III region encoding an EF hand protein expressed in a monocytic lineage. *Biochem Biophys Res Commun* 1996; **224**: 855–862.
- 58 Ito D, Imai Y, Ohsawa K, Nakajima K, Fukuchi Y, Kohsaka S. Microglia-specific localisation of a novel calcium binding protein, Iba1. *Brain Res Mol Brain Res* 1998; **57**: 1–9.
- 59 Ohsawa K, Imai Y, Kanazawa H, Sasaki Y, Kohsaka S. Involvement of Iba1 in membrane ruffling and phagocytosis of macrophages/microglia. *J Cell Sci* 2000; **113**(Pt 17): 3073–3084.
- 60 Walker DG, Lue LF. Immune phenotypes of microglia in human neurodegenerative disease: challenges to detecting microglial polarization in human brains. *Alzheimers Res Ther* 2015; **7**: 56.
- 61 Lue LF, Schmitz CT, Serrano G, Sue LI, Beach TG, Walker DG. TREM2 protein expression changes correlate with Alzheimer's disease neurodegenerative pathologies in post-mortem temporal cortices. *Brain Pathol* 2015; **25**: 469–480.
- 62 Nielsen HM, Ek D, Avdic U, Orbjorn C, Hansson ONetherlands Brain Banket *et al*. NG2 cells, a new trail for Alzheimer's disease mechanisms? *Acta Neuropathol Commun* 2013; **1**: 7.
- 63 Rangaraju S, Gearing M, Jin LW, Levey A. Potassium channel Kv1.3 is highly expressed by microglia in human Alzheimer's disease. *J Alzheimers Dis* 2015; **44**: 797–808.
- 64 Rivera EJ, Goldin A, Fulmer N, Tavares R, Wands JR, de la Monte SM. Insulin and insulin-like growth factor expression and function deteriorate with progression of Alzheimer's disease: link to brain reductions in acetylcholine. *J Alzheimers Dis* 2005; **8**: 247–268.
- 65 Tang SC, Lathia JD, Selvaraj PK, Jo DG, Mughal MR, Cheng A *et al*. Toll-like receptor-4 mediates neuronal apoptosis induced by amyloid beta-peptide and the membrane lipid peroxidation product 4-hydroxynonenal. *Exp Neurol* 2008; **213**: 114–121.
- 66 Walker DG, Whetzel AM, Serrano G, Sue LI, Beach TG, Lue LF. Association of CD33 polymorphism rs3865444 with Alzheimer's disease pathology and CD33 expression in human cerebral cortex. *Neurobiol Aging* 2015; **36**: 571–582.
- 67 Lastres-Becker I, Innamorato NG, Jaworski T, Rabano A, Kugler S, Van Leuven F *et al*. Fractalkine activates NRF2/NFE2L2 and heme oxygenase 1 to restrain tauopathy-induced microgliosis. *Brain* 2014; **137**(Pt 1): 78–91.
- 68 Satoh J, Kino Y, Asahina N, Takitani M, Miyoshi J, Ishida T *et al*. TMEM119 marks a subset of microglia in the human brain. *Neuropathology* 2016; **36**: 39–49.
- 69 Bachstetter AD, Van Eldik LJ, Schmitt FA, Neltner JH, Ighodaro ET, Webster SJ *et al*. Disease-related microglia heterogeneity in the hippocampus of Alzheimer's disease, dementia with Lewy bodies, and hippocampal sclerosis of aging. *Acta Neuropathol Commun* 2015; **3**: 32.
- 70 Griciuc A, Serrano-Pozo A, Parrado AR, Lesinski AN, Asselin CN, Mullin K *et al*. Alzheimer's disease risk gene CD33 inhibits microglial uptake of amyloid beta. *Neuron* 2013; **78**: 631–643.
- 71 Magistri M, Velmeshev D, Makhmutova M, Faghihi MA. Transcriptomics profiling of Alzheimer's disease reveal neurovascular defects, altered amyloid-beta homeostasis, and deregulated expression of long noncoding RNAs. *J Alzheimers Dis* 2015; **48**: 647–665.
- 72 Davies DS, Ma J, Jegathees T, Goldsby C. Microglia show altered morphology and reduced arborization in human brain during aging and Alzheimer's disease. *Brain Pathol* 2016; **27**: 795–808.
- 73 Ekonomou A, Savva GM, Brayne C, Forster G, Francis PT, Johnson M *et al*. Stage-specific changes in neurogenic and glial markers in Alzheimer's disease. *Biol Psychiatry* 2015; **77**: 711–719.
- 74 Lee C, Low CY, Francis PT, Attems J, Wong PT, Lai MK *et al*. An isoform-specific role of FynT tyrosine kinase in Alzheimer's disease. *J Neurochem* 2016; **136**: 637–650.
- 75 Marlatt MW, Bauer J, Aronica E, van Haastert ES, Hoozemans JJ, Joels M *et al*. Proliferation in the Alzheimer hippocampus is due to microglia, not astroglia, and occurs at sites of amyloid deposition. *Neural Plast* 2014; **2014**: 693851.
- 76 Sanchez-Mejias E, Navarro V, Jimenez S, Sanchez-Mico M, Sanchez-Varo R, Nunez-Diaz C *et al*. Soluble phospho-tau from Alzheimer's disease hippocampus drives microglial degeneration. *Acta Neuropathol* 2016; **132**: 897–916.
- 77 Streit WJ, Braak H, Xue QS, Bechmann I. Dystrophic (senescent) rather than activated microglial cells are associated with tau pathology and likely precede neurodegeneration in Alzheimer's disease. *Acta Neuropathol* 2009; **118**: 475–485.
- 78 Fiala M, Liu QN, Sayre J, Pop V, Brahmandam V, Graves MC *et al*. Cyclooxygenase-2-positive macrophages infiltrate the Alzheimer's disease brain and damage the blood-brain barrier. *Eur J Clin Invest* 2002; **32**: 360–371.
- 79 Arnold SE, Han LY, Clark CM, Grossman M, Trojanowski JQ. Quantitative neuro-histological features of frontotemporal degeneration. *Neurobiol Aging* 2000; **21**: 913–919.
- 80 DeLuca GC, Joseph A, George J, Yates RL, Hamard M, Hofer M *et al*. Olfactory pathology in central nervous system demyelinating diseases. *Brain Pathol* 2015; **25**: 543–551.
- 81 Doorn KJ, Goudriaan A, Blits-Huizinga C, Bol JG, Rozemuller AJ, Hoogland PV *et al*. Increased amoeboid microglial density in the olfactory bulb of Parkinson's and Alzheimer's patients. *Brain Pathol* 2014; **24**: 152–165.
- 82 Hultman K, Strickland S, Norris EH. The APOE varepsilon4/varepsilon4 genotype potentiates vascular fibrin(ogen) deposition in amyloid-laden vessels in the brains of Alzheimer's disease patients. *J Cereb Blood Flow Metab* 2013; **33**: 1251–1258.
- 83 Perez-Nieves BG, Stein TD, Tai HC, Dols-Icardo O, Scotton TC, Barroeta-Espar I *et al*. Dissecting phenotypic traits linked to human resilience to Alzheimer's pathology. *Brain* 2013; **136**(Pt 8): 2510–2526.
- 84 Serrano-Pozo A, Mielke ML, Gomez-Isla T, Betensky RA, Growdon JH, Frosch MP *et al*. Reactive glia not only associates with plaques but also parallels tangles in Alzheimer's disease. *Am J Pathol* 2011; **179**: 1373–1384.
- 85 Van Everbroeck B, Dewulf E, Pals P, Lubke U, Martin JJ, Cras P. The role of cytokines, astrocytes, microglia and apoptosis in Creutzfeldt-Jakob disease. *Neurobiol Aging* 2002; **23**: 59–64.
- 86 Arnold SE, Trojanowski JQ, Gur RE, Blackwell P, Han LY, Choi C. Absence of neurodegeneration and neural injury in the cerebral cortex in a sample of elderly patients with schizophrenia. *Arch Gen Psychiatry* 1998; **55**: 225–232.
- 87 Alvarez MI, Rivas L, Lacruz C, Toledano A. Astroglial cell subtypes in the cerebella of normal adults, elderly adults, and patients with Alzheimer's disease: a histological and immunohistochemical comparison. *Glia* 2015; **63**: 287–312.
- 88 Falke E, Han LY, Arnold SE. Absence of neurodegeneration in the thalamus and caudate of elderly patients with schizophrenia. *Psychiatry Res* 2000; **93**: 103–110.
- 89 Lue LF, Walker DG, Rogers J. Modeling microglial activation in Alzheimer's disease with human postmortem microglial cultures. *Neurobiol Aging* 2001; **22**(6): 945–956.
- 90 Doens D, Fernandez PL. Microglia receptors and their implications in the response to amyloid beta for Alzheimer's disease pathogenesis. *J Neuroinflamm* 2014; **11**: 48.
- 91 Shan L, Bossers K, Unmehopa U, Bao AM, Swaab DF. Alterations in the histaminergic system in Alzheimer's disease: a postmortem study. *Neurobiol Aging* 2012; **33**: 2585–2598.
- 92 Becher B, Antel JP. Comparison of phenotypic and functional properties of immediately ex vivo and cultured human adult microglia. *Glia* 1996; **18**: 1–10.
- 93 Licastro F, Mallory M, Hansen LA, Masliah E. Increased levels of alpha-1-antichymotrypsin in brains of patients with Alzheimer's disease correlate with activated astrocytes and are affected by APOE 4 genotype. *J Neuroimmunol* 1998; **88**: 105–110.
- 94 Masliah E, Mallory M, Hansen L, Alford M, Albright T, Terry R *et al*. Immunoreactivity of CD45, a protein phosphotyrosine phosphatase, in Alzheimer's disease. *Acta Neuropathol* 1991; **83**: 12–20.

- 95 Colton CA, Mott RT, Sharpe H, Xu Q, Van Nostrand WE, Vitek MP. Expression profiles for macrophage alternative activation genes in AD and in mouse models of AD. *J Neuroinflamm* 2006; **3**: 27.
- 96 Kwiatek-Majkusiak J, Dickson DW, Tacik P, Aoki N, Tomasiuk R, Kozirowski D et al. Relationships between typical histopathological hallmarks and the ferritin in the hippocampus from patients with Alzheimer's disease. *Acta Neurobiol Exp* 2015; **75**: 391–398.
- 97 Meadowcroft MD, Connor JR, Yang QX. Cortical iron regulation and inflammatory response in Alzheimer's disease and APPSWE/PS1DeltaE9 mice: a histological perspective. *Front Neurosci* 2015; **9**: 255.
- 98 Ohgami T, Kitamoto T, Shin RW, Kaneko Y, Ogomori K, Tateishi J. Increased senile plaques without microglia in Alzheimer's disease. *Acta Neuropathol* 1991; **81**: 242–247.
- 99 Mochizuki A, Peterson JW, Mufson EJ, Trapp BD. Amyloid load and neural elements in Alzheimer's disease and nondemented individuals with high amyloid plaque density. *Exp Neurol* 1996; **142**: 89–102.
- 100 Fukumoto H, Asami-Odaka A, Suzuki N, Iwatsubo T. Association of A beta 40-positive senile plaques with microglial cells in the brains of patients with Alzheimer's disease and in non-demented aged individuals. *Neurodegeneration* 1996; **5**: 13–17.
- 101 DiPatre PL, Gelman BB. Microglial cell activation in aging and Alzheimer disease: partial linkage with neurofibrillary tangle burden in the hippocampus. *J Neuropathol Exp Neurol* 1997; **56**: 143–149.
- 102 Bertram L, Lange C, Mullin K, Parkinson M, Hsiao M, Hogan MF et al. Genome-wide association analysis reveals putative Alzheimer's disease susceptibility loci in addition to APOE. *Am J Hum Genet* 2008; **83**: 623–632.
- 103 Hollingworth P, Harold D, Sims R, Gerrish A, Lambert JC, Carrasquillo MM et al. Common variants at ABCA7, MS4A6A/MS4A4E, EPHA1, CD33 and CD2AP are associated with Alzheimer's disease. *Nat Genet* 2011; **43**: 429–435.
- 104 Malik M, Simpson JF, Parikh I, Wilfred BR, Fardo DW, Nelson PT et al. CD33 Alzheimer's risk-altering polymorphism, CD33 expression, and exon 2 splicing. *J Neurosci* 2013; **33**: 13320–13325.
- 105 Takahashi K, Rochford CD, Neumann H. Clearance of apoptotic neurons without inflammation by microglial triggering receptor expressed on myeloid cells-2. *J Exp Med* 2005; **201**: 647–657.
- 106 Steubl D, Roos M, Hettwer S, Angermann S, Wen M, Schmauder C et al. Comparison of peritoneal low-molecular-weight-protein-removal in CCPD and CAPD patients based on C-terminal agrin fragment clearance. *Kidney Blood Press Res* 2016; **41**: 175–185.
- 107 Roussos P, Katsel P, Fam P, Tan W, Purohit DP, Haroutunian V. The triggering receptor expressed on myeloid cells 2 (TREM2) is associated with enhanced inflammation, neuropathological lesions and increased risk for Alzheimer's dementia. *Alzheimers Dement* 2015; **11**: 1163–1170.
- 108 Itoh Y, Yamada M, Suematsu N, Matsushita M, Otomo E. An immunohistochemical study of centenarian brains: a comparison. *J Neurol Sci* 1998; **157**: 73–81.
- 109 Paulus W, Bancher C, Jellinger K. Microglial reaction in Pick's disease. *Neurosci Lett* 1993; **161**: 89–92.
- 110 Yamada M, Itoh Y, Sodeyama N, Suematsu N, Otomo E, Matsushita M et al. Senile dementia of the neurofibrillary tangle type: a comparison with Alzheimer's disease. *Dement Geriatr Cogn Disord* 2001; **12**: 117–126.
- 111 Sheng JG, Jones RA, Zhou XQ, McGinness JM, Van Eldik LJ, Mrak RE et al. Interleukin-1 promotion of MAPK-p38 overexpression in experimental animals and in Alzheimer's disease: potential significance for tau protein phosphorylation. *Neurochem Int* 2001; **39**: 341–348.
- 112 Sheng JG, Mrak RE, Griffin WS. Microglial interleukin-1 alpha expression in brain regions in Alzheimer's disease: correlation with neuritic plaque distribution. *Neuropathol Appl Neurobiol* 1995; **21**: 290–301.
- 113 Sheng JG, Griffin WS, Royston MC, Mrak RE. Distribution of interleukin-1-immunoreactive microglia in cerebral cortical layers: implications for neuritic plaque formation in Alzheimer's disease. *Neuropathol Appl Neurobiol* 1998; **24**: 278–283.
- 114 Mannoji H, Yeger H, Becker LE. A specific histochemical marker (lectin Ricinus communis agglutinin-1) for normal human microglia, and application to routine histopathology. *Acta Neuropathol* 1986; **71**: 341–343.
- 115 Mackenzie IR, Hao C, Munoz DG. Role of microglia in senile plaque formation. *Neurobiol Aging* 1995; **16**: 797–804.
- 116 Sheffield LG, Marquis JG, Berman NE. Regional distribution of cortical microglia parallels that of neurofibrillary tangles in Alzheimer's disease. *Neurosci Lett* 2000; **285**: 165–168.
- 117 Kravitz E, Gaisler-Salomon I, Biegan A. Hippocampal glutamate NMDA receptor loss tracks progression in Alzheimer's disease: quantitative autoradiography in postmortem human brain. *PLoS ONE* 2013; **8**: e81244.
- 118 Marutle A, Gillberg PG, Bergfors A, Yu W, Ni R, Nennesmo I et al. (3)H-deprenyl and (3)H-PIB autoradiography show different laminar distributions of astroglia and fibrillar beta-amyloid in Alzheimer brain. *J Neuroinflamm* 2013; **10**: 90.
- 119 Knezevic D, Mizrahi R. Molecular imaging of neuroinflammation in Alzheimer's disease and mild cognitive impairment. *Prog Neuropsychopharmacol Biol Psychiatry* 2017.
- 120 Pey P, Pearce RK, Kalaitzakis ME, Griffin WS, Gentleman SM. Phenotypic profile of alternative activation marker CD163 is different in Alzheimer's and Parkinson's disease. *Acta Neuropathol Commun* 2014; **2**: 21.
- 121 Shefer VF. Hippocampal pathology as a possible factor in the pathogenesis of senile dementias. *Neurosci Behav Physiol* 1977; **8**: 236–239.
- 122 Pelvig DP, Pakkenberg H, Regeur L, Oster S, Pakkenberg B. Neocortical glial cell numbers in Alzheimer's disease. A stereological study. *Dement Geriatr Cogn Disord* 2003; **16**: 212–219.
- 123 Cribbs DH, Berchtold NC, Perreau V, Coleman PD, Rogers J, Tenner AJ et al. Extensive innate immune gene activation accompanies brain aging, increasing vulnerability to cognitive decline and neurodegeneration: a microarray study. *J Neuroinflamm* 2012; **9**: 179.
- 124 Durrenberger PF, Fernando FS, Kashefi SN, Bonnert TP, Seilhean D, Nait-Oumesmar B et al. Common mechanisms in neurodegeneration and neuroinflammation: a BrainNet Europe gene expression microarray study. *J Neural Transm* 2015; **122**: 1055–1068.
- 125 Li X, Long J, He T, Belshaw R, Scott J. Integrated genomic approaches identify major pathways and upstream regulators in late onset Alzheimer's disease. *Sci Rep* 2015; **5**: 12393.
- 126 Podtelezchnikov AA, Tanis KQ, Nebozhyn M, Ray WJ, Stone DJ, Loboda AP. Molecular insights into the pathogenesis of Alzheimer's disease and its relationship to normal aging. *PLoS ONE* 2011; **6**: e29610.
- 127 Seyfried NT, Dammer EB, Swarup V, Nandakumar D, Duong DM, Yin L et al. A multi-network approach identifies protein-specific co-expression in asymptomatic and symptomatic Alzheimer's disease. *Cell Syst* 2017; **4**: 60–72 e64.
- 128 Su JH, Deng G, Cotman CW. Bax protein expression is increased in Alzheimer's brain: correlations with DNA damage, Bcl-2 expression, and brain pathology. *J Neuropathol Exp Neurol* 1997; **56**: 86–93.
- 129 McGeer PL, Akiyama H, Itagaki S, McGeer EG. Immune system response in Alzheimer's disease. *Can J Neurol Sci* 1989; **16**(4 Suppl): 516–527.
- 130 Bayer TA, Buslei R, Havas L, Falkai P. Evidence for activation of microglia in patients with psychiatric illnesses. *Neurosci Lett* 1999; **271**: 126–128.
- 131 Togo T, Akiyama H, Kondo H, Ikeda K, Kato M, Iseki E et al. Expression of CD40 in the brain of Alzheimer's disease and other neurological diseases. *Brain Res* 2000; **885**: 117–121.
- 132 Bowser R, Reilly S. Expression of FAC1 in activated microglia during Alzheimer's disease. *Neurosci Lett* 1998; **253**: 163–166.
- 133 Rogers J, Lubner-Narod J, Styren SD, Civin WH. Expression of immune system-associated antigens by cells of the human central nervous system: relationship to the pathology of Alzheimer's disease. *Neurobiol Aging* 1988; **9**: 339–349.
- 134 Yoshiyama Y, Arai K, Oki T, Hattori T. Expression of invariant chain and pro-cathepsin L in Alzheimer's brain. *Neurosci Lett* 2000; **290**: 125–128.
- 135 Wiendl H, Feger U, Mittelbronn M, Jack C, Schreiner B, Stadelmann C et al. Expression of the immune-tolerogenic major histocompatibility molecule HLA-G in multiple sclerosis: implications for CNS immunity. *Brain* 2005; **128**(Pt 11): 2689–2704.
- 136 Christie RH, Chung H, Rebeck GW, Strickland D, Hyman BT. Expression of the very low-density lipoprotein receptor (VLDL-r), an apolipoprotein-E receptor, in the central nervous system and in Alzheimer's disease. *J Neuropathol Exp Neurol* 1996; **55**: 491–498.
- 137 Dickson DW, Sinicropi S, Yen SH, Ko LW, Mattiace LA, Bucala R et al. Glycation and microglial reaction in lesions of Alzheimer's disease. *Neurobiol Aging* 1996; **17**: 733–743.
- 138 Perlmutter LS, Scott SA, Barron E, Chui HC. MHC class II-positive microglia in human brain: association with Alzheimer lesions. *J Neurosci Res* 1992; **33**: 549–558.
- 139 Tooyama I, Kimura H, Akiyama H, McGeer PL. Reactive microglia express class I and class II major histocompatibility complex antigens in Alzheimer's disease. *Brain Res* 1990; **523**: 273–280.
- 140 An L, Sato H, Konishi Y, Walker DG, Beach TG, Rogers J et al. Expression and localization of lactotransferrin messenger RNA in the cortex of Alzheimer's disease. *Neurosci Lett* 2009; **452**: 277–280.
- 141 Christie RH, Freeman M, Hyman BT. Expression of the macrophage scavenger receptor, a multifunctional lipoprotein receptor, in microglia associated with senile plaques in Alzheimer's disease. *Am J Pathol* 1996; **148**: 399–403.
- 142 Sasaki A, Yamaguchi H, Ogawa A, Sugihara S, Nakazato Y. Microglial activation in early stages of amyloid beta protein deposition. *Acta Neuropathol* 1997; **94**: 316–322.
- 143 Styren SD, Civin WH, Rogers J. Molecular, cellular, and pathologic characterization of HLA-DR immunoreactivity in normal elderly and Alzheimer's disease brain. *Exp Neurol* 1990; **110**: 93–104.

- 144 McGeer PL, Kawamata T, Walker DG, Akiyama H, Tooyama I, McGeer EG. Microglia in degenerative neurological disease. *Glia* 1993; **7**: 84–92.
- 145 Lopez-Gonzalez I, Schluter A, Aso E, Garcia-Esparcia P, Ansoleaga B, LL F *et al*. Neuroinflammatory signals in Alzheimer disease and APP/PS1 transgenic mice: correlations with plaques, tangles, and oligomeric species. *J Neuropathol Exp Neurol* 2015; **74**: 319–344.
- 146 Wu CK, Thal L, Pizzo D, Hansen L, Masliah E, Geula C. Apoptotic signals within the basal forebrain cholinergic neurons in Alzheimer's disease. *Exp Neurol* 2005; **195**: 484–496.
- 147 Jellinger KA, Paulus W, Grundke-Iqbal I, Riederer P, Youdim MB. Brain iron and ferritin in Parkinson's and Alzheimer's diseases. *J Neural Transm Park Dis Dement Sect* 1990; **2**: 327–340.
- 148 Zeineh MM, Chen Y, Kitzler HH, Hammond R, Vogel H, Rutt BK. Activated iron-containing microglia in the human hippocampus identified by magnetic resonance imaging in Alzheimer disease. *Neurobiol Aging* 2015; **36**: 2483–2500.
- 149 van Duijn S, Nabuurs RJ, van Duinen SG, Natta R. Comparison of histological techniques to visualize iron in paraffin-embedded brain tissue of patients with Alzheimer's disease. *J Histochem Cytochem* 2013; **61**: 785–792.
- 150 Grundke-Iqbal I, Fleming J, Tung YC, Lassmann H, Iqbal K, Joshi JG. Ferritin is a component of the neuritic (senile) plaque in Alzheimer dementia. *Acta Neuropathol* 1990; **81**: 105–110.
- 151 Yamada T, Yoshiyama Y, Sato H, Seiki M, Shinagawa A, Takahashi M. White matter microglia produce membrane-type matrix metalloprotease, an activator of gelatinase A, in human brain tissues. *Acta Neuropathol* 1995; **90**: 421–424.
- 152 Akiyama H, Ikeda K, Katoh M, McGeer EG, McGeer PL. Expression of MRP14, 27E10, interferon-alpha and leukocyte common antigen by reactive microglia in postmortem human brain tissue. *J Neuroimmunol* 1994; **50**: 195–201.
- 153 Yamada T, Horisberger MA, Kawaguchi N, Moroo I, Toyoda T. Immunohistochemistry using antibodies to alpha-interferon and its induced protein, MxA, in Alzheimer's and Parkinson's disease brain tissues. *Neurosci Lett* 1994; **181**: 61–64.
- 154 Yamada T, Yoshiyama Y, Kawaguchi N, Ichinose A, Iwaki T, Hirose S *et al*. Possible roles of transglutaminases in Alzheimer's disease. *Dement Geriatr Cogn Disord* 1998; **9**: 103–110.
- 155 Yamada T, Miyazaki K, Koshikawa N, Takahashi M, Akatsu H, Yamamoto T. Selective localization of gelatinase A, an enzyme degrading beta-amyloid protein, in white matter microglia and in Schwann cells. *Acta Neuropathol* 1995; **89**: 199–203.
- 156 Scott SA, Johnson SA, Zarow C, Perlmuter LS. Inability to detect beta-amyloid protein precursor mRNA in Alzheimer plaque-associated microglia. *Exp Neurol* 1993; **121**: 113–118.
- 157 Dickson DW, Farlo J, Davies P, Crystal H, Fuld P, Yen SH. Alzheimer's disease. A double-labeling immunohistochemical study of senile plaques. *Am J Pathol* 1988; **132**: 86–101.
- 158 Wirths O, Hillmann A, Pradier L, Hartig W, Bayer TA. Oligomeric pyroglutamate amyloid-beta is present in microglia and a subfraction of vessels in patients with Alzheimer's disease: implications for immunotherapy. *J Alzheimers Dis* 2013; **35**: 741–749.
- 159 Togo T, Akiyama H, Iseki E, Kondo H, Ikeda K, Kato M *et al*. Occurrence of T cells in the brain of Alzheimer's disease and other neurological diseases. *J Neuroimmunol* 2002; **124**: 83–92.
- 160 Yamada T, Yamanaka I. Microglial localization of alpha-interferon receptor in human brain tissues. *Neurosci Lett* 1995; **189**: 73–76.
- 161 Akiyama H, Ikeda K, Kondo H, Kato M, McGeer PL. Microglia express the type 2 plasminogen activator inhibitor in the brain of control subjects and patients with Alzheimer's disease. *Neurosci Lett* 1993; **164**: 233–235.
- 162 Yamada T, Tsujioka Y, Taguchi J, Takahashi M, Tsuboi Y, Moroo I *et al*. Melanotransferrin is produced by senile plaque-associated reactive microglia in Alzheimer's disease. *Brain Res* 1999; **845**: 1–5.
- 163 Liu Y, Walter S, Stagi M, Cherny D, Letiembre M, Schulz-Schaeffer W *et al*. LPS receptor (CD14): a receptor for phagocytosis of Alzheimer's amyloid peptide. *Brain* 2005; **128**(Pt 8): 1778–1789.
- 164 Guillemain GJ, Brew BJ, Noonan CE, Takikawa O, Cullen KM. Indoleamine 2,3 dioxygenase and quinolinic acid immunoreactivity in Alzheimer's disease hippocampus. *Neuropathol Appl Neurobiol* 2005; **31**: 395–404.
- 165 Akiyama H, Nishimura T, Kondo H, Ikeda K, Hayashi Y, McGeer PL. Expression of the receptor for macrophage colony stimulating factor by brain microglia and its upregulation in brains of patients with Alzheimer's disease and amyotrophic lateral sclerosis. *Brain Res* 1994; **639**: 171–174.
- 166 Bryan KJ, Zhu X, Harris PL, Perry G, Castellani RJ, Smith MA *et al*. Expression of CD74 is increased in neurofibrillary tangles in Alzheimer's disease. *Mol Neurodegener* 2008; **3**: 13.
- 167 Minnash P, Yamamoto Y, Ohkubo I, Nishi K. Demonstration of puromycin-sensitive alanyl aminopeptidase in Alzheimer disease brain. *Leg Med* 2003; **5** (Suppl 1): S285–S287.
- 168 Haga S, Akai K, Ishii T. Demonstration of microglial cells in and around senile (neuritic) plaques in the Alzheimer brain. An immunohistochemical study using a novel monoclonal antibody. *Acta Neuropathol* 1989; **77**: 569–575.
- 169 Du Yan S, Zhu H, Fu J, Yan SF, Roher A, Tourtellotte WW *et al*. Amyloid-beta peptide-receptor for advanced glycation endproduct interaction elicits neuronal expression of macrophage-colony stimulating factor: a proinflammatory pathway in Alzheimer disease. *Proc Natl Acad Sci USA* 1997; **94**: 5296–5301.
- 170 Carrano A, Hoozemans JJ, van der Vies SM, Rozemuller AJ, van Horsen J, de Vries HE. Amyloid Beta induces oxidative stress-mediated blood-brain barrier changes in capillary amyloid angiopathy. *Antioxid Redox Signal* 2011; **15**: 1167–1178.
- 171 Wong A, Luth HJ, Deuther-Conrad W, Dukic-Stefanovic S, Gasic-Milenkovic J, Arendt T *et al*. Advanced glycation endproducts co-localize with inducible nitric oxide synthase in Alzheimer's disease. *Brain Res* 2001; **920**: 32–40.
- 172 Rozemuller AJ, Eikelenboom P, Theeuwes JW, Jansen Steur EN, de Vos RA. Activated microglial cells and complement factors are unrelated to cortical Lewy bodies. *Acta Neuropathol* 2000; **100**: 701–708.
- 173 Ryu JK, McLarnon JG. A leaky blood-brain barrier, fibrinogen infiltration and microglial reactivity in inflamed Alzheimer's disease brain. *J Cell Mol Med* 2009; **13**(9A): 2911–2925.
- 174 Xia MQ, Qin SX, Wu LJ, Mackay CR, Hyman BT. Immunohistochemical study of the beta-chemokine receptors CCR3 and CCR5 and their ligands in normal and Alzheimer's disease brains. *Am J Pathol* 1998; **153**: 31–37.
- 175 Hoozemans JJ, Rozemuller AJ, Janssen I, De Groot CJ, Veerhuis R, Eikelenboom P. Cyclooxygenase expression in microglia and neurons in Alzheimer's disease and control brain. *Acta Neuropathol* 2001; **101**: 2–8.
- 176 Arends YM, Duyckaerts C, Rozemuller JM, Eikelenboom P, Hauw JJ. Microglia, amyloid and dementia in Alzheimer disease. A correlative study. *Neurobiol Aging* 2000; **21**: 39–47.
- 177 Ricciarelli R, D'Abramo C, Zingg JM, Giliberto L, Markesbery W, Azzi A *et al*. CD36 overexpression in human brain correlates with beta-amyloid deposition but not with Alzheimer's disease. *Free Radic Biol Med* 2004; **36**: 1018–1024.
- 178 Walker DG, Lue LF, Beach TG. Increased expression of the urokinase plasminogen-activator receptor in amyloid beta peptide-treated human brain microglia and in AD brains. *Brain Res* 2002; **926**: 69–79.
- 179 Lyoo CH, Ikawa M, Liow JS, Zoghbi SS, Morse CL, Pike VW *et al*. Cerebellum can serve as a pseudo-reference region in Alzheimer disease to detect neuroinflammation measured with PET radioligand binding to translocator protein. *J Nucl Med* 2015; **56**: 701–706.
- 180 Green PS, Mendez AJ, Jacob JS, Crowley JR, Growdon W, Hyman BT *et al*. Neuronal expression of myeloperoxidase is increased in Alzheimer's disease. *J Neurochem* 2004; **90**: 724–733.
- 181 Heneka MT, Sastre M, Dumitrescu-Ozimek L, Hanke A, Dewachter I, Kuperi C *et al*. Acute treatment with the PPARgamma agonist pioglitazone and ibuprofen reduces glial inflammation and Abeta1-42 levels in APPV7171 transgenic mice. *Brain* 2005; **128**(Pt 6): 1442–1453.
- 182 Mackenzie IR, Munoz DG. Nonsteroidal anti-inflammatory drug use and Alzheimer-type pathology in aging. *Neurology* 1998; **50**: 986–990.
- 183 Lucas SM, Rothwell NJ, Gibson RM. The role of inflammation in CNS injury and disease. *Br J Pharmacol* 2006; **147**(Suppl 1): S232–S240.
- 184 Najjar S, Pearlman DM, Alper K, Najjar A, Devinsky O. Neuroinflammation and psychiatric illness. *J Neuroinflamm* 2013; **10**: 43.
- 185 Trepanier MO, Hopperton KE, Mizrahi R, Mechawar N, Bazinet RP. Postmortem evidence of cerebral inflammation in schizophrenia: a systematic review. *Mol Psychiatry* 2016; **21**: 1009–1026.
- 186 Zotova E, Barambe V, Cheaveau M, Morgan W, Holmes C, Harris S *et al*. Inflammatory components in human Alzheimer's disease and after active amyloid-beta42 immunization. *Brain* 2013; **136**(Pt 9): 2677–2696.
- 187 Paquet C, Amin J, Mouton-Liger F, Nasser M, Love S, Gray F *et al*. Effect of active Abeta immunotherapy on neurons in human Alzheimer's disease. *J Pathol* 2015; **235**: 721–730.
- 188 Lyck L, Dalmau I, Chemnitz J, Finsen B, Schroder HD. Immunohistochemical markers for quantitative studies of neurons and glia in human neocortex. *J Histochem Cytochem* 2008; **56**: 201–221.
- 189 Torres EM, Meldrum A, Kirik D, Dunnett SB. An investigation of the problem of two-layered immunohistochemical staining in paraformaldehyde fixed sections. *J Neurosci Methods* 2006; **158**: 64–74.
- 190 Ghatak A, Combs CK. Iba1 immunoreactivity is enhanced following an antigen retrieval treatment with EDTA, pH 6.0. *MethodsX* 2014; **1**: 269–274.
- 191 Kashon ML, Ross GW, O'Callaghan JP, Miller DB, Petrovitch H, Burchfiel CM *et al*. Associations of cortical astrogliosis with cognitive performance and dementia status. *J Alzheimers Dis* 2004; **6**: 595–604.
- 192 Muramori F, Kobayashi K, Nakamura I. A quantitative study of neurofibrillary tangles, senile plaques and astrocytes in the hippocampal subdivisions and entorhinal cortex in Alzheimer's disease, normal controls and non-Alzheimer neuropsychiatric diseases. *Psychiatry Clin Neurosci* 1998; **52**: 593–599.

- 193 Akiyama H, Mori H, Saido T, Kondo H, Ikeda K, McGeer PL. Occurrence of the diffuse amyloid beta-protein (Abeta) deposits with numerous Abeta-containing glial cells in the cerebral cortex of patients with Alzheimer's disease. *Glia* 1999; **25**: 324–331.
- 194 Bouvier DS, Jones EV, Quesseveur G, Davoli MA, Ferreira AT, Quirion R *et al*. High resolution dissection of reactive glial nets in Alzheimer's disease. *Sci Rep* 2016; **6**: 24544.
- 195 Irie-Sasaki J, Sasaki T, Penninger JM. CD45 regulated signaling pathways. *Curr Top Med Chem* 2003; **3**: 783–796.
- 196 Kaneko Y, Kitamoto T, Tateishi J, Yamaguchi K. Ferritin immunohistochemistry as a marker for microglia. *Acta Neuropathol* 1989; **79**: 129–136.
- 197 Wlodarczyk A, Cedile O, Jensen KN, Jasson A, Mony JT, Khorrooshi R *et al*. Pathologic and protective roles for microglial subsets and bone marrow- and blood-derived myeloid cells in central nervous system inflammation. *Front Immunol* 2015; **6**: 463.
- 198 Dinarello CA. Immunological and inflammatory functions of the interleukin-1 family. *Annu Rev Immunol* 2009; **27**: 519–550.
- 199 Hutchins KD, Dickson DW, Rashbaum WK, Lyman WD. Localization of microglia in the human fetal cervical spinal cord. *Brain Res Dev Brain Res* 1992; **66**: 270–273.
- 200 Beckers L, Ory D, Geric I, Declercq L, Koole M, Kassiou M *et al*. Increased expression of translocator protein (TSPO) marks pro-inflammatory microglia but does not predict neurodegeneration. *Mol Imaging Biol* 2017.
- 201 Bonsack Ft Alleyne CH Jr, Sukumari-Ramesh S. Augmented expression of TSPO after intracerebral hemorrhage: a role in inflammation? *J Neuroinflammation* 2016; **13**: 151.



This work is licensed under a Creative Commons Attribution-NonCommercial-ShareAlike 4.0 International License. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in the credit line; if the material is not included under the Creative Commons license, users will need to obtain permission from the license holder to reproduce the material. To view a copy of this license, visit <http://creativecommons.org/licenses/by-nc-sa/4.0/>

© The Author(s) 2018

Supplementary Information accompanies the paper on the Molecular Psychiatry website (<http://www.nature.com/mp>)