

ARTICLE

Received 24 Jan 2015 | Accepted 13 May 2015 | Published 25 Jun 2015

DOI: 10.1038/ncomms8464

OPEN

IL-1 receptor antagonist-deficient mice develop autoimmune arthritis due to intrinsic activation of IL-17-producing CCR2⁺ V γ 6⁺ $\gamma\delta$ T cells

Aoi Akitsu^{1,2,3,4,5}, Harumichi Ishigame^{1,†}, Shigeru Kakuta^{1,†}, Soo-hyun Chung^{1,5}, Satoshi Ikeda¹, Kenji Shimizu^{1,5}, Sachiko Kubo^{1,5}, Yang Liu^{1,†}, Masayuki Umemura⁶, Goro Matsuzaki⁶, Yasunobu Yoshikai⁷, Shinobu Saijo^{1,†} & Yoichiro Iwakura^{1,2,4,5}

Interleukin-17 (IL-17)-producing $\gamma\delta$ T ($\gamma\delta$ 17) cells have been implicated in inflammatory diseases, but the underlying pathogenic mechanisms remain unclear. Here, we show that both CD4⁺ and $\gamma\delta$ 17 cells are required for the development of autoimmune arthritis in IL-1 receptor antagonist (IL-1Ra)-deficient mice. Specifically, activated CD4⁺ T cells direct $\gamma\delta$ T-cell infiltration by inducing CCL2 expression in joints. Furthermore, IL-17 reporter mice reveal that the V γ 6⁺ subset of CCR2⁺ $\gamma\delta$ T cells preferentially produces IL-17 in inflamed joints. Importantly, because IL-1Ra normally suppresses IL-1R expression on $\gamma\delta$ T cells, IL-1Ra-deficient mice exhibit elevated IL-1R expression on V γ 6⁺ cells, which play a critical role in inducing them to produce IL-17. Our findings demonstrate a pathogenic mechanism in which adaptive and innate immunity induce an autoimmune disease in a coordinated manner.

¹Laboratory of Molecular Pathogenesis, Center for Experimental Medicine and Systems Biology, The Institute of Medical Science, The University of Tokyo, Tokyo 108-8639, Japan. ²Department of Biophysics and Biochemistry, Graduate School of Science, The University of Tokyo, Tokyo 113-0032, Japan. ³Research Fellow of the Japan Society for the Promotion of Science (JSPS), Tokyo 102-0083, Japan. ⁴Core Research for Evolutional Science and Technology (CREST), Japan Science and Technology Agency, Saitama 332-0012, Japan. ⁵Division of Experimental Animal Immunology, Center for Animal Disease Models, Research Institute for Biomedical Sciences, Tokyo University of Science, Chiba 278-0022, Japan. ⁶Tropical Biosphere Research Center, University of the Ryukyus, Okinawa 903-0213, Japan. ⁷Research Center for Prevention of Infectious Diseases, Medical Institute of Bioregulation, Kyushu University, Fukuoka 812-8582, Japan. † Present addresses: RIKEN Center for Integrative Medical Sciences, Kanagawa 230-0045, Japan (H.I.); Department of Biomedical Science, Graduate School of Agricultural and Life Sciences, The University of Tokyo, Tokyo 113-8657, Japan (S.K.); Renji Hospital Clinical Stem Cell Research Center, Shanghai Jiao Tong University School of Medicine, Shang Hai 200127, China (Y.L.); Department of Molecular Immunology, Medical Mycology Research Center, Chiba University, Chiba 263-8522, Japan (S.S.). Correspondence and requests for materials should be addressed to Y.I. (email: iwakura@rs.tus.ac.jp).

Interleukin (IL)-17 plays important roles in the development of autoimmune diseases, such as rheumatoid arthritis and psoriasis, by inducing expression of proinflammatory cytokines and chemokines, recruiting neutrophils and activating T cells and B cells^{1,2}. Although helper CD4⁺ T (Th17) cells are well-known producers of IL-17 that contribute to the development of autoimmune diseases, recent studies showed that innate immune cells and innate-like cells are also important sources of IL-17 in local inflammatory tissues^{3,4}. Mouse autoimmune disease models have revealed that IL-17-producing $\gamma\delta$ T ($\gamma\delta 17$) cells are an important innate source of IL-17 (refs 5–13). In collagen-induced arthritis, experimental autoimmune encephalomyelitis and psoriasis-like skin inflammation, the synergy between $\gamma\delta 17$ and $\alpha\beta$ T cells is important for disease development^{5,6,11,14}, but it remains unclear how $\gamma\delta 17$ cells induce tissue-specific inflammation.

$\gamma\delta 17$ cells share many characteristics with Th17 cells. However, in contrast to Th17 cells in which differentiation in the periphery is required for IL-17 production, the functional potential of $\gamma\delta 17$ cells is already determined during intrathymic development^{15–17}. These $\gamma\delta$ thymocytes, which express the transcription factor ROR γ t and the signature cytokine receptor IL-23R¹⁸, leave the thymus as functionally committed cells¹⁹. Therefore, $\gamma\delta$ T cells produce IL-17 directly following stimulation with IL-1 β and IL-23 without T cell receptor (TCR) stimulation in the periphery^{5,13}. Although the expression of IL-23R on $\gamma\delta 17$ cells is constitutive⁵, expression of IL-1R in the periphery is tissue-type dependent²⁰. *Il1r1*^{-/-} mice or anti-IL-1R monoclonal antibody (mAb) treatment abrogates IL-17 production in $\gamma\delta$ T cells^{11,20}, suggesting that IL-1R expression plays a critical role in IL-17 production. However, the regulatory mechanism of IL-1R expression remains unclear. In addition to ROR γ t, transcription factors such as Blk²¹, Hes-1 (ref. 22), NF κ B²³, SOX4 and SOX13 (ref. 24) are also required for $\gamma\delta 17$ cell development.

In mice, the TCR γ locus consists of seven *V γ* (*V γ 1–V γ 7*) genes that are closely correlated with effector functions, although in most strains *V γ 3* is a pseudogene²⁵. IL-17 is produced by *V γ 4*⁺ and *V γ 6*⁺ $\gamma\delta$ T cells²⁶ (Heilig and Tonegawa's nomenclature²⁷). Although overall gene expression patterns are similar between these two subsets¹⁸, each subset has distinct features. *V γ 6*⁺ $\gamma\delta$ T cells express the invariant *V γ 6/V δ 1* TCR, develop only in the late-embryonic thymus and preferentially localize in the uterus, vagina, lung, dermis and peritoneal cavity^{28,29}. On the other hand, *V γ 4*⁺ $\gamma\delta$ T cells have a more diverse TCR repertoire, and develop in both foetal and adult thymus. Subsequently, they circulate in blood and reside in the dermis and secondary lymphoid organs³⁰. However, the differences between the pathogenic roles of *V γ 6*⁺ and *V γ 4*⁺ $\gamma\delta 17$ cells, particularly the contribution of *V γ 6*⁺ $\gamma\delta 17$ cells to inflammatory diseases, remain unclear.

The mechanism of $\gamma\delta 17$ cell migration into inflammatory sites is also poorly understood. Recent gene array analysis showed that the expression of chemokine receptors such as CCR6, CCR2 and CXCR6 is upregulated in $\gamma\delta 17$ thymocytes¹⁸. Moreover, CCR6 expression is often used as a marker of $\gamma\delta 17$ cells¹⁷. However, it remains unknown whether these chemokine receptors function in $\gamma\delta 17$ cell migration.

IL-1 receptor antagonist (IL-1Ra, gene symbol: *Il1rn*) is an endogenous inhibitor of IL-1 activity that competes with IL-1 α and IL-1 β for IL-1R binding. IL-1Ra-deficient (*Il1rn*^{-/-}) mice spontaneously develop arthritis in an IL-17- and T-cell-dependent manner^{31–33}, suggesting that excess IL-1 signalling caused by IL-1Ra deficiency induces IL-17 production from T cells and the development of arthritis. Here, we found that $\gamma\delta 17$ cells are the main producers of IL-17 in joints of *Il1rn*^{-/-} mice, and that recruitment of CCR2⁺ $\gamma\delta$ T cells to the joints via induction of CCL2 by CD4⁺ T cells is important for the

development of arthritis. Both *V γ 6*⁺ and *V γ 4*⁺ $\gamma\delta$ T cells were recruited to the joints, but only the *V γ 6*⁺ subset efficiently produced IL-17, because *Il1rn*^{-/-} *V γ 6*⁺ cells intrinsically expressed high levels of IL-1R due to the loss of down-regulation of IL-1R expression by IL-1Ra. These observations provide novel insights into the pathogenic mechanism underlying the development of autoimmune arthritis in *Il1rn*^{-/-} mice.

Results

$\gamma\delta$ T cells mainly produce IL-17 in *Il1rn*^{-/-} mouse joints. We analysed IL-17-producing cells in arthritic *Il1rn*^{-/-} mice. Both $\gamma\delta 17$ cell and Th17 cell populations were elevated in draining lymph nodes (LNs) of affected joints (Fig. 1a,b), whereas proportions of IL-17⁺CD8⁺ T cells and IL-17⁺DX5⁺ T cells were unchanged (Supplementary Fig. 1). Notably, most joint-infiltrating IL-17-producing cells were $\gamma\delta$ T cells, whereas Th17 cells were rare (Fig. 1c). Moreover, immunofluorescence staining revealed that IL-17 in the joints was primarily expressed in $\gamma\delta$ T cells (Fig. 1d). These observations suggest that $\gamma\delta 17$ cells play a pathogenic role in the development of arthritis in *Il1rn*^{-/-} mice.

$\gamma\delta$ and CD4⁺ T cells are involved in arthritis development. To analyse the pathogenic roles of $\gamma\delta$ T cells and CD4⁺ T cells, we injected either anti- $\gamma\delta$ TCR or anti-CD4 mAb into *Il1rn*^{-/-} mice before onset of disease. Both antibody treatments significantly suppressed the incidence of arthritis (Fig. 2a,b). Although the severity scores, determined by the swelling and ankylosing changes of the affected ankles of antibody-treated mice, were similar to those of untreated mice (Supplementary Fig. 2a,b), the microscopic histological scores significantly decreased following treatment with anti- $\gamma\delta$ TCR mAb (Fig. 2c,d). Anti- $\gamma\delta$ TCR mAb treatment depleted 70–90% of $\gamma\delta$ T cells at early time points, but the population gradually recovered (Supplementary Fig. 2c), probably due to the development of antibodies against this mAb. Nonetheless, the anti- $\gamma\delta$ TCR mAb treatment greatly decreased the $\gamma\delta 17$ population (Fig. 2e; Supplementary Fig. 2d), suggesting that anti- $\gamma\delta$ TCR mAb effectively depleted $\gamma\delta 17$ cells, at least at earlier times. IL-17 production in the $\gamma\delta$ TCR⁻ population was not elevated in these mice (Supplementary Fig. 2d). Most populations of CD4⁺ T cells, including Th17 cells, remained depleted in *Il1rn*^{-/-} mice on day 11 after anti-CD4 mAb treatment (Fig. 2f; Supplementary Fig. 2e). These results suggest that both $\gamma\delta$ T cells and CD4⁺ T cells are involved in the development of arthritis in *Il1rn*^{-/-} mice. However, arthritis developed normally in *Tcrd*^{-/-}*Il1rn*^{-/-} mice (Supplementary Fig. 2f), and IL-17-producing CD4⁻CD8⁻ $\gamma\delta$ TCR⁻ T cells were increased in the LNs and joints of these mice (Supplementary Fig. 2g).

Both $\gamma\delta 17$ and CD4⁺ T cells collaborate to develop arthritis.

Next, we directly examined the pathogenic role of $\gamma\delta$ T cells and CD4⁺ T cells by adoptive cell transfer. $\gamma\delta$ T cells derived from *Cd4*^{-/-}*Il1rn*^{-/-} mice and/or CD4⁺ T cells derived from *Tcrd*^{-/-}*Il1rn*^{-/-} mice were transferred into *scid/scid* mice. We found that *scid/scid* mice that received transfer of whole-*Il1rn*^{-/-} T cells or a mixture of $\gamma\delta$ and CD4⁺ T cells developed arthritis, whereas those that received transfer of $\gamma\delta$ or CD4⁺ T cells alone did not (Fig. 2g). Instead, development of inflammation was observed in other organs, such as the colon and dermis, when $\gamma\delta$ T cells alone were transferred (Supplementary Fig. 2h). When total T cells or a mixture of $\gamma\delta$ and CD4⁺ T cells were transferred, $\gamma\delta 17$ cells were present in arthritic joints (Fig. 2h); however, no $\gamma\delta$ T cells were located in the joints when $\gamma\delta$ T cells were transferred alone. When CD4⁺ T cells were transferred, no Th17 cells were observed in the joints, although





Figure 1 | $\gamma\delta$ T cells are the main source of IL-17 in the inflamed joints of *Il1rn*^{-/-} mice. (a-c) Flow cytometry of LN cells from WT ($n = 3$) and arthritic *Il1rn*^{-/-} mice ($n = 3$) (a) or joint-infiltrating cells from WT (pool of five mice) and arthritic *Il1rn*^{-/-} mice (pool of two mice) (c). All cells were stimulated with P/I for 5 h, and then stained for intracellular IL-17. Numbers refer to percent cells in CD3 ϵ ⁺ cells. Quantification of IL-17⁺ $\gamma\delta$ TCR⁺ and IL-17⁺ CD4⁺ in CD3 ϵ ⁺ cells are shown (b). ** $P < 0.01$ versus WT mice (unpaired Student's *t*-test). Data show mean \pm s.e.m. (d) Frozen sections of arthritic joints of *Il1rn*^{-/-} mice were co-stained with anti-IL-17 Ab (green), anti- $\gamma\delta$ TCR Ab (red) and 4,6-diamidino-2-phenylindole (DAPI) (blue). Sections were observed under a fluorescence microscope (top and middle panels, scale bars, 100 μ m), and a confocal microscope (bottom panels, scale bars, 10 μ m). B, bone. IL-17⁺ $\gamma\delta$ TCR⁺ cells are shown by white arrows. Similar results were obtained in six other *Il1rn*^{-/-} mice. All data except d are representative of > 5 independent experiments.

IL-17⁻ CD4⁺ T cells were found (Fig. 2h). In contrast to the joints, $\gamma\delta 17$ and Th17 cells were observed in LNs of *scid/scid* mice that received transfer of $\gamma\delta$ and CD4⁺ T cells, respectively

(Supplementary Fig. 2i). Thus, *Il1rn*^{-/-} CD4⁺ T cells are required for the recruitment of $\gamma\delta$ T cells to the joints, and $\gamma\delta$ T cells are required for the production of IL-17.

Joint-infiltrating $\gamma\delta 17$ cells predominantly express CCR2. To analyse $\gamma\delta 17$ cells in joints, we generated an IL-17 reporter (*Il17^{g/g}*) mouse, in which IRES-eGFP was inserted into the *Il17a* locus without affecting IL-17 production (Supplementary Fig. 3a,b). Similar to *Il1rn^{-/-}* mice, *Il17^{g/g}Il1rn^{-/-}* mice spontaneously developed arthritis (Supplementary Fig. 3c). Green

fluorescent protein (GFP) expression correlated with intracellular IL-17 expression in joint-infiltrating cells from *Il17^{g/g}Il1rn^{-/-}* mice after phorbol myristate acetate and ionomycin (P/I) stimulation (Supplementary Fig. 3d), indicating that GFP expression accurately reflects IL-17 expression. Consistent with intracellular IL-17 staining in *Il1rn^{-/-}* mice (Fig. 1c), joint GFP



expression was mostly limited to $\gamma\delta$ T cells (Supplementary Fig. 3e). GFP expression was detectable without any stimulation in joints, whereas only few were detected in LNs (Supplementary Fig. 3f,d), suggesting that $\gamma\delta 17$ cells were activated in joints.

Several chemokine receptors, including CCR2, CXCR6 and CCR6, were expressed on joint-infiltrating $\gamma\delta$ T cells in *Il1rn*^{-/-} mice (Supplementary Fig. 4a). In particular, almost all $\gamma\delta$ T cells expressed CCR2, but not CCR5, CXCR4 or CCR9. Using these *Il17^{g/g}Il1rn*^{-/-} mice, we found that about 95% of $\gamma\delta 17$ cells expressed CCR2, and that relatively large proportions of $\gamma\delta 17$ cells also expressed CXCR6 and CCR6 (Fig. 3a).

Next, we examined chemokine expression in joints. Consistent with our previous microarray analysis³⁴, expression of *Ccl2* (ligand for CCR2), *Cxcl16* (ligand for CXCR6), *Ccl5* (ligand for CCR5) and *Cxcl12* (ligand for CXCR4) was significantly elevated in joints of *Il1rn*^{-/-} mice relative to wild-type (WT) mice, whereas expression of *Ccl20* (ligand for CCR6) was unchanged (Fig. 3b). Moreover, we detected moderate levels of $\gamma\delta$ T-cell infiltration in the non-arthritis joints of *Il17a*^{-/-}*Il1rn*^{-/-} mice³². In these mice, $\gamma\delta$ T cells expressed high levels of CCR2, but not CXCR6 (Fig. 3c), indicating that the accumulation of CCR2⁺ $\gamma\delta$ T cells is not a result of inflammation. These observations suggest that the CCL2-CCR2 axis is involved in the recruitment of $\gamma\delta$ T cells into joints.

***Il1rn*^{-/-} CD4⁺ cells induce *Ccl2* expression in joints.** CCR2⁺ $\gamma\delta$ T cells accumulated preferentially in arthritic joints, but not other organs, of *Il1rn*^{-/-} mice (Fig. 3d; Supplementary Fig. 4b). Consistent with this, expression of *Ccl2* (Fig. 3e), but not *Ccl20* (Supplementary Fig. 4c), was elevated in joints of *Il1rn*^{-/-} mice. Moreover, expression of *Ccl2*, but not *Cxcl16*, was specifically elevated in joints of *scid/scid* mice when *Il1rn*^{-/-} CD4⁺ T cells, but not $\gamma\delta$ T cells, were transferred (Fig. 3f,g). Thus, activated CD4⁺ T cells induce *Ccl2* expression in synovial resident cells and cause migration of CCR2⁺ $\gamma\delta$ T cells.

CCL2 recruits $\gamma\delta$ T cells to joints and induces arthritis. To determine whether elevated CCL2 expression in joints causes $\gamma\delta 17$ cell accumulation and disease development in *Il1rn*^{-/-} mice, we injected anti-CCL2 mAb into *Il17^{g/g}Il1rn*^{-/-} mice and analysed joint-infiltrating $\gamma\delta$ T cells. Anti-CCL2 mAb treatment significantly suppressed development of arthritis in *Il17^{g/g}Il1rn*^{-/-} mice (Fig. 4a; Supplementary Fig. 5), suggesting an important role for CCL2 in pathogenesis. The frequencies of

$\gamma\delta$ T cells and CCR2⁺ $\gamma\delta$ T cells were significantly reduced in joints of non-arthritis mice after mAb treatment (Fig. 4b-e), indicating that CCL2 is responsible for recruitment of CCR2⁺ $\gamma\delta$ T cells into joints. In mAb-treated non-arthritis mice, only a few GFP⁺ $\gamma\delta$ T cells were detected (Fig. 4d,f), suggesting a pathogenic role for joint-infiltrating $\gamma\delta 17$ cells in development of arthritis. Notably, when GFP⁺ $\gamma\delta$ T cells were gated, CCR2 was expressed at similar levels even in cells from anti-CCL2 mAb-treated non-arthritis mouse joints (Fig. 4g,h), suggesting that CCR2 expression in $\gamma\delta 17$ cells is required for the development of arthritis. On the other hand, the correlation between CXCR6 expression and disease development suggests that CXCR6 expression is a result of inflammation (Fig. 4i,j). These data indicate that CCR2⁺ $\gamma\delta$ T cells migrate into joints in response to high levels of CCL2 in *Il1rn*^{-/-} mouse joints, leading to development of arthritis.

IL-1Ra suppresses IL-1R expression on $\gamma\delta$ T cells. Next, we analysed the mechanism of IL-17 induction in $\gamma\delta$ T cells. Consistent with a previous report³⁵, IL-23 alone or IL-23 plus IL-1 β induced IL-17 production from magnetic-activated cell sorting (MACS)-purified splenic $\gamma\delta$ T cells (purity: 80%) (Fig. 5a). However, MACS-purified *Il1a*^{-/-}*Il1b*^{-/-} $\gamma\delta$ T cells or fluorescence-activated cell sorting (FACS)-purified WT $\gamma\delta$ T cells (purity: >99%) did not respond to IL-23 alone, and IL-1 β was required to induce IL-17 (Fig. 5a,b), suggesting that IL-1 β is essential for the IL-17 induction in $\gamma\delta$ T cells. Consistent with these results, the development of arthritis was completely suppressed in *Il1b*^{-/-}*Il1rn*^{-/-} mice (Supplementary Table 1).

We then analysed the effects of IL-1 β and IL-23 on expression of genes encoding transcription factors characteristic of Th17 cells, such as ROR γ t, ROR α , I κ B ζ and BATF. ROR γ t expression in $\gamma\delta$ T cells was drastically increased by addition of IL-1 β and IL-23 together, but only marginally by IL-1 β or IL-23 alone (Supplementary Fig. 6a). A similar effect was observed on ROR α expression. I κ B ζ and BATF were induced by IL-1 β alone, and the increase in their expression was synergistically enhanced by the addition of IL-23.


IL-23 remarkably increased IL-1R expression on $\gamma\delta$ T cells (Fig. 5c,d; Supplementary Fig. 6b). On the other hand, IL-23R was expressed on unstimulated $\gamma\delta$ T cells from WT, *Il1a*^{-/-}*b*^{-/-} and *Il1rn*^{-/-} mice, and its expression was not enhanced by the addition of IL-1 β (Fig. 5e). IL-1 β or IL-23 signalling alone was not sufficient for induction of IL-17; instead, synergistic

Figure 2 | Collaboration between CD4⁺ T cells and $\gamma\delta 17$ cells is important for the development of arthritis. (a,b) Suppression of arthritis development in *Il1rn*^{-/-} mice after treatment with anti- $\gamma\delta$ TCR (a) or anti-CD4 (b) mAb. Non-arthritis *Il1rn*^{-/-} mice at the age of 4 weeks were injected on days 0, 3, 7 and 10 (ages of 28, 31, 35 and 38 days) with anti- $\gamma\delta$ TCR mAb (■, n = 10) or isotype-matched hamster IgG (◇, n = 10) (a), or with anti-CD4 mAb (■, n = 10) or isotype-matched rat IgG (◇, n = 9) (b). *P < 0.05 versus control IgG, assessed by the χ^2 -test. (c,d) Non-arthritis *Il1rn*^{-/-} mice at the age of 20 days were injected with anti- $\gamma\delta$ TCR mAb or hamster IgG every 3 days (ages of 20, 23 and 26 days), and mice were killed at the age of 27 days. Representative haematoxylin and eosin-stained sections of ankle joints in non-treated WT mouse (left column, n = 6) and *Il1rn*^{-/-} mouse treated with control hamster IgG (middle column, n = 6) or α - $\gamma\delta$ TCR mAb (right column, n = 5) are shown. Synovial cell proliferation and inflammatory cell infiltration (arrows), bone erosion (arrowheads), fibrin clots (*) and pannus formation (★) in control *Il1rn*^{-/-} mouse (middle column) are indicated. Scale bars, 100 μ m. Tib, tibia; Tal, talus; Cal, calcaneum; Nav, navicular bone; Cun, cuneiform bone (c). (d) The means of histological scores are shown. *P < 0.05 versus *Il1rn*^{-/-} mice treated with hamster IgG. (e,f) Flow cytometry of LN cells from antibody-treated *Il1rn*^{-/-} mice. Cells were collected from *Il1rn*^{-/-} mice, 8 days after the first injection with anti- $\gamma\delta$ TCR mAb or hamster IgG-treated (e) or 11 days after the first treatment with anti-CD4 mAb or rat IgG (f) or from non-treated WT mice. Cells were stimulated with P/I for 5 h, and then stained for intracellular IL-17. Numbers refer to percentages in CD3e⁺ cells. (a-f) Data are representative of two independent experiments. (g,h) *scid/scid* mice at the age of 4 weeks were transferred with $\gamma\delta$ T cells from *Cd4*^{-/-}*Il1rn*^{-/-} mice (■, n = 8), CD4⁺ T cells from *Tcrd*^{-/-}*Il1rn*^{-/-} mice (◇, n = 9), Thy1.2⁺ T cells (whole-T cells) from *Il1rn*^{-/-} mice (▲, n = 8) or $\gamma\delta$ T cells from *Cd4*^{-/-}*Il1rn*^{-/-} mice plus CD4⁺ T cells from *Tcrd*^{-/-}*Il1rn*^{-/-} mice (×, n = 13). Incidence of arthritis is shown (g). Flow cytometry of the joint-infiltrating cells of *scid/scid* mice after 18 weeks of transfer, or age-matched and non-treated WT or *Il1rn*^{-/-} mice (h). Cells were stimulated with P/I for 5 h, and then stained for intracellular IL-17. Numbers refer to percentage in CD3e⁺ cells. Data are pooled from g or representative of (h) two independent experiments.

activation by IL-1 β and IL-23 was required (Fig. 5f; Supplementary Fig. 6c).

IL-1R expression induced by IL-23 was suppressed by the addition of exogenous IL-1Ra, and IL-1R expression was elevated

on *Il1rn*^{-/-} $\gamma\delta$ T cells (Fig. 5c,d), indicating that IL-23 and IL-1Ra reciprocally regulates IL-17 production by regulating IL-1R expression. Consistent with the elevation in IL-1R expression, IL-17 production in response to IL-23 and IL-1 β



stimulation was higher in *Il1rn*^{-/-} $\gamma\delta$ T cells than in WT $\gamma\delta$ T cells (Fig. 5f; Supplementary Fig. 6c). *Il1rn*^{-/-} $\gamma\delta$ T cells were less sensitive than WT $\gamma\delta$ T cells to inhibition of IL-1R expression by IL-1 β , and the maximum levels of IL-1R expression induced by IL-1 β and IL-23 on WT $\gamma\delta$ T cells were lower than those on *Il1rn*^{-/-} $\gamma\delta$ T cells (Fig. 5g). These results indicate that IL-1R expression is abnormally elevated in *Il1rn*^{-/-} $\gamma\delta$ T cells, resulting in higher IL-17 production in these cells (Fig. 5h). Thus, these results suggest that IL-1Ra is not merely an antagonist of IL-1R for the binding of IL-1 α and IL-1 β , but is also an important regulator of IL-1R expression on the cell surface.

V γ 6⁺ subset is the main IL-17 producer in *Il1rn*^{-/-} joints.

To identify the $\gamma\delta$ subset responsible for IL-17 production in *Il1rn*^{-/-} mouse joints, we examined the V γ subset in joint-infiltrating $\gamma\delta$ T cells. Proportions of V γ subsets were analysed using V γ TCR-specific antibodies except for V γ 6, because no anti-V γ 6 antibody was available. We found that $\gamma\delta$ T cells in the joints consisted of only two major populations: V γ 4⁺ and an 'other' population that was stained by none of antibodies against V γ 1, V γ 2, V γ 4, V γ 5 or V γ 7 TCR (Supplementary Fig. 7a). We assumed this 'other' V γ subset represented V γ 6⁺ cells, because only V γ 6⁺ cells are predicted to be unstained by all of these antibodies. $\gamma\delta$ T-cell subsets in LNs were heterogeneous, but only the V γ 4⁺ and putative V γ 6⁺ subsets were capable of producing IL-17 (Supplementary Fig. 7b).

Next, we analysed V γ subsets in $\gamma\delta$ 17 cells using *Il17^{g/g}Il1rn*^{-/-} mice. An average of 80% of joint-infiltrating GFP⁺ $\gamma\delta$ T cells were unstained by anti-V γ 4 and anti-V γ 1/2 antibodies (putative V γ 6⁺ cells), and only 20% of the population was V γ 4⁺ (Fig. 6a,b). Furthermore, mean fluorescence intensity of GFP was significantly higher in V γ 4⁻ cells than in V γ 4⁺ cells (Fig. 6c). Furthermore, the putative V γ 6⁺ cell population of IL-17⁺ $\gamma\delta$ T cells (63%) was larger than the V γ 4⁺ cell population (37%), as estimated by intracellular IL-17 staining (Supplementary Fig. 7c). The proportions of V γ 4⁺ cells in IL-17⁺ and in GFP⁺ $\gamma\delta$ T cells were slightly elevated following P/I stimulation (Supplementary Fig. 7c,d), probably because GFP as well as IL-17 expression is elevated after P/I stimulation or cell activation^{23,36,37}, and most V γ 6⁺ cells were already activated in *Il1rn*^{-/-} $\gamma\delta$ T cells *in vivo*.

We also examined V γ subset composition by measuring V γ messenger RNA (mRNA) expression using reverse transcription (RT)-PCR. V γ 6 mRNA was preferentially expressed in FACS-sorted GFP⁺ $\gamma\delta$ T cells in joints from *Il17^{g/g}Il1rn*^{-/-} mice, whereas other V γ mRNAs were also detected in LN $\gamma\delta$ T cells (Fig. 6d), consistent with the results of FACS analyses. V δ 1 mRNA was exclusively detected in GFP⁺ $\gamma\delta$ T cells in joints

(Fig. 6e), suggesting that these cells are of the canonical V γ 6/V δ 1 $\gamma\delta$ T-cell subset³⁸. Thus, V γ 6⁺ $\gamma\delta$ T cells are the major producers of IL-17 in joints of *Il1rn*^{-/-} mice.

Il1rn^{-/-} V γ 6⁺ cells highly express IL-1R intrinsically.

Because both V γ 6⁺ and V γ 4⁺ cells were present in joints with high levels of CCR2 expression on the surface (Fig. 7a), we asked why IL-17 was preferentially produced by V γ 6⁺ cells in *Il1rn*^{-/-} mouse joints. IL-1R expression on $\gamma\delta$ T cells was greatly elevated in both LNs and joints of *Il1rn*^{-/-} mice (Fig. 7b), and most GFP⁺ $\gamma\delta$ T cells in *Il17^{g/g}Il1rn*^{-/-} mouse joints were IL-1R⁺ (Fig. 7c). This IL-1R⁺ population was mostly V γ 6⁺ (Fig. 7d). The mean fluorescence intensity of IL-1R was also significantly higher in V γ 6⁺ cells than in V γ 4⁺ cells (Fig. 7e). This elevated IL-1R expression was already present in newborn thymic $\gamma\delta$ T cells from *Il1rn*^{-/-} mice (Fig. 7f), and was detected on V γ 6⁺ (V γ 4⁻, V γ 1/2⁻, V γ 5⁻ and V γ 7⁻) cells (Fig. 7g). Notably, IL-1R expression on $\gamma\delta$ T cells in newborn thymus correlated with CCR2 expression (Fig. 7h). IL-1 β mRNA expression was observed even in joints of WT mice, and was greatly elevated in joints of *Il1rn*^{-/-} mice (Fig. 7i). IL-23 mRNA expression in joints was also significantly higher in *Il1rn*^{-/-} mice than in WT mice (Fig. 7j). These observations suggest that joint-infiltrating CCR2⁺V γ 6⁺ *Il1rn*^{-/-} $\gamma\delta$ T cells, which intrinsically express high levels of IL-1R, preferentially produce IL-17 in response to IL-1 β and IL-23, resulting in development of arthritis.

Discussion

Here we showed that both CD4⁺ T cells and $\gamma\delta$ 17 cells are important for development of arthritis in *Il1rn*^{-/-} mice, because antibody-mediated depletion of either $\gamma\delta$ T or CD4⁺ T cells suppressed the development of arthritis. Furthermore, upon adoptive transfer, only a mixture of $\gamma\delta$ T and CD4⁺ T cells induced arthritis in *scid/scid* mice. $\gamma\delta$ 17 cells localized in joints of *scid/scid* mice when $\gamma\delta$ T cells were transferred along with CD4⁺ T cells, whereas $\gamma\delta$ T cells were not detected in joints when $\gamma\delta$ T cells were transferred alone. These observations suggest that $\gamma\delta$ T cells alone cannot distribute into joints, and CD4⁺ T cells are required for the localization of $\gamma\delta$ T cells.

Anti- $\gamma\delta$ TCR mAb injection significantly suppressed not only the incidence of arthritis but also the histological severity score, indicating that $\gamma\delta$ T cells are involved in development of arthritis. It was recently reported that treatment with anti- $\gamma\delta$ TCR mAb results in internalization of $\gamma\delta$ TCR rather than $\gamma\delta$ T-cell depletion³⁹. In our hands, however, the $\gamma\delta$ T-cell population was greatly diminished by treatment with this antibody, indicating that the $\gamma\delta$ 17 population is actually depleted by this antibody.

Figure 3 | CCR2⁺ $\gamma\delta$ 17 cells predominantly accumulate in *Il1rn*^{-/-} mouse joints. (a) Expression of chemokine receptors and GFP on joint-infiltrating cells from WT ($n=3$, pool of two mice each), *Il1rn*^{-/-} ($n=3$) and *Il17^{g/g}Il1rn*^{-/-} ($n=3$) mice at 16 weeks of age. Numbers represent percentages in CD3 ϵ ⁺ $\gamma\delta$ TCR⁺ cells. Numbers in parentheses represent percentages in GFP⁺ CD3 ϵ ⁺ $\gamma\delta$ TCR⁺ cells. (b) Quantitative PCR (qPCR) analysis of the transcripts of CCL2 (*Ccl2*), CXCL16 (*Cxcl16*), CCL20 (*Ccl20*), CCL5 (*Ccl5*) and CXCL12 (*Cxcl12*) in joints of WT ($n=4$) or *Il1rn*^{-/-} ($n=4$) mice. Values are shown relative to those in WT mice. * $P<0.05$; ** $P<0.01$; NS, not significant (versus WT mice) (unpaired Student's *t*-test). (c) Flow cytometry of the joint-infiltrating $\gamma\delta$ T cells from WT, arthritic *Il1rn*^{-/-} and non-arthritic *Il17a^{-/-}Il1rn*^{-/-} mice. Numbers represent percentages in CD3 ϵ ⁺ $\gamma\delta$ TCR⁺ cells. (d) Contents of CCR2⁺ cells in CD3 ϵ ⁺ $\gamma\delta$ TCR⁺ cells in the thymus (Thy), mesenteric LNs (mLN), draining LNs (dLN), spleens (SP), blood and joints of WT ($n=3$) and *Il1rn*^{-/-} ($n=3$) mice at the age of 14 weeks. * $P<0.05$; NS (versus WT mice) (unpaired Student's *t*-test). (e) Levels of CCL2 mRNA in whole tissue were measured by qPCR in various tissues from WT ($n=5$) and *Il1rn*^{-/-} ($n=5$) mice at the age of 8 weeks. Values are shown relative to those in WT mouse thymus total RNA. ** $P<0.01$; NS (versus WT mice) (unpaired Student's *t*-test). (f,g) qPCR analysis of mRNA for CCL2 (f) and CXCL16 (g) in joints of *scid/scid* mice after 24 weeks of transfer of CD4⁺ T cells ($n=4$) or $\gamma\delta$ T cells ($n=3$), or age-matched *scid/scid* mice ($n=3$). Values are shown relative to those in control *scid/scid* mice. * $P<0.05$; NS versus control *scid/scid* mice (unpaired Student's *t*-test). mRNA expression was normalized to that of *Gapdh* (b,e-g). All data except a and c show mean \pm s.e.m. Data are representative of two (c-g) or >3 (a,b) independent experiments.




Figure 4 | CCL2 is essential for $\gamma\delta$ T-cell infiltration into joints and arthritis development. (a) Incidence (left) and severity scores of arthritis in affected mice (right) in *Il17^{g/g}Il1m*^{-/-} mice after anti-CCL2 mAb treatment. Non-arthritic *Il17^{g/g}Il1m*^{-/-} mice at the age of 21 days were injected with anti-CCL2 mAb (■, $n=9$) or PBS (◇, $n=11$) every 3 days (ages of 21, 24, 27 and 30 days). * $P<0.05$ versus treatment with PBS, assessed by the χ^2 -test. Data represent a pool of two independent experiments. (b–j) Flow cytometry of joint-infiltrating cells from *Il17^{g/g}Il1m*^{-/-} mice 11 days after the first injection. Cells were collected from PBS-treated arthritic (A) mice ($n=5$) and anti-CCL2 mAb-treated non-arthritic (NA) ($n=3$) or arthritic (A) ($n=2$) mice. Age-matched and non-treated WT ($n=3$) and arthritic *Il1m*^{-/-} ($n=3$) mice were used as controls. Numbers refer to percentage in CD3 ϵ ⁺ cells (b), in CD3 ϵ ⁺ $\gamma\delta$ TCR⁺ cells (d) or in GFP⁺CD3 ϵ ⁺ $\gamma\delta$ TCR⁺ cells (g,i). The average proportions of $\gamma\delta$ T cells in CD3 ϵ ⁺ cells (c), CCR2⁺ (e) or GFP⁺ (f) cells among $\gamma\delta$ TCR⁺ cells, or CCR2⁺ (h) or CXCR6⁺ (j) cells among GFP⁺ $\gamma\delta$ TCR⁺ cells, are shown. * $P<0.05$; *** $P<0.001$; NS, not significant (versus *Il17^{g/g}Il1m*^{-/-} mice treated with PBS) (unpaired Student's t -test). Data show mean \pm s.e.m. Data (b–j) are representative of two independent experiments.




Figure 5 | IL-23 induces expression of IL-1R on the surface of γδ T cells, whereas IL-1Ra suppresses its expression. (a,b) Concentrations of IL-17 in culture supernatants of magnetic-activated cell sorting (MACS)-purified (about 80%) (a) or FACS-purified (>99% pure) (b) splenic γδ T cells from pools of 16 WT mice (a,b) or 16 *Il1a*^{-/-}*b*^{-/-} mice (b) stimulated for 3 days with medium only, IL-1β, IL-23 or IL-23 plus IL-1β, without γδ TCR stimulation. IL-17 was detected by enzyme-linked immunosorbent assay (ELISA). ****P*<0.001 versus WT mice (a); ****P*<0.001 versus medium only (b) (unpaired Student's *t*-test). (c-h) FACS-purified γδ T cells from pooled spleens of WT, *Il1r1*^{-/-}, *Il1a*^{-/-}*b*^{-/-} or *Il1m*^{-/-} mice (11-16 mice each) were stimulated for 3 days with medium only, IL-1β, IL-23, IL-23 plus IL-1β or IL-23 plus IL-1Ra. Flow cytometry of γδ T cells stained for surface IL-1R (c) and IL-23R (e) are shown. Quantification of IL-1R⁺ γδ T cells is indicated in d and g. Concentrations of IL-17 in culture supernatants were determined by ELISA (f,h). **P*<0.05; ***P*<0.01; ****P*<0.001 (versus WT mice) (unpaired Student's *t*-test). Numbers in parentheses indicate the concentration of cytokines (ng ml⁻¹). Representative data (c,e) and mean ± s.e.m. (a,b,d,f,g,h) of triplicate cultures are shown. All data are representative of two or three independent experiments.




Figure 6 | The Vγ6⁺ γδ T-cell subset is the major source of IL-17 in *Il1rn*^{-/-} mouse joints. (a–c) GFP expression in joint-infiltrating γδ T cells from WT, *Il1rn*^{-/-} and *Il17^{g/g}Il1rn*^{-/-} mice at 16 weeks of age. Numbers refer to percentage in CD3ε⁺ γδ TCR⁺ cells (a, left). Quantification of GFP⁺ Vγ subsets in CD3ε⁺ γδ TCR⁺ cells is shown (a, right). Numbers in the pie chart show the percentages of the indicated Vγ subset among GFP⁺ γδ TCR⁺ CD3ε⁺ cells, and represent the average of three mice (b). Mean fluorescence intensity of GFP in Vγ6⁺ (Vγ4⁻) or Vγ4⁺ cells is shown in c. **P* < 0.05, ****P* < 0.001 (versus Vγ4⁺ cells) (unpaired Student's *t*-test). Data show the mean ± s.e.m. of three mice. (d,e) Vγ (d) and Vδ (e) gene expression in γδ T cells. CD3ε⁺ γδ TCR⁺ (whole γδ T) cells from LNs (first row) or from joints (second row) of *Il1rn*^{-/-} mice and GFP⁺ CD3ε⁺ γδ TCR⁺ (GFP⁺ γδ T) (third row) or CD3ε⁺ γδ TCR⁻ (non-γδ T) (fourth row) cells from joints of *Il17^{g/g}Il1rn*^{-/-} mice, were sorted on a FACSaria, and Vγ and Vδ gene expression was analysed by RT-PCR.


Importantly, the γδ17 population was significantly reduced without a compensatory increase in IL-17⁺ cells in the γδ TCR⁻ population, suggesting that the reduction of γδ17 cells was not the result of replacement of γδ17 cells by hypothetical 'invisible γδ T cells'³⁹. A reduction in IL-17 production following anti-γδ TCR mAb treatment was also reported by another group⁴⁰. Although we examined the effect of γδ T-cell deficiency using *Tcrd*^{-/-} mice, we could not find any effect of this gene

mutation on the development of arthritis. Interestingly, we found IL-17-producing CD4⁻CD8⁻γδTCR⁻ T cells were increased in the joints of *Tcrd*^{-/-} *Il1rn*^{-/-} mice, suggesting that these cells may substitute for the deficiency of γδ T cells to produce IL-17. We are now further characterizing these cells.

In *Il17^{g/g}Il1rn*^{-/-} mice, most of the joint-infiltrating γδ17 cells expressed CCR2, in association with elevated expression of CCL2 in the joints. Antibody-mediated blockade of CCL2 in

Il17^g/Il1rn^{-/-} mice reduced infiltration of CCR2⁺ GFP⁺ $\gamma\delta$ T cells in joints and suppressed the development of arthritis, suggesting that CCR2⁺ $\gamma\delta$ 17 cell accumulation in the joints, which is critical for the development of arthritis, is mediated by the CCL2-CCR2 interaction. The CCL2-CCR2 axis-mediated $\gamma\delta$ T-cell migration has also been reported as a protection

mechanism against tumours⁴¹. Regarding CCR2 expression, a subset of $\gamma\delta$ T cells acquires effector functions and expresses IL-17 during intrathymic development^{15,16}, and these thymic $\gamma\delta$ 17 subsets already express some chemokine receptors such as CCR2, CCR6 and CXCR6 (refs 17,18). In contrast to $\gamma\delta$ 17 cells in *Il1rn^{-/-}* mice, pathogenic Th17 cells expressing CCR6 are



recruited to inflammatory sites, such as joints of SKG mice⁴² and the central nervous system of an experimental autoimmune encephalomyelitis model^{43,44}, via the CCR6–CCL20 interaction. Because we did not observe a significant increase in CCL20 expression in joints of *Il1rn*^{-/-} mice relative to WT mice, the CCR6–CCL20 axis may not be important for the $\gamma\delta$ 17 cell migration into joints, even though >50% of joint-infiltrating GFP⁺ $\gamma\delta$ T cells in *Il17^{g/g}Il1rn*^{-/-} mice also expressed CCR6. CXCR6 was also expressed on joint-infiltrating $\gamma\delta$ 17 cells in *Il17^{g/g}Il1rn*^{-/-} mice. However, because this chemokine receptor was not expressed on joint-infiltrating $\gamma\delta$ T cells in non-arthritic *Il17a*^{-/-}*Il1rn*^{-/-} mice, CXCR6 may not be involved in recruitment of $\gamma\delta$ T cells, at least under non-inflammatory conditions.

Importantly, *Il1rn*^{-/-} CD4⁺ T-cell transfer induced CCL2 expression in joints of recipient *scid/scid* mice, suggesting that CD4⁺ T cells directed the migration of CCR2⁺ $\gamma\delta$ T cells. We rarely detected Th17 cells in the inflamed joints, indicating that IL-17 production from CD4⁺ T cells is not required for pathogenesis in *Il1rn*^{-/-} mice, at least for induction of local inflammation. Therefore, our results show that CD4⁺ T cells direct the tissue specificity of inflammation, and $\gamma\delta$ 17 cell-derived IL-17 elicits local inflammation and arthritis in *Il1rn*^{-/-} mice. Although the importance of $\gamma\delta$ 17 cells in the development of arthritis in the collagen-induced arthritis model was already suggested by Ito *et al.*¹³, the mechanism how these $\gamma\delta$ T cells are distributed to joints has not been elucidated. In this report, we first clarified the mechanism how $\gamma\delta$ T cells are recruited to the inflammatory site. Actually, when $\gamma\delta$ T cells alone were transferred to *scid/scid* mice, development of inflammation was observed in other organs, such as the colon and dermis, suggesting the importance of CD4⁺ T cells for the tissue-specific distribution of $\gamma\delta$ T cells. We showed previously that T cells from *Il1rn*^{-/-} mice are hyper-reactive due to the overexpression of CD40L and OX40 on the cell surface, and consequently lose tolerance against self-antigens^{32,33}.

A combination of IL-1 β and IL-23, but not IL-1 β or IL-23 alone, induces IL-17 in $\gamma\delta$ T cells without TCR engagement^{11,13,20,35}. In this report, we showed that IL-23 is required for the induction of IL-1R on $\gamma\delta$ T cells, and IL-1 β is essential for the induction of IL-17. However, IL-1 β alone could not induce IL-17 production in *Il1rn*^{-/-} $\gamma\delta$ T cells, even though these cells expressed IL-1R, consistent with a report that IL-1 β alone does not induce IL-17 in peritoneum- and lung-derived $\gamma\delta$ T cells expressing high levels of IL-1R²⁰. These observations suggest that IL-23 plays roles, other than upregulating IL-1R, in the induction of IL-17 expression in $\gamma\delta$ T cells. In this context, expression of

IL-17 signature transcription factors such as *Rorc*, *Rora*, *Nfkbz* and *Batf* was increased by IL-1 β and IL-23 together.

Moreover, IL-23-induced IL-1R expression on $\gamma\delta$ T cells was suppressed by exogenous IL-1Ra, suggesting that IL-1Ra not only antagonizes IL-1 α and IL-1 β for IL-1R binding, but also regulates cell-surface expression of IL-1R. Consistent with this, IL-1R expression on $\gamma\delta$ T cells was intrinsically augmented in *Il1rn*^{-/-} mice, making IL-17 production by these *Il1rn*^{-/-} $\gamma\delta$ T cells hyper-sensitive to the action of IL-1 β and IL-23. *Il1rn*^{-/-} $\gamma\delta$ T cells induced arthritis upon transfer into *scid/scid* mice. Because IL-1R expression levels induced by IL-1 β and IL-23 were much higher on *Il1rn*^{-/-} $\gamma\delta$ T cells than on WT $\gamma\delta$ T cells, endogenous low levels of IL-1Ra in the recipients could not suppress IL-1R expression on *Il1rn*^{-/-} $\gamma\delta$ T cells. Thus, the strict control of IL-1R expression by IL-1Ra is important for regulation of IL-17 production in $\gamma\delta$ T cells.

Children with homozygous point mutations or deletions of the *IL1RN* gene develop life-threatening severe inflammatory diseases with prominent involvement of bone and skin after birth^{45,46}. The population of IL-17-producing cells is markedly elevated in inflamed skin, and a large amount of IL-1 β is produced in mononuclear cells, as in *Il1rn*^{-/-} mice^{45,46}. Although the pathogenic roles of IL-17 in local inflammation in these patients remain unknown, $\gamma\delta$ T cells may be involved in the inflammation. Our observation that exogenous IL-1Ra suppresses IL-1R expression on $\gamma\delta$ T cells may partly explain why treatment with the recombinant IL-1Ra anakinra completely resolves symptoms in these affected children^{45,46}.

Using *Il17^{g/g}* mice, we showed that IL-17 was preferentially produced in V γ 6⁺ $\gamma\delta$ T cells of *Il1rn*^{-/-} mouse joints, although both CCR2⁺ V γ 6⁺ cells and CCR2⁺ V γ 4⁺ cells were localized in the joints. These results suggest that IL-17 production capacity differs in these two $\gamma\delta$ T-cell subsets. We found that IL-1R expression in joints is much higher on V γ 6⁺ cells than on V γ 4⁺ cells, explaining why IL-17 is induced preferentially in V γ 6⁺ cells in *Il1rn*^{-/-} mice. This elevated IL-1R expression was observed even in newborn thymus. Notably in this regard, different transcription factors are required for the development of V γ 6⁺ and V γ 4⁺ cell subset²⁴, and V γ 6⁺ cells acquire the innate capacity to produce IL-17 in the embryonic thymus⁴⁷. Therefore, during embryonic development, the level of IL-1R is probably intrinsically higher in V γ 6⁺ $\gamma\delta$ T cells than in V γ 4⁺ cells.

V γ 6⁺ $\gamma\delta$ 17 cells are thought to be a tissue-resident, long-lived, self-renewing population⁴⁷. Although our CCL2 inhibition and transfer experiments suggested that LN V γ 6⁺ $\gamma\delta$ T cells capable of producing IL-17 were recruited to joints (where they induced

Figure 7 | IL-17-producing *Il1rn*^{-/-} V γ 6⁺ $\gamma\delta$ T cells intrinsically express high levels of IL-1R. (a) Flow cytometry of joint-infiltrating cells from *Il1rn*^{-/-} mice ($n = 3$) for the expression of CCR2 and V γ 4. Numbers refer to percentage in CD3 ϵ ⁺ $\gamma\delta$ TCR⁺ cells. Numbers in parentheses represent percentage in V γ 6⁺ (V γ 4⁻) or V γ 4⁺ cells (left). Quantification of CCR2⁺ cells in V γ 6⁺ (V γ 4⁻) or V γ 4⁺ cells is shown (right). NS, not significant versus V γ 4⁺ cells (unpaired Student's *t*-test). **(b)** Flow cytometry of IL-1R expression in LNs and joint-infiltrating cells from WT or *Il1rn*^{-/-} mice. Numbers refer to percentages in CD3 ϵ ⁺ cells. Numbers in parenthesis represent percentages in CD3 ϵ ⁺ $\gamma\delta$ TCR⁺ cells (left). The percentage of IL-1R⁺ cells among CD3 ϵ ⁺ $\gamma\delta$ TCR⁺ cells is indicated on the right. ** $P < 0.01$; *** $P < 0.001$ (versus WT mice) (unpaired Student's *t*-test). **(c)** FACS analysis of IL-1R and GFP expression on joint-infiltrating CD3 ϵ ⁺ $\gamma\delta$ TCR⁺ cells from WT, *Il1rn*^{-/-} and *Il17^{g/g}Il1rn*^{-/-} mice. Numbers in parentheses represent percentage in GFP⁺ $\gamma\delta$ TCR⁺ cells. **(d,e)** Flow cytometry of IL-1R expression in joint-infiltrating cells from *Il1rn*^{-/-} mice. Numbers refer to percentages in CD3 ϵ ⁺ $\gamma\delta$ TCR⁺ cells. Numbers in parentheses represent percentages in V γ 6⁺ (V γ 4⁻) or V γ 4⁺ cells (**d**, left). Quantification of IL-1R⁺ cells (**d**, right) and mean fluorescence intensity of IL-1R in V γ 6⁺ or V γ 4⁺ cells are indicated (**e**). * $P < 0.05$; *** $P < 0.001$ (versus V γ 4⁺ cells) (unpaired Student's *t*-test). **(f-h)** Flow cytometry of IL-1R expression in $\gamma\delta$ T cells in newborn thymus (within the first day after birth) from WT and *Il1rn*^{-/-} mice. IL-1R expression on $\gamma\delta$ T cells (**f**) and on different V γ subsets (**g**), and CCR2 and IL-1R expression on $\gamma\delta$ T cells (**h**) are shown. Numbers refer to percentage in CD3 ϵ ⁺ cells (**f,h**, top), or in CD3 ϵ ⁺ $\gamma\delta$ TCR⁺ cells (**g,h**, bottom). Numbers in parentheses represent percentage in $\gamma\delta$ TCR⁺ cells (**f,h**, top) or in IL-1R⁺ $\gamma\delta$ TCR⁺ cells (**h**, bottom). **(i,j)** qPCR analysis of the transcripts for IL-1 β (*Il1b*) (**i**) or IL-23 (*Il23p19*) (**j**) in cells from joints, draining LNs (dLN), mesenteric LNs (mLN) and spleens (SP) from WT or *Il1rn*^{-/-} mice. mRNA levels were normalized against *Gapdh*, and values are shown relative to mLN of WT mice. * $P < 0.05$; NS, not significant (versus WT mice) (unpaired Student's *t*-test). Data show mean \pm s.e.m. All data are representative of two or three independent experiments.

arthritis) via CCR2–CCL2 interaction, development of arthritis was not completely suppressed by treatment with anti-CCL2 mAb. Therefore, we could not completely exclude the possibility that these tissue-resident $V\gamma 6^+ \gamma\delta 17$ cells are also involved in development of arthritis.

$\gamma\delta 17$ cells have been observed in the peripheral tissues of several inflammatory diseases^{11,48,49}, and the involvement of $\gamma\delta 17$ cells is suggested in some autoimmune models^{5,7,11–13,24}. Innate immune cell-derived IL-17 has also been implicated in the pathogenesis of intestinal inflammatory diseases⁴. However, the mechanisms by which these IL-17-producing $\gamma\delta$ T cells or innate immune cells are recruited to the inflammatory sites remain largely unknown. In this report, we showed that $CD4^+$ T cells determine the tissue specificity, and $CD4^+$ T-cell-induced CCL2 recruits $CCR2^+ V\gamma 6^+ \gamma\delta$ T cells to the joints. Therefore, our model may explain how IL-17-producing $\gamma\delta$ T cells or innate immune cells infiltrate local tissues and induce inflammation in other autoimmune diseases⁵⁰. Taken together, our findings provide important insight into the immunological mechanisms driving tissue-specific autoimmune diseases, and may provide a clue for the development of novel therapies.

Methods

Mice. All mice, except C.B.-17 *scid/scid* mice (CLEA Japan, Tokyo, Japan), were on the BALB/cA genetic background. *Il1rn*^{-/-} mice were produced as described^{31,51}, and backcrossed to BALB/cA (CLEA Japan) for nine generations. *Il17^{g/g}* mice were generated as shown in Supplementary Fig. 3a, and backcrossed to BALB/cA mice for eight generations. *Il17^{g/g}Il1rn*^{-/-} mice were produced by crossing *Il17^{g/g}* mice with *Il1rn*^{-/-} mice. *Tcrd*^{-/-} mice were generated by Itoharu *et al.*⁵². *Cd4*^{-/-} mice were a gift from Dr. Fujii (RIKEN, Kanagawa, Japan). These mice were backcrossed to the BALB/cA mice for eight generations. *Il1b*^{-/-} and *Il1a*^{-/-b} mice were produced as described⁵¹. *Il1r1*^{-/-} mice were obtained from Immunex Corporation. *Il17a*^{-/-Il1rn}^{-/-} mice were generated as described³². Sex- and age-matched mice, usually at 8–12 weeks of age, were used for each experiment. In some experiments, newborn mice and younger (3–4 weeks old) or elder (24 weeks old) mice were also used as described in the figure legends. All mice were kept under specific pathogen-free conditions in environmentally controlled clean rooms at the Center for Experimental Medicine and Systems Biology, The Institute of Medical Science, The University of Tokyo, and Institute for Biomedical Sciences, Tokyo University of Science. All experiments were approved by the institutional ethical committees for animal experiments and the committees for gene manipulation experiments.

Clinical assessment of arthritis. Development of arthritis was monitored by macroscopic evaluation as described previously³¹. Briefly, each paw was graded as follows: 0, no change; 1, mild swelling; 2, obvious swelling; 3, severe swelling and ankylosis changes (maximum 12 points for individual mice).

Histological assessment of arthritis. Whole-animal joints were fixed in 10% formalin in 0.001 M phosphate buffer (pH 7.2), decalcified in 10% EDTA and embedded in paraffin. Serial sections (4 μ m) were stained with haematoxylin and eosin. Each joint was graded on a scale of 0–3: 0, normal; 1, thickening and proliferation of the synovial lining, with slight inflammatory cell infiltration; 2, grade-1 changes plus extensive synoviocyte proliferation and severe inflammatory cell infiltration; 3, grade-2 changes plus pannus formation and bone erosion. Histological score of the ankle joint was estimated from the sum grade of both ankle joints (maximum 6 points for individual mice).

Immunofluorescence staining. For immunohistochemical analysis, hindlimbs were embedded in Super Cryoembedding Medium (Leica Microsystems Japan, Tokyo, Japan), and frozen sections (5 μ m) were generated. The sections were fixed in cold acetone for 5 min and blocked with 4% bovine serum albumin (Sigma, St Louis, MO, USA) and 5% goat serum in PBS. Antibodies used were as follows: 2 μ g ml⁻¹ rabbit anti-mouse IL-17A polyclonal Ab (Abcam, ab9565-100, Cambridge, UK), 5 μ g ml⁻¹ hamster anti-mouse $\gamma\delta$ TCR mAb (GL3, BD Pharmingen, San Diego, CA, USA), 2.5 μ g ml⁻¹ Alexa Fluor 488-goat anti-rabbit-IgG (Life Technologies, Carlsbad, CA, USA), and 2.5 μ g ml⁻¹ Cy3-goat anti-hamster IgG (Jackson ImmunoResearch, West Grove, PA, USA). Nuclei were stained with 0.5 μ g ml⁻¹ 4,6-diamidino-2-phenylindole. The slides were visualized on a fluorescence microscope (Keyence, Osaka, Japan) and on an Olympus FV1000 Confocal Microscope, operated by the FluoView software (Olympus, Tokyo, Japan).

Isolation of joint-infiltrating cells. Ankle joints were cut out and digested with 2.4 mg ml⁻¹ hyaluronidase (Sigma), 1 mg ml⁻¹ collagenase (Sigma) in RPMI 1640 (Sigma) plus 10% foetal bovine serum (FBS) for 1 h at 37 °C. The cells were filtered through a cell strainer with a 70- μ m nylon mesh (Becton Dickinson, Franklin Lakes, NJ, USA) and washed with RPMI 1640 plus 10% FBS.

Cell transfer. Purified $\gamma\delta$ T cells (8×10^5 cells), $CD4^+$ T cells (2×10^7 cells) or $Thy1.2^+$ T cells (2×10^7 cells) from LN (axillary, brachial, inguinal, popliteal and cervical) cells were suspended in 200 μ l sterile PBS and then injected intravascularly into *scid/scid* mice.

Antibody treatment. Non-arthritic *Il1rn*^{-/-} mice (4 weeks old) received twice weekly intraperitoneal injections of 400 μ g anti-TCR $\gamma\delta$ mAb (UC7-13D5 (ref. 53), purified from hybridoma culture supernatant) or 250 μ g anti-CD4 mAb (GK1.5, purified from hybridoma culture supernatant provided by Dr Nariuchi (The University of Tokyo)). Isotype-matched Armenian hamster IgG (400 μ g per mouse; Innovate Research, Noida, India) was used as a control for UC7-13D5. Isotype-matched rat IgG (250 μ g per mouse; Thermo, Waltham, MA, USA) was used as a control for GK1.5. A dose of 200 μ g anti-CCL2 mAb (123616; R&D Systems, Minneapolis, MN, USA), 200 μ g rat IgG2b isotype control (Fitzgerald, Sudbury Road Acton, MA, USA) or PBS alone was intraperitoneally injected into non-arthritic *Il17^{g/g}Il1rn*^{-/-} mice at ages of 21, 24, 27 and 30 days.

Cell isolation. To examine LN cells by flow cytometry, axillary, brachial, inguinal and popliteal LNs were collected. To obtain FACS-purified $\gamma\delta$ T cells, LN and/or spleen cells were first incubated with biotin-conjugated anti-mouse $\gamma\delta$ TCR mAb (GL3; #13-5711, eBioscience, San Diego, CA, USA) (1/100 diluted) and then with MicroBeads conjugated to anti-biotin mAb (#130-090-485, Miltenyi Biotec, Bergisch Gladbach, Germany) (1/8 diluted). Labelled cells were positively selected twice using autoMACS (Miltenyi Biotec). To purify whole-T cells and $CD4^+$ T cells, LN and/or spleen cells were stained with MicroBeads conjugated to anti-mouse $Thy1.2$ mAb (#130-049-101, Miltenyi Biotec) and $CD4$ mAb (#130-049-201, Miltenyi Biotec), respectively (1/8 diluted), and separated using an autoMACS. MACS-purified $\gamma\delta$ T ($\gamma\delta$ TCR⁺ CD3 ϵ ⁺) cells and GFP⁺ $\gamma\delta$ T (GFP⁺ $\gamma\delta$ TCR⁺ CD3 ϵ ⁺) cells were further purified using a FACSARIA (Becton Dickinson). FITC-anti-mouse $\gamma\delta$ TCR mAb (UC7-13D5; eBioscience), PE-anti-mouse $\gamma\delta$ TCR mAb (GL3; BioLegend, San Diego, CA, USA), and APC/Cy7-anti-mouse CD3 ϵ mAb (145-2C11; BioLegend) (1/100 diluted) were used for labelling. The efficiency of these cell purifications was determined by flow cytometry.

Cell culture. For analysis of flow cytometry and concentrations of IL-17 in culture supernatants, cells from LNs and/or spleens were cultured for 72 h in RPMI 1640 containing 10% FBS with or without IL-1 β (10 ng ml⁻¹) (PeproTech, Rocky Hill, NJ, USA), IL-23 (10 ng ml⁻¹) (R&D Systems), IL-1Ra (200 ng ml⁻¹) (R&D Systems) or cytokines at the indicated concentrations (ng ml⁻¹) in the absence of $\gamma\delta$ TCR stimulation. For analysis of mRNA expression, $\gamma\delta^+$ CD3 ϵ^+ or $\gamma\delta^-$ CD3 ϵ^+ T cells from spleens were purified using a FACSARIA (Becton Dickinson) and cultured for 48 h with or without IL-1 β (10 ng ml⁻¹) and IL-23 (10 ng ml⁻¹).

Flow cytometry. Intracellular cytokine staining was performed as described previously⁵⁴ after stimulation with 50 ng ml⁻¹ phorbol myristate acetate (Sigma), 500 ng ml⁻¹ ionomycin (Sigma) and 2 μ M monensin (Sigma) for 5 h. For staining of cell-surface molecules, cells were first treated with anti-mouse CD16/CD32 mAbs (2.4G2, purified from hybridoma culture supernatant) in staining buffer (HBSS containing 2% FCS and 0.1% sodium azide) to block FcR binding, and then stained with antibodies (1/100 diluted). A list of all antibodies used in the study is shown in Supplementary Table 2. For intracellular cytokine staining, cells were fixed with 4% paraformaldehyde and treated with a permeabilization buffer (0.1% saponin (Sigma) in staining buffer), and then incubated with antibodies (1/100 diluted) against intracellular cytokine. 7-Aminoactinomycin D (Sigma) was used to stain dead cells. Cells were analysed on a FACSCanto II system (Becton Dickinson), and data were analysed with FlowJo software (Tree Star).

Measurement of cytokine. Concentrations of IL-17 in culture supernatants were determined by enzyme-linked immunosorbent assay using commercially available kits (Ready-Set-Go, eBioscience).

RT-PCR and real-time PCR. To prepare single cells for RNA expression analysis, draining LN, mesenteric LN and spleen were digested with 200 U ml⁻¹ collagenase (Sigma) for 30 min. Joints were digested with 200 U ml⁻¹ collagenase (Sigma) and 2.4 mg ml⁻¹ hyaluronidase (Sigma). Next, these cells were filtered through a cell strainer with a 70- μ m nylon mesh (Becton Dickinson). RNA from joint-, dLN-, mLN- and spleen-derived single cells and RNA from cultured cells were extracted using the GenElute Mammalian Total RNA Miniprep Kit (Sigma). Tissue total RNA from joint, draining LN, spleen, thymus, colon, lung and liver was extracted using the Sepasol reagent (Nacalai Tesque, Kyoto, Japan). All RNA were denatured in the presence of an oligo dT primer and reverse transcribed using the High

Capacity cDNA Reverse Transcription Kit (Applied Biosystems, Foster City, CA, USA). Quantitative real-time PCR was performed with a SYBR Green qPCR kit (Invitrogen, Carlsbad, CA, USA) and an iCycler system (Bio-Rad, Hercules, CA, USA) with the sets of primers described in Supplementary Table 3.

Statistics. Unless otherwise specified, all results are shown as means and s.e.m. Unpaired Student's *t*-tests were used to statistically analyse all results, except that Mann-Whitney's *U*-tests were used to evaluate disease severity and χ^2 -tests were used to evaluate disease incidence. Differences were considered significant at $P < 0.05$.

References

- Iwakura, Y., Ishigame, H., Saijo, S. & Nakae, S. Functional specialization of interleukin-17 family members. *Immunity* **34**, 149–162 (2011).
- Kellner, H. Targeting interleukin-17 in patients with active rheumatoid arthritis: rationale and clinical potential. *Ther. Adv. Musculoskelet. Dis.* **5**, 141–152 (2013).
- Cua, D. J. & Tato, C. M. Innate IL-17-producing cells: the sentinels of the immune system. *Nat. Rev. Immunol.* **10**, 479–489 (2010).
- Akitsu, A., Kakuta, S., Saijo, S. & Iwakura, Y. Rag2-deficient IL-1 receptor antagonist-deficient mice are a novel colitis model in which innate lymphoid cell-derived IL-17 is involved in the pathogenesis. *Exp. Anim.* **63**, 235–246 (2014).
- Sutton, C. E. *et al.* Interleukin-1 and IL-23 induce innate IL-17 production from gammadelta T cells, amplifying Th17 responses and autoimmunity. *Immunity* **31**, 331–341 (2009).
- Petermann, F. *et al.* $\gamma\delta$ T cells enhance autoimmunity by restraining regulatory T cell responses via an interleukin-23-dependent mechanism. *Immunity* **33**, 351–363 (2010).
- Roark, C. L. *et al.* Exacerbation of collagen-induced arthritis by oligoclonal, IL-17-producing gamma delta T cells. *J. Immunol.* **179**, 5576–5583 (2007).
- Park, S.-G. *et al.* T regulatory cells maintain intestinal homeostasis by suppressing $\gamma\delta$ T cells. *Immunity* **33**, 791–803 (2010).
- Gray, E. E., Suzuki, K. & Cyster, J. G. Cutting edge: identification of a motile IL-17-producing T cell population in the dermis. *J. Immunol.* **186**, 6091–6095 (2011).
- Sumaria, N. *et al.* Cutaneous immunosurveillance by self-renewing dermal gammadelta T cells. *J. Exp. Med.* **208**, 505–518 (2011).
- Cai, Y. *et al.* Pivotal role of dermal IL-17-producing $\gamma\delta$ T cells in skin inflammation. *Immunity* **35**, 596–610 (2011).
- Gray, E. E. *et al.* Deficiency in IL-17-committed V γ 4(+) $\gamma\delta$ T cells in a spontaneous Sox13-mutant CD45.1(+) congenic mouse strain provides protection from dermatitis. *Nat. Immunol.* **14**, 584–592 (2013).
- Ito, Y. *et al.* Gamma/delta T cells are the predominant source of interleukin-17 in affected joints in collagen-induced arthritis, but not in rheumatoid arthritis. *Arthritis Rheum.* **60**, 2294–2303 (2009).
- Roark, C. L. *et al.* A canonical V γ 4V δ 4 + $\gamma\delta$ T cell population with distinct stimulation requirements which promotes the Th17 response. *Immunol. Res.* **55**, 217–230 (2012).
- Shibata, K. *et al.* Identification of CD25 + gamma delta T cells as fetal thymus-derived naturally occurring IL-17 producers. *J. Immunol.* **181**, 5940–5947 (2008).
- Ribot, J. C. *et al.* CD27 is a thymic determinant of the balance between interferon-gamma- and interleukin 17-producing gammadelta T cell subsets. *Nat. Immunol.* **10**, 427–436 (2009).
- Haas, J. D. *et al.* CCR6 and NK1.1 distinguish between IL-17A and IFN-gamma-producing gammadelta effector T cells. *Eur. J. Immunol.* **39**, 3488–3497 (2009).
- Narayan, K. *et al.* Intrathymic programming of effector fates in three molecularly distinct $\gamma\delta$ T cell subtypes. *Nat. Immunol.* **13**, 511–518 (2012).
- Prinz, I., Silva-Santos, B. & Pennington, D. J. Functional development of $\gamma\delta$ T cells. *Eur. J. Immunol.* **43**, 1988–1994 (2013).
- Duan, J., Chung, H., Troy, E. & Kasper, D. L. Microbial colonization drives expansion of IL-1 receptor 1-expressing and IL-17-producing gamma/delta T cells. *Cell Host Microbe* **7**, 140–150 (2010).
- Laird, R. M., Laky, K. & Hayes, S. M. Unexpected role for the B cell-specific Src family kinase B lymphoid kinase in the development of IL-17-producing T cells. *J. Immunol.* **185**, 6518–6527 (2010).
- Shibata, K. *et al.* Notch-Hes1 pathway is required for the development of IL-17-producing $\gamma\delta$ T cells. *Blood* **118**, 586–593 (2011).
- Powolny-Budnicka, I. *et al.* RelA and RelB transcription factors in distinct thymocyte populations control lymphotoxin-dependent interleukin-17 production in $\gamma\delta$ T cells. *Immunity* **34**, 364–374 (2011).
- Malhotra, N. *et al.* A network of high-mobility group box transcription factors programs innate interleukin-17 production. *Immunity* **38**, 681–693 (2013).
- Pereira, P. *et al.* Rearrangement and expression of V γ 1, V γ 2 and V γ 3 TCR γ genes in C57BL/6 mice. *Int. Immunol.* **8**, 83–90 (1996).
- O'Brien, R. L. & Born, W. K. gammadelta T cell subsets: a link between TCR and function? *Semin. Immunol.* **22**, 193–198 (2010).
- Heilig, J. S. & Tonegawa, S. Diversity of murine gamma genes and expression in fetal and adult T lymphocytes. *Nature* **322**, 836–840 (1986).
- Cai, Y. *et al.* Differential developmental requirement and peripheral regulation for dermal V γ 4 and V γ 6T17 cells in health and inflammation. *Nat. Commun.* **5**, 3986 (2014).
- Itohara, S. *et al.* Homing of a gamma delta thymocyte subset with homogeneous T-cell receptors to mucosal epithelia. *Nature* **343**, 754–757 (1990).
- Bonneville, M., O'Brien, R. L. & Born, W. K. Gammadelta T cell effector functions: a blend of innate programming and acquired plasticity. *Nat. Rev. Immunol.* **10**, 467–478 (2010).
- Horai, R. *et al.* Development of chronic inflammatory arthropathy resembling rheumatoid arthritis in interleukin 1 receptor antagonist-deficient mice. *J. Exp. Med.* **191**, 313–320 (2000).
- Nakae, S. *et al.* IL-17 production from activated T cells is required for the spontaneous development of destructive arthritis in mice deficient in IL-1 receptor antagonist. *Proc. Natl Acad. Sci. USA* **100**, 5986–5990 (2003).
- Horai, R. *et al.* TNF-alpha is crucial for the development of autoimmune arthritis in IL-1 receptor antagonist-deficient mice. *J. Clin. Invest.* **114**, 1603–1611 (2004).
- Fujikado, N., Saijo, S. & Iwakura, Y. Identification of arthritis-related gene clusters by microarray analysis of two independent mouse models for rheumatoid arthritis. *Arthritis Res. Ther.* **8**, R100 (2006).
- Lalor, S. J. *et al.* Caspase-1-processed cytokines IL-1beta and IL-18 promote IL-17 production by gammadelta and CD4 T cells that mediate autoimmunity. *J. Immunol.* **186**, 5738–5748 (2011).
- Nakamura, R. *et al.* Tyk2-signaling plays an important role in host defense against Escherichia coli through IL-23-induced IL-17 production by gammadelta T cells. *J. Immunol.* **181**, 2071–2075 (2008).
- Hayes, S. M. & Laird, R. M. Genetic requirements for the development and differentiation of interleukin-17-producing $\gamma\delta$ T cells. *Crit. Rev. Immunol.* **32**, 81–95 (2012).
- Lafaille, J. J., DeCloux, A., Bonneville, M., Takagaki, Y. & Tonegawa, S. Junctional sequences of T cell receptor gamma delta genes: implications for gamma delta T cell lineages and for a novel intermediate of V-(D)-J joining. *Cell* **59**, 859–870 (1989).
- Koenecke, C. *et al.* In vivo application of mAb directed against the $\gamma\delta$ TCR does not deplete but generates 'invisible' $\gamma\delta$ T cells. *Eur. J. Immunol.* **39**, 372–379 (2009).
- Shibata, K., Yamada, H., Hara, H., Kishihara, K. & Yoshikai, Y. Resident Vdelta1 + gammadelta T cells control early infiltration of neutrophils after Escherichia coli infection via IL-17 production. *J. Immunol.* **178**, 4466–4472 (2007).
- Lança, T. *et al.* Protective role of the inflammatory CCR2/CCL2 chemokine pathway through recruitment of type 1 cytotoxic $\gamma\delta$ T lymphocytes to tumor beds. *J. Immunol.* **190**, 6673–6680 (2013).
- Hirota, K. *et al.* Preferential recruitment of CCR6-expressing Th17 cells to inflamed joints via CCL20 in rheumatoid arthritis and its animal model. *J. Exp. Med.* **204**, 2803–2812 (2007).
- Reboldi, A. *et al.* C-C chemokine receptor 6-regulated entry of TH-17 cells into the CNS through the choroid plexus is required for the initiation of EAE. *Nat. Immunol.* **10**, 514–523 (2009).
- Arima, Y. *et al.* Regional neural activation defines a gateway for autoreactive T cells to cross the blood-brain barrier. *Cell* **148**, 447–457 (2012).
- Reddy, S. *et al.* An autoinflammatory disease due to homozygous deletion of the IL1RN locus. *N. Engl. J. Med.* **360**, 2438–2444 (2009).
- Aksentjevich, I. *et al.* An autoinflammatory disease with deficiency of the interleukin-1-receptor antagonist. *N. Engl. J. Med.* **360**, 2426–2437 (2009).
- Haas, J. D. *et al.* Development of interleukin-17-producing $\gamma\delta$ T cells is restricted to a functional embryonic wave. *Immunity* **37**, 48–59 (2012).
- Kenna, T. J. *et al.* Enrichment of circulating interleukin-17-secreting interleukin-23 receptor-positive $\gamma\delta$ T cells in patients with active ankylosing spondylitis. *Arthritis Rheum.* **64**, 1420–1429 (2012).
- Caccamo, N. *et al.* Differentiation, phenotype, and function of interleukin-17-producing human V γ 9V δ 2 T cells. *Blood* **118**, 129–138 (2011).
- Hueber, A. J. *et al.* Mast cells express IL-17A in rheumatoid arthritis synovium. *J. Immunol.* **184**, 3336–3340 (2010).
- Horai, R. *et al.* Production of mice deficient in genes for interleukin (IL)-1alpha, IL-1beta, IL-1alpha/beta, and IL-1 receptor antagonist shows that IL-1beta is crucial in turpentine-induced fever development and glucocorticoid secretion. *J. Exp. Med.* **187**, 1463–1475 (1998).
- Itohara, S. *et al.* T cell receptor delta gene mutant mice: independent generation of alpha beta T cells and programmed rearrangements of gamma delta TCR genes. *Cell* **72**, 337–348 (1993).
- Houlden, B. A. *et al.* A TCR gamma delta cell recognizing a novel TL-encoded gene product. *Cold Spring Harb. Symp. Quant. Biol.* **54**(Pt 1): 45–55 (1989).
- Komiyama, Y. *et al.* IL-17 plays an important role in the development of experimental autoimmune encephalomyelitis. *J. Immunol.* **177**, 566–573 (2006).

Acknowledgements

We thank S. Fujii (RIKEN, Kanagawa, Japan) for providing us with *Cd4*^{-/-} mice. Y. Ishii and A. Fujita (FACS Core Laboratory, The Institute of Medical Science, The University of Tokyo, Tokyo, Japan) for cell sorting; H. Saito (The University of Tokyo) for studentship support; K. Shibata (Kyushu University, Fukuoka, Japan) for reading the manuscript and for valuable discussion; A. Seno for technical assistance in designing the figures; and all members of our laboratory for excellent animal care. This work was supported by CREST (Y.I.), Grants-in-Aid from the Ministry of Education, Culture, Sports, Science and Technology of Japan (Y.I.), the Scientific Technique Research Promotion Program for Agriculture, Forestry, Fisheries and Food Industry (Y.I.) and the JSPS (A.A.).

Author contributions

A.A. made the principal contribution to all aspects of this work. H.I., S.K. and S.S. gave advice and technical support. S.-H.C., S.I. and K.S. provided technical support. S.K. and Y.L. provided technical support for the generation of *Il17^{2/8}* mice. M.U. and G.M. provided anti-*V γ 7* mAb. Y.Y. provided anti- $\gamma\delta$ TCR mAb-producing hybridoma and *Tcrd*^{-/-} mice. Y.I. organized and supervised the project and edited the draft manuscript.

Additional information

Supplementary Information accompanies this paper at <http://www.nature.com/naturecommunications>

Competing financial interests: The authors declare no competing financial interests.

Reprints and permission information is available online at <http://npg.nature.com/reprintsandpermissions/>

How to cite this article: Akitsu, A. *et al.* IL-1 receptor antagonist-deficient mice develop autoimmune arthritis due to intrinsic activation of IL-17-producing CCR2⁺ V γ 6⁺ $\gamma\delta$ T cells. *Nat. Commun.* 6:7464 doi: 10.1038/ncomms8464 (2015).



This work is licensed under a Creative Commons Attribution 4.0 International License. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in the credit line; if the material is not included under the Creative Commons license, users will need to obtain permission from the license holder to reproduce the material. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>