



Cognitive functioning throughout adulthood and illness stages in individuals with psychotic disorders and their unaffected siblings

Eva Velthorst^{1,2,3} · Josephine Mollon⁴ · Robin M. Murray⁵ · Lieuwe de Haan^{3,6} · Inez Myin Germeys⁷ · David C. Glahn⁴ · Celso Arango⁸ · Els van der Ven^{9,10} · Marta Di Forti^{11,12} · Miguel Bernardo¹³ · Sinan Guloksuz^{10,14} · Philippe Delespaul^{10,15} · Gisela Mezquida¹³ · Silvia Amoretti¹³ · Julio Bobes¹⁶ · Pilar A. Saiz¹⁶ · María Paz García-Portilla¹⁶ · José Luis Santos^{16,17,18} · Estela Jiménez-López^{16,19} · Julio Sanjuan²⁰ · Eduardo J. Aguilar²⁰ · Manuel Arrojo²¹ · Angel Carracedo^{22,23} · Gonzalo López¹⁶ · Javier González-Peñas¹⁶ · Mara Parellada⁸ · Cem Atbaşoğlu²⁴ · Meram Can Saka²⁴ · Alp Üçok²⁵ · Köksal Alptekin^{26,27} · Berna Akdede²⁶ · Tolga Binbay²⁶ · Vesile Altinyazar²⁸ · Halis Ulaş²⁶ · Berna Yalınçetin²⁷ · Güvem Gümüş-Akay^{29,30} · Burçin Cihan Beyaz³¹ · Haldun Soygür³² · Eylem Şahin Cankurtaran³³ · Semra Ulusoy Kaymak³⁴ · Nadja P. Maric³⁵ · Marina M. Mihaljevic³⁵ · Sanja Andric Petrovic³⁶ · Tijana Mirjanic³⁷ · Cristina Marta Del-Ben³⁸ · Laura Ferraro⁵ · Charlotte Gayer-Anderson³⁹ · Peter B. Jones⁴⁰ · Hannah E. Jongsma^{40,41} · James B. Kirkbride⁴¹ · Caterina La Cascia⁴² · Antonio Lasalvia⁴³ · Sarah Tosato⁴³ · Pierre-Michel Llorca⁴⁴ · Paulo Rossi Menezes⁴⁵ · Craig Morgan^{39,46} · Diego Quattrone¹¹ · Marco Menchetti⁴⁷ · Jean-Paul Selten^{10,48} · Andrei Szöke⁴⁹ · Iliaria Tarricone⁵⁰ · Andrea Tortelli⁵¹ · Philip McGuire⁵ · Lucia Valmaggia⁵² · Matthew J. Kempton⁵ · Mark van der Gaag⁵³ · Anita Riecher-Rössler⁵⁴ · Rodrigo A. Bressan⁵⁵ · Neus Barrantes-Vidal^{56,57} · Barnaby Nelson^{58,59} · Patrick McGorry^{58,60} · Chris Pantelis⁶⁰ · Marie-Odile Krebs⁶¹ · Stephan Ruhrmann⁶² · Gabriele Sachs⁶³ · Bart P. F. Rutten¹⁰ · Jim van Os^{5,10,64} · Behrooz Z. Alizadeh^{65,66} · Therese van Amelsvoort¹⁰ · Agna A. Bartels-Velthuis⁶⁶ · Richard Bruggeman^{66,67} · Nico J. van Beveren^{68,69,70} · Jurjen J. Luyckx^{64,71,72} · Wiepke Cahn^{64,73} · Claudia J. P. Simons^{10,74} · Rene S. Kahn^{1,75} · Frederike Schirmbeck^{3,6} · Ruud van Winkel^{10,76} · EU-GEI High Risk Study · Abraham Reichenberg^{1,2}

Received: 19 August 2020 / Revised: 3 November 2020 / Accepted: 18 November 2020 / Published online: 7 January 2021
© The Author(s), under exclusive licence to Springer Nature Limited 2021

Abstract

Important questions remain about the profile of cognitive impairment in psychotic disorders across adulthood and illness stages. The age-associated profile of familial impairments also remains unclear, as well as the effect of factors, such as symptoms, functioning, and medication. Using cross-sectional data from the EU-GEI and GROUP studies, comprising 8455 participants aged 18 to 65, we examined cognitive functioning across adulthood in patients with psychotic disorders ($n = 2883$), and their unaffected siblings ($n = 2271$), compared to controls ($n = 3301$). An abbreviated WAIS-III measured verbal knowledge, working memory, visuospatial processing, processing speed, and IQ. Patients showed medium to large deficits across all functions (ES range = -0.45 to -0.73 , $p < 0.001$), while siblings showed small deficits on IQ, verbal knowledge, and working memory (ES = -0.14 to -0.33 , $p < 0.001$). Magnitude of impairment was not associated with participant age, such that the size of impairment in older and younger patients did not significantly differ. However, first-episode patients performed worse than prodromal patients (ES range = -0.88 to -0.60 , $p < 0.001$). Adjusting for cannabis use, symptom severity, and global functioning attenuated impairments in siblings, while deficits in patients remained statistically significant, albeit reduced by half (ES range = -0.13 to -0.38 , $p < 0.01$). Antipsychotic medication also accounted for around half of the impairment in patients (ES range = -0.21 to -0.43 , $p < 0.01$). Deficits in verbal knowledge, and working memory may specifically index familial, i.e., shared genetic and/or shared environmental, liability for psychotic disorders. Nevertheless, potentially modifiable illness-related factors account for a significant portion of the cognitive impairment in psychotic disorders.

These authors contributed equally: Eva Velthorst, Josephine Mollon

Members of the EU-GEI High Risk Study are listed below
Acknowledgements.

Supplementary information The online version of this article (<https://doi.org/10.1038/s41380-020-00969-z>) contains supplementary material, which is available to authorized users.

✉ Eva Velthorst
eva.velthorst@mssm.edu

Extended author information available on the last page of the article

Introduction

Cognitive impairment is a common feature of schizophrenia and related psychotic disorders. Some even argue that schizophrenia should be conceptualized as a cognitive rather than psychotic illness, with cognitive dysfunction representing *the* core component of the disorder [1, 2]. Indeed, the DSM-V emphasizes the importance of assessing cognitive functioning alongside the five symptom domains (delusions, hallucinations, disorganized speech, disorganized, behavior, negative symptoms) [3], and in ICD-11, cognitive symptoms are listed alongside positive, negative, depressive, manic, and psychomotor symptoms [4]. However, despite more than a century of research on cognitive functioning in schizophrenia-related disorders, important knowledge gaps remain.

First, the profile of cognitive impairment across the lifespan remains poorly characterized, and its relationship with illness stages is unclear. Evidence suggests that patients with schizophrenia-related disorders experience some degree of cognitive decline over their lifetime, with the largest decline occurring during the years prior and up to the first few years after onset [5]. After illness onset, both cross-sectional [6] and longitudinal [7] evidence suggests at least some stabilization of impairment. However, there is also evidence for decline after onset [8], a second ‘peak’ in decline during later, chronic illness stages [9], and increased risk of dementia in very-late onset schizophrenia [10]. Efforts to examine the profile of specific cognitive functions across adulthood have also yielded mixed findings [6, 8]. Delineating the profile of these functions across adulthood and illness stages may reveal critical functions and periods for detection and intervention.

Second, cognitive decline has not been fully considered as an age-associated process, rather than in relation to stage of illness (i.e., premorbid, first-episode, chronic) [5]. Similarly, most studies have examined early [7] or late adulthood [9], without being able to trace cognitive functioning across the entirety of adulthood. While evidence suggests that cognitive decline during the first two decades of life reflects a failure to keep up with developmental norms rather than loss of cognitive capacity [11, 12], studies have not yet delineated the nature of age-associated processes throughout adulthood. The importance of considering cognitive functioning beyond adolescence is highlighted by the fact that brain and cognitive development continue well into the third decade of life [13].

Third, studies have yet to examine the full age-associated profile of familial deficits. Substantial evidence suggests that relatives of patients with schizophrenia-related disorders also show some degree of cognitive impairment [14, 15], at least in part due to shared heritable genetic mechanisms. The genetic underpinnings of cognitive impairments in schizophrenia-related disorders have been demonstrated by studies showing overlap between schizophrenia polygenic risk scores (PRS) and cognitive

performance [16], as well as educational attainment [17]. However, despite continued evidence for familial cognitive impairments, it remains unclear whether siblings show greater impairments in certain domains and whether the age-associated pattern of cognitive impairments resembles that of patients. A detailed evaluation of the familiarity of cognitive deficits across adulthood and cognitive domains may provide important additional etiological clues about the genetic and neurobiological underpinnings of cognitive impairments in schizophrenia-related disorders.

Lastly, it remains unknown whether illness-related factors such as symptom severity, global functioning, and anti-psychotic medication influence cognitive impairments differentially throughout adulthood. The age-associated influence of cannabis use on cognitive impairments is similarly unclear, despite the role cannabis may play in the emergence of psychotic symptoms [18]. The potentially moderating effect of sex on age-associated cognitive processes in psychotic disorders also remains largely unexplored. Examining these factors across adulthood may advance understanding of the etiology of cognitive impairment, as well as its clinical significance.

Using the largest sample of patients with schizophrenia-related disorders, non-psychotic siblings, and controls to date, we examined cognitive impairments across adulthood. We used cross-sectional data on general and specific functions from 8,455 individuals aged 18 to 65 from the Genetic Risk and Outcome of Psychosis (GROUP) and EUropean network of national schizophrenia networks studying Gene-Environment Interactions (EU-GEI) studies, which comprise prodromal (i.e., converted to a schizophrenia-related disorder during the study), first-episode and established illness stages. We examined whether: (1) cognitive impairments in patients differ by age category (i.e., very early-, early- and mid-adulthood) and/or stage of illness (i.e., prodromal stage, first-episode, established stage), (2) siblings of these patients show a similar pattern of impairment, and (3) this impairment is influenced by socioeconomic status, education, sex, symptom severity, global functioning, antipsychotic medication, and cannabis use.

Methods

Sample

Data were collected in 30 centers across 13 countries (UK, Netherlands, Spain, France, Italy, Serbia, Turkey, Austria, Switzerland, Germany, Australia, Denmark, Brazil), and were part of the baseline assessment for the EU-GEI study, which ran from May 1st 2010 to April 30th 2015 [19, 20], or the GROUP study, which ran from April 2004 to December 2013 [21]. Ethical approval was granted in each center and all

participants provided written informed consent. Of the combined dataset of 10,136 individuals, 685 (21.6%) patients, 259 (11.0%) siblings and 334 (10.0%) controls did not complete cognitive testing, leaving a total of 8858 individuals (3341 controls, 2347 siblings, 3,170 patients). Patients were either in the prodromal (i.e., had converted to a schizophrenia-related disorder during the study period), first-episode or established stage of illness, and were excluded if an organic cause was the primary reason for their psychotic symptoms. Control participants were excluded if they had a past or current diagnosis of any schizophrenia-related disorder. All participants had to have adequate language skills local to each center in order to undergo cognitive testing. Other exclusion and inclusion criteria for individual studies/work packages covering the different illness stages are described in the supplement.

Measures

Cognition

An abbreviated WAIS-III [22], comprising the information, arithmetic, block design, and digit symbol coding subtests, was used to measure performance in the domains of verbal knowledge, working memory visuospatial processing, and processing speed, respectively. Each WAIS subtest taps into many different abilities and the domains mentioned herein are simplified. In *GROUP*, all items of each subtest were administered. In *EU-GEI*, the digit symbol coding was administered for the standard time, along with every second item of the block design and arithmetic subtests, and every third item of the information subtest [23]. Raw scores were then multiplied by two (arithmetic and block design) or three (information). This abbreviated WAIS-III version was developed for *EU-GEI* and provides a reliable approximation of IQ and the four subtests [23].

Sociodemographic characteristics

Age, sex, ethnicity, years of education, and parental socioeconomic status (SES) were obtained (Table 1). In *EU-GEI*, parental SES (i.e., occupation level) was obtained using an amended version of the Medical Research Council Socioeconomic Schedule (MRC SDS) [24], and in *GROUP* using a comparable scale. Current cannabis use was ascertained in *GROUP* using the Composite International Diagnostic Interview (CIDI) [25] and in *EUGEI* using the Cannabis Experiences Questionnaire (CEQ) [26].

Clinical characteristics

Diagnoses were obtained using the Operational Criteria Checklist algorithm (OPCRIT) [27]. OPCRIT shows high interrater reliability generally [28] and in our study, after

training ($\kappa = 0.7$). Illness duration and current antipsychotic medication use (yes/no) were assessed using the abbreviated Nottingham Onset Schedule (NOS) [29]. Symptom severity and general functioning were measured using the Global Assessment of Functioning (GAF) symptom and disability scales [30], respectively.

Statistical analyses

Group differences in sample characteristics were examined using χ^2 , t - and Mann–Whitney- U -tests. Raw scores on the digit symbol coding, block design, information and arithmetic subtests, and the sum of these subtests (i.e., raw IQ) were z transformed. Thus, β values throughout represent standardized effect sizes (ES), with values 0.2, 0.5, and 0.8 indicating small, medium, and large ESs, respectively. These z scores were used in all statistical analyses, and are plotted by age separately for each country in sFigs. 1–5.

Age-associated group differences in cognitive functioning

Multilevel linear models (MLMs) were fitted to account for the hierarchical structure of the data (i.e., with random intercepts for country, center, and family). Based on age distributions (sFig. 6) and nonlinear relationships between age and cognitive functioning (Fig. 1), we categorized individuals into approximately equal-sized age groups: 18–25, 26–39, and 40–65 years, representing very early-, early- and mid-adulthood, respectively (Table 1). Effects of interest were group main effects and the interaction between group and age category. A statistically significant group main effect would indicate a difference in cognitive performance between patients and/or siblings compared to controls. A statistically significant group-by-age interaction would indicate that group differences in cognitive performance differ by participants' age. Sex and ethnicity were entered as covariates in all models.

Illness stage and duration

The effect of illness stage was examined using MLMs as described above, but with prodromal patients set as the reference. Illness stage was based on study (i.e., prodrome study = prodromal stage, first-episode study = first-episode, course studies = established stage), except for individuals in the course studies with an illness duration of less than 2 years ($n = 314$), who were considered first-episode. Illness duration (measured in years from illness onset) was entered into MLMs as a continuous effect of interest.

Sociodemographic and illness-related factors

We entered sociodemographic and illness-related factors (current cannabis use; symptom severity i.e., GAF

Table 1 Sample characteristics.

	Prodrome ^a N = 56		First-episode ^b N = 865		Established-stage ^c N = 1962		Siblings ^d N = 2271		Controls ^c N = 3,301		Group differences
	n/mean	%/SD	n/mean	%/SD	n/mean	%/SD	n/mean	%/SD	n/mean	%/SD	
Male	32	57.1	530	61.3	1406	71.7	996	43.7	1609	48.7	a=c,d; b=c,d,e;
<i>Missing</i>	–	–	–	–	–	–	1	0.0	–	–	c=d,e; d≠e
Age	22.80	4.33	30.72	10.46	30.46	8.57	31.60	9.14	34.23	11.68	a=b,c,d,e; b=d,e; c≠d,
<i>Missing</i>	–	–	–	–	–	–	3	0.0	2	0.0	e; d≠e
Age category											
18–25	46	82.1	350	40.5	660	33.6	701	30.9	921	27.9	a=b,c,d,e; b=c,d,e;
26–39	10	17.9	346	40.0	953	48.6	1102	48.6	1327	40.2	c=d,e; d≠e
40–65	–	–	169	19.5	349	17.9	465	20.5	1051	31.9	
White ethnicity	36	64.3	556	64.3	1721	88.9	2036	89.9	2953	89.78	a=c,d,e; b=c,d,e
<i>Missing</i>	–	–	–	–	27	1.4	6	0.3	12	0.4	
Illness duration in years	–	–	0.67	2.86	6.27	6.02	–	–	–	–	b=c
<i>Missing</i>	–	–	298	34.5	485	24.72	–	–	–	–	
Antipsychotics	9	25.0	816	96.3	1745	93.7	–	–	–	–	a=b,c; b=c
<i>Missing</i>	20	35.7	18	2.1	99	5.0	–	–	–	–	
Paternal SES											
Higher	16	35.6	193	25.2	52	24.3	59	26.0	412	28.1	a=b,c,d,e; b=d,e
Intermediate	17	37.8	181	23.6	64	29.9	77	33.9	419	28.6	
Lower	9	20.0	369	48.1	97	45.3	90	39.7	625	42.6	
Unemployed	3	6.7	24	3.1	1	0.5	1	0.4	11	0.8	
<i>Missing</i>	11	18.3	98	11.3	1748 ^a	89.1 ^a	2044 ^a	90.0 ^a	1834 ^a	55.6 ^a	
Country											
UK	22	39.3	155	17.9	–	–	5	0.2	324	9.8	–
Netherlands	5	8.9	176	20.4	998	50.9	1012	44.6	755	22.9	
Austria	3	5.4	–	–	–	–	–	–	–	–	
Switzerland	5	8.9	–	–	–	–	–	–	–	–	
Spain	2	3.6	185	21.4	385	19.6	538	23.7	637	19.3	
Turkey	–	–	–	–	528	26.9	565	24.9	949	28.8	
Serbia	–	–	–	–	51	2.6	50	2.2	36	1.1	
Australia	3	5.4	–	–	–	–	–	–	15	0.5	
Germany	5	8.9	–	–	–	–	–	–	–	–	
France	7	12.5	87	10.1	–	–	6	0.3	144	4.4	
Brazil	–	–	162	18.7	–	–	88	3.9	280	8.5	
Denmark	4	7.1	–	–	–	–	–	–	–	–	
Italy	–	–	100	11.6	–	–	7	0.3	161	4.9	
Years of education	14.36	3.19	13.10	4.22	12.14	3.98	13.33	4.30	13.60	4.25	a=c,d,e; b=c,d,e; c≠d
<i>Missing</i>	9	13.8	18	2.1	98	5.0	62	2.7	44	1.3	
Current cannabis use	15	27.8	189	22.1	205	14.7	203	12.5	304	10.5	a=c,d,e; b=c,d,e;
<i>Missing</i>	2	3.3	9	1.0	566	28.8	642	28.3	414	12.5	c≠e; d≠e
GAF symptoms	51.65	8.25	47.61	17.50	54.60	16.54	82.46	12.15	85.31	9.90	a=d,e; b=c,d,e;
<i>Missing</i>	4	6.2	38	4.4	139	7.1	1097	46.8	609	18.4	c=d,e; d≠e
GAF disability	52.14	10.27	51.82	16.53	54.32	16.82	84.91	11.34	86.60	9.10	a=d,e; b=c,d,e;
<i>Missing</i>	–	–	37	4.3	143	7.3	1037	45.7	608	18.4	c=d,e; d≠e
Scaled cognition scores											
IQ	97.32	14.64	84.86	18.06	88.74	16.93	97.19	16.60	98.88	18.06	a=b,c; b=c,d,e;
											c=d,e; d≠e
Information	10.50	3.39	8.77	3.80	9.56	3.67	9.69	3.52	10.19	3.55	a=b; b=c,d,e; c≠e; d≠e
Arithmetic	9.67	3.35	7.70	3.34	8.84	3.23	9.82	3.25	10.08	3.19	a=b; b=c,d,e;
											c=d,e; d≠e
Block design	9.64	3.27	7.82	3.64	8.72	3.36	9.89	3.06	9.84	3.29	a=b; b=c,d,e;
											c=d,e; d≠e
Symbol coding	8.70	2.76	6.68	2.91	5.99	3.43	9.04	3.66	9.10	4.06	a=b,c; b=c,d,e; c≠d,e

^aMissingness due to comparable measures not being available in the GROUP study.

≠Significantly different at $p < 0.05$.




Fig. 1 Loess curves of z scores against age for each groups. Lines represent the cognitive performance Z-scores (Y-axis) for controls (red), siblings (green) and patients (blue) by age (X-axis).

symptoms; global functioning i.e., GAF disability; illness duration; parental SES; years of education; antipsychotic medication) as covariates into separate MLMs to examine whether each of these factors influenced group and group-by-age effects.

Sex-related differences

We fitted MLMs separately in males and females to examine potential sex differences in group and group-by-age effects. In order to formally test for sex-related

differences, we also entered sex-by-group, and three-way interactions between sex, group, and age into MLMs on the whole sample.

Sensitivity analyses

Sensitivity analyses were conducted to determine whether cognitive patterns were similar in patients with non-affective and affective psychosis. To rule out any potential bias due to the inclusion of patients without a participating sibling (37.4% of sample), we repeated the main analyses including only patients with a participating sibling. We also analyzed controls and siblings with high GAF disability scores (80+, controls: $n = 2193$; siblings: $n = 834$) and low GAF disability scores (<80, controls: $n = 400$; siblings: $n = 405$) separately to examine potential bias from missing GAF data.

An adjusted p -value threshold of 0.005 ($0.05 \div 10$ (5 cognitive subtests \times 2 statistical models for (1) main effects and (2) interaction effects)) was used in all models to account for multiple comparisons. Statistical analyses were performed in Stata 15.1 [31]. The R [32] package *ggplot2* [33] was used to create graphics.

Results

Sample characteristics

Table 1 shows sample characteristics. The small number of participants below the age of 18 were excluded ($n = 179$). Thus, the final sample comprised 2883 patients, 2271 siblings, and 3301 controls. Of this sample, 1805 (62.6%) patients had at least one participating sibling (range = 1–5; median = 1).

Older participants showed lower scores than younger participants

Across all groups, IQ ($\beta = -0.42$, $z = -12.01$, $SE = 0.04$, $p < 0.001$), block design ($\beta = -0.45$, $z = -11.39$, $SE = 0.04$, $p < 0.001$), and digit symbol coding ($\beta = -0.42$, $z = -12.15$, $SE = 0.04$, $p < 0.001$) was significantly associated with participant age, such that participants in mid-adulthood performed worse than participants in very-early-adulthood (Fig. 2, Table 2).

Patients showed substantial cognitive impairments that were not associated with participant age

Patients showed medium to large deficits across all cognitive measures (Fig. 2, Table 2). Large deficits were seen on IQ ($\beta = -0.73$, $z = -20.39$, $SE = 0.04$, $p < 0.001$), and digit symbol coding ($\beta = -0.71$, $z = -20.30$, $SE = 0.03$,

$p < 0.001$). Medium deficits were observed on information ($\beta = -0.45$, $z = -11.05$, $SE = 0.04$, $p < 0.001$), arithmetic ($\beta = -0.66$, $z = -15.87$, $SE = 0.04$, $p < 0.001$), and block design ($\beta = -0.45$, $z = -11.05$, $SE = 0.04$, $p < 0.001$). No group-by-age interactions reached statistical significance, suggesting no differential association between cognitive performance and participant age in patients and controls. Namely, older participants scored worse than younger participants in both groups, and the magnitude of difference between older and younger participants was not significantly different between groups (Table 2).

First-episode patients performed worse than patients in other illness-stages

First-episode patients performed worse than prodromal patients on IQ ($\beta = -0.88$, $z = -4.75$, $SE = 0.19$, $p < 0.001$), information ($\beta = -0.60$, $z = -3.06$, $SE = 0.19$, $p = 0.002$), arithmetic ($\beta = -0.61$, $z = -3.11$, $SE = 0.20$, $p = 0.002$), block design ($\beta = -0.82$, $z = -4.07$, $SE = 0.20$, $p < 0.001$), and slightly worse than established stage patients on information ($\beta = -0.16$, $z = -3.72$, $SE = 0.04$, $p < 0.001$). Established stage patients performed worse than prodromal patients on IQ ($\beta = -0.80$, $z = -4.26$, $SE = 0.19$, $p < 0.001$), arithmetic ($\beta = -0.58$, $z = -2.89$, $SE = 0.20$, $p = 0.004$), block design ($\beta = -0.70$, $z = -3.42$, $SE = 0.20$, $p = 0.001$), and digit symbol coding ($\beta = -0.66$, $z = 3.65$, $SE = 0.18$, $p < 0.001$). All differences remained after adjusting for age. Illness duration showed no statistically significant effects on cognition.


Siblings showed small cognitive impairments that were smaller in older than younger participants

Compared to controls, siblings showed small deficits on IQ ($\beta = -0.14$, $z = -3.65$, $SE = 0.04$, $p < 0.001$), information ($\beta = -0.33$, $z = -7.64$, $SE = 0.04$, $p < 0.001$) and arithmetic ($\beta = -0.23$, $z = -4.97$, $SE = 0.05$, $p < 0.001$) (Fig. 2, Table 2). These deficits were smaller in mid-adulthood than in early-adulthood, reflected by significant group-by-age interactions on IQ ($\beta = 0.16$, $z = 2.88$, $SE = 0.06$, $p = 0.004$), information ($\beta = 0.22$, $z = 3.47$, $SE = 0.06$, $p = 0.001$) and block design ($\beta = 0.20$, $z = 3.15$, $SE = 0.06$, $p = 0.002$) (Table 2). Thus, while older participants scored worse than younger participants in both groups, the magnitude of difference between older and younger participants was smaller in siblings than controls.

Sociodemographic and illness-related factors influenced cognitive impairments

Figure 3 shows differences in effect sizes of group main effects for models adjusting for each covariate, compared to

Fig. 2 Cognitive z scores at each age category by group. Cognitive scores for controls aged 18–25 set to 0 i.e., as reference category.



the unadjusted model. Adjusting for parental SES statistically attenuated deficits in verbal knowledge, and working memory in siblings, but effect sizes remained small (sTable 1, sFig. 7). Adjusting for current cannabis use (sTables 2, sFig. 8) and years of education (sTables 3, sFig. 9) attenuated the small IQ deficit in siblings. Adjusting for symptom severity (GAF symptoms) (sTable 4, sFig. 10) and global functioning (GAF disability) (sTable 5, sFig. 11) attenuated the IQ and arithmetic deficits in siblings, and the

block design impairment in patients. Information, arithmetic, and symbol coding deficits in patients remained statistically significant, but were reduced by more than half when adjusting for these factors (sTables 4 and 5). Interestingly, when entering both global functioning and symptom severity into the same model, only functioning remained statistically associated with cognition ($z = 9.84$, $SE = 0.001$, $p < 0.001$), while symptoms were not ($z = 1.59$, $SE = 0.001$, $p = 0.11$). This finding suggests that the

Table 2 Age, group and group-by-interaction effects on cognitive measures, adjusting for sex and ethnicity.

Measure	Age effect					Group effect					Group-by-age interactionVery-early- to early-adulthood					Group-by-age interactionEarly- to mid-adulthood				
	Age	β	z	SE	p	Group	β	z	SE	p	β	z	SE	p	β	z	SE	p	Δ ES	
IQ	18-25	-	-	-	-	Controls	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	26-39	-0.02	-0.73	0.03	0.46	Siblings	-0.14	-3.65	0.04	<0.001	0.08	1.72	0.05	0.09	0.16	2.88	0.06	0.004	0.06	-0.32
	40-65	-0.42	-12.01	0.04	<0.001	Patients	-0.73	-20.39	0.04	<0.001	0.03	0.56	0.05	0.57	0.08	1.56	0.05	0.12	0.05	0.34
Information	18-25	-	-	-	-	Controls	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	26-39	0.09	2.37	0.04	0.02	Siblings	-0.33	-7.64	0.04	<0.001	0.12	2.17	0.05	0.03	0.22	3.47	0.06	0.001	0.06	0.04
	40-65	0.03	0.77	0.04	0.43	Patients	-0.46	-11.24	0.04	<0.001	0.06	1.26	0.05	0.21	0.12	1.97	0.06	0.05	0.05	0
Arithmetic	18-25	-	-	-	-	Controls	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	26-39	0.02	0.41	0.04	0.68	Siblings	-0.23	-4.97	0.05	<0.001	0.06	2.24	0.06	0.03	0.11	1.65	0.07	0.10	0.06	-0.11
	40-65	-0.09	-2.32	0.04	0.02	Patients	-0.66	-15.87	0.04	<0.001	0.05	1.80	0.05	0.07	0.14	2.24	0.06	0.03	0.06	-0.06
Block design	18-25	-	-	-	-	Controls	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	26-39	-0.06	-1.45	0.04	0.14	Siblings	-0.09	-1.95	0.04	0.05	0.08	1.41	0.05	0.16	0.20	3.15	0.06	0.002	0.06	-0.28
	40-65	-0.45	-11.39	0.04	<0.001	Patients	-0.45	-11.05	0.04	<0.001	0.01	0.22	0.05	0.82	0.07	1.11	0.06	0.27	0.06	-0.39
Symbol coding	18-25	-	-	-	-	Controls	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	26-39	-0.03	-0.99	0.03	0.32	Siblings	-0.04	-0.94	0.04	0.35	0.02	0.48	0.05	0.63	0.04	0.73	0.06	0.47	0.06	-0.37
	40-65	-0.42	-12.15	0.03	<0.001	Patients	-0.71	-20.30	0.03	<0.001	0	-0.02	0.04	0.99	0.03	0.61	0.05	0.54	0.05	-0.35

Since cognitive test scores were z transformed, β coefficients correspond to standardized ESs (values 0.2, 0.5 and 0.8 indicate small, medium and large ESs, respectively). Bolded estimates signify a statistical significant difference between the cognitive score of patients or siblings with the control group ($p < 0.005$).

association between functioning and cognition may account for most of the association between symptoms and cognition. Finally, adjusting for antipsychotic medication attenuated the information and block design impairments in patients, and reduced the magnitude of deficits in IQ, arithmetic, and digit symbol coding by about half (sTable 6, sFig. 12). sTable 7 shows correlations between socio-demographic and illness-related factors and all cognitive measures.

Cognitive impairments were comparable in male and female patients but not male and female siblings


Compared to controls, both male and female patients showed medium to large impairments across all cognitive measures (sTables 8 and 9). Accordingly, sex-by-group interactions did not reveal any sex-related differences in patients on any cognitive measure, such that cognitive impairments were comparable in male and female patients. On the other hand, male siblings showed a smaller deficit on information ($\beta = -0.21$, $z = -3.34$, $SE = 0.06$, $p = 0.001$), than female siblings ($\beta = -0.40$, $z = -6.72$, $SE = 0.06$, $p < 0.001$), and the deficit on arithmetic did not reach significance in male siblings ($\beta = -0.12$, $z = -1.83$, $SE = 0.07$, $p = 0.07$), but did in female siblings ($\beta = -0.30$, $z = -4.94$, $SE = 0.06$, $p < 0.001$) (sTables 8 and 9, sFigs. 13 and 14). Accordingly, significant sex-by-group interactions on information ($\beta = -0.11$, $z = -2.46$, $SE = 0.05$, $p = 0.014$) and arithmetic ($\beta = -0.13$, $z = -0.05$, $SE = 2.66$, $p = 0.008$), confirmed that female siblings showed greater deficits on these domains than male siblings.

Group-by-age interactions no longer reached significance in male siblings, but female siblings showed a significant group-by-age interaction on information between early- and mid-adulthood ($\beta = 0.27$, $z = 3.25$, $SE = 0.08$, $p = 0.001$), suggesting a differential association between cognitive performance and participant age in female siblings and controls (sTables 8 and 9). Namely, while older participants scored worse than younger participants across groups, the magnitude of difference between older and younger participants was smaller in female siblings than controls. However, none of the sex-by-group-by-age interactions reached statistical significance, suggesting negligible sex-related differences in age-associated effects on cognitive functioning.

Sensitivity analyses

Sensitivity analyses comparing patients with non-affective and affective psychosis, patients with and without participating siblings, and participants with high and low GAF scores revealed similar patterns overall (sResults).

Fig. 3 Difference in effect size of group effect in models adjusting for each covariate compared to main model. Positive differences in effect size indicate reduced difference between controls and patients/siblings and negative differences indicate increased difference between controls and patients/siblings.



Discussion

In this large, cross-sectional patient-sibling-control study, patients with psychotic disorders showed large, widespread cognitive impairments, which were not associated with participant age. However, first-episode patients showed larger deficits than prodromal patients, and a slightly larger verbal knowledge deficit compared to established stage patients. Siblings showed small deficits on IQ, and measures of verbal knowledge and working memory, which were attenuated when adjusting for cannabis use, symptom severity, global functioning, and education. These findings add to current knowledge in several important ways.

First, effects of participant age on the magnitude of cognitive impairments were minimal. Examining cognitive raw scores throughout adulthood revealed that older patients showed lower scores than younger patients, but also that the same was true of controls, such that the magnitude of impairment remained stable. However, magnitude of impairment did differ by illness stage, with first-episode patients showing much larger deficits than prodromal patients. In line with previous evidence [34], impairments in the oldest siblings were smaller than impairments in the youngest siblings, which may be because older siblings have passed the critical period for psychosis-onset, while younger siblings have not and may still be at risk for psychotic disorders. Alternatively, controls may experience greater age-associated cognitive decline later in adulthood, while siblings and patients, who already experienced decline earlier in adulthood, may show relative stabilization or even normalization [6]. Our results should also be considered in the context of the well-documented Flynn effect [35], whereby cognitive performance and IQ in any fixed age group improves over time due to improvements in education, nutrition, etc. Thus, while our finding of lower

cognitive scores in older participants compared to younger participants may reflect age-associated cognitive decline [36, 37], the Flynn effect may also account for this finding. Conversely, recent data suggest a reversal of the Flynn effect [38–40], which may partly explain our finding of smaller impairments in older siblings compared to younger siblings. Moreover, while we made considerable efforts to recruit a well-matched representative sample using quota sampling methods, we cannot rule out selection bias. For example, participation in research studies is associated with better cognitive functioning and this bias may be more pronounced in later adulthood. Similarly, since lower IQ is associated with earlier mortality [41], older individuals with more pronounced cognitive impairment may be less likely to participate. Future longitudinal studies that are able to prospectively follow individuals throughout adulthood are needed to determine the profile and underlying mechanisms of age-associated cognitive processes in psychotic disorders. Overall, our findings support previous evidence that most of the cognitive deficit associated with psychotic disorders is already apparent at illness onset [7].

Second, including a large sample of siblings with similar genetic and environmental predispositions as patients, but without the potentially confounding effects of illness-related factors, provides important insights into the familiarity of cognitive impairments [14]. Specifically, while patients showed medium to large deficits across all measures, siblings showed small deficits on IQ, verbal knowledge, and working memory, but not on processing speed and visuospatial processing. This latter finding contrasts with a meta-analytic finding of large processing speed deficits in first-degree relatives [42], likely because most studies in this meta-analysis combined data from parents and siblings, while we only examined siblings, who are younger. Nevertheless, our findings suggest that verbal

knowledge and working memory deficits may specifically index familial, i.e., shared genetic and/or environmental, liability for psychotic disorders. Accordingly, a recent meta-analysis of cognition in first-degree relatives of patients with schizophrenia also reported the largest deficits in IQ and verbal measures [15]. The notion of verbal impairments as a familial marker for psychotic illness is in line with evidence that verbal deficits emerge early [11, 12]. Non-verbal impairments, on the other hand, emerge over time [12], increase throughout the early illness stage [11], but remain stable thereafter [8]. Thus, shared genetic and/or environmental factors may lead to deficits in verbal abilities in individuals at familial risk for psychosis, while additional risk factors, possibly interacting with these verbal deficits, may lead to emerging nonverbal deficits and psychotic illness. It is important to note that we examined only two cognitive tests requiring verbal skills, and shared genetic liability may depend on the subtest measured. For example, genomic loci that jointly influence schizophrenia and verbal-numerical reasoning have been identified [43], but a recent study showed no association between schizophrenia PRSs and a verbal reading test [44].

Third, adjusting for both symptom severity and global functioning attenuated the IQ and working memory deficits in siblings, and reduced cognitive deficits in patients by half. Yet our findings also suggest that the association between symptoms and cognition is confounded by functioning. Thus, while the two GAF subscales are highly correlated ($r=0.69$) and adjusting for each subscale reduced deficits by similar magnitudes, it may be important to disentangle the effects of these factors. These findings are in line with evidence of the significant impact of cognition on functional outcomes [45], as well as the lack of a strong association between cognition and symptom severity [46]. Interestingly, siblings outperformed controls in visuospatial ability and processing speed after adjustment for symptoms and functioning, suggesting a potentially protective mechanism. Deficits in working memory, visuospatial, and processing speed abilities may therefore be ameliorated by improving functioning levels. Impairments in both patients and siblings were also reduced, albeit to a lesser extent, when adjusting for education. Interestingly, this reduction was slightly more pronounced on the verbal knowledge and working memory tests, suggesting that these deficits may partly reflect an impairment in the ability to learn in standard educational settings. While the relationship between cognitive impairment and factors such as educational attainment and functioning are difficult to discern due to reverse causality, future studies that are able to disentangle whether they act as mediators, moderators, or lie on the causal pathway between cognition and psychosis will provide important insights. The finding that female siblings showed larger deficits than male siblings is also intriguing,

and highlights the need for further examination of sex-specific genetic risk factors [47]. Finally, adjusting for current cannabis use had a negligible impact on patient impairments, but attenuated IQ deficits in siblings. These findings are in line with evidence of a minimal association between cannabis use and cognitive functioning in psychotic disorders [48], as well as more severe symptomatology in sibling cannabis users [49].

This study has some limitations. First, our findings require replication in longitudinal samples since we used cross-sectional data. We also cannot rule out age-associated effects on cognition in early life or late adulthood since our youngest and oldest participants were 18 and 65, respectively. Moreover, while the large age range is a strength, cohort effects should be considered. Nevertheless, the current findings are in line with longitudinal results from the same sample [50]. Second, one limitation inherent to large cohorts is the tradeoff between breadth and depth. While we examined a number of covariates, future studies that are able to examine anti-psychotic dosage, type and adherence, comorbidities, such as anxiety and depression, and positive and negative psychotic symptoms are needed. Comorbidity is the rule rather than the exception in psychotic disorders [51], making it difficult to disentangle the effects of psychotic versus other psychiatric symptoms. The reduction in power due to missing data when adjusting for covariates also warrants consideration, and effect sizes should be considered alongside statistical significance. Third, while our sensitivity analyses eliminate certain sources of bias, others cannot be ruled out. Individuals with better functioning may be more likely to participate in research studies, although the reverse is also possible. Fourth, our findings regarding specific cognitive domains require replication using larger test batteries that are able to cover each domain in greater detail. Relatedly, abbreviated tests, such as those administered in EUGEI, may both over-estimate (reduced fatigue) and underestimate (less attenuated learning) cognitive functioning, especially in individuals of low ability [23]. However, our data show normal distribution of IQ and subtest scaled scores across all groups (see sTable 10).

In conclusion, using a large, cross-sectional sample of patients with psychotic disorders, their siblings and controls, we found that patients showed substantial and widespread cognitive impairments, while siblings showed smaller verbal knowledge and working memory impairments. Moreover, effects of age and illness stage (beyond the first episode) on these impairments were minimal, while illness-related factors accounted for much of the impairment in siblings, and around half of the patient deficit. Thus, our findings suggest that most of the cognitive impairment associated with psychotic disorders is already apparent at illness onset, highlighting the importance of early cognitive remediation intervention efforts. Therapeutic efforts targeting illness-related factors, such as symptoms and

functioning, which account for a significant portion of the cognitive impairment, could also have substantial benefits. Finally, deficits in verbal knowledge and working memory may specifically index familial liability and could be useful targets for studies aimed at elucidating the heritable neurobiological mechanisms underlying psychotic disorders.

Acknowledgements The European Community's Seventh Framework Programme under grant agreement No. HEALTH-F2-2010-241909 (EU-GEI). The Geestkracht programme of the Dutch Health Research Council (Zon-Mw)(GROUP). Dr. Velthorst is supported by The Seaver Foundation. Dr. Pantelis was supported by a NHMRC Senior Principal Research Fellowship (ID: 1105825), a NHMRC Program Grant (ID: 1150083). The Melbourne arm of the study was supported by a grant from the Australian National Health & Medical Research Council (NHMRC-EU grant ID: 567215). The French cohort was supported by the French Ministry grant (PHRC ICAAR - AOM07-118). The Spanish sample was supported by the Spanish Ministry of Science and Innovation, Instituto de Salud Carlos III (SAM16PE07CP1, PI16/02012, PI19/024), Madrid Regional Government (B2017/BMD-3740 AGES-CM-2), Fundación Familia Alonso and Fundación Alicia Koplowitz. The Brazilian sample was supported by FAPESP-Brazil (grant 2012/05178-0). Additional support was provided by a Medical Research Council Fellowship to Dr. Kempton (grant MR/J008915/1). Dr. Kirkbride was supported by the National Institute for Health Research University College London Hospital Biomedical Research Centre. Dr. Nelson was supported by an NHMRC Senior Research Fellowship (1137687). We would like to thank the EU-GEI WP2 Group not mentioned in main author list: Kathryn Hubbard, Stephanie Beards, Simona A. Stilo, Pedro Cuadrado, José Juan Rodríguez Solano, David Fraguas, Álvaro Andreu-Bernabeu, Gonzalo López, Bibiana Cabrera, Juan Nacher, Javier Costas, Mario Matteis [8], Marta Rapado-Castro, Emiliano González, Covadonga M. Díaz-Caneja [8], Emilio Sánchez, Manuel Durán-Cutilla, Nathalie Franke, Fabian Termorshuizen, Daniella van Dam, Elles Messchaert, Marion Leboyer [4], Franck Schürhoff, Stéphane Jamain, Grégoire Baudin, Aziz Ferchiou, Baptiste Pignon, Jean-Romain Richard, Thomas Charpeaud, Anne-Marie Tronche, Flora Frijda, Giovanna Marrazzo, Crocettarachele Sartorio, Fabio Seminerio, Camila Marcelino Loureiro, Rosana Shuhama, Mirella Ruggeri, Chiara Bonetto, Doriana Cristofalo, Domenico Berardi, Marco Seri, Elena Bonora, Anastasios Nougus, Giuseppe D'Andrea, Laura Ferraro, Giada Tripoli, Ulrich Reininghaus, Enrique García Bernardo, Laura Roldán, Esther Lorente-Rovira, Ma Soledad Olmeda, Daniele La Barbera, Cristina M. Del-Ben, Lucia Sideli. Study funders contributed to the salaries of the research workers employed, but did not participate in the study design, data analyses, data interpretation, or writing of the manuscript. All authors had full access to the study data and had final responsibility for the decision to submit for publication.

EU-GEI High Risk Study Maria Calem⁵, Stefania Tognin⁵, Gemma Modinos⁵, Sara Pisani⁵, Tamar C. Kraan³, Daniella S. van Dam³, Nadine Burger⁵³, G. Paul Amminger⁵⁸, Athena Politis⁵⁸, Joanne Goodall⁵⁸, Stefan Borgwardt⁷⁷, Erich Studerus⁷⁷, Ary Gadelha⁵⁵, Elisa Brietzke⁷⁸, Gracielle Asevedo⁵⁵, Elson Asevedo⁵⁵, Andre Zugman⁵⁵, Tecelli Domínguez-Martínez⁷⁹, Manel Monsonet⁸⁰, Paula Cristóbal-Narváez⁸⁰, Anna Racioppi⁵⁶, Thomas R. Kwapił⁸¹, Mathilde Kazes⁶¹, Claire Daban⁶¹, Julie Bourgin⁶¹, Olivier Gay⁶¹, Céilia Mam-Lam-Fook⁶¹, Dorte Nordholm⁸², Lasse Rander⁸², Kristine Krakauer⁸², Louise Birkedal Glenthøj⁸², Birte Glenthøj⁸³, Dominika Gebhard⁶², Julia Arnhold⁸⁴, Joachim Klosterkötter⁶², Iris Lasser⁶³, Bernadette Winklbauer⁶³

⁷⁷University of Basel, Basel, Switzerland; ⁷⁸Department of Psychiatry, Escola Paulista de Medicina, Universidade Federal de São Paulo (UNIFESP), São Paulo, Brazil; ⁷⁹CONACYT-Dirección de

Investigaciones Epidemiológicas y Psicosociales, Instituto Nacional de Psiquiatría Ramón de la Fuente Muñiz, México, México; ⁸⁰Departament de Psicologia Clínica i de la Salut (Universitat Autònoma de Barcelona), Barcelona, Spain; ⁸¹Department of Psychology, University of Illinois at Urbana-Champaign, Champaign, IL, USA; ⁸²Mental Health Center Copenhagen and Center for Clinical Intervention and Neuropsychiatric Schizophrenia Research, CINS, Mental Health Center Glostrup, Mental Health Services in the Capital Region of Copenhagen, University of Copenhagen, Copenhagen, Denmark; ⁸³Centre for Neuropsychiatric Schizophrenia Research (CNSR) & Centre for Clinical Intervention and Neuropsychiatric Schizophrenia Research (CINS), Mental Health Centre Glostrup, University of Copenhagen, Glostrup, Denmark; ⁸⁴Psyberlin, Berlin, Germany

Compliance with ethical standards

Conflict of interest Dr. Arango. has been a consultant to or has received honoraria or grants from Acadia, Angelini, Gedeon Richter, Janssen Cilag, Lundbeck, Minerva, Otsuka, Roche, Sage, Servier, Shire, Schering Plough, Sumitomo Dainippon Pharma, Sunovion and Takeda. Dr. Pantelis has received honoraria for talks at educational meetings and has served on an advisory board for Lundbeck, Australia Pty Ltd. Dr. Bernardo has been a consultant for, received grant/research support and honoraria from, and been on the speakers/advisory board of ABBiotics, Adamed, Angelini, Casen Recordati, Eli Lilly, Ferrer, Forum Pharmaceuticals, Janssen-Cilag, Lundbeck, Menarini, Otsuka, Takeda and Somatics. MO Krebs received financial support from Janssen, Otsuka Lundbeck alliance, Eisai for educational activities. All other authors report no financial relationships with commercial interests.

Publisher's note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

References

- Kahn RS, Keefe RS. Schizophrenia is a cognitive illness: time for a change in focus. *JAMA Psy.* 2013;70:1107–12.
- Elvevag B, Goldberg TE. Cognitive impairment in schizophrenia is the core of the disorder. *Critical Rev Neurobiol.* 2000;14:1–21.
- American Psychiatric Association. Diagnostic and statistical manual of mental disorders (DSM-5®). Washington, D.C.: American Psychiatric Association; 2013.
- World Health Organization. International classification of diseases, 11th revision (ICD-11). ICD-11 is here Geneva: World Health Organization; 2018.
- Becker H, Nieman D, Wiltink S, Dingemans P, Van de Fliert J, Velthorst E, et al. Neurocognitive functioning before and after the first psychotic episode: does psychosis result in cognitive deterioration? *Psych Med.* 2010;40:1599–606.
- Mollon J, Mathias SR, Knowles EE, Rodrigue A, Koenis MM, Pearlson GD, et al. Cognitive impairment from early to middle adulthood in patients with affective and nonaffective psychotic disorders. *Psychol Med.* 2019;50:48–57.
- Bozikas VP, Andreou C. Longitudinal studies of cognition in first episode psychosis: a systematic review of the literature. *Aust NZealand J Psych.* 2011;45:93–108.
- Zanelli J, Mollon J, Sandin S, Morgan C, Dazzan P, Pilecka I, et al. Cognitive change in schizophrenia and other psychoses in the decade following the first episode. *Am J Psychiatry.* 2019; 176:811–9.
- Harvey PD. What is the evidence for changes in cognition and functioning over the lifespan in patients with schizophrenia? *J Clin Psychiatry.* 2014;75:34–38.

10. Kodesh A, Goldberg Y, Rotstein A, Weinstein G, Reichenberg A, Sandin S, et al. Risk of dementia and death in very-late-onset schizophrenia-like psychosis: a national cohort study. *Schizophrenia Res.* 2020;223:220–6.
11. Mollon J, David AS, Zammit S, Lewis G, Reichenberg A. Course of cognitive development from infancy to early adulthood in the psychosis spectrum. *JAMA Psychiatry.* 2018;75:270–9.
12. Reichenberg A, Caspi A, Harrington H, Houts R, Keefe RS, Murray RM, et al. Static and dynamic cognitive deficits in childhood preceding adult schizophrenia: a 30-year study. *Am J Psych.* 2010;167:160–9.
13. Lebel C, Walker L, Leemans A, Phillips L, Beaulieu C. Microstructural maturation of the human brain from childhood to adulthood. *Neuroimage.* 2008;40:1044–55.
14. Snitz BE, MacDonald III AW, Carter CS. Cognitive deficits in unaffected first-degree relatives of schizophrenia patients: a meta-analytic review of putative endophenotypes. *Schizophr Bull.* 2005;32:179–94.
15. Agnew-Blais J, Seidman LJ. Neurocognition in youth and young adults under age 30 at familial risk for schizophrenia: a quantitative and qualitative review. *Cogn Neuropsychiatry.* 2013;18:44–82.
16. van Os J, van der Steen Y, Islam MA, Guloksuz S, Rutten BP, Simons CJ, et al. Evidence that polygenic risk for psychotic disorder is expressed in the domain of neurodevelopment, emotion regulation and attribution of salience. *Psychol Med.* 2017;47:2421–37.
17. Sorensen HJ, Debost JC, Agerbo E, Benros ME, McGrath JJ, Mortensen PB, et al. Polygenic risk scores, school achievement, and risk for schizophrenia: a danish population-based study. *Biol Psychiatry.* 2018;84:684–91.
18. Kuepper R, van Os J, Lieb R, Wittchen H-U, Höfler M, Henquet C. Continued cannabis use and risk of incidence and persistence of psychotic symptoms: 10 year follow-up cohort study. *Bmj.* 2011;342:d738.
19. EU-GEI. Identifying gene-environment interactions in schizophrenia: contemporary challenges for integrated, large-scale investigations. *Schizophrenia Bulletin.* 2014;40:729–36.
20. Gayer-Anderson C, Jongsma HE, Di Forti M, Quattrone D, Velthorst E, de Haan L, et al. The European Network of National Schizophrenia networks studying gene-environment interactions (EU-GEI): incidence and first-episode case-control programme. *Social Psych Psych Epidemiol.* 2020;55:645–57.
21. Korver N, Quee PJ, Boos HB, Simons CJ, de Haan L, Investigators G. Genetic Risk and Outcome of Psychosis (GROUP), a multi site longitudinal cohort study focused on gene-environment interaction: objectives, sample characteristics, recruitment and assessment methods. *Int J Methods Psych Res.* 2012;21:205–21.
22. Blyler CR, Gold JM, Iannone VN, Buchanan RW. Short form of the WAIS-III for use with patients with schizophrenia. *Schizophrenia Res.* 2000;46:209–15.
23. Velthorst E, Levine SZ, Henquet C, de Haan L, van Os J, Myin-Germeys I, et al. To cut a short test even shorter: reliability and validity of a brief assessment of intellectual ability in schizophrenia—a control-case family study. *Cog Neuropsych.* 2013;18:574–93.
24. Mallet J. MRC sociodemographic schedule. Section of Social Psychiatry, Institute of Psychiatry; 1997.
25. WHO. Composite international diagnostic interview. Geneva, Switzerland: World Health Organization; 1990.
26. Di Forti M, Morgan C, Dazzan P, Pariante C, Mondelli V, Marques TR, et al. High-potency cannabis and the risk of psychosis. *Br J Psych.* 2009;195:488–91.
27. McGuffin P, Farmer A, Harvey I. A polydiagnostic application of operational criteria in studies of psychotic illness: development and reliability of the OPCRIT system. *Arch Gen Psych.* 1991;48:764–70.
28. Craddock N, Asherson P, Owen MJ, Williams J, McGuffin P, Farmer AE. Concurrent validity of the OPCRIT diagnostic system: comparison of OPCRIT diagnoses with consensus best-estimate lifetime diagnoses. *Br J Psych.* 1996;169:58–63.
29. Singh SP, Cooper JE, Fisher HL, Tarrant CJ, Lloyd T, Banjo J, et al. Determining the chronology and components of psychosis onset: the Nottingham Onset Schedule (NOS). *Schizophrenia Res.* 2005;80:117–30.
30. Jones SH, Thornicroft G, Coffey M, Dunn G. A brief mental health outcome scale: reliability and validity of the Global Assessment of Functioning (GAF). *Br J Psych.* 1995;166:654–9.
31. StataCorp. Stata Statistical Software: Release 15. College Station, TX: StataCorp; 2017. <https://www.stata.com/features/documentation/>. Accessed on 1 March 2018.
32. R-Core-Team. A language and environment for statistical computing. Vienna, Austria: R Foundation for Statistical Computing; 2014. <http://www.R-project.org>.
33. Wickham H. *ggplot2: Elegant graphics for data analysis*. New York: Springer-Verlag; 2016.
34. Zalesky A, Pantelis C, Cropley V, Fornito A, Cocchi L, McAdams H, et al. Delayed development of brain connectivity in adolescents with schizophrenia and their unaffected siblings. *JAMA Psychiatry.* 2015;72:900–8.
35. Flynn JR. *What is intelligence?: Beyond the Flynn effect*. Cambridge, UK: Cambridge University Press; 2007.
36. Salthouse TA. When does age-related cognitive decline begin? *Neurobiol Aging.* 2009;30:507–14.
37. Deary IJ, Corley J, Gow AJ, Harris SE, Houlihan LM, Marioni RE, et al. Age-associated cognitive decline. *Br Med Bull.* 2009;92:135–52.
38. Dutton E, van der Linden D, Lynn R. The negative Flynn Effect: a systematic literature review. *Intelligence.* 2016;59:163–9.
39. Teasdale TW, Owen DR. A long-term rise and recent decline in intelligence test performance: the Flynn Effect in reverse. *Pers Ind Diff.* 2005;39:837–43.
40. Bratsberg B, Rogeberg O. Flynn effect and its reversal are both environmentally caused. *Proc Nat Acad Sci.* 2018;115:6674–8.
41. Sachs GA, Carter R, Holtz LR, Smith F, Stump TE, Tu W, et al. Cognitive impairment: an independent predictor of excess mortality: a cohort study. *Ann Intern Med.* 2011;155:300–8.
42. Dickinson D. Digit symbol coding and general cognitive ability in schizophrenia: worth another look? *Br J Psychiatry.* 2008;193:354–6.
43. Smeland OB, Frei O, Kauppi K, Hill WD, Li W, Wang Y, et al. Identification of genetic loci jointly influencing schizophrenia risk and the cognitive traits of verbal-numerical reasoning, reaction time, and general cognitive function. *JAMA Psychiatry.* 2017;74:1065–75.
44. Shafee R, Nanda P, Padmanabhan JL, Tandon N, Alliey-Rodriguez N, Kalapurakkel S, et al. Polygenic risk for schizophrenia and measured domains of cognition in individuals with psychosis and controls. *Transl Psychiatry.* 2018;8:78.
45. Fett AK, Viechtbauer W, Dominguez MD, Penn DL, van Os J, Krabbendam L. The relationship between neurocognition and social cognition with functional outcomes in schizophrenia: a meta-analysis. *Neurosci Biobehav Rev.* 2011;35:573–88.
46. Gold JM. Cognitive deficits as treatment targets in schizophrenia. *Schizophr Res.* 2004;72:21–28.
47. Goldstein JM, Cherkertzian S, Tsuang MT, Petryshen TL. Sex differences in the genetic risk for schizophrenia: History of the evidence for sex-specific and sex-dependent effects. *Am J Med Gen Part B: Neuropsych Genetics.* 2013;162:698–710.
48. Loberg EM, Hugdahl K. Cannabis use and cognition in schizophrenia. *Front Hum Neurosci.* 2009;3:53.

49. van Winkel R, Investigators G. Further evidence that cannabis moderates familial correlation of psychosis-related experiences. *PLoS One*. 2015;10:e0137625.
50. Van Haren NEM, Van Dam DS, Stellato RK. Genetic risk and outcome of psychosis (group) investigators. Change in IQ in schizophrenia patients and their siblings: a controlled longitudinal study. *Psychol Med*. 2019;49:2573–81.
51. Knowles EEM, Mathias SR, Pearson GD, Barrett J, Mollon J, Denbow D, et al. Clinical correlates of subsyndromal depression in African American individuals with psychosis: The relationship with positive symptoms and comorbid substance dependence. *Schizophr Res*. 2019;206:333–46.

Affiliations

Eva Velthorst^{1,2,3} · Josephine Mollon⁴ · Robin M. Murray⁵ · Lieuwe de Haan^{3,6} · Inez Myin Germeys⁷ · David C. Glahn⁴ · Celso Arango⁸ · Els van der Ven^{9,10} · Marta Di Forti^{11,12} · Miguel Bernardo¹³ · Sinan Guloksuz^{10,14} · Philippe Delespaul^{10,15} · Gisela Mezquida¹³ · Silvia Amoretti¹³ · Julio Bobes¹⁶ · Pilar A. Saiz¹⁶ · María Paz García-Portilla¹⁶ · José Luis Santos^{16,17,18} · Estela Jiménez-López^{16,19} · Julio Sanjuan²⁰ · Eduardo J. Aguilar²⁰ · Manuel Arrojo²¹ · Angel Carracedo^{22,23} · Gonzalo López¹⁶ · Javier González-Peñas¹⁶ · Mara Parellada⁸ · Cem Atbaşoğlu²⁴ · Meram Can Saka²⁴ · Alp Üçok²⁵ · Köksal Alptekin^{26,27} · Berna Akdede²⁶ · Tolga Binbay²⁶ · Vesile Altınyazar²⁸ · Halis Ulaş²⁶ · Berna Yalınçetin²⁷ · Güvem Gümüş-Akay^{29,30} · Burçin Cihan Beyaz³¹ · Haldun Soygür³² · Eylem Şahin Cankurtaran³³ · Semra Ulusoy Kaymak³⁴ · Nadja P. Maric³⁵ · Marina M. Mihaljevic³⁵ · Sanja Andric Petrovic³⁶ · Tijana Mirjanic³⁷ · Cristina Marta Del-Ben³⁸ · Laura Ferraro⁵ · Charlotte Gayer-Anderson³⁹ · Peter B. Jones⁴⁰ · Hannah E. Jongsma^{40,41} · James B. Kirkbride⁴¹ · Caterina La Cascia⁴² · Antonio Lasalvia⁴³ · Sarah Tosato⁴³ · Pierre-Michel Llorca⁴⁴ · Paulo Rossi Menezes⁴⁵ · Craig Morgan^{39,46} · Diego Quattrone¹¹ · Marco Menchetti⁴⁷ · Jean-Paul Selten^{10,48} · Andrei Szöke⁴⁹ · Iliaria Tarricone⁵⁰ · Andrea Tortelli⁵¹ · Philip McGuire⁵ · Lucia Valmaggia⁵² · Matthew J. Kempton⁵ · Mark van der Gaag⁵³ · Anita Riecher-Rössler⁵⁴ · Rodrigo A. Bressan⁵⁵ · Neus Barrantes-Vidal^{56,57} · Barnaby Nelson^{58,59} · Patrick McGorry^{58,60} · Chris Pantelis⁶⁰ · Marie-Odile Krebs⁶¹ · Stephan Ruhrmann⁶² · Gabriele Sachs⁶³ · Bart P. F. Rutten¹⁰ · Jim van Os^{5,10,64} · Behrooz Z. Alizadeh^{65,66} · Therese van Amelsvoort¹⁰ · Agna A. Bartels-Velthuis⁶⁶ · Richard Bruggeman^{66,67} · Nico J. van Beveren^{68,69,70} · Jurjen J. Luyckx^{64,71,72} · Wiepke Cahn^{64,73} · Claudia J. P. Simons^{10,74} · Rene S. Kahn^{1,75} · Frederike Schirmbeck^{3,6} · Ruud van Winkel^{10,76} · EU-GEI High Risk Study · Abraham Reichenberg^{1,2}

¹ Department of Psychiatry, Icahn School of Medicine at Mount Sinai, New York, NY, USA

² Seaver Center for Research and Treatment, Icahn School of Medicine at Mount Sinai, New York, NY, USA

³ Department of Psychiatry, Amsterdam UMC, Amsterdam, The Netherlands

⁴ Department of Psychiatry, Boston Children's Hospital, Harvard Medical School, Boston, MA, USA

⁵ Department of Psychosis Studies, Institute of Psychiatry, Psychology and Neuroscience, King's College London, London, England, UK

⁶ Arkin Institute for Mental Health, Amsterdam, The Netherlands

⁷ Department of Neurosciences, Psychiatry Research Group, Center for Contextual Psychiatry, KU Leuven, Leuven, Belgium

⁸ Department of Child and Adolescent Psychiatry, Institute of Psychiatry and Mental Health, Hospital General Universitario Gregorio Marañón, School of Medicine, Universidad Complutense, Instituto de Investigación Sanitaria Gregorio Marañón (IiSGM), Spanish Mental Health Research Network (CIBERSAM), Madrid, Spain

⁹ Mailman School of Public Health, Columbia University, New York, NY, USA

¹⁰ Department of Psychiatry and Neuropsychology, School for Mental Health and Neuroscience, Maastricht University Medical Centre, Maastricht, The Netherlands

¹¹ Department of Social Genetics and Developmental Psychiatry, Institute of Psychiatry, Psychology and Neuroscience, King's College London, London, England, UK

¹² South London and Maudsley NHS Mental Health Foundation Trust, London, UK

¹³ Barcelona Clinic Schizophrenia Unit, Neuroscience Institute, Hospital Clinic of Barcelona, University of Barcelona, IDIBAPS, Spanish Mental Health Research Network (CIBERSAM), Barcelona, Spain

¹⁴ Department of Psychiatry, Yale School of Medicine, New Haven, CT, USA

¹⁵ Mondriaan Mental Health Care, Heerlen/Maastricht, The Netherlands

¹⁶ Faculty of Medicine and Health Sciences - Psychiatry, Universidad de Oviedo, Mental Health Services of Principado de Asturias (SESPA), Instituto de Investigación Sanitaria del Principado de Asturias (ISPA), INEUROPA, CIBERSAM, Oviedo, Spain

¹⁷ Neurobiological Research Group, Institute of Technology, Universidad de Castilla-La Mancha, Cuenca, Spain

- 18 Department of Psychiatry, Hospital “Virgen de la Luz”, Cuenca, Spain
- 19 Universidad de Castilla-La Mancha, Health and Social Research Center, Cuenca, Spain
- 20 Department of Psychiatry, Hospital Clínico Universitario de Valencia, INCLIVA, CIBERSAM, School of Medicine, Universidad de Valencia, Valencia, Spain
- 21 Department of Psychiatry, Instituto de Investigación Sanitaria (IDIS), Complejo Hospitalario Universitario de Santiago de Compostela, Santiago de Compostela, Spain
- 22 Grupo de Medicina Xenómica, Fundación Pública Galega de Medicina Xenómica, Instituto de Investigación Sanitaria de Santiago de Compostela (IDIS), Galician Health Service (SERGAS), Santiago de Compostela, Spain
- 23 Centro de Investigación en Red de Enfermedades Raras (CIBERER) y Centro Nacional de Genotipado (CEGEN-PRB3), Center for Research in Molecular Medicine and Chronic Diseases (CIMUS), Universidad de Santiago de Compostela, Santiago de Compostela, Spain
- 24 Department of Psychiatry, School of Medicine, Ankara University, Ankara, Turkey
- 25 Department of Psychiatry, Faculty of Medicine, Istanbul University, Istanbul, Turkey
- 26 Department of Psychiatry, Dokuz Eylül University, School of Medicine, Izmir, Turkey
- 27 Department of Neuroscience, Health Sciences Institute, Dokuz Eylül University, Izmir, Turkey
- 28 Department of Psychiatry, Adnan Menderes University, School of Medicine, Aydın, Turkey
- 29 Department of Physiology, School of Medicine, Ankara University, Ankara, Turkey
- 30 Brain Research Center, Ankara University, Ankara, Turkey
- 31 Department of Psychology, Middle East Technical University, Ankara, Turkey
- 32 Turkish Federation of Schizophrenia Associations, Ankara, Turkey
- 33 Private Practice, Ankara, Turkey
- 34 Ankara City Hospital Psychiatry Clinic, Ankara, Turkey
- 35 Faculty of Medicine, University of Belgrade & Institute of Mental Health, Belgrade, Serbia
- 36 Faculty of Medicine, University of Belgrade & Clinic for Psychiatry, Belgrade, Serbia
- 37 Special Hospital for Psychiatric Disorders Kovin, Kovin, Serbia
- 38 Ribeirão Preto Medical School, University of São Paulo, São Paulo, Brazil
- 39 Department of Health Service and Population Research, Institute of Psychiatry, King’s College London, London, UK
- 40 Department of Psychiatry, University of Cambridge, Cambridge, England, UK
- 41 PsyLife Group, Division of Psychiatry, UCL, London, England, UK
- 42 Department of Biomedicine, Neuroscience and Advanced Diagnostics, University of Palermo, Palermo, Italy
- 43 Section of Psychiatry, Department of Neuroscience, Biomedicine and Movement, University of Verona, Verona, Italy
- 44 Centre Hospitalier Universitaire de Clermont-Ferrand, Clermont-Ferrand, France
- 45 Department of Preventive Medicine, Faculdade de Medicina, Universidade of São Paulo, São Paulo, Brazil
- 46 ESRC Centre for Society and Mental Health, King’s College London, London, UK
- 47 Department of Biomedical and Neuromotor Sciences, University of Bologna, Bologna, Italy
- 48 Rivierduinen Institute for Mental Health Care, Leiden, The Netherlands
- 49 INSERM, U955 Créteil, France
- 50 Department of Medical and Surgical Sciences, Psychiatry Unit, Alma Mater Studiorum Università di Bologna, Bologna, Italy
- 51 Etablissement Public de Santé Maison Blanche, Paris, France
- 52 Department of Psychology, Institute of Psychiatry, Psychology and Neuroscience, King’s College London, London, UK
- 53 Amsterdam Public Mental Health Research Institute, Department of Clinical Psychology, Faculty of Behavioural and Movement Sciences, Vrije Universiteit Amsterdam, Amsterdam, The Netherlands
- 54 Medizinische Fakultät, Universität Basel, Basel, Switzerland
- 55 LiNC-Lab Interdisciplinar Neurociências Clínicas, Depto Psiquiatria, Escola Paulista de Medicina, Universidade Federal de São Paulo (UNIFESP), São Paulo, Brazil
- 56 Departament de Psicologia Clínica i de la Salut, Universitat Autònoma de Barcelona, Barcelona, Spain
- 57 Fundació Sanitària Sant Pere Claver, Spanish Mental Health Research Network (CIBERSAM), Barcelona, Spain
- 58 Orygen, Parkville, VIC, Australia
- 59 Centre for Youth Mental Health, The University of Melbourne, Parkville, VIC, Australia
- 60 Melbourne Neuropsychiatry Centre, Department of Psychiatry, University of Melbourne & Melbourne Health, Carlton South, VIC, Australia
- 61 University of Paris, GHU-Paris, Sainte-Anne, C’JAAD, Inserm U1266, Institut de Psychiatrie (CNRS 3557), Paris, France
- 62 Department of Psychiatry and Psychotherapy, Faculty of Medicine and University Hospital, University of Cologne, Cologne, Germany
- 63 Department of Psychiatry and Psychotherapy, Medical University of Vienna, Vienna, Austria

- ⁶⁴ Department of Psychiatry, Brain Centre Rudolf Magnus, University Medical Center Utrecht, Utrecht University, Utrecht, The Netherlands
- ⁶⁵ Department of Epidemiology, University of Groningen, University Medical Center Groningen, Groningen, The Netherlands
- ⁶⁶ University of Groningen, University Medical Center Groningen, University Center for Psychiatry, Rob Giel Research Center, Groningen, The Netherlands
- ⁶⁷ Department of Clinical and Developmental Neuropsychology, University of Groningen, Groningen, The Netherlands
- ⁶⁸ Antes Center for Mental Health Care, Rotterdam, The Netherlands
- ⁶⁹ Department of Psychiatry, Erasmus MC, Rotterdam, The Netherlands
- ⁷⁰ Department of Neuroscience, Erasmus MC, Rotterdam, The Netherlands
- ⁷¹ Department of Translational Neuroscience, UMC Utrecht Brain Center, University Medical Center Utrecht, Utrecht University, Utrecht, The Netherlands
- ⁷² Outpatient Second Opinion Clinic, GGNet Mental Health, Apeldoorn, The Netherlands
- ⁷³ Altrecht Science, Altrecht Mental Health Care, Utrecht, The Netherlands
- ⁷⁴ GGzE Institute for Mental Health Care, Eindhoven, The Netherlands
- ⁷⁵ VISN 2 Mental Illness Research, Education and Clinical Center (MIRECC), James J. Peters Department of Veterans Affairs Medical Center, New York, NY, USA
- ⁷⁶ Department of Neurosciences, Psychiatry Research Group, Center for Clinical Psychiatry, KU Leuven, Leuven, Belgium