

REVIEW ARTICLE

Open Access

Establishing a clinical service to prevent psychosis: What, how and when? Systematic review

Gonzalo Salazar de Pablo^{1,2}, Andrés Estradé^{1,3}, Marcello Cutroni⁴, Olivier Andlauer^{5,6} and Paolo Fusar-Poli^{1,4,7,8}

Abstract

The first rate-limiting step to successfully translate prevention of psychosis in to clinical practice is to establish specialised Clinical High Risk for Psychosis (CHR-P) services. This study systematises the knowledge regarding CHR-P services and provides guidelines for translational implementation. We conducted a PRISMA/MOOSE-compliant (PROSPERO-CRD42020163640) systematic review of Web of Science to identify studies until 4/05/2020 reporting on CHR-P service configuration, outreach strategy and referrals, service user characteristics, interventions, and outcomes. Fifty-six studies (1998–2020) were included, encompassing 51 distinct CHR-P services across 15 countries and a catchment area of 17,252,666 people. Most services (80.4%) consisted of integrated multidisciplinary teams taking care of CHR-P and other patients. Outreach encompassed active (up to 97.6%) or passive (up to 63.4%) approaches: referrals came mostly (90%) from healthcare agencies. CHR-P individuals were more frequently males (57.2%). Most (70.6%) services accepted individuals aged 12–35 years, typically assessed with the CAARMS/SIPS (83.7%). Baseline comorbid mental conditions were reported in two-third (69.5%) of cases, and unemployment in one third (36.6%). Most services provided up to 2-years (72.4%), of clinical monitoring (100%), psychoeducation (81.1%), psychosocial support (73%), family interventions (73%), individual (67.6%) and group (18.9%) psychotherapy, physical health interventions (37.8%), antipsychotics (87.1%), antidepressants (74.2%), anxiolytics (51.6%), and mood stabilisers (38.7%). Outcomes were more frequently ascertained clinically (93.0%) and included: persistence of symptoms/comorbidities (67.4%), transition to psychosis (53.5%), and functional status (48.8%). We provide ten practical recommendations for implementation of CHR-P services. Health service knowledge summarised by the current study will facilitate translational efforts for implementation of CHR-P services worldwide.

Introduction

The clinical high risk for psychosis (CHR-P) paradigm¹ represents one of the most established preventive approaches in clinical psychiatry². It originated in Australia around 25 years ago³ and since then, it has progressively gained importance⁴. CHR-P individuals

are young and accumulate risk factors for the disorders^{5–7}, that lead to functional impairments⁸ and attenuated psychotic symptoms⁹. Because of these features, these individuals seek help¹⁰ at specialised CHR-P mental health services. The detection¹¹, prognostic assessment¹² and preventive treatment^{13–16} in CHR-P individuals¹⁵ have the potential to maximize the benefits of early interventions in psychosis^{17,18}. A recent evidence-based summary by the European College of Neuropsychopharmacology Network for the Prevention of Mental Disorders and Mental Health Promotion¹⁹ indicated that the first rate-limiting step to prevent psychosis is to establish specialised CHR-P services²⁰. Accordingly, several CHR-P services have been implemented worldwide, as recently mapped by the

Correspondence: Paolo Fusar-Poli (paolo.fusar-poli@kcl.ac.uk)

¹Early Psychosis: Interventions and Clinical-detection (EPIC) Lab, Department of Psychosis Studies, Institute of Psychiatry, Psychology & Neuroscience, King's College London, London, UK

²Institute of Psychiatry and Mental Health, Department of Child and Adolescent Psychiatry, Hospital General Universitario Gregorio Marañón School of Medicine, Universidad Complutense, Instituto de Investigación Sanitaria Gregorio Marañón (IISGM), CIBERSAM, Madrid, Spain

Full list of author information is available at the end of the article

These authors contributed equally: Gonzalo Salazar de Pablo, Andrés Estradé

Endnote reference file: CHR-P clinical services.enl

© The Author(s) 2021



Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons license and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>.

International Early Psychosis Association (IEPA²¹: <https://iepa.org.au/list-a-service>).

Despite these progresses, health service research in this field has been fragmented to the point that the characteristics (“what”) of a CHR-P service per se are poorly defined. As CHR-P services expand globally²¹, it becomes essential to synthesize the core CHR-P health service features that have been implemented in real-world scenarios. While a CHR-P clinic can be broadly defined as a “multidisciplinary community mental health service that provides treatment and support to people at high risk of developing psychosis” (page 16 from NHS England²²), this definition remains elusive. Similarly, there is no clear guidance on “how” to integrate different service components. The three main models for delivering CHR-P services include the “stand-alone”, “hub and spoke”, and “integrated” models²². While the standalone model works independently from other more generic community mental health teams, in the “hub and spoke” model, dedicated team workers (“spokes”) are based within more generic community teams to route patients needing more intensive services to the central “hub”²³. In an integrated model, the CHR-P service is completely integrated into the community mental health care. In addition, these models can be combined within broad mental health services enhancing transitional primary care platforms across adolescents and young adults²⁴. The additional limitation of knowledge is that the timing (“when”) for preventive approaches, which is reflected by CHR-P entry age criteria is uncertain. While this has been typically set for young people aged 8–40⁴ years, more recent lifespan-inclusive approaches for those under the age of 25 (0–25 years)²⁵ models have been piloted.

While previous systematic reviews have addressed these issues for services taking care of patients with a first episode of psychosis^{26,27}, CHR-P research has remained mostly “academic” and did not systematically address real-world service characteristics such as: service configuration, outreach strategy and referrals, service user characteristics, interventions, and outcomes. The current systematic review summarizes, for the first time, evidence on these domains to inform the real-world implementation (i.e., what, how, and when) of CHR-P clinical services worldwide.

Methods

This study (study protocol: PROSPERO CRD42020163640) was conducted in accordance with PRISMA²⁸ (eTable 1) checklist.

Search strategy and selection criteria

A multistep systematic literature search strategy was used to identify relevant articles by two independent researchers (GSP, AE). First, the Web of Science database (Clarivate

Analytics) was searched, incorporating the Web of Science Core Collection, BIOSIS Citation Index, KCI-Korean Journal Database, MEDLINE, Russian Science Citation Index, and SciELO Citation Index as well as Cochrane Central Register of Reviews, and Ovid/PsychINFO databases, as well as the OpenGrey database (for grey literature) from inception until 4th May 2020, with no restrictions on language. The following search terms were applied: (“risk” OR “prodrom*” OR “ultra-high risk” OR “clinical high risk” OR “attenuat*” OR “high risk” OR “genetic high risk” OR “risk syndrome” OR “at risk mental state” OR “at-risk mental state” OR “ARMS” OR “risk of progression” OR “schizophrenia” OR “schizoaffective disorder” OR “schizophreniform disorder”) AND (“psychosis”) AND (“prevention” OR “intervention” OR “early intervention” OR “referral” OR “assessment” OR “service” OR “clinical service” OR “psychiatric service” OR “implementation” OR “care pathways”). The references of the articles identified in previous reviews and relevant commentaries and the references from the included studies were manually searched to identify additional relevant records. Abstracts were screened, and potential full texts were assessed against inclusion and exclusion criteria.

The inclusion criteria were a) being an original study published in international databases or in the grey literature, b) describing clinical services for individuals in a CHR-P state as defined according to established instruments: Comprehensive Assessment of At-Risk Mental States (CAARMS³), Structured Interview for Psychosis-risk Syndromes (SIPS^{29,30}), Bonn Scale for the Assessment of Basic Symptoms (BSABS³¹), Basel Screening Instrument for Psychosis (BSIP³²), Schizophrenia Proneness Instrument³³ - Adult (SPI-A) and Child and Youth (SPI-CY) version -, Positive and Negative Syndrome Scale (PANSS³⁴), Scale for the Assessment of Negative Symptoms (SANS³⁵), Brief Psychiatric Rating Scale (BPRS³⁶) and Early Recognition Inventory (ERIRAOS³⁷), c) providing information on any of the following: service configuration, outreach strategy and referrals, service user characteristics, interventions and outcomes, d) providing relevant information without any restrictions on language, sex, age, or ethnicity. The exclusion criteria were a) non-original studies such as abstracts, conference proceedings, study protocols, reviews, guidelines, b) studies with a primary research focus (e.g., research networks) and lacking description of CHR-P clinical services, c) studies describing clinical services for conditions other than the CHR-P or services without a CHR-P component, d) national or regional survey studies with aggregate data and lacking a service-specific description.

Descriptive measures and data extraction

Independent researchers (GSP, AE, MC) extracted data from the included studies; discrepancies were resolved

through consensus, consulting a senior researcher (PFP). The variables included were those necessary to describe “what, how and when” to implement CHR-P services. These variables were grouped according to health service domains previously established (beyond general data such as first author, year of publication, name of the CHR-P service, country)³⁸: (i) service configuration: continent, service set-up date, population in the catchment area, type of service, professionals involved, (ii) outreach strategy and referrals: outreach activities—measured using an adapted version of the Longitudinal Youth-At-Risk Study (LYRIKS) study³⁹ classification—, referral sources (iii) service user characteristics: sociodemographic characteristics, CHR-P assessment, CHR-P subgroups (defined as in previous studies)⁴⁰, minimum and maximum age inclusion criteria and service use age range, comorbidities and employment (iv) interventions: type of intervention (non-pharmacological vs. psychopharmacological) and duration of service provision and (v) outcomes: type of outcomes monitored and outcome instruments. Furthermore, we reported quality assessment (see below).

Data analysis

Systematic review

All the studies were systematically summarized in tables reporting on various health service domains: service configuration, outreach strategy, and referrals (Table 1), service user characteristics (Table 2), interventions and outcomes (Table 3). We complement this with descriptive analysis of common operational and clinical challenges. An online tool (<https://www.maptive.com>) was used to create a graphical representation of the geographical distribution of the CHR-P services included in the review.

Quality assessment

We adapted the mixed Methods Appraisal Tool (MMAT)^{41,42} questions for non-randomized clinical studies due to the heterogeneity expected in the included studies to assess the quality of the included studies (eMethods 1), considering the content and characteristics of the studies according to our inclusion criteria.

Results

Database

The literature search yielded 12,130 citations, which were screened for eligibility. Two hundred and twenty-one full-text articles were evaluated for eligibility, and 165 were excluded. In total, 49 studies reporting information on individual CHR-P services (eTable 2), and seven multisite studies (eTable 3) were selected (PRISMA, Fig. 1). All CHR-P services (100%) used validated assessment instruments and no studies were excluded for this reason. The final pool of 56 included studies were published between the years 1998 and 2020. The total sample

Table 1 Service configuration (above); outreach strategy and referrals (below).

	Number of services (%)
Service configuration	
Continent	51
Europe	30 (58.8)
North America	13 (25.5)
Australia	4 (7.8)
Asia	3 (5.9)
South America	1 (2.0)
Service set-up date	50
1991–1999	6 (12.0)
2000–2009	31 (62.0)
2010–2019	13 (26.0)
Population in the catchment area	35
Combined total population	17,252,666
Average total pop. per service	492,933
Type of service	51
Integrated CHR-P service	41 (80.4)
Standalone CHR-P service	10 (19.6)
Hub and spoke CHR-P service	0
Professionals involved (not mutually exclusive)	30
Psychiatrist ^a	30 (100)
Clinical psychologist or counsellor ^b	23 (76.7)
Case manager/care coordinator ^c	15 (50.0)
Nurse ^d	15 (50.0)
Occupational therapist/social worker/educator	12 (40.0)
Research personnel ^e	3 (10.0)
Neuropsychologist	2 (6.7)
General practitioner	1 (3.3)
Exercise physiologist	1 (3.3)
Outreach strategy and referrals	
Outreach activities (not mutually exclusive)	41
Active approaches	
Workshops	
General workshops	40 (97.6)
Targeting healthcare professionals	35 (85.4)
Targeting education professionals ^f	20 (48.8)
Targeting community organisations	14 (34.1)

Table 1 continued

	Number of services (%)
Service promotion to NGOs and community services ^g	14 (34.1)
Service promotion to social and governmental services ^h	8 (19.5)
Service promotion to family members	2 (4.9)
General public awareness campaigns	15 (36.6)
Passive approaches	
Dedicated online site	23 (63.4)
Print and other media ⁱ	22 (53.7)
Referral sources (not mutually exclusive)	40
Outpatient or community mental health services ^j	36 (90.0)
General healthcare ^k	30 (75.0)
Education organisations or services ^l	26 (65.0)
Self	24 (60.0)
Family, relatives or friends	24 (60.0)
Inpatient mental health services ^m	17 (42.5)
A&E departments	9 (22.5)
Social services & welfare	7 (17.5)
Government organisations ⁿ	6 (15.0)
Community organisations ^o	5 (12.5)
Early Intervention for Psychosis services	5 (12.5)

For footnotes see the supplementary section (eResults 1).

A&E Accident and emergency Departments, NGO non-governmental organization.

encompassed 51 distinct CHR-P clinical services, from 41 different regions, across 15 countries (Fig. 2).

Most multisite studies reported on collaborative networks of clinical CHR-P services, including the Pan-London Network for Psychosis-Prevention (PNP³⁸), the Early Detection and Intervention for the Prevention of Psychosis Program (EDIPPP^{43–45}), and the Swiss Early Psychosis Project (SWEPP⁴⁶). Two additional multisite studies report on five centres operating under the Italian Departments⁴⁷ and six CHR-P services in Canada⁴⁸.

Service configuration

The CHR-P services were located mostly in Europe (58.8%), followed by North America (25.5%), Australia (7.8%), Asia (5.9%), and South America (2.0%; Fig. 2 and Table 1). The first program to be implemented was the Personal Assessment and Crisis Evaluation (PACE) clinic in 1994, in Melbourne⁴⁹, and the most recent one the City & Hackney At-Risk Mental State Service (HEADS UP) in

Table 2 Service user characteristics.

	Number of services (%)
Sociodemographic characteristics	
Sample size	33
<50	12 (39.4)
50–100	6 (18.2)
>100	15 (45.5)
Sex	43 (CHR-P individuals)
Male (frequency %)	57.2
CHR-P Assessment (not mutually exclusive)	37
CAARMS	18 (48.6)
SIPS	13 (35.1)
ERlraos-CL	6 (16.2)
BSIP	2 (5.4)
BSABS/SPI-A/SPI-CY	1 (2.7)
Min. age inclusion criteria	48
Between 8 and 6 years	2 (4.2)
12 years	12 (25)
Between 13 and 15 years	15 (31.3)
Between 16 and 17 years	13 (27.1)
18 years or older	6 (12.5)
Max. age inclusion criteria	49
18 years	4 (8.2)
Between 24 and 29 years	14 (28.6)
Between 30 and 35 years	24 (49.0)
Between 40 and 56 years	4 (8.2)
65 years or older	3 (6.1)
Service users age range	51
Children and adolescents only (<18)	2 (3.9)
Adolescents only (12–18)	2 (3.9)
Adolescents and young adults (12–35)	36 (70.6)
Children, adolescents and adults (8–40)	1 (2.0)
Adolescents and adults (≥12) ^a	5 (9.8)
Young adults (18–35)	4 (7.8)
Adults only (≥18)	1 (2.0)
	CHR-P individuals (%)
Diagnostic subgroups (not mutually exclusive)	17 ^b
APS	82.6
BLIPS	10.7
GRD	8.5

Table 2 continued

	Number of services (%)
Comorbidity (not mutually exclusive) ^c	31 ^b
Any DSM/ICD comorbid disorder	69.5
Depressive disorders	42.3
Bipolar disorder	15.5
Persistent depressive disorder	6.7
History of suicide attempts	10.5
Anxiety disorders	24.1
Social phobia	5.9
Obsessive compulsive disorder	5.1
Adjustment disorder	11.6
Any personality disorder	15.5
Schizotypal personality disorder	11.0
Substance use disorders	12.4
Employment	13 ^b
Unemployment rates	36.6

^aIncludes two services that enrolled “adolescents and adults” without further specification.

^bNumber of services providing data for the service user characteristics as % CHR-P individuals.

^cDiagnosis according to DSM or ICD criteria established using either structured interviews or clinical interviews.

APS attenuated psychosis symptoms, BLIPS brief limited intermittent psychotic symptoms, BSABS Bonn scale for the assessment of basic symptoms, BSIP Basel screening instrument for psychosis, CAARMS comprehensive assessment of at-risk mental states, ERIRAS-CL early recognition inventory retrospective assessment of symptoms checklist, GRD genetic risk and deterioration, SIPS structured interview for psychosis-risk syndromes, SPI-A schizophrenia proneness instrument (adults version), SPI-CY schizophrenia proneness instrument (child and youth version).

2015, in London³⁸. 62.0% CHR-P services were set up from 2000 to 2009 (Table 1). Population in the catchment area covered a total of 17,252,666 people, with an average of 492,933 people (SD: 396,997, Table 1). Most services (80.4%) consisted of teams integrated into the community mental health care. Standalone CHR-P services were less frequent (19.6%) and there were no hub and spoke services. CHR-P clinical services involve a wide range of professionals, the most frequent ones being psychiatrists (100%), who were involved in all the services. Other professionals include clinical psychologists or counsellors (76.7%), case managers/care coordinators (50%), and nurses (50%).

Outreach strategy and referrals

Outreach activities and audiences were highly variable (Table 1). Within active strategies, workshops for referral sources were the most frequent (97.6%), often targeting healthcare professionals (85.4%), educational professionals (48.8%), or community organisations (34.1%). Services also approached NGOs and community services

Table 3 Interventions (above) and outcomes (below).

	Number of services (%)
Interventions	
Non-pharmacological interventions (not mutually exclusive)	37
Clinical monitoring	37 (100.0)
Psychoeducation	30 (81.1%)
Case management and psychosocial support ^a	27 (73.0)
Family intervention ^b	27 (73.0)
CBT-based individual intervention ^c	25 (67.6)
Other individual psychotherapeutic intervention ^d	24 (64.9)
Physical health interventions ^e	14 (37.8)
Group social or therapeutic interventions ^f	7 (18.9)
Pharmacological interventions (not mutually exclusive)	31
Antipsychotic medication	27 (87.1)
Antidepressants	23 (74.2)
Anxiolytics	16 (51.6)
Mood stabilizers	12 (38.7)
Omega-3 fatty acids	3 (9.7)
Duration of service provision	29
6 months	1 (3.4)
12 months	7 (24.1)
24 months	13 (44.8)
36 months	3 (10.3)
60 months or more	5 (17.2)
Outcomes	
Type of outcomes (not mutually exclusive)	43
Persistence of symptoms/comorbidities	29 (67.4)
Transition to psychosis	23 (53.5)
Functional status	21 (48.8)
Remission	18 (41.9)
Physical health outcomes	13 (30.2)
Service users' satisfaction	11 (25.6)
Hospitalisation	8 (18.6)
Mortality	6 (13.9)
Outcome instruments (not mutually exclusive)	43
Clinical interviews	40 (93.0)

Table 3 continued

	Number of services (%)
Psychometric instruments	16 (37.2)
CAARMS	10 (23.2)
SIPS	5 (11.6)
Electronic health records	7 (16.3)

For footnotes see the supplementary section (eResults 2).

(34.1%), and less frequently social and governmental services (19.5%) and family members of mental health patients (4.9%). About one-third of CHR-P services (36.6%) implemented general public awareness campaigns including TV or radio appearances^{50,51}, theatre adverts, high school art contests, and sponsors for minor league sports teams⁴⁵. More than half of services (63.4%) implemented either a dedicated online site for service promotion, or have elaborated printed and other media materials (53.7%), such as information brochures and leaflets^{50–52}, posters⁵³, articles in professional journals and local newspapers^{51,54}, presentations in scientific conferences⁴⁶, newsletters⁵¹, and promotional videos^{43,55}. Most CHR-P services received young people with a putative risk of psychosis from health-related organizations, including both outpatient or community mental health services (90.0%) and general healthcare services (75.0%). Education organisations are also frequent referral sources (65.0%), followed by self (60.0%), family or relatives (60.0%), inpatient mental health services (42.5%), and accident and emergency departments (22.5%). Other referral sources were reported in less than 20% of CHR-P services.

Service user characteristics

The total sample size of service users was of 5637 CHR-P individuals, ranging from 4⁵⁶ to 467⁵⁷ individuals: most of them were males (% of CHR-P females = 42.8, see Table 2).

CHR-P status was most frequently assessed using the Comprehensive Assessment of At-Risk Mental States (CAARMS) (48.6%), followed by the Structured Interview for Psychosis-risk Syndromes (SIPS) (35.1%) and the Early Recognition Inventory retrospective assessment of symptoms checklist (ERiraos-CL) (16.2%; Table 2). The Basel screening instrument for psychosis (BSIP) and basic symptoms instruments were infrequently used (<6% of services). Most services provided treatment starting in adolescence, from the ages 12 to 17 (83.4%). The most frequent minimum inclusion age range was 13–15 years (31.3%); only two services reported the inclusion of

children from the age of 8 and 6^{46,48,58}. Most services accepted users until 30–35 years (49.0%) or 24–29 years (28.6%). A few services (6.1%) accepted service-users of 65 years or older^{56,59}. The most frequent age range (70.6% of services) was 12 and 35 years.

82.6% of the CHR-P individuals fulfilled APS criteria, 10.7% fulfilled BLIPS criteria and 8.5% fulfilled GRD criteria (not mutually exclusive). Baseline comorbid mental disorders were reported in 69.5% CHR-P individuals. Mood disorders were the most common: depressive disorders (42.3%), bipolar disorders (15.5%), and persistent depressive disorder (6.7%). Anxiety disorders were also frequent (24.1%), including social phobia (5.9%) and obsessive–compulsive disorder (OCD) (5.1%). Adjustment disorder appeared in 11.6% of CHR-P individuals. Comorbid personality disorders were present in 15.5% of CHR-P individuals, particularly schizotypal personality disorder (11.0%). Substance use disorders were present in 12.4% of CHR-P individuals. Past history of suicide attempts was present in 10.5% of CHR-P subjects. Unemployment rate (i.e., neither work nor study) was observed in 36.6% in CHR-P individuals in clinical services.

Interventions

Across non-pharmacological interventions, clinical monitoring was the most common intervention and was carried out in all the services (100%). Other common interventions were psychoeducation (81.1%), case management and psychosocial support (73.0%) and family interventions (73.0%) (Table 3). Cognitive behavioural therapy-based interventions and any other type of individual psychotherapeutic intervention (encompassing individual motivational interviewing⁶⁰ or sessions⁴⁷, supportive counselling^{38,44,47,49,52,61–71}, relaxation training⁶¹, cognitive remediation⁵⁵, solution focused brief therapy⁵⁵, social skills training^{65,70}, substance misuse work⁷⁰, and psychotherapy NOS^{72,73}) were provided by about two-thirds of the services (67.6% and 64.9%, respectively). Physical health interventions and group psychotherapy sessions were more infrequent (37.8% and 18.9%, respectively).

Most (87.1%) services employed low-dose antipsychotic (AP) medication, although not as the first-line intervention but only following worsening of symptoms or functioning^{52,74,75}. Other interventions employed by CHR-P services included antidepressants (74.2%), anxiolytics (51.6%), mood stabilisers (38.7%). Three services (9.7%) reported the use of omega-3 fatty acids^{38,66,71,76,77}. Most services provided care for 24 months (44.8%) or 12 months (24.1%). Three services (10.3%) provided 36 months of clinical follow-up. Extended service provision of 60 months or more were reported in 17.2% of services.





Table 4 Ten simple recommendations for real-world implementation of CHR-P service.

Service configuration	
1	Implement a standalone community service (“what”)
2	Train a multidisciplinary team (psychiatrists, clinical psychologists or counsellors, case managers and nurses) (“what”)
Outreach strategy and referrals	
3	Adopt active and passive outreach, primarily targeting healthcare agencies (“how”)
4	Ensure adequate risk enrichment during the recruitment (“how”)
CHR-P service user characteristics	
5	Define CHR-P through established psychometric instruments (not in general population) (“how”)
6	Implement a transitional and transdiagnostic service across adolescents and young adults (“when”)
Interventions	
7	Offer needs-based interventions and psychological interventions (“how”)
8	Titrate the intervention according to the characteristics and risk profile ^a as well as the values and preferences of the individuals (“how”)
Outcomes	
9	Collect information and target recovery, physical health outcomes, service users’ satisfaction, functioning and quality of life (“how”)
10	Extend clinical monitoring for outcomes for at least three years (“how”)

^aCHR-P subgroups BLIPS > APS > GRD, severity of attenuated positive and negative symptoms, and level of functioning.

Outcomes

The outcomes most frequently evaluated in the CHR-P services were persistence of symptoms/comorbidities (67.4%), transition to psychosis (53.5%), functional status (48.8%), and remission (41.9%). Physical health outcomes (30.2%), service users’ satisfaction (25.6%), hospitalisation (18.6%) and mortality (13.9%) were less frequently evaluated.

Outcomes were most commonly evaluated with standard clinical interviews with the service users (93.0%), and more infrequently with psychometric instruments (37.2%). In the latter case, the CAARMS (23.2%) and SIPS (11.6%) were more frequently employed. About 16.3% of CHR-P services evaluated outcomes via electronic health records.

Quality of the included studies

Study quality scores ranged from 1 to 5. The overall mean quality score for included studies reporting on individual services was 3.8 (moderately high quality) on the MMAT scale, with a SD of 1.3 (eTable 2).

Discussion

To our knowledge, this is the first systematic review to comprehensively summarize the evidence from real-world implementation of CHR-P clinical services. This review encompasses 56 studies describing a total of 51 services for CHR-P individuals.

Consistent with recent surveys of CHR-P services^{21,47,78}, there is great diversity in how clinical services have been implemented in real-world scenarios, across all aspects of

service delivery: 1) service configuration, 2) outreach strategy and referrals, 3) service user characteristics, 4) interventions and 5) outcomes. We discuss these points, while also mentioning common challenges. The evidence summarised will then be used to operationalise ten empirical recommendations for overcoming these challenges and facilitating the real-world implementation of CHR-P services (Table 4).

In terms of service configuration, several CHR-P clinical services have been implemented across—at least—15 countries (Fig. 2), covering a catchment area of over 17 M people. Following a period of rapid expansion (2000–2009), new CHR-P services continue to emerge^{38,76}. At present, CHR-P services spread across most continents²¹, although they are mostly established in high-income countries. While most CHR-P services are configured as integrated services (80.4%), standalone models of care (19.6%) seem to be associated with high levels of service efficiency²⁷. For example, CHR-P standalone services had dropout rates in the range of 12–19.2%^{50,77,79–82} compared to 25.4% in integrated services⁵². One possible explanation is that in integrated models of care, healthcare resources are typically diverted towards more severe service users (e.g., first-episode vs. CHR-P patients)³⁸. In line with this notion, the actual caseload of CHR-P individuals was minimal ($n = 4$ out of 239 clients) in some integrated services⁵⁶, and more severe patients had more frequent contacts with these services⁷³. Another issue is that standalone services may be physically located outside general psychiatric services, which is preferable to reduce stigmatisation risks^{77,81}.

Service users and their relatives were generally more satisfied with standalone CHR-P services, particularly with clinical contact being outside traditional mental health settings⁶⁷. Conversely, family disengagement was the most significant barrier (71.4%) in integrated services⁶⁰. Likewise, primary care clinicians favoured standalone models of care because of the superior accessibility of the services⁶⁷. Standalone services are more costly to set up in the first year but deliver highest economic savings in the longer term⁶⁹, mainly associated with the improved outcome of the disorder⁶⁸. These considerations are of relevance given that poor financial support and lack of adequate infrastructures are frequently cited barriers for the establishment of standalone CHR-P services outside mental healthcare²¹. Future health service research is expected to consolidate these speculations, as well as to test the efficiency of innovative models of care. For example, although there were no hub and spoke services, this organization design, which arranges service delivery assets into a network, may be particularly promising in this field and fit well with the youth friendly mental health reform which is undergoing in several countries⁸³. Based on this evidence we recommend to preferably implement standalone services (Table 4).

This review also indicates that the CHR-P clinical services are essentially multidisciplinary, reflecting the complexity of the psychopathological assessment and case formulation⁸⁴. Based on the most frequent professionals involved in CHR-P services, we recommend a minimum team encompassing psychiatrists, clinical psychologists, or counsellors, case managers and nurses (Table 4). Because multidisciplinary work requires adequate articulation and training of staff, a core associated recommendation is to ensure adequate training⁸⁵. National surveys have found lack of specialised training in evidence-based interventions to cause dismal across staff^{78,86}. Ensuring proper training is particularly challenging for non-academic services, with less resources and limited organizational support⁴⁸.

In terms of outreach strategy, school, mental health, and physical health practitioners were the core targets of community outreach⁴³. We have confirmed high heterogeneity across two main strategies: active (up to 97.6%) and passive (up to 63.4%) outreach. The first strategy involved active efforts to organise workshops more frequently targeting healthcare professionals (85.4%), or service promotion activities in the community (up to 34.1% of CHR-P services) and implementing general public awareness campaigns (36.6%). The second strategy involved passive approaches such as a dedicated online site (63.4%), or printed and other media materials (53.7%). This heterogeneity is likely to reflect diverse culturally sensitive approaches across CHR-P services that led to variable pathways to care. In terms of referrals, most

CHR-P services received young people with a putative risk of psychosis from health-related organisations such as mental health services (90.0%) and general healthcare services (75.0%). Implementing an outreach to promote referrals of CHR-P individuals is challenging. In the lack of clear guidance, there is high risk of inefficient use of resources (e.g., staff) and inappropriate referrals that eventually do not meet CHR-P criteria. For example, some CHR-P services reported a high number of inappropriate referrals following intense media campaigns, switching to more focused outreach strategies^{49,51,61,79,87–90}. At times of financial constraints, the core outreach activities and referral targets summarised in the current study can be used as benchmark to maximise the efficiency of resources when implementing a new CHR-P service. There are also empirical constraints. For example, difficulties in recruiting participants is the most difficult challenge in countries where the CHR-P paradigm is starting to be implemented⁷⁶ and in culturally diverse catchment areas⁴³. Even in countries with an established CHR-P network like the UK, increasing numbers of referrals following the implementation of new national policies resulted in more dedicated CHR-P services that were needed to manage the referrals⁵⁹. Finally, the type of outreach and referrals determine the accumulation of established risk factors for psychosis^{5,7,91}, thus influencing the level of psychosis risk among individuals recruited for undergoing a CHR-P assessment (also termed as pretest risk enrichment)^{92,93}. For example, individuals sampled from inpatient units may have accumulated more risk factors for psychosis and therefore present with a higher level of psychosis compared to those sampled from the community. This level of risk enrichment^{93,94}, substantially impacts the clinical utility of CHR-P instruments¹². Accordingly, intense outreach strategies mainly targeting the general population end up diluting the level of pretest psychosis risk⁹³, and therefore impeding a clinically meaningful identification of CHR-P individuals^{11,95}. In line with recent psychometric guidances^{12,20}, we recommend CHR-P outreach to primarily target healthcare agencies to promote referrals from these sources (Table 4). Community outreach and recruitment from the general public should be considered only if adequate risk enrichment strategies can be implemented (for a detailed review see ref. 11). For example, pre-screening approaches can increase pretest risk enrichment among referrals²¹ and was employed by some services⁴⁷.

In terms of service users characteristics, we confirmed that males were relatively more represented than females, in line with the epidemiological gender distribution of psychosis risk⁶. Currently, the vast majority (83.7%) of CHR-P services employ the CAARMS or the SIPS, while basic symptoms instruments failed to enter clinical

practice at large scale. This suggests that the harmonisation of these two instruments could deliver a widely used gold standard assessment measure for clinical practice. A rapid response to referrals⁶² and flexibility with time and setting of assessments⁶⁷ have been found to improve engagement with CHR-P services.

Age intake is a core implicit criterion (along with the help-seeking behaviour) defining the CHR-P state^{4,10,20}. The most frequently applied age range (70.6% of services) was of 12–35 years, in line with epidemiological research indicating that the peak of risk is between 15 and 35 years⁶. Empirical research confirms that CHR-P psychometric assessment (e.g., the CAARMS) is valid in young people aged 12 years upwards⁶⁴. This finding also confirms the transitional nature of the CHR-P paradigm that cuts across adults and children and adolescent mental health services²⁵. Accordingly, most services provided treatment starting in adolescence (between 13 and 15 years). Conversely, only a few services accepted users beyond 40 years^{56,59,96}. The requirement of extending the assessment and care of emerging psychosis in the older people, introduced by national guidelines such as the Access and Waiting Time Standards in the UK²² is against the evidence that CHR-P instruments are valid up to 40 years⁴. Furthermore, it conflicts with recent mental health reforms that are lowering—as opposed to increasing—the age threshold for preventive approaches to those aged from 0 to 25 years²⁵. Based on these findings we recommend that CHR-P services ascertain the at-risk status through the CAARMS or SIPS in both adolescents and young adults (Table 4). This review also indicated that presentation to CHR-P services was associated with frequent comorbid mental health conditions (in particular mood and anxiety disorders^{97,98}) in two-thirds (69.5%) of the individuals, coupled with past history of suicide attempts in about one in ten (10.5%) and unemployment in about one third (36.6%) of cases. We further observed regional heterogeneity in clinical presentation: substance misuse was more prevalent in Western services^{38,51,54,60,65}, while non-existent in Japan⁵². These findings recommend that CHR-P services should adopt a broader “transdiagnostic” approach^{99–101}, which is cutting across several psychopathological dimensions (Table 4), given that psychosis onset can occur from preceding mood dysregulation¹⁰² or substance abuse. This recommendation is also relevant for current operationalisations of at-risk syndromes, which require formulating a differential diagnosis between psychosis risk and other psychopathological dimensions such as the SIPS or the DSM-5-APS². Although psychotic experiences are frequent in the general population^{103,104}, clinical attenuated psychotic symptoms are infrequent and not normally distributed. Only 0.3% of the general young population meet DSM-5-APS criteria^{2,105}.

In terms of interventions, most services (72.4%) provided care for 2 years or less (see outcomes below), with some exceptions^{38,48,52,54,55,62,63,72,74,81,87,106}, encompassing clinical monitoring (100%), psychoeducation (81.1%), psychosocial support (73%), family interventions (73%), CBT-based individual interventions (67.6%), group psychotherapy (18.9%), physical health interventions (37.8%), antipsychotics (87.1%), antidepressants (74.2%), anxiolytics (51.6%), and mood stabilisers (38.7%). It appears that CHR-P clinical services currently provide a wide range of psychosocial and biological interventions to meet the clinical needs of CHR-P service users. Clinical monitoring, case management and targeted case management are essential elements of preventive treatment²², based on the principles of social psychiatry and the importance of engaging CHR-P individuals with healthcare services¹⁰⁷. These often included psychoeducation and informing patients about their risk, as done in other preventive approaches in medicine⁸⁴. Despite current guidelines recommending psychological interventions (such as cognitive behavioural therapy) as first-line treatment, about one-third of CHR-P services did not provide them. Evidence to favour psychotherapy over other types of interventions in this population is currently uncertain^{13,15,16,20}. Conversely, antipsychotic treatment, which is discouraged by current treatment guidelines, was frequently considered, although typically at low dosages and only when the symptoms were deteriorating. This is consistent with data from global and national surveys of CHR-P services^{21,78} that report frequent use of antipsychotic drugs. The relatively frequent use of anxiolytics, antidepressants, and mood stabilisers—which is not considered by current guidelines—can index the transdiagnostic nature of the CHR-P state with frequent affective and anxiety comorbidities. The variety in provision of treatments likely reflects the high clinical heterogeneity of this population and the lack of clear treatment guidelines stratified on their individual needs. For example, current guidelines are not stratified across CHR-P subgroups. Individuals with brief psychotic episodes may be defined through research-based operationalisations, such as brief and limited intermittent psychotic symptoms (BLIPS) or standard psychiatric classifications including “Acute and Transient Psychotic Disorder” as per ICD-11 or DSM-5 “Brief Psychotic Disorder”. There is diagnostic and prognostic overlap across these definitions of brief psychotic episodes^{108,109}. Individuals with brief psychotic episodes have the highest risk of developing psychosis²⁰—especially when recurrent or presenting with seriously disorganizing or dangerous features—^{108,110}. They also display poor clinical outcomes and do not engage with the recommended cognitive behavioural therapy^{40,108}, leaving them with unmet need for care¹¹⁰. Stratification across these clinical subgroups has been proposed in recent revisions

of the CHR-P paradigm^{1,20} and should be considered in future clinical guidelines²¹. Because the uncertainty of current evidence is high, we align with the recent recommendations of the European College of Neuropsychopharmacology Prevention of Mental Disorders and Mental Health Promotion Network¹⁹ to still offer needs-based interventions and psychological interventions, titrating the intervention according to the characteristics and risk profile (i.e., transition risk, symptom severity, and functional impairment)²⁰ as well as the values and preferences of the CHR-P individuals (Table 4)²⁰. For example, it seems important to individualise physical health and lifestyle interventions on the needs presented by each service user^{60,76}.

In terms of outcomes, surprisingly, persistence of symptoms/comorbidities (67.4%) was measured more frequently than transition to psychosis (53.5%), functional status (48.8%), and remission (41.9%). This likely reflects the efforts of CHR-P services to treat comorbid conditions, aiming for improving recovery, functioning, and quality of life²⁰. At the same time, other outcomes such as physical health were collected in only about a third (30.2%) of CHR-P services¹¹¹. CHR-P individuals accumulate genetic and environmental risk factors²⁰, including cardiometabolic risk factors as decreased physical activity¹¹² and high rates of substance use¹¹², including tobacco¹¹², alcohol¹¹², and cannabis¹¹³. Thus, more attention should be paid to recording the physical health of CHR-P individuals in clinical services¹¹⁴. Another domain of improvement includes a more frequent monitoring of service users' satisfaction, which is pivotal to higher engagement and decreased drop-out rates. Furthermore CHR-P services should also more extensively monitor healthcare utilisation (e.g., hospital admissions)¹¹⁵ and broad outcomes such as mortality rates to better characterise the overall burden of this condition¹¹⁶. Future research is needed to standardise a core outcome set for CHR-P research and therefore facilitate collaborative efforts. These initiative should also indicate the assessment measures to be employed to monitor outcomes. Currently, clinical outcomes in CHR-P services are most commonly evaluated with standard clinical interviews (93.0%), and psychometric instruments are more infrequently used (37.2% of cases, most frequently CAARMS or SIPS to evaluate transition to psychosis)²⁰. In the future, monitoring broad health outcomes in CHR-P services could leverage electronic health records that can provide real-world, real-time valuable clinical information^{11,117–120} and that are being increasingly implemented in healthcare providers. As noted above, duration of care including clinical monitoring is currently limited to, most frequently (44.8%), 2 years. However, accumulating evidence has clearly indicated that although the risk of psychosis onset peaks within 2 years¹²¹, it can increase

in the longer term at least until 3–4 years^{40,122,123}. In addition, non-transitioning CHR-P individuals can continue to experience functional impairment and symptomatology at 6-years⁹⁷. This confirms that a 2-year service provision is insufficient²¹. As such, we recommend clinical monitoring for outcomes to be implemented for at least 3 years (Table 4). Flexible follow-up after this timepoint can help make more efficient use of clinical resources, while tailoring interventions to users' needs¹²⁴. For example, the clinical follow-up can be extended if service users are still symptomatic or present socio-occupational difficulties^{55,62}. Finally, CHR-P services should be prepared to collect information and target outcomes other than psychosis such as recovery, physical health outcomes, service users' satisfaction, functioning, and quality of life^{20,124}. Harmonisation of core outcome set for CHR-P services is a clinical research priority for the future. Several national and regional networks of CHR-P services started to emerge during the decade of 2010–2019 (e.g., EUGEL, PRONIA, PSYSCAN, NAPLS, PNC, HARMONY, PRONET, and STEP) and may facilitate this enterprise, allowing services to leverage best practices and expertise, increasing lobby capacity and enhancing collaborative efforts^{38,44,46}. International clinical research infrastructures have also been developed such as the European College of Neuropsychopharmacology Network for the Prevention of Mental Disorders and Mental Health Promotion (ECNP PMD-MHP)¹⁹. These initiatives will introduce several innovations in the CHR-P field, encompassing personalised prediction of outcomes and individualised interventions, digital screening for improving detection of psychosis risk and enhancement of transdiagnostic research capability within CHR-P services (e.g., preventive interventions for bipolar risk)¹²⁵.

Limitations

The main limitation of this study is that health service information was scattered across services, and that there are no established standards to measure the core domains. This limited the capacity to quantitatively compare the different services with meta-analyses. Future harmonisation efforts in terms of CHR-P healthcare research would be extremely valuable. The database was nonetheless large and sufficiently powered to analyse different factors including service configuration, outreach strategy and referrals, CHR-P service user characteristics, interventions, and outcomes. Another limitation is the limited knowledge provided about the long-term outcomes. Furthermore, our results are based on data from the literature that has been published. However, some clinical services may be running but not publishing details about service configuration, outreach strategy and referrals,

service user characteristics, interventions, and outcomes. At the same time, we hope that our review will stimulate the establishment of a global network of CHR-P services with shared clinical research infrastructures²¹. Finally, a considerable amount of studies were carried out in relatively small samples, with only 45.5% services^{39,50–52,59,62,68,69,73,75,80,82,87–90,106,126–128} including more than 100 CHR-P individuals.

Conclusions

Health service knowledge summarised by the current study will facilitate translational efforts for implementation of CHR-P services worldwide.

Acknowledgements

This study was supported by the King's College London Confidence in Concept award from the Medical Research Council (MRC) (MC_PC_16048) to Dr. Fusar-Poli. Dr. Fusar-Poli is supported by the PSYSCAN project through the European commission. Dr. Salazar de Pablo is supported by the Alicia Koplowitz Foundation. The views expressed are those of the authors and not necessarily those of the NHS, the NIHR or the Department of Health and Social Care.

Author details

¹Early Psychosis: Interventions and Clinical-detection (EPIC) Lab, Department of Psychosis Studies, Institute of Psychiatry, Psychology & Neuroscience, King's College London, London, UK. ²Institute of Psychiatry and Mental Health, Department of Child and Adolescent Psychiatry, Hospital General Universitario Gregorio Marañón School of Medicine, Universidad Complutense, Instituto de Investigación Sanitaria Gregorio Marañón (IISGM), CIBERSAM, Madrid, Spain. ³Department of Clinical and Health Psychology, Catholic University, Montevideo, Uruguay. ⁴Department of Brain and Behavioral Sciences, University of Pavia, Pavia, Italy. ⁵Heads UP Service, East London NHS Foundation Trust, London, UK. ⁶Centre for Psychiatry, Wolfson Institute of Preventive Medicine, Barts and the London School of Medicine and Dentistry, Queen Mary University of London, London, UK. ⁷National Institute for Health Research, Maudsley Biomedical Research Centre, South London and Maudsley NHS Foundation Trust, London, UK. ⁸OASIS Service, South London and Maudsley NHS Foundation Trust, London, UK

Conflict of interest

Dr. Fusar-Poli has received grants from Lundbeck and personal fees from Menarini, Lundbeck, and Angelini. No other disclosures reported.

Publisher's note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Supplementary Information accompanies this paper at (<https://doi.org/10.1038/s41398-020-01165-x>).

Received: 11 July 2020 Revised: 27 November 2020 Accepted: 3 December 2020

Published online: 13 January 2021

References

- Fusar-Poli, P. The clinical high-risk state for psychosis (CHR-P), version II. *Schizophr. Bull.* **43**, 44–47 (2017).
- Salazar de Pablo, G., Catalan, A. & Fusar-Poli, P. Clinical validity of DSM-5 attenuated psychosis syndrome: advances in diagnosis, prognosis, and treatment. *JAMA Psychiatry* **77**, 311–320 (2019).
- Yung, A. R. et al. Mapping the onset of psychosis: the comprehensive assessment of at-risk mental states. *Aust. N. Z. J. Psychiatry* **39**, 964–971 (2005).
- Fusar-Poli, P. et al. The psychosis high-risk state: A comprehensive state-of-the-art review. *Jama Psychiatry* **70**, 107–120 (2013).
- Fusar-Poli, P. et al. Deconstructing vulnerability for psychosis: meta-analysis of environmental risk factors for psychosis in subjects at ultra high-risk. *Eur. Psychiatry* **40**, 65–75 (2017).
- Radua, J. et al. What causes psychosis? An umbrella review of risk and protective factors. *World Psychiatry* **17**, 49–66 (2018).
- Oliver, D. et al. What causes the onset of psychosis in individuals at clinical high risk? A meta-analysis of risk and protective factors. *Schizophr. Bull.* **46**, 110–120 (2019).
- Fusar-Poli, P. et al. Disorder, not just state of risk: meta-analysis of functioning and quality of life in people at high risk of psychosis. *Br. J. Psychiatry* **207**, 198–206 (2015).
- Fusar-Poli, P., Raballo, A. & Parnas, J. What is an attenuated psychotic symptom? On the importance of the context. *Schizophr. Bull.* **43**, 687–692 (2017).
- Falkenberg, I. et al. Why are help-seeking subjects at ultra-high risk for psychosis help-seeking? *Psychiatry Res.* **228**, 808–815 (2015).
- Fusar-Poli, P., Sullivan, S. A., Shah, J. L. & Uhlhaas, P. J. Improving the detection of individuals at clinical risk for psychosis in the community, primary and secondary care: an integrated evidence-based approach. *Front. Psychiatry* **10**, 774 (2019).
- Fusar-Poli, P. et al. At risk or not at risk? A meta-analysis of the prognostic accuracy of psychometric interviews for psychosis prediction. *World Psychiatry* **14**, 322–332 (2015).
- Davies, C. et al. Lack of evidence to favor specific preventive interventions in psychosis: a network meta-analysis. *World Psychiatry* **17**, 196–209 (2018).
- Devoe, D. J., Farris, M. S., Townes, P. & Addington, J. Attenuated psychotic symptom interventions in youth at risk of psychosis: a systematic review and meta-analysis. *Early Interv. Psychiatry* **13**, 3–17 (2019).
- Fusar-Poli, P. et al. Preventive treatments for psychosis: umbrella review (Just the Evidence). *Front. Psychiatry* **10**, 764 (2019).
- Davies, C. et al. Efficacy and acceptability of interventions for attenuated positive psychotic symptoms in individuals at clinical high risk of psychosis: a network meta-analysis. *Front. Psychiatry* **9**, 187 (2018).
- Correll, C. U. et al. Comparison of early intervention services vs treatment as usual for early-phase psychosis: a systematic review, meta-analysis, and meta-regression. *JAMA Psychiatry* **75**, 555–565 (2018).
- Fusar-Poli, P., McGorry, P. D. & Kane, J. M. Improving outcomes of first-episode psychosis: an overview. *World Psychiatry* **16**, 251–265 (2017).
- Fusar-Poli, P. et al. European college of neuropsychopharmacology network on the prevention of mental disorders and mental health promotion (ECNP PMD-MHP). *Eur. Neuropsychopharmacol.* **29**, 1301–1311 (2019).
- Fusar-Poli, P. et al. Prevention of psychosis: advances in detection, prognosis, and intervention. *JAMA Psychiatry* **77**, 755–765 (2020).
- Kotlicka-Antczak, M. et al. Worldwide implementation of clinical services for the prevention of psychosis: the IEPA early intervention in mental health survey. *Early Interv. Psychiatry* (2020).
- The National Collaborating Centre for Mental Health and the National Institute for Health and Care Excellence. *Implementing the Early Intervention in Psychosis Access and Waiting Time Standard: Guidance*. (NHS, England, 2016).
- Elrod, J. K. & Fortenberry, J. L. The hub-and-spoke organization design: an avenue for serving patients well. *BMC Health Serv. Res.* **17**, 457 (2017).
- Rickwood, D. et al. Australia's innovation in youth mental health care: the headspace centre model. *Early Interv. Psychiatry* **13**, 159–166 (2019).
- Fusar-Poli, P. Integrated mental health Services for the developmental period (0 to 25 Years): a critical review of the evidence. *Front. Psychiatry* **10**, 355 (2019).
- Bird, V. et al. Early intervention services, cognitive-behavioural therapy and family intervention in early psychosis: systematic review. *Br. J. Psychiatry* **197**, 350–356 (2010).
- Behan, C., Masterson, S. & Clarke, M. Systematic review of the evidence for service models delivering early intervention in psychosis outside the stand-alone centre. *Early Interv. Psychiatry* **11**, 3–13 (2017).
- Moher, D., Liberati, A., Tetzlaff, J., Altman, D. G. & Group, P. Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. *BMJ* **339**, b2535 (2009).
- Fusar-Poli, P. et al. Towards a standard psychometric diagnostic interview for subjects at ultra high risk of psychosis: CAARMS versus SIPS. *Psychiatry J.* **2016**, 7146341 (2016).

30. McGlashan, T. W. B. & Woods, S. *The Psychosis-Risk Syndrome: Handbook for Diagnosis and Follow-up*. (Oxford University, Oxford, 2010).
31. Vollmer-Larsen, A., Handest, P. & Parnas, J. Reliability of measuring anomalous experience: the Bonn Scale for the assessment of basic symptoms. *Psychopathology* **40**, 345–348 (2007).
32. Riecher-Rössler, A. et al. The Basel screening instrument for psychosis (BSIP): development, structure, reliability and validity. *Fortschr. Neurol. Psychiatr.* **76**, 207–216 (2008).
33. Fux, L., Walger, P., Schimmelmann, B. G. & Schultze-Lutter, F. The schizophrenia proneness instrument, child and youth version (SPI-CY): practicability and discriminative validity. *Schizophr. Res.* **146**, 69–78 (2013).
34. Kay, S. R., Fiszbein, A. & Opler, L. A. The positive and negative syndrome scale (PANSS) for schizophrenia. *Schizophr. Bull.* **13**, 261–276 (1987).
35. Andreasen, N. C. The scale for the assessment of negative symptoms (SANS): conceptual and theoretical foundations. *Br. J. Psychiatry* **155**, 49–58 (1989).
36. Overall, J. & Gorham, D. The Brief Psychiatric Rating Scale (BPRS): recent developments in ascertainment and scaling. *Psychopharmacol. Bull.* **24**, 97–99 (1988).
37. Haefner, H., Bechdolf, A., Klosterkotter, J. & Maurer, K. *Early Detection and Intervention in Psychosis. A Practice Handbook*. (Schattauer, Stuttgart, 2011).
38. Fusar-Poli, P. et al. Pan-London network for psychosis-prevention (PNP). *Front. Psychiatry* **10**, 707 (2019).
39. Mitter, N., Nah, G. Q. R., Bong, Y. L., Lee, J. & Chong, S. A. Longitudinal youth-at-risk study (LYRIKS): outreach strategies based on a community-engaged framework. *Early Interv. Psychiatry* **8**, 298–303 (2014).
40. Fusar-Poli, P. et al. Heterogeneity of psychosis risk within individuals at clinical high risk: A meta-analytical stratification. *Jama Psychiatry* **73**, 113–120 (2016).
41. Hong, Q. N., Gonzalez-Reyes, A. & Pluye, P. Improving the usefulness of a tool for appraising the quality of qualitative, quantitative and mixed methods studies, the Mixed Methods Appraisal Tool (MMAT). *J. Eval. Clin. Pr.* **24**, 459–467 (2018).
42. Hong, Q. N. et al. Improving the content validity of the mixed methods appraisal tool: a modified e-Delphi study. *J. Clin. Epidemiol.* **111**, 49–59.e41 (2019).
43. Lynch, S. et al. Early detection, intervention and prevention of psychosis program: community outreach and early identification at six US sites. *Psychiatr. Serv.* **67**, 510–516 (2016).
44. McFarlane, W. et al. Early detection, intervention, and prevention of psychosis program: rationale, design, and sample description. *Adolesc. Psychiatry* **2**, 112–124 (2012).
45. Ruff, A., McFarlane, W., Downing, D., Cook, W. & Woodberry, K. A community outreach and education model for early identification of mental illness in young people. *Adolesc. Psychiatry* **2**, 140–145 (2012).
46. Simon, A. E., Theodoridou, A., Schimmelmann, B., Schneider, R. & Conus, P. The Swiss Early Psychosis Project SWEPP: a national network. *Early Interv. Psychiatry* **6**, 106–111 (2012).
47. Cocchi, A. et al. Early intervention in psychosis: a feasibility study financed by the Italian Center on Control of Maladies. *Early Interv. Psychiatry* **9**, 163–171 (2015).
48. Bertulies-Esposito, B. et al. Où en sommes-nous? An overview of successes and challenges after 30 Years of early intervention services for psychosis in Quebec. *Can. J. Psychiatry* 706743719895193 (2020).
49. Phillips, L. et al. The PACE Clinic: identification and management of young people at “Ultra” high risk of psychosis. *J. Psychiatr. Pr.* **8**, 255–269 (2002).
50. McFarlane et al. Portland identification and early referral: a community-based system for identifying and treating youths at high risk of psychosis. *Psychiatr. Serv.* **61**, 512–515 (2010).
51. Schultze-Lutter, F., Ruhmann, S. & Klosterkotter, J. Early detection of psychosis—establishing a service for persons at risk. *Eur. Psychiatry* **24**, 1–10 (2009).
52. Katsura, M. et al. A naturalistic longitudinal study of at-risk mental state with a 24 year follow-up at a specialized clinic setting in Japan. *Schizophr. Res.* **158**, 32–38 (2014).
53. Carr, V. et al. A risk factor screening and assessment protocol for schizophrenia and related psychosis. *Aust. N. Z. J. Psychiatry* **34**, S170–S180. (2000).
54. Riecher-Rössler, A. et al. The Basel early-detection-of-psychosis (FEPSY)-study—design and preliminary results. *Acta Psychiatr. Scand.* **115**, 114–125 (2007).
55. Rao, S. et al. Support for wellness achievement programme (SWAP): a service for individuals with at-risk mental state in Singapore. *Ann. Acad. Med. Singap.* **42**, 552–555 (2013).
56. Penno, S. J., Hamilton, B. & Petrakis, M. Early intervention in psychosis: Health of the Nation Outcome Scales (HoNOS) outcomes from a five-year prospective study. *Arch. Psychiatr. Nurs.* **31**, 553–560 (2017).
57. Geros, H. et al. Migrant status and identification as ultra-high risk for psychosis and transitioning to a psychotic disorder. *Acta Psychiatr. Scand.* **141**, 52–59 (2020).
58. Schultze-Lutter, F., Hubl, D., Schimmelmann, B. G. & Michel, C. Age effect on prevalence of ultra-high risk for psychosis symptoms: replication in a clinical sample of an early detection of psychosis service. *Eur. Child Adolesc. Psychiatry* **26**, 1401–1405 (2017).
59. Adamson, V. et al. Implementing the access and waiting time standard for early intervention in psychosis in the United Kingdom: an evaluation of referrals and post-assessment outcomes over the first year of operation. *Early Interv. Psychiatry* **12**, 979–986 (2018).
60. Coates, D. et al. The psychiatric, psychosocial and physical health profile of young people with early psychosis: data from an early psychosis intervention service. *Child Youth Serv.* **40**, 93–115 (2019).
61. Yung, A. R. et al. Prediction of psychosis—a step towards indicated prevention of schizophrenia. *Br. J. Psychiatry* **172**, 14–20 (1998).
62. Pruessner, M. et al. The Clinic for Assessment of Youth at Risk (CAYR): 10 years 812 of service delivery and research targeting the prevention of psychosis in 813 Montreal, Canada. *Early Interv. Psychiatry* **11**, 177–184 (2017).
63. Kollias, C. et al. Early psychosis intervention outpatient service of the 1st Psychiatric University Clinic in Athens: 3 Years of experience. *Early Interv. Psychiatry* **12**, 491–496 (2018).
64. Spada, G. et al. Identifying children and adolescents at ultra high risk of psychosis in Italian neuropsychiatry services: a feasibility study. *Eur. Child Adolesc. Psychiatry* **25**, 91–106 (2016).
65. Meneghelli, A., Cocchi, A. & Preti, A. ‘Programma2000’: a multi-modal pilot programme on early intervention in psychosis underway in Italy since 1999. *Early Interv. Psychiatry* **4**, 97–103 (2010).
66. Leanza, L. et al. Predictors of study drop-out and service disengagement in patients at clinical high risk for psychosis. *Soc. Psychiatry Psychiatr. Epidemiol.* **55**, 539–548 (2020).
67. Broome, M. R. et al. Outreach and support in south London (OASIS): implementation of a clinical service for prodromal psychosis and the at risk mental state. *Eur. Psychiatry* **20**, 372–378 (2005).
68. Fusar-Poli, P., Byrne, M., Badger, S., Valmaggia, L. R. & McGuire, P. K. Outreach and support in South London (OASIS), 2001–2011: ten years of early diagnosis and treatment for young individuals at high clinical risk for psychosis. *Eur. Psychiatry* **28**, 315–326 (2013).
69. Valmaggia, L. R. et al. Economic impact of early intervention in people at high risk of psychosis. *Psychol. Med.* **39**, 1617–1626 (2009).
70. Tiffin, P. A. & Hudson, S. An early intervention in psychosis service for adolescents. *Early Interv. Psychiatry* **1**, 212–218 (2007).
71. Kwon, J. S., Byun, M. S., Lee, T. Y. & An, S. K. Early intervention in psychosis: insights from Korea. *Asian J. Psychiatr.* **5**, 98–105 (2012).
72. Theodoridou, A. et al. Early recognition of high risk of bipolar disorder and psychosis: an overview of the ZInEP “Early Recognition” Study. *Front. Public Health* **2**, 166 (2014).
73. Conrad, A. M. et al. Ten-year audit of clients presenting to a specialised service for young people experiencing or at increased risk for psychosis. *BMC Psychiatry* **14**, 318 (2014).
74. Pelizza, L. et al. The “Reggio Emilia At-Risk Mental States” program: a diffused, “liquid” model of early intervention in psychosis implemented in an Italian Department of Mental Health. *Early Interv. Psychiatry* **13**, 1513–1524 (2019).
75. Leuci, E., Quattrone, E., Pellegrini, P. & Pelizza, L. The “Parma-Early Psychosis” program: general description and process analysis after 5 years of clinical activity. *Early Interv. Psychiatry* **14**, 356–364 (2019).
76. Gaspar, P. A. et al. Early psychosis detection program in Chile: a first step for the South American challenge in psychosis research. *Early Interv. Psychiatry* **13**, 328–334 (2019).
77. Kotlicka-Antczak, M., Pawelczyk, T., Rabe-Jablonska, J. & Pawelczyk, A. PORT (Programme of Recognition and Therapy): the first Polish recognition and treatment programme for patients with an at-risk mental state. *Early Interv. Psychiatry* **9**, 339–342 (2015).
78. Stain, H., Mawn, L., Common, S., Pilton, M. & Andrew, T. Research and practice for ultra-high risk for psychosis: a national survey of early intervention in psychosis services in England. *Eur. Psychiatry* **41**, S191–S192 (2017).
79. Yung, A. R. et al. Can we predict the onset of first-episode psychosis in a high-risk group? *Int. Clin. Psychopharmacol.* **13**, S23–S30 (1998).

80. Green, C. E. L., McGuire, P. K., Ashworth, M. & Valmaggia, L. R. Outreach and support in South London (OASIS). Outcomes of non-attenders to a service for people at high risk of psychosis: the case for a more assertive approach to assessment. *Psychol. Med.* **41**, 243–250 (2011).
81. Kotlicka-Antczak, M. et al. Polish individuals with an at-risk mental state: demographic and clinical characteristics. *Early Interv. Psychiatry* **12**, 391–399 (2018).
82. Power, P. et al. Lambeth early onset (LEO) and outreach & support in South London (OASIS) service. *Early Interv. Psychiatry* **1**, 97–103 (2007).
83. Mei, C. et al. Global research priorities for youth mental health. *Early Interv. Psychiatry* **14**, 3–13 (2020).
84. Fusar-Poli, P., Davies, C. & Bonoldi, I. A case of a college student presenting with mild mental health problems. *JAMA Psychiatry* **75**, 1298–1299 (2018).
85. Schultze-Lutter, F. et al. EPA guidance on the early detection of clinical high risk states of psychoses. *Eur. Psychiatry* **30**, 405–416 (2015).
86. Cocchi, A. et al. Implementation and development of early intervention in psychosis services in Italy: a national survey promoted by the Associazione Italiana Interventi Precoci nelle Psicosi. *Early Interv. Psychiatry* **12**, 37–44 (2018).
87. Nelson, B. et al. Long-term follow-up of a group at ultra high risk ("Prodromal") for psychosis the PACE 400 study. *Jama Psychiatry* **70**, 793–802 (2013).
88. Yung, A. R. et al. PACE: a specialised service for young people at risk of psychotic disorders. *Med. J. Aust.* **187**, S43–S46 (2007).
89. Yung, A. R. et al. Testing the Ultra High Risk (prodromal) criteria for the prediction of psychosis in a clinical sample of young people. *Schizophr. Res.* **84**, 57–66 (2006).
90. Yung, A. R. et al. Validation of "prodromal" criteria to detect individuals at ultra high risk of psychosis: 2 year follow-up. *Schizophr. Res.* **105**, 10–17 (2008).
91. Davies, C. et al. Prenatal and perinatal risk and protective factors for psychosis: a systematic review and meta-analysis. *Lancet Psychiatry* **7**, 399–410 (2020).
92. Fusar-Poli, P. et al. Why transition risk to psychosis is not declining at the OASIS ultra high risk service: the hidden role of stable pretest risk enrichment. *Schizophr. Res.* **192**, 385–390 (2018).
93. Fusar-Poli, P. et al. The dark side of the moon: meta-analytical impact of recruitment strategies on risk enrichment in the clinical high risk state for psychosis. *Schizophr. Bull.* **42**, 732–743 (2016).
94. Fusar-Poli, P. et al. Deconstructing pretest risk enrichment to optimize prediction of psychosis in individuals at clinical high risk. *JAMA Psychiatry* **73**, 1260–1267 (2016).
95. Oliver, D., Radua, J., Reichenberg, A., Uher, R. & Fusar-Poli, P. Psychosis polyrisk score (PPS) for the detection of individuals at-risk and the prediction of their outcomes. *Front. Psychiatry* **10**, 174 (2019).
96. Quijada, Y., Tizon, J. L., Artigue, J. & Parra, B. At-risk mental state (ARMS) detection in a community service center for early attention to psychosis in Barcelona. *Early Interv. Psychiatry* **4**, 257–262 (2010).
97. Rutigliano, G. et al. Persistence or recurrence of non-psychotic comorbid mental disorders associated with 6-year poor functional outcomes in patients at ultra high risk for psychosis. *J. Affect. Disord.* **203**, 101–110 (2016).
98. Fusar-Poli, P., Nelson, B., Valmaggia, L., Yung, A. R. & McGuire, P. K. Comorbid depressive and anxiety disorders in 509 individuals with an at-risk mental state: impact on psychopathology and transition to psychosis. *Schizophr. Bull.* **40**, 120–131 (2014).
99. Fusar-Poli, P. et al. Transdiagnostic psychiatry: a systematic review. *World Psychiatry* **18**, 192–207 (2019).
100. Fusar-Poli, P. et al. Development and validation of a clinically based risk calculator for the transdiagnostic prediction of psychosis. *JAMA Psychiatry* **74**, 493–500 (2017).
101. Fusar-Poli, P. TRANSD recommendations: improving transdiagnostic research in psychiatry. *World Psychiatry* **18**, 361–362 (2019).
102. Mishara, A. L. & Fusar-Poli, P. The phenomenology and neurobiology of delusion formation during psychosis onset: Jaspers, Truman symptoms, and aberrant salience. *Schizophr. Bull.* **39**, 278–286 (2013).
103. Moriyama, T. S. et al. Differences between self-reported psychotic experiences, clinically relevant psychotic experiences, and attenuated psychotic symptoms in the general population. *Front. Psychiatry* **10**, 782 (2019).
104. Schimmelmänn, B. G., Michel, C., Martz-Iimgartinger, A., Linder, C. & Schultze-Lutter, F. Age matters in the prevalence and clinical significance of ultra-high-risk for psychosis symptoms and criteria in the general population: findings from the BEAR and BEARS-kid studies. *World Psychiatry* **14**, 189–197 (2015).
105. Schultze-Lutter, F., Michel, C., Ruhrmann, S. & Schimmelmänn, B. G. Prevalence and clinical significance of DSM-5-attenuated psychosis syndrome in adolescents and young adults in the general population: the Bern Epidemiological At-Risk (BEAR) Study. *Schizophr. Bull.* **40**, 1499–1508 (2014).
106. Conrad, A. M. et al. Utility of risk-status for predicting psychosis and related outcomes: evaluation of a 10-year cohort of presenters to a specialised early psychosis community mental health service. *Psychiatry Res.* **247**, 336–344 (2017).
107. Dimitrakopoulos, S., Kollias, C., Stefanis, N. C. & Kontaxakis, V. Early psychotic experiences: interventions, problems and perspectives. *Psychiatrike Psychiatriki* **26**, 45–54 (2015).
108. Fusar-Poli, P. et al. Diagnostic and prognostic significance of brief limited intermittent psychotic symptoms (BLIPS) in individuals at ultra high risk. *Schizophr. Bull.* **43**, 48–56 (2017).
109. Fusar-Poli, P. et al. Prognosis of brief psychotic episodes: a meta-analysis. *JAMA Psychiatry* **73**, 211–220 (2016).
110. Fusar-Poli, P. et al. Unmet needs for treatment in 102 individuals with brief and limited intermittent psychotic symptoms (BLIPS): implications for current clinical recommendations. *Epidemiol. Psychiatr. Sci.* **29**, e67 (2019).
111. Carney, R., Cotter, J., Bradshaw, T. & Yung, A. R. Examining the physical health and lifestyle of young people at ultra-high risk for psychosis: a qualitative study involving service users, parents and clinicians. *Psychiatry Res.* **255**, 87–93 (2017).
112. Carney, R., Cotter, J., Bradshaw, T., Firth, J. & Yung, A. R. Cardiometabolic risk factors in young people at ultra-high risk for psychosis: a systematic review and meta-analysis. *Schizophr. Res.* **170**, 290–300 (2016).
113. Carney, R., Cotter, J., Firth, J., Bradshaw, T. & Yung, A. R. Cannabis use and symptom severity in individuals at ultra high risk for psychosis: a meta-analysis. *Acta Psychiatr. Scand.* **136**, 5–15 (2017).
114. Yung, A. R. et al. The reality of at risk mental state services: a response to recent criticisms. *Psychol. Med.* 1–7 (2019).
115. Yoviene Sykes, L. A. et al. Predictive validity of conversion from the clinical high risk syndrome to frank psychosis. *Schizophr. Res.* **216**, 184–191 (2020).
116. Chang, C. K. et al. All-cause mortality among people with serious mental illness (SMI), substance use disorders, and depressive disorders in southeast London: a cohort study. *BMC Psychiatry* **10**, 77 (2010).
117. Oliver D. et al. Real-world implementation of precision psychiatry: transdiagnostic risk calculator for the automatic detection of individuals at-risk of psychosis. *Schizophr. Res.* **10**, 109 (2020).
118. Wang T. et al. Implementation of a real-time psychosis risk detection and alerting system based on electronic health records using CogStack. *J. Vis. Exp.* **15**, <https://doi.org/10.3791/60794> (2020).
119. Fusar-Poli, P. et al. Transdiagnostic risk calculator for the automatic detection of Individuals at risk and the prediction of psychosis: second replication in an Independent National Health Service trust. *Schizophr. Bull.* **45**, 562–570 (2019).
120. Rutigliano, G. et al. Long term outcomes of acute and transient psychotic disorders: the missed opportunity of preventive interventions. *Eur. Psychiatry* **52**, 126–133 (2018).
121. Kempton, M. J., Bonoldi, I., Valmaggia, L., McGuire, P. & Fusar-Poli, P. Speed of psychosis progression in people at ultra-high clinical risk: a complementary meta-analysis. *JAMA Psychiatry* **72**, 622–623 (2015).
122. Fusar-Poli, P. et al. Predicting psychosis: meta-analysis of transition outcomes in individuals at high clinical risk. *Arch. Gen. Psychiatry* **69**, 220–229 (2012).
123. Fusar-Poli, P. et al. Long-term validity of the At Risk Mental State (ARMS) for predicting psychotic and non-psychotic mental disorders. *Eur. Psychiatry* **42**, 49–54 (2017).
124. Schmidt, S. J. et al. EPA guidance on the early intervention in clinical high risk states of psychoses. *Eur. Psychiatry* **30**, 388–404 (2015).
125. Fusar-Poli, P. et al. Semistructured interview for bipolar at risk states (SIBARS). *Psychiatry Res.* **264**, 302–309 (2018).
126. Lee, J. et al. The Longitudinal Youth at Risk Study (LYRIKS)—an Asian UHR perspective. *Schizophr. Res.* **151**, 279–283 (2013).
127. Tay, S. A. et al. Support for Wellness Achievement Programme (SWAP): clinical and demographic characteristics of young people with at-risk mental state in Singapore. *Early Interv. Psychiatry* **9**, 516–522 (2015).
128. Fusar-Poli, P. et al. Antidepressant, antipsychotic and psychological interventions in subjects at high clinical risk for psychosis: OASIS 6-year naturalistic study. *Psychol. Med.* **45**, 1327–1339 (2015).