

Effects of paleogeographic changes and CO₂ variability on northern mid-latitude temperature gradients in the Cretaceous

Received: 28 February 2023

Accepted: 15 August 2023

Published online: 25 August 2023

 Check for updatesKaushal Gianchandani¹✉, Sagi Maor¹, Ori Adam¹,
Alexander Farnsworth^{2,3}, Hezi Gildor¹, Daniel J. Lunt² & Nathan Paldor¹

The Cretaceous ‘greenhouse’ period (~145 to ~66 million years ago, Ma) in Earth’s history is relatively well documented by multiple paleoproxy records, which indicate that the meridional sea surface temperature (SST) gradient increased (non-monotonically) from the Valanginian (~135 Ma) to the Maastrichtian (~68 Ma). Changes in atmospheric CO₂ concentration, solar constant, and paleogeography are the primary drivers of variations in the spatio-temporal distribution of SST. However, the particular contribution of each of these drivers (and their underlying mechanisms) to changes in the SST distribution remains poorly understood. Here we use data from a suite of paleoclimate simulations to compare the relative effects of atmospheric CO₂ variability and paleogeographic changes on mid-latitude SST gradient through the Cretaceous. Further, we use a fundamental model of wind-driven ocean gyres to quantify how changes in the Northern Hemisphere paleogeography weaken the circulation in subtropical ocean gyres, leading to an increase in extratropical SSTs.

Analysis of oxygen isotopes in foraminifera tests ($\delta^{18}\text{O}$) and the index of tetraethers that quantifies the archaeal lipid distributions in marine sediments (TEX₈₆) indicate that the sea surface temperature (SST) during the Cretaceous period (~145 to ~66 million years ago, Ma) ranged between 28–35 °C in low latitudes to 11–22 °C in high latitudes (first and third quartiles of LOESS smoothed data)¹. A similar trend is observed during the Paleocene (~66 to ~56 Ma) and Eocene (~56 to ~34 Ma) epochs of the Paleogene period (~66 to ~23 Ma) when the tropical and high latitude temperatures were between 31–33 °C and 11–14 °C, respectively². The persistence of such high globally averaged SST and low pole-to-tropics temperature gradient during the Cretaceous–Paleocene–Eocene (CPE) is attributed predominantly to the strong greenhouse effect ensuing from the higher partial pressure of CO₂ ($p\text{CO}_2$) in the atmosphere^{3,4}. However, in addition to changes in $p\text{CO}_2$, the spatiotemporal variability in SST can also stem from

tectonically driven changes in paleogeography, including surface topography, ocean bathymetry, and the arrangement of continents^{5–7}. Specifically, paleogeographic alterations during the Cretaceous have been demonstrated to dampen seasonal variations and trigger widespread flooding, ultimately leading to moderation of the pole-to-tropics temperature gradient^{8,9}. The rearrangement of continents during the period has also been associated with shifts in the deep-water formation sites^{7,10}, alternations in the oceanic overturning circulation¹¹, and variations in mid-Cretaceous deep-sea ventilation¹². Furthermore, the paleogeographic features associated with certain ages in the Early and Late Cretaceous have been found to be more conducive to glacial events than the mid-Cretaceous¹³. Moreover, changes in Antarctic orography¹⁴ and the opening of the Drake passage¹⁵ could have both contributed to the variability in meridional SST gradients during the Paleocene–Eocene (PE).

¹Fredy & Nadine Herrmann Institute of Earth Sciences, Hebrew University of Jerusalem, Edmond J. Safra Campus, Givat Ram, Jerusalem 9190401, Israel.

²School of Geographical Sciences and Cabot Institute, University of Bristol, Bristol BS8 1SS, UK. ³State Key Laboratory of Tibetan Plateau Earth System, Environment and Resources (TPESER), Institute of Tibetan Plateau Research, Chinese Academy of Sciences, Beijing 100101, China.

✉ e-mail: kaushal.g@mail.huji.ac.il

Although previous studies have investigated the effect of different paleogeographies corresponding to multiple ages in the CPE on variations in the pole-to-tropics temperature gradient and the ocean circulation^{10–12,16}, the effect of paleogeographic changes on the surface ocean circulation specifically has not been examined before. Surface ocean gyres contained between two meridional continental boundaries are prominent features of the present-day ocean. The meridional volume (mass) transport (ψ) associated with these gyres redistributes heat in the ocean and is critical to regulating the climate on Earth's surface.

Henry Stommel was the first to formulate a simple mathematical model comprising five parameters that capture the fundamental characteristics of the circulation in such gyres¹⁷. While the linear dependence of ψ on the amplitude of zonally averaged wind stress (τ) is evident from the analytical solution of Stommel's model, discerning its dependence on the zonal and meridional extents of the ocean basin (L_x and L_y , respectively) is rather non-trivial. It should be noted that in Stommel's model, the meridional boundaries of the basin are co-located with the latitudes of zero wind-stress curl. However, in the real ocean, the two do not necessarily coincide in general, and the extent of the surface gyres is constrained only by the latitudes corresponding to the vanishing wind-stress curl. Previous studies established that both ψ and the poleward heat transport (\mathcal{H}) associated with a surface gyre increase with the horizontal aspect ratio ($\frac{L_y}{L_x}$) of the ocean basin containing the gyre^{18–20} (hereafter referred to as the 'gyral basin'). Thus, the aspect ratio of any gyral basin is governed by the curl of the overlying wind stress and the relative position of the zonal boundaries set by the continents. Based on these first-order constraints, it was recently postulated that the large meridional SST gradients observed during some geologic periods may have resulted from the smaller ψ (and consequently a smaller \mathcal{H}) associated with the surface gyres contained in basins with small aspect ratios²⁰.

Here we use data generated from two ensembles of paleoclimate simulations carried out using the HadCM3L (specifically, HadCM3LB-M2.1aD) model^{6,7} to better constrain the relative contribution of changes in $p\text{CO}_2$ and paleogeographic effects on the SST distribution during the CPE. In particular, we develop a simple model that isolates the role of paleogeographically driven changes in extratropical gyral circulation and the corresponding effect on the evolution of pole-to-tropics SST gradient. The model is then applied to the HadCM3L data to show that paleogeography-driven reduction in ψ (and consequently \mathcal{H}) associated with the surface ocean gyres may have been a key factor contributing to the variability in meridional SST gradients observed during the CPE^{1,2}.

Results

Numerical simulations

All simulations necessary for this work were carried out using the UK Met Office coupled atmosphere-ocean model, HadCM3L (resolution: 3.75° in longitude \times 2.5° in latitude), which includes multiple climate feedbacks, including vegetation feedbacks^{6,21}. The atmospheric CO_2 concentration in the two ensembles is kept constant at 560 ppmv and 1120 ppmv (i.e., $\times 2$ and $\times 4$ preindustrial atmospheric CO_2 concentration) while the geography and the solar constant are varied. The chosen values of atmospheric CO_2 provide an idealized framework that allows us to isolate the effect of CO_2 versus paleogeography. In addition, this range of CO_2 values captures the maximum variation (530 ppmv to 840 ppmv) in the long-term trend of LOESS smoothed data on estimated atmospheric CO_2 concentration during the CPE²². The range does not capture the very lowest values of CO_2 seen in some proxies in the latest Cretaceous²³, but these low values are inconsistent with relatively high reconstructed global mean temperatures at this time²⁴. The geographies corresponding to different geological ages are constrained by data aggregated from lithological, tectonic, fossil, and

deep-sea studies⁶. The solar constant in both ensembles increases monotonically from the Berriasian age (-143 Ma) to the Priabonian age (-36 Ma) by 0.9% (see Methods). In practice, the relatively small magnitude of the forcing associated with the increase in solar constant is dwarfed by changes in $p\text{CO}_2$ and paleogeography between simulations and their relative impact on SST distributions.

Wind-driven gyral circulation in the CPE

The mid-latitude gyral basin in the northern paleo-Pacific (hereafter referred to as the North Pacific) retains an approximately rectangular geometry – similar to Stommel's idealized ocean – during the CPE, despite considerable plate tectonic activity^{25,26}. Variations in the North Pacific's circulation are expected to strongly affect the climate during the CPE, since for these -110 million years much of the planet's sea-water was in the large paleo-Pacific^{27,28}.

The L_x of the mid-latitude gyral basin in the North Pacific, i.e., the typical longitudinal distance between continental margins (see Methods), remained nearly constant during the CPE (-13,000 km, Supplementary Fig. 1). On the other hand, L_y of this basin, which is defined as the distance between the two latitudes where the wind-stress curl over the surface vanishes, decreased (non-monotonically) from -3900 km to -2800 km and then increased to -3600 km (Fig. 1a). These variations in L_y determine the variations in the aspect ratio and hence the value of ψ in this gyral basin. This is illustrated by comparing ψ in the longer Valanginian gyral basin (-135 Ma; Fig. 1b), which is double that of the -30% narrower Maastrichtian gyral basin (-68 Ma; Fig. 1c). The ocean during the Valanginian is also characterized by a much weaker SST gradient, which is consistent with a stronger \mathcal{H} compared to the Maastrichtian.

Variability in meridional SST gradients and its dependence on atmospheric CO_2 and paleogeography

To examine the variability in meridional SST gradients during the CPE, we first consider the SST gradients from the Early to the Late Cretaceous and subsequently focus on the trend from the Paleocene to the Eocene. Specifically, we compare the Maastrichtian to the Valanginian (rather than the Berriasian, given the paucity of the proxy-derived SST data in this age¹) in the Cretaceous and the Priabonian to the Danian age (-63 Ma) in the PE.

In the HadCM3L simulations carried out for an atmospheric CO_2 concentration of 560 ppmv, the meridional gradient of the zonally averaged SST (over all longitudes, ΔSST_y) calculated between 20°N and 50°N increases from 12.5°C in the Valanginian to 15.7°C in the Maastrichtian, whereas, ΔSST_y decreases from 16.5°C in the Danian to 15.5°C in the Priabonian. The same trend persists for an atmospheric CO_2 concentration of 1120 ppmv, ΔSST_y increases from 11.1°C to 13.9°C during the Cretaceous and decreases from 14.3°C to 13.7°C during the PE. Figure 2a illustrates the temporal trend in ΔSST_y for the two atmospheric CO_2 concentrations. In both simulation ensembles, the temporal variation in ΔSST_y from the Valanginian to the Maastrichtian is -3°C even though the global mean SST varies only slightly (standard deviation of less than 1°C about the ensemble averages, see Supplementary Fig. 3). Alternatively, the temporal variation in ΔSST_y from the Danian to the Priabonian (-0.8°C) is comparable to the standard deviation about the global mean SST for the ensembles. Furthermore, comparing the two simulation ensembles shows that during the CPE, a doubling of $p\text{CO}_2$ decreases ΔSST_y between 0.7°C (observed in the Albian age, -106 Ma) and 2.6°C (observed in the Selandian age, -60 Ma), resulting in an average decrease of 1.8°C in ΔSST_y .

The long-term trend of LOESS smoothed data on estimated atmospheric CO_2 concentration suggests that a decrease in atmospheric CO_2 from -700 ppmv in the Valanginian to -570 ppmv in the Maastrichtian²² can cause a decrease in global mean SST by -1.5°C (based on a typical climate sensitivity parameter of -4.5°C per CO_2




Fig. 1 | Changes in the meridional extent of the mid-latitude gyral basin (L_y) in the Northern paleo-Pacific during the Cretaceous and Paleogene periods for an atmospheric CO_2 concentration of 560 ppmv (solid blue curve) and 1120 ppmv (dashed red curve). **a Change in L_y during the Early-Late Cretaceous, the Paleocene, and the Eocene epochs. **b, c** Sea surface temperature ($^{\circ}\text{C}$, color) and gyral streamlines (Sv, contour; $1 \text{ Sv} = 10^6 \text{ m}^3 \text{ s}^{-1}$) in the ocean during the Valanginian**

(-135 Ma) and the Maastrichtian (-68 Ma) ages for an atmospheric CO_2 concentration of 1120 ppmv. Dotted lines mark the latitudes where the wind-stress curl is zero. For completeness, Supplementary Fig. 2 shows the sea surface temperature fields and streamlines in the ocean during the Valanginian and the Maastrichtian for an atmospheric CO_2 concentration of 560 ppmv.

doubling⁷) and an increase in ΔSST_y by -0.6°C (since a doubling of atmospheric CO_2 decreases ΔSST_y by 1.8°C a decrease in CO_2 from -700 ppmv to -570 ppmv, i.e., a reduction by a factor of 0.8, is expected to increase ΔSST_y by $\frac{\log(0.8)}{\log(2)} \times -1.8 = 0.6^{\circ}\text{C}$). Similarly, a decrease in atmospheric CO_2 from -590 ppmv in the Danian to -530 ppmv in the Priabonian²² is expected to cause a 0.7°C decrease in global mean SST and an 0.3°C increase in ΔSST_y . These estimates are consistent with the inverse relation between ΔSST_y and global mean SST observed across multiple climate states in Earth's history with varied paleogeographies². It should be noted that changes in atmospheric CO_2 may have had a more pronounced effect on ΔSST_y since this study does not account for the short-term fluctuations in $p\text{CO}_2$ trend or the lowest values of CO_2 inferred from some proxies²³.

Comparing the typical variation in ΔSST_y from the change in $p\text{CO}_2$ with paleogeography-driven variation in ΔSST_y indicates that paleogeographic changes had a greater effect on the mid-latitude temperature gradient during the Cretaceous than the long-term changes in $p\text{CO}_2$. This is also true for the PE. However, since the overall ΔSST_y variation during the PE is comparable to the standard deviation in the ensemble-averaged global mean SST, the analysis that follows only focuses on the Cretaceous.

The differences between the long-term SST gradients in the simulations and estimates based on paleotemperature reconstructions may be attributed to the limited spatial distribution of available proxies. Paleotemperature reconstructions based on $\delta^{18}\text{O}$ and TEX_{86} proxies suggest that meridional SST gradients first decreased from -10 – -17°C in the Valanginian to -3 – -5°C in the Aptian (-119 Ma) and subsequently increased through the middle-Late Cretaceous to reach a maximum of -19 – -21°C in the Maastrichtian¹. However, the trend in this proxy-derived meridional SST gradient is only expected to capture the long-term trends in Equator-to-pole temperature difference. It does not capture the mid-latitude temperature gradient in the Northern Hemisphere (the focus of this study) since no data exists between 30°N and 80°N and the data for the high latitude paleotemperature originates from sites located between 48°S and 54°S paleolatitudes. Nevertheless, the estimated ΔSST_y variation resulting

from the combined effect of reduction in atmospheric CO_2 concentration and paleogeographic changes from the Valanginian to the Maastrichtian (-3.6°C) is within 55% of the variation estimated from proxies (-6.5°C).

Role of gyral circulation in determining the meridional SST gradients

The decrease in ΔSST_y resulting from an increase in $p\text{CO}_2$ of the model atmosphere can be attributed to polar amplification^{22, 29–32}, whereas the increase in ΔSST_y from the Early to Late Cretaceous in the two ensembles of simulations is related to paleogeography-driven changes. Paleogeography can influence many factors that could in turn affect polar amplification (e.g., cloud distribution and other local feedbacks); here, we focus on its influence on oceanic heat transport (\mathcal{H}). Figure 2b shows a strong anti-correlation between ΔSST_y and the maximal volumetric (mass) transport (ψ) in mid-latitudes, which indicates that Cretaceous basins with smaller L_y contain surface gyres with smaller ψ . Thus, the weakening of mid-latitude \mathcal{H} (Supplementary Fig. 4), caused by the weakened mid-latitude ocean gyres contributes substantially to the increase of ΔSST_y in the middle-Late Cretaceous. To quantify this effect, we assume that:

1. To first order, ΔSST_y decreases linearly with \mathcal{H} and with the poleward heat transport associated with the atmospheric meridional overturning circulation (MOC).
2. The heat transport associated with the atmospheric MOC is proportional to \mathcal{H}^{33} .
3. The ocean circulation, to first order, is similar to the circulation in a 'Stommel-like' gyre. As such, due to rapid advection, there is negligible loss of heat by warm water along the western boundary. In contrast, the water mass loses much of its heat while it traverses the northern and eastern edges of the basin at a much lower speed. Thus, SST along the southern and western boundaries is high, whereas SST along the northern and eastern boundaries is low. Further, the East-West SST gradient is proportional to ΔSST_y .




Fig. 2 | Meridional temperature gradient and maximal volumetric (mass) transport in the mid-latitude northern paleo-Pacific. **a** Meridional gradient of zonal-mean SST (ΔSST_y) between 20°N and 50°N for an atmospheric CO₂ concentration of 560 ppmv (blue curve) and 1120 ppmv (red curve). The best-fit curves for the two atmospheric CO₂ concentrations (dashed blue and red curves) are calculated using ΔSST_y and the maximal volume (mass) transport (ψ_{max}). The shaded region shows the range between which the best-fit curve can vary with 1 σ error in κ and ΔSST_y^{Rad} . **b** Decrease in ψ_{max} from the Early to Late Cretaceous and the subsequent increase from the Paleocene to the Eocene for the two atmospheric CO₂ concentrations.

Based on these assumptions, we find (see Methods):

$$\Delta SST_y = \frac{\Delta SST_y^{Rad}}{1 + (\kappa \times \psi_{max} \times \Delta SST_y^{Rad})} \quad (1)$$

where ΔSST_y^{Rad} is the meridional gradient of SST at radiative equilibrium ($H = 0$), ψ_{max} is the maximal gyral volume (mass) transport in the mid-latitude North Pacific, and κ is an empirical constant. The above equation is developed based on the paleogeographic considerations relevant to the Cretaceous, but we expect it to be also applicable to the PE since the majority (>60%) of the volumetric (mass) transport in the ocean's surface is concentrated in the North Pacific until the Priabonian. Performing a regression analysis, based on this equation, on ΔSST_y and ψ_{max} estimated from the HadCM3L model data for the 19 ages considered here yields $\Delta SST_y^{Rad} = 22.6 \pm 1.4$ °C and $\kappa = (3.9 \pm 0.5) \times 10^{-4} \text{ Sv}^{-1} \text{ °C}^{-1}$ ($1 \text{ Sv} = 10^6 \text{ m}^3 \text{ s}^{-1}$) for an

atmospheric CO₂ of 560 ppmv, and $\Delta SST_y^{Rad} = 18.0 \pm 0.9$ °C and $\kappa = (3.7 \pm 0.6) \times 10^{-4} \text{ Sv}^{-1} \text{ °C}^{-1}$ for an atmospheric CO₂ of 1120 ppmv. The best-fit curves (Fig. 2a, dashed curves) illustrate that the reduction in ψ_{max} explains -80% and -75% of the variance in ΔSST_y from the Berriasian to the Priabonian in the simulation ensemble with an atmospheric CO₂ of 560 ppmv and 1120 ppmv, respectively. This underscores the role of L_y in the increase in ΔSST_y during the Cretaceous and in the subsequent decrease in ΔSST_y from the Paleocene to the Eocene.

Though our regression analysis captures the general trend in the evolution of ΔSST_y during the CPE for both simulation ensembles, we observe a steep increase in ΔSST_y between -130 Ma and -100 Ma for the simulations with an atmospheric CO₂ of 1120 ppmv (Fig. 2a, solid red curve). This steep increase lies outside the range in which the best-fit curve is expected to vary based on one standard deviation variation in κ and ΔSST_y^{Rad} . It stems from an abrupt decrease in the SST at 50°N between the Barremian age (-128 Ma) and the Albian age (Supplementary Fig. 5). However, examining the precise climatic variations and feedback associated with paleogeographic changes that led to this abrupt cooling between the Barremian and the Albian are beyond the scope of this study.

The HadCM3L simulations show that geologic ages in the Cretaceous with a large L_y (e.g., the Valanginian) are characterized by a wide Equator-to-pole extent of the ocean basin (\tilde{L}_y) (Fig. 3a, b), while geologic ages with a small L_y (e.g., the Maastrichtian) are marked by a narrower \tilde{L}_y (Fig. 3c, d). Moreover, like L_y , \tilde{L}_y also decreased from the Early to Late Cretaceous (Supplementary Fig. 6). The correlation between L_y and \tilde{L}_y can be attributed to the increase in land area in the polar North Pacific since the mid-Cretaceous (Supplementary Fig. 7), because the increased land area amplifies the damping effect (friction) over the surface, which reduces the meridional distance between the zonal wind extremes (thus reducing the distance between the latitudes where the wind-stress curl over the surface vanishes). However, multiple other features of the atmosphere-ocean system not considered here can affect L_y , e.g., meridional extent and/or intensity of the Hadley cell, the intensity of the Walker circulation, and exchange of water masses between ocean basins³⁴. For instance, L_y and \tilde{L}_y do not coevolve during the PE (Supplementary Fig. 6), which could be related to one or more of these features. Exploring a causal relation between changes in these climatic features and L_y on geologic timescales is left for future study.

Discussion

In summary, we find that a large portion of the increase in ΔSST_y observed in the Stommel-like mid-latitude North Pacific basin during the Cretaceous results from the decrease in poleward ocean heat transport. This decrease in heat transport is a consequence of a weakened intensity of the gyral circulation, i.e., a reduction in the volumetric/mass transport associated with the gyre. This weakening results from tectonically driven changes in Northern Hemisphere paleogeography, which reduces the horizontal aspect ratio of the gyral basin. The anti-correlation between oceanic heat transport and ΔSST_y persists in the PE as well, and an increase in oceanic heat transport during these epochs is accompanied by a decrease in ΔSST_y . However, this decrease is of the order of the standard deviation of the ensemble-averaged global mean SST. We note that in contrast to the Northern Hemisphere, the two ensembles of simulations do not show a discernible trend in ΔSST_y variations in the Southern Hemisphere, likely because of the break-up of the Antarctica–Australia landmass during this period – demonstrating the limitations of the theory proposed here.

We also find that fluctuations in atmospheric CO₂ concentration can substantially affect the ΔSST_y during the Cretaceous and estimate the magnitude of ΔSST_y variation based on the long-term trend of atmospheric CO₂. However, there is a considerable uncertainty in the




Fig. 3 | Zonal-mean wind speed and temperature distribution in the northern paleo-Pacific Ocean for an atmospheric CO₂ concentration of 1120 ppmv in the Valanginian (~142 Ma) and Maastrichtian (~68 Ma) ages. Panels a and c show the zonally averaged winds in the Valanginian and the Maastrichtian, respectively; the dotted lines denote the latitudes where the wind-stress curl vanishes and L_y is the meridional distance between them. Panels b and d show the temperature fields

(°C, color) and streamlines (Sv, contours) of the surface gyre during the two ages; \tilde{L}_y is the typical Equator-to-pole extent of the basin. The yellow-purple loops depict the meridional extent of the atmospheric meridional overturning circulations. The latitude where the winds change from Easterlies to Westerlies marks the poleward extent of the atmospheric overturning cell⁴⁰.

estimated atmospheric CO₂ levels^{23,35} in general and, thus by extension, in the magnitude of ΔSST_y variation we attribute to pCO_2 fluctuations during the period. In addition to changes in atmospheric CO₂ concentration and paleogeography, ΔSST_y can also be non-trivially affected by several other drivers in the climate system, e.g., fluctuations in deep-sea temperatures^{22,36}, opening/closing/deepening of oceanic gateways^{10,11,37} and variations in the extent of marine ice cover⁸.

The temporal variation in meridional temperature gradients obtained from the HadCM3L model for both the simulation ensembles agrees with the proxy-inferred increase in temperature gradient from the Early to the Late Cretaceous¹. Since the atmospheric CO₂ and continental arrangement do not coevolve in the current set of simulations, we stop short of comparing the model temperature gradients with proxy data. This should be addressed in future work using data from multiple climate models in which atmospheric CO₂ and geography are varied simultaneously. Additionally, numerical simulations that use a hierarchy of coupled atmosphere–ocean models with idealized continental configurations can further advance our understanding of how paleogeography affects ΔSST_y . Nevertheless, our analysis is based on foundational models of atmospheric and oceanic circulation and first-order assumptions to quantify the characteristics of a given paleoclimate – which are simple and robust. It may, therefore, also explain the effects of changes in paleogeography on variations in ΔSST_y in other geologic periods.

Methods

Identifying the meridional and zonal extents of the mid-latitude gyral basin in the North Pacific

To determine the typical meridional and zonal extents of the gyral basin in the mid-latitude North Pacific, we construct a ‘trapezoidal’ domain in the ocean. The meridional margins of the gyral basin are given by the latitudes, ϕ_1 and ϕ_2 , where the zonally averaged wind calculated by the HadCM3L model reaches an extremum. These extrema correspond to the points where the curl of the zonally averaged wind stress vanishes. Subsequently, we determine four typical

longitudes, λ_1 – λ_4 , based on the continental boundaries between the latitudes ϕ_1 and ϕ_2 such that the region enclosed between the vertices P (ϕ_1, λ_1), Q (ϕ_1, λ_2), R (ϕ_2, λ_3) and S (ϕ_2, λ_4) denotes the ‘gyral basin’. The meridional extent of the gyral basin (L_y) is defined as the latitudinal distance between the edges PQ and RS and the zonal extent (L_x) is defined as the cosine(latitude) weighted average of the lengths of PQ and RS. Supplementary Fig. 1 illustrates the gyral basins during different geologic ages in the Cretaceous and the Paleogene. We note that both L_y and L_x depend on the grid resolution. Despite the rather stringent definition of L_y and L_x and their dependence on the grid resolution, the correlation between L_y and ΔSST_y is very strong, which further demonstrates the robustness of our analysis.

Relation between meridional sea surface temperature gradients and the volume (mass) transport in the ocean

Poleward volume (mass) and heat transport associated with the atmospheric meridional overturning circulation and the gyral circulation in the ocean’s surface lower the meridional sea surface temperature gradient (ΔSST_y), which is driven by differential solar heating. We assume that ΔSST_y is given by (see assumption 1, main text):

$$\Delta SST_y = \Delta SST_y^{Rad} (1 - c_O \mathcal{H}_O - c_A \mathcal{H}_A) \quad (2)$$

where ΔSST_y^{Rad} is the meridional gradient of SST at radiative equilibrium when ocean and atmosphere heat transport are zero ($\mathcal{H}_O = \mathcal{H}_A = 0$), and c_O and c_A are the appropriate empirical constants. Assuming that atmosphere and ocean heat transport are proportional to each other (on average, and not at every latitude)³³, we may write $c_A \mathcal{H}_A = (\gamma - 1) c_O \mathcal{H}_O$, where $\gamma > 1$ is an arbitrary constant (see assumption 2, main text). Equation (2) can then be rewritten as:

$$\Delta SST_y = \Delta SST_y^{Rad} (1 - \gamma c_O \mathcal{H}_O). \quad (3)$$

We further assume that the gyral heat transport can be written as $\mathcal{H}_O = c_p \rho \psi_{max} \Delta SST_x$, where ΔSST_x is the East-West SST gradient³⁸,

yielding:

$$\Delta SST_y = \Delta SST_y^{Rad} \left(1 - \gamma c_o c_p \rho \psi_{\max} \Delta SST_x \right). \quad (4)$$

Given the well-constrained geometry of a Stommel-like gyre, we substitute $\Delta SST_x = c_{xy} \Delta SST_y$ (see assumption 3, main text) to obtain:

$$\Delta SST_y = \frac{\Delta SST_y^{Rad}}{1 + \left(\gamma c_o c_p \rho \frac{1}{c_{xy}} \psi_{\max} \Delta SST_y^{Rad} \right)}, \quad (5)$$

where $0 < c_{xy} \leq 1$ is another constant. Substituting $\kappa = \gamma c_o c_p \rho \frac{1}{c_{xy}}$, yields Eq. (1).

Coupled atmosphere–ocean model description

The data analyzed in this study was generated from two ensembles of simulations carried out using the UK Met Office coupled atmosphere–ocean model, HadCM3L, which includes multiple climate feedbacks, including vegetation feedbacks^{6,21}. The model has a horizontal resolution of 3.75° in longitude \times 2.5° in latitude. The atmosphere and the ocean are divided into 19 and 20 vertical levels, respectively. The CO₂ in the model atmosphere is kept fixed at 560 ppmv and 1120 ppmv for the two ensembles of simulations, while the paleogeography and solar constant are varied. The chosen range of atmospheric CO₂ captures the maximum variation (530 ppmv to 840 ppmv) in the long-term trend of LOESS smoothed data on estimated atmospheric CO₂ concentration during the CPE²². The geographies corresponding to different geological ages are developed by Getech Plc using methods described in ref. 39, and are constrained by geological data aggregated from lithological, tectonic, fossil, and deep-sea studies. The geographies were originally produced at a resolution of 0.5° in both latitude and longitude, which were subsequently used to generate geographies at model resolution. Additionally, some smoothing was applied to ensure model stability. A more detailed description is available in Section 2.1 of ref. 6. Given that the variation in solar constant was quite small (a monotonic increase of -0.9% from the Berriasian age in the Cretaceous to the Priabonian age in the Paleogene), we assume that for a given $p\text{CO}_2$ in the model atmosphere, any changes in the spatial distribution of SSTs between simulations arise primarily from the changing paleogeography during the CPE and the atmosphere–ocean feedbacks associated with it.

Data availability

The simulations used in this study are identical to those described in ref. 7, and the full details are available therein. The data from the model simulations are available from https://www.paleo.bristol.ac.uk/unmodel/scripts/papers/Farnsworth_et_al_2019.html.

Code availability

The data generated from the model simulations was analyzed using standard Python packages.

References

- O'Brien, C. L. et al. Cretaceous sea-surface temperature evolution: constraints from TEX₈₆ and planktonic foraminiferal oxygen isotopes. *Earth Sci. Rev.* **172**, 224–247 (2017).
- Gaskell, D. E. et al. The latitudinal temperature gradient and its climate dependence as inferred from foraminiferal $\delta^{18}\text{O}$ over the past 95 million years. *Proc. Natl Acad. Sci. USA* **119**, e2111332119 (2022).
- Wang, Y. et al. Paleo-CO₂ variation trends and the Cretaceous greenhouse climate. *Earth Sci. Rev.* **129**, 136–147 (2014).
- Anagnostou, E. et al. Changing atmospheric CO₂ concentration was the primary driver of early Cenozoic climate. *Nature* **533**, 380–384 (2016).
- Godd ris, Y., Donnadi u, Y., le Hir, G., Lefebvre, V. & Nardin, E. The role of palaeogeography in the Phanerozoic history of atmospheric CO₂ and climate. *Earth Sci. Rev.* **128**, 122–138 (2014).
- Lunt, D. J. et al. Palaeogeographic controls on climate and proxy interpretation. *Climate* **12**, 1181–1198 (2016).
- Farnsworth, A. et al. Climate sensitivity on geological timescales controlled by nonlinear feedbacks and ocean circulation. *Geophys. Res. Lett.* **46**, 9880–9889 (2019).
- Donnadi u, Y., Pierrehumbert, R., Jacob, R. & Fluteau, F. Modelling the primary control of paleogeography on Cretaceous climate. *Earth Planet Sci. Lett.* **248**, 426–437 (2006).
- Fluteau, F., Ramstein, G., Besse, J., Guiraud, R. & Masse, J. P. Impacts of palaeogeography and sea level changes on Mid-Cretaceous climate. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* **247**, 357–381 (2007).
- Ladant, J.-B. et al. Paleogeographic controls on the evolution of Late Cretaceous ocean circulation. *Climate* **16**, 973–1006 (2020).
- Poulsen, C. J., Barron, E. J., Arthur, M. A. & Peterson, W. H. Response of the Mid-Cretaceous global oceanic circulation to tectonic and CO₂ forcings. *Paleoceanography* **16**, 576–592 (2001).
- Donnadi u, Y., Puc eat, E., Moiroud, M., Guillocheau, F. & Deconinck, J.-F. A better-ventilated ocean triggered by Late Cretaceous changes in continental configuration. *Nat. Commun.* **7**, 10316 (2016).
- Ladant, J.-B. & Donnadi u, Y. Palaeogeographic regulation of glacial events during the Cretaceous supergreenhouse. *Nat. Commun.* **7**, 12771 (2016).
- Heinemann, M., Jungclauss, J. H. & Marotzke, J. Warm Paleocene/Eocene climate as simulated in ECHAM5/MPI-OM. *Climate* **5**, 785–802 (2009).
- Toumoulin, A. et al. Quantifying the effect of the Drake passage opening on the Eocene Ocean. *Paleoceanogr. Palaeoclimatol.* **35**, e2020PA003889 (2020).
- Shellito, C. J., Lamarque, J.-F. & Sloan, L. C. Early Eocene Arctic climate sensitivity to $p\text{CO}_2$ and basin geography. *Geophys. Res. Lett.* **36**, L09707 (2009).
- Stommel, H. The westward intensification of wind-driven ocean currents. *Trans. Am. Geophys. Union* **29**, 202 (1948).
- Bye, J. A. T. & Veronis, G. A correction to the sverdrup transport. *J. Phys. Oceanogr.* **9**, 649–651 (1979).
- Bye, J. A. T. & Veronis, G. Poleward heat flux by an ocean gyre. *Dyn. Atmos. Oceans* **4**, 101–114 (1980).
- Gianchandani, K., Gildor, H. & Paldor, N. On the role of domain aspect ratio in the westward intensification of wind-driven surface ocean circulation. *Ocean Sci.* **17**, 351–363 (2021).
- Valdes, P. J. et al. The BRIDGE HadCM3 family of climate models: HadCM3@Bristol v1.0. *Geosci. Model Dev.* **10**, 3715–3743 (2017).
- Valdes, P. J., Scotese, C. R. & Lunt, D. J. Deep ocean temperatures through time. *Climate* **17**, 1483–1506 (2021).
- Foster, G. L., Royer, D. L. & Lunt, D. J. Future climate forcing potentially without precedent in the last 420 million years. *Nat. Commun.* **8**, 14845 (2017).
- Scotese, C. R., Song, H., Mills, B. J. W. & van der Meer, D. G. Phanerozoic paleotemperatures: the earth's changing climate during the last 540 million years. *Earth Sci. Rev.* **215**, 103503 (2021).
- Seton, M. et al. Global continental and ocean basin reconstructions since 200Ma. *Earth Sci. Rev.* **113**, 212–270 (2012).
- Lawver, L. A., Grantz, A. & Gahagan, L. M. Plate kinematic evolution of the present Arctic region since the Ordovician. in *Tectonic Evolution of the Bering Shelf-Chukchi Sea-Artic Margin and Adjacent Landmasses* (eds Miller, E. L., Grantz, A. & Klemperer, S. L.) **360** (Geological Society of America, 2002).
- Hay, W. Cretaceous Oceans and Ocean Modeling. in *Cretaceous Ocean Redbeds* 243–271 <https://doi.org/10.2110/sepmp.091.233> (SEPM (Society for Sedimentary Geology) 2009).

28. Barron, E. J. & Peterson, W. H. The Cenozoic ocean circulation based on ocean General Circulation Model results. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* **83**, 1–28 (1991).
 29. Manabe, S. & Wetherald, R. T. The effects of doubling the CO₂ Concentration on the Climate of A General Circulation Model. *J. Atmos. Sci.* **32**, 3–15 (1975).
 30. Holland, M. M. & Bitz, C. M. Polar amplification of climate change in coupled models. *Clim. Dyn.* **21**, 221–232 (2003).
 31. Stuecker, M. F. et al. Polar amplification dominated by local forcing and feedbacks. *Nat. Clim. Chang.* **8**, 1076–1081 (2018).
 32. Cavalheiro, L. et al. Impact of global cooling on Early Cretaceous high pCO₂ world during the Weissert Event. *Nat. Commun.* **12**, 5411 (2021).
 33. Held, I. M. The partitioning of the poleward energy transport between the tropical ocean and atmosphere. *J. Atmos. Sci.* **58**, 943–948 (2001).
 34. Adam, O., Farnsworth, A. & Lunt, D. J. Modality of the tropical rain belt across models and simulated climates. *J. Clim.* <https://doi.org/10.1175/JCLI-D-22-0521.1> (2022).
 35. Rae, J. W. B. et al. Atmospheric CO₂ over the past 66 million years from marine archives. *Annu. Rev. Earth Planet Sci.* **49**, 609–641 (2021).
 36. Cramer, B. S., Miller, K. G., Barrett, P. J. & Wright, J. D. Late Cretaceous–Neogene trends in deep ocean temperature and continental ice volume: reconciling records of benthic foraminiferal geochemistry ($\delta^{18}\text{O}$ and Mg/Ca) with sea level history. *J. Geophys. Res. Oceans* **116**, C12023 (2011).
 37. Poulsen, C. J., Gendaszek, A. S. & Jacob, R. L. Did the rifting of the Atlantic Ocean cause the Cretaceous thermal maximum? *Geology* **31**, 115–118 (2003).
 38. Czaja, A. & Marshall, J. The partitioning of poleward heat transport between the atmosphere and ocean. *J. Atmos. Sci.* **63**, 1498–1511 (2006).
 39. Markwick, P. J. & Valdes, P. J. Palaeo-digital elevation models for use as boundary conditions in coupled ocean–atmosphere GCM experiments: a Maastrichtian (late Cretaceous) example. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* **213**, 37–63 (2004).
 40. Adam, O. et al. The TropD software package (v1): standardized methods for calculating tropical-width diagnostics. *Geosci. Model Dev.* **11**, 4339–4357 (2018).
- A.F. acknowledges the Chinese Academy of Sciences Visiting Professorship for Senior International Scientists grant 2021FSE0001.

Author contributions

K.G. conceived the initial idea, developed the analytical model, and applied it to the HadCM3L data. O.A., H.G., and N.P. contributed to the development of the analytical model, and S.M. contributed to the application of the analytical model to the HadCM3L data. A.F. and D.J.L. carried out the HadCM3L paleoclimate simulations. K.G. drafted the initial version of the paper, and all other authors contributed to the text.

Competing interests

The authors declare no competing interests.

Additional information

Supplementary information The online version contains supplementary material available at <https://doi.org/10.1038/s41467-023-40905-7>.

Correspondence and requests for materials should be addressed to Kaushal Gianchandani.

Peer review information *Nature Communications* thanks Wolf Dumman and the other anonymous reviewer(s) for their contribution to the peer review of this work. A peer review file is available.

Reprints and permissions information is available at <http://www.nature.com/reprints>

Publisher's note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

© The Author(s) 2023

Acknowledgements

K.G. and H.G. were supported by the Joint National Natural Science Foundation of China–Israel Science Foundation (research grant number 2547/17) and by the U.S.–Israel Binational Science Foundation (BSF grant number 2018152). D.J.L. and A.F. acknowledge Leverhulme grant RPG-2019-365; D.J.L. acknowledges NERC grant NE/X000222/1;