

SCIENTIFIC REPORTS



OPEN

Seed Pelleting with Calcium Peroxide Improves Crop Establishment of Direct-seeded Rice under Waterlogging Conditions

Junhao Mei¹, Weiqin Wang¹, Shaobing Peng¹ & Lixiao Nie^{1,2}

Poor crop establishment of direct-seeded rice (DSR) is one of the major constraints to wide adoption of DSR, particularly in areas prone to flooding after sowing or where fields are not level. Seed pelleting is an effective, practical and facile technique to enhance crop establishment under unfavorable environmental conditions. To evaluate the effects of seed pelleting on rice germination, seedling growth and associated metabolic events under waterlogging stress, various seed pelleting treatments including formulae, pelleting times (the weight ratio of pelleting agents: rice seeds = 1:1~7:1 (w/w) and CaO₂ contents were tested in series of experiments. Naked seeds were maintained for comparison as a control. Pelleting treatments with CaO₂ significantly increased seed germination and seedling growth of DSR under waterlogging conditions compared with pelleted seeds without CaO₂ and naked seeds. The optimum weight ratios of CaO₂ to dry seeds were found to be in the range of 0.6:1–1:1 based on seed germination and seedling growth performance under waterlogging conditions. Under waterlogging conditions, high seed germination percentage and vigorous seedling growth of DSR due to seed pelleting with CaO₂ was associated with an increase in α -amylase activity, but decrease in alcohol dehydrogenase (ADH) and pyruvate decarboxylase (PDC) activities in pelleted seeds.

Rice (*Oryza sativa* L.), a staple food for more than half of the world's population, is grown on 163.2 million hectares, with an average annual production of 740.9 million tones across the globe¹. China is the main producer of rice, contributing more than 29% of total global rice production². Therefore, the stability of China's rice production is an indispensable part of the global food security³. In China, traditional transplanted flooded rice is the major planting pattern of rice and nearly 50% of the rice is grown under such conditions⁴. However, in recent years, depleting water resources, labor shortage and increasing labor cost have threatened the sustainability and productivity of transplanted flooded rice systems, suggesting that traditionally transplanted rice is no longer suitable for sustainable development in China^{5,6}.

Taking the advantages of saving water and labor and increasing system productivity, DSR has been believed to be an optimal option for rice production⁷. Tao *et al.* reported that direct-seeded rice was a feasible and sustainable alternative to transplanted rice in Central China based on comparable or even higher yield performances and greater water use efficiency, as well as lower global warming potential and conducive to mechanization⁸. Many countries have shifted from traditional transplanted flooded rice to DSR^{9,10}. At present, 23% of rice is direct-seeded globally¹¹. In China, the planting area for direct-seeded rice is increasing rapidly. By 2012, the percentage of direct-seeded rice area to total rice planting area in China increased to 28%⁴.

Nevertheless, poor crop establishment remains a major obstacle for large-scale adoption of DSR, particularly in areas prone to flooding after sowing or where fields are not level^{12–14}. Globally, the flood-prone ecosystem

¹National Key Laboratory of Crop Genetic Improvement, MOA Key Laboratory of Crop Ecophysiology and Farming System in the Middle Reaches of the Yangtze River, College of Plant Science and Technology, Huazhong Agricultural University, Wuhan, Hubei, 430070, China. ²Hubei Collaborative Innovation Center for Grain Industry, Yangtze University, Jingzhou, Hubei, 434023, China. Correspondence and requests for materials should be addressed to L.N. (email: nielixiao@mail.hzau.edu.cn)




Figure 1. Germination dynamics of pelleted seeds in different formulae with various pelleting times and CaO_2 contents under waterlogging conditions (1.5 cm water layer) in Experiment 1. (a) Formula 1. (b) Formula 2. (c) Formula 3. (d) Formula 4. CK: naked seeds. CP: Calcium peroxide. CP1: Seeds were pelleted by pelleting agents with 10% of CaO_2 at a pelleting time of 1. CP3: Seeds were pelleted by pelleting agents with 10% of CaO_2 at a pelleting time of 3. CP5: Seeds were pelleted by pelleting agents with 10% of CaO_2 at a pelleting time of 5. CP7: Seeds were pelleted by pelleting agents with 10% of CaO_2 at a pelleting time of 7. NCP1: Seeds were pelleted by pelleting agents without 10% of CaO_2 at a pelleting time of 1. NCP3: Seeds were pelleted by pelleting agents without 10% of CaO_2 at a pelleting time of 3. NCP5: Seeds were pelleted by pelleting agents without 10% of CaO_2 at a pelleting time of 5. NCP7: Seeds were pelleted by pelleting agents without 10% of CaO_2 at a pelleting time of 7. Seed germination data were recorded from 3 DAS until constant at 8 DAS. Error bars indicate standard error ($n = 3$).

accounts for about 9% of total rice lands, but for many countries in South and Southeast Asia, flood-prone rice can represent more than 25% of total rice lands, and the submergence duration varies from 1 to 10 days¹⁵. In China, 1/10 of the land areas, 33 million ha of farmland, 70% of the agricultural production are threatened by flood stress¹⁶. The waterlogging after rice seeding would result in seed floating, or the low seed germination rates, as well as uneven and irregular emergence of rice¹⁷. Furthermore, prolonged duration of flooding would result in wilting and death of the seeds and seedlings, inadequate crop stands, and eventually the reduction of grain yield^{18,19}.

In recent years, various strategies have been employed to improve abiotic stress tolerance during seed germination. Seed pelleting is an effective, practical and facile technique to enhance rapid and uniform emergence, high seedling vigor, and better yields in many field crops particularly under unfavorable environmental conditions²⁰. Pelleting is one type of seed coating technology that is defined as the deposition of a layer of inert materials that may obscure the original shape and size of the seed, resulting in the shape and the general size of the raw seed changed, and a substantial weight increase and improved plantability^{21,22}. Ryu *et al.* reported that seed germination and root colonization of sesame were significantly improved when the sesame seeds were pelleted with a combination of clay, vermiculite and talc; consequently the grain yield was increased by 10.2%²³. Previous research demonstrated that seed pelleting with ammonium molybdate and ferrous sulfate in rhizobium inoculated seeds improved growth parameters of soybean and yield²⁴. Asagi *et al.* indicated that pelleted rice seeds with clay or diatomaceous clay increased germination percentage by 43% and 26%, respectively, compared with naked seeds²⁵. Guan *et al.* pelleted tobacco seeds using the combined materials with a superabsorbent polymer, poly hydrogel, and salicylic acid enhanced drought-tolerance and significantly improved seed germination, as well as seedling growth²⁶. Other studies found that seeds pelleting with calcium peroxide (CaO_2) promoted

Formula	Clay (g)	Talc (g)	Attapulgate (g)	Bentonite (g)	Total weight (g)
1	66.7	33.3	—	—	100
2	—	71.4	28.6	—	100
3	50	50	—	—	100
4	30	60	—	10	100

Table 1. Formulae of rice seed-pelleting agents used in the experiments. In experiment 1, Formulae 1–4 were adopted. In experiment 2, Formulae 1 and 2 were adopted. In experiment 3, Formulae 1 and 2 were adopted. In field experiment, Formulae 1 and 2 were adopted.

germination of rice directly sown into flooded soil, and improved plant growth^{27–30}. However, little work has been done regarding the comparative performance of different seed pelleting formulae in enhancing the submergence tolerance of rice and to explore the mechanisms of pelleting-induced stress tolerance. The objectives of the present study were (1) to examine the effects of various seed pelleting formulae with various pelleting times (the weight ratio of pelleting agents: rice seeds = 1:1~7:1 (w/w)) on seed emergence performance under waterlogging conditions during seed germination and (2) to unravel the physiological and biochemical changes in rice seedlings under the influence of seed pelleting and waterlogging stress to obtain a better understanding of pelleting-induced mechanisms.

Results

Experiment 1: Rice seed germination and seedling growth between pelleted and naked rice seeds under waterlogging conditions. *Seed germination.* There were significant ($p \leq 0.05$) variations in rice seed germination percentages among different seed pelleting treatments under waterlogging conditions (Fig. 1). The seed germination percentages were significantly higher in the pelleting treatments with 10% of CaO₂ than those pelleting treatments without CaO₂ and naked seeds under waterlogging conditions. Furthermore, the seed germination percentage was increased as the pelleting times increased (Fig. 1). Nevertheless, seed pelleting without CaO₂ was not effective in rice seed emergence under waterlogging conditions. At 8 DAS, the maximum seed germination percentages were above 80% in all formulae, when pelleted with 10% of CaO₂ at a pelleting time of 7 under waterlogging conditions. While the seed germination percentage of naked seeds was only 31.1%.

Seedling growth. Pelleting treatments with CaO₂ significantly enhanced rice seedling growth under waterlogging conditions compared with pelleted seeds without CaO₂ and naked seeds. Moreover, seedling growth parameters were increased as the pelleting times increased (Table 2). The highest plant growth parameters were observed in pelleting treatments with 10% of CaO₂ at a pelleting time of 7 (Table 2). On average, seed pelleting with CaO₂ treatments significantly increased the shoot length, root length, shoot fresh weight, and root fresh weight by 92.99%, 33.49%, 119.39%, and 87.93%, respectively, compared with naked seeds control. While seed pelleting treatments without CaO₂ did not consistently increase seedling growth parameters compared to the naked seed control. No significant differences in seedling growth parameters were observed among different formulae, as well as pelleting treatments without CaO₂ and naked seed control.

Experiment 2: Effects of different CaO₂ contents and pelleting times on seed germination and early seedling growth of pelleted rice under waterlogging conditions. *Seed germination.* All of the pelleting treatments with CaO₂ significantly enhanced rice seed emergence of rice compared with pelleted without CaO₂ and naked seed control (Fig. 2). When the weight ratio of pelleting agents to rice seed was 1:1, an increase in CaO₂ content significantly increased the germination percentage under waterlogging conditions. The maximum germination rates at 8 DAS were 87.8% and 84.4% in formulae 1 and 2, respectively, when the CaO₂ content in the pelleting agents was increased to 60%. When the pelleting times were 3:1 or 5:1, the highest seed germination percentages were observed at a CaO₂ content of 20%. When the pelleting time was increased to 7:1, the germination percentages of pelleted seeds decreased with the increase in content of CaO₂, and the highest germination rates at 8 DAS were 86.7% and 83.3%, respectively, at 10% CaO₂ content in formulae 1 and 2.

Seedling growth. All of the pelleting treatments with CaO₂ significantly enhanced rice seedling growth compared with pelleted without CaO₂ and naked seed control in both formulae 1 and 2 (Table 3). The highest seedling growth parameters were found at 60% of CaO₂ content when the pelleting time was 1:1; at 20% of CaO₂ content when the pelleting times was 3:1 or 5:1; at 10% of CaO₂ content when the pelleting times was 7:1 in both formulae 1 and 2. The shoot length, root length and their fresh weights increased with the increase in CaO₂ content, but these parameters were decreased when the weight ratio of CaO₂ to rice seed was over 1:1. When the pelleting time was increased to 7:1, seedling growth parameters showed continuing downward trends as the content of CaO₂ increased in formulae 1 and 2 under waterlogging conditions.

The optimum weight ratio of CaO₂ to dry seeds was determined based on the highest germination percentage and the best seedling growth, which was achieved under a certain CaO₂ content at a certain pelleting time. In this study, the optimum weight ratio of CaO₂ to dry seeds was at 0.6:1–1:1 (a CaO₂ content × a pelleting time). Adverse effects on seed germination and seedling growth was observed when the ratio was over 1:1.

Experiment 3: The mechanisms underlying the waterlogging tolerance during seed germination and seedling growth induced by rice seed pelleting with CaO₂. *Seed germination and Seedling*

Treatment	CaO ₂ content	Pelleting time	Shoot length	Root length	Shoot FW	Root FW
			(cm)	(cm)	(mg seedling ⁻¹)	(mg seedling ⁻¹)
Naked seed		0	4.89 ± 0.10 efg	6.50 ± 0.31 bc	14.67 ± 0.59 d	8.08 ± 0.92 e
Pelleting Formula 1	0	1:1	4.28 ± 0.35 g	6.36 ± 0.23 c	13.64 ± 0.97 d	9.68 ± 0.85 de
	0	3:1	4.64 ± 0.30 fg	6.03 ± 0.96 c	14.10 ± 0.89 d	7.72 ± 0.68 e
	0	5:1	5.37 ± 0.16 e	6.43 ± 0.39 c	15.02 ± 0.64 d	8.67 ± 0.79 e
	0	7:1	5.22 ± 0.29 ef	6.40 ± 0.58 c	16.33 ± 1.10 d	8.93 ± 0.62 e
	10%	1:1	8.14 ± 0.19 d	7.16 ± 0.95 bc	29.80 ± 0.68 c	11.64 ± 1.16 cd
	10%	3:1	9.23 ± 0.30 c	8.13 ± 0.49 ab	29.96 ± 1.11 c	13.26 ± 0.30 bc
	10%	5:1	10.42 ± 0.20 b	9.42 ± 0.29 a	35.21 ± 0.94 b	15.12 ± 0.75 b
Naked seed		0	4.89 ± 0.10 d	6.50 ± 0.31 de	14.67 ± 0.59 d	8.08 ± 0.92 c
Pelleting Formula 2	0	1:1	4.53 ± 0.35 d	6.16 ± 0.30 ef	13.60 ± 1.47 d	8.24 ± 1.15 c
	0	3:1	5.12 ± 0.19 d	5.48 ± 0.18 f	13.84 ± 0.75 d	6.88 ± 0.80 c
	0	5:1	5.00 ± 0.12 d	5.90 ± 0.25 ef	14.03 ± 0.44 d	8.30 ± 0.42 c
	0	7:1	6.42 ± 0.25 c	6.55 ± 0.55 de	21.05 ± 0.80 c	12.60 ± 0.80 b
	10%	1:1	7.23 ± 0.42 c	7.15 ± 0.45 d	26.88 ± 0.68 b	13.60 ± 0.61 b
	10%	3:1	8.89 ± 0.42 b	8.13 ± 0.42 c	27.59 ± 2.29 b	13.71 ± 0.84 b
	10%	5:1	10.25 ± 0.20 a	10.10 ± 0.23 a	34.67 ± 1.76 a	17.12 ± 1.29 a
Naked seed		0	4.89 ± 0.10 de	6.50 ± 0.31 b	14.67 ± 0.59 e	8.08 ± 0.92 de
Pelleting Formula 3	0	1:1	4.85 ± 0.21 de	5.56 ± 0.30 b	13.59 ± 1.30 e	8.06 ± 0.23 de
	0	3:1	4.60 ± 0.26 e	5.66 ± 0.50 b	13.49 ± 1.66 e	6.56 ± 0.59 e
	0	5:1	4.92 ± 0.24 de	6.44 ± 0.47 b	15.04 ± 1.54 de	8.63 ± 1.44 cde
	0	7:1	5.76 ± 0.42 d	6.22 ± 0.75 b	19.03 ± 1.35 d	10.87 ± 1.09 cd
	10%	1:1	7.61 ± 0.48 c	6.64 ± 0.41 b	26.18 ± 2.92 c	11.96 ± 2.14 bc
	10%	3:1	8.47 ± 0.34 c	9.22 ± 0.25 a	27.60 ± 1.08 c	15.73 ± 0.54 a
	10%	5:1	9.95 ± 0.56 b	9.72 ± 0.61 a	33.11 ± 0.59 b	15.22 ± 1.29 ab
Naked seed		0	4.89 ± 0.10 e	6.50 ± 0.31 bc	14.67 ± 0.59 c	8.08 ± 0.92 c
Pelleting Formula 4	0	1:1	4.85 ± 0.19 e	5.47 ± 0.27 c	13.91 ± 1.69 c	6.91 ± 0.16 c
	0	3:1	4.66 ± 0.20 e	5.38 ± 0.31 c	13.47 ± 0.70 c	6.95 ± 0.16 c
	0	5:1	5.39 ± 0.36 de	6.59 ± 0.38 bc	15.60 ± 0.92 c	9.82 ± 0.61 bc
	0	7:1	5.87 ± 0.03 d	7.57 ± 0.67 b	17.93 ± 2.67 c	11.79 ± 0.21 b
	10%	1:1	6.98 ± 0.68 c	7.12 ± 0.61 b	23.25 ± 2.18 b	12.37 ± 1.79 b
	10%	3:1	7.75 ± 0.35 c	7.64 ± 0.13 b	26.08 ± 1.87 b	12.77 ± 1.13 b
	10%	5:1	10.42 ± 0.05 b	9.88 ± 0.69 a	36.02 ± 0.88 a	17.40 ± 1.27 a
		10%	11.31 ± 0.33 a	9.97 ± 0.54 a	37.93 ± 1.72 a	16.89 ± 1.38 a
Formula			ns	ns	ns	ns
CaO ₂ content			**	**	**	**
Pelleting time			**	**	**	**
Formula × CaO ₂ content			*	ns	ns	ns
Formula × Pelleting time			ns	ns	ns	ns
CaO ₂ content × Pelleting time			**	**	**	*
Formula × CaO ₂ content × Pelleting time			ns	ns	ns	ns

Table 2. The effect of different seed pelleting formulae with various pelleting times on seedling growth parameters of direct-seeded rice under waterlogging conditions (1.5 cm water layer) at 8 DAS in Experiment 1. Pelleting time is the weight ratio of pelleting agents to rice seeds. Within a column under each pelleting formula, means followed by different letters are significantly different at a 0.05 probability level according to a least significant difference (LSD) test. The naked seeds control were not involved in three-way ANOVA. FW: fresh weight. ** and * denote significance at the 0.01 and 0.05 probability level, respectively. ns: non-significant.

growth. Results revealed that seed pelleting treatments at a pelleting time of 3:1 with 20% of CaO₂ significantly ($p \leq 0.05$) enhanced seed germination and seedling growth compared with naked seeds under waterlogging conditions, while there were no significant differences in seed germination and seedling growth between pelleted seed treatments and the naked seed control under normal condition (Fig. 3 and Table 4). The germination percentages in the two pelleted treatments under waterlogging conditions and three treatments under normal conditions were above 86.9%, which was significantly higher than naked seeds (45.6%) under waterlogging conditions (Fig. 3).




Figure 2. Germination dynamics of pelleted seeds in two formulae with various pelleting times and CaO_2 contents under waterlogging conditions (1.5 cm water layer) in Experiment 2. Seeds were pelleted in Formula 1 (a (pelleting time of 1), c (pelleting time of 3), e (pelleting time of 5), g (pelleting time of 7)) and Formula 2 (b (pelleting time of 1), d (pelleting time of 3), f (pelleting time of 5), h (pelleting time of 7)). CK: naked seeds. F + 0% CaO_2 : Seeds were pelleted by pelleting agents without CaO_2 . F + 10% CaO_2 : Seeds were pelleted by pelleting agents with 10% of CaO_2 . F + 20% CaO_2 : Seeds were pelleted by pelleting agents with 20% of CaO_2 . F + 40% CaO_2 : Seeds were pelleted by pelleting agents with 40% of CaO_2 . F + 60% CaO_2 : Seeds were pelleted by pelleting agents with 60% of CaO_2 . Seed germination data were recorded from 3 DAS until constant at 8 DAS. Error bars indicate standard error ($n=3$).

Treatment	Pelleting time	CaO ₂ content	Shoot length (cm)	Root length (cm)	Shoot FW (mg seedling ⁻¹)	Root FW (mg seedling ⁻¹)
Naked seed	0		1.92 ± 0.06 j	2.41 ± 0.37 g	6.21 ± 0.27 k	3.61 ± 1.31 gh
Pelleting Formula 1	1:1	0	2.25 ± 0.30 j	2.01 ± 0.91 gh	6.80 ± 0.66 k	3.79 ± 1.32 fgh
	1:1	10%	4.21 ± 0.48 gh	6.17 ± 0.40 cd	14.15 ± 1.80 i	8.63 ± 2.17 de
	1:1	20%	8.00 ± 0.64 c	8.38 ± 1.03 ab	27.22 ± 2.57 cd	17.02 ± 1.42 abc
	1:1	40%	9.22 ± 0.80 b	8.88 ± 0.61 ab	28.39 ± 2.29 cd	18.66 ± 0.47 a
	1:1	60%	10.54 ± 0.63 a	9.58 ± 0.80 a	33.84 ± 2.39 ab	18.36 ± 5.05 ab
	3:1	0	2.47 ± 0.34 ij	3.65 ± 1.14 fg	8.64 ± 1.44 k	4.78 ± 1.40 fg
	3:1	10%	7.83 ± 1.04 c	8.16 ± 1.46 ab	26.67 ± 2.98 cd	14.07 ± 1.68 c
	3:1	20%	9.96 ± 0.71 ab	8.35 ± 0.77 ab	35.18 ± 0.71 a	19.55 ± 0.58 a
	3:1	40%	6.59 ± 0.44 de	7.26 ± 0.97 bcd	25.22 ± 0.85 de	14.57 ± 0.90 c
	3:1	60%	5.05 ± 0.22 fg	5.66 ± 0.80 de	22.29 ± 0.72 ef	10.24 ± 0.79 d
	5:1	0	2.89 ± 0.22 ij	4.25 ± 1.42 ef	10.03 ± 1.24 jk	5.50 ± 2.49 efg
	5:1	10%	7.75 ± 0.43 cd	7.56 ± 0.86 bc	29.95 ± 0.53 bc	15.32 ± 0.85 bc
	5:1	20%	10.40 ± 0.18 a	8.23 ± 0.41 ab	34.77 ± 0.57 a	19.82 ± 2.18 a
	5:1	40%	5.82 ± 0.06 ef	4.21 ± 0.87 ef	20.30 ± 0.89 fg	7.09 ± 0.89 def
	5:1	60%	4.28 ± 0.24 gh	2.39 ± 0.81 g	16.30 ± 0.07 hi	4.81 ± 0.23 fg
	7:1	0	2.59 ± 0.16 lj	2.43 ± 1.82 g	8.34 ± 0.91 k	3.18 ± 1.74 gh
	7:1	10%	10.25 ± 0.33 ab	9.34 ± 1.10 a	34.60 ± 2.47 a	18.72 ± 3.72 a
	7:1	20%	7.90 ± 0.15 c	7.24 ± 0.69 bcd	26.90 ± 0.58 cd	14.93 ± 1.86 c
7:1	40%	5.03 ± 0.11 fg	3.19 ± 1.82 fg	18.10 ± 0.90 gh	5.87 ± 2.92 efg	
7:1	60%	3.61 ± 0.19 hi	0.62 ± 0.36 h	13.84 ± 0.73 lj	1.44 ± 0.75 h	
Naked seed	0		1.92 ± 0.06 g	2.41 ± 0.37 gh	6.21 ± 0.27 k	3.61 ± 1.31 jk
Pelleting Formula 2	1:1	0	1.74 ± 0.19 g	1.29 ± 0.90 h	5.04 ± 1.02 k	2.01 ± 0.37 k
	1:1	10%	5.12 ± 0.21 cd	5.19 ± 1.20 de	15.49 ± 1.18 i	9.93 ± 2.52 f
	1:1	20%	7.11 ± 0.23 b	7.48 ± 1.15 bc	21.73 ± 1.02 f	14.62 ± 0.11 de
	1:1	40%	9.43 ± 0.58 a	8.25 ± 1.09 abc	28.26 ± 1.85 cd	16.94 ± 2.44 cd
	1:1	60%	10.02 ± 0.17 a	8.62 ± 0.60 ab	31.46 ± 0.79 ab	20.96 ± 2.30 ab
	3:1	0	3.64 ± 0.78 ef	4.11 ± 1.54 ef	11.74 ± 1.56 j	5.56 ± 2.50 ijk
	3:1	10%	5.69 ± 0.66 c	6.77 ± 1.94 cd	19.75 ± 1.31 fg	11.56 ± 3.96 ef
	3:1	20%	9.60 ± 0.18 a	8.92 ± 0.43 ab	31.08 ± 0.49 bc	19.62 ± 1.31 abc
	3:1	40%	7.12 ± 0.33 b	7.90 ± 0.09 abc	25.03 ± 0.92 d	15.22 ± 0.42 d
	3:1	60%	5.31 ± 0.14 cd	4.93 ± 0.89 e	18.92 ± 0.84 fgh	9.59 ± 1.01 fg
	5:1	0	3.03 ± 0.21 f	4.13 ± 0.82 ef	9.63 ± 0.62 j	5.89 ± 0.68 hij
	5:1	10%	7.43 ± 0.51 b	8.42 ± 0.71 abc	26.09 ± 2.30 de	16.24 ± 4.63 cd
	5:1	20%	10.28 ± 0.32 a	9.43 ± 0.44 a	33.13 ± 0.53 ab	22.91 ± 0.86 a
	5:1	40%	5.32 ± 0.53 cd	4.63 ± 2.08 e	19.37 ± 0.87 fg	9.18 ± 3.22 fgh
	5:1	60%	4.61 ± 0.26 de	2.63 ± 1.07 fg	17.01 ± 0.99 ghi	4.61 ± 2.14 jk
	7:1	0	3.22 ± 0.21 f	4.34 ± 1.22 ef	11.46 ± 0.84 j	6.10 ± 0.81 g hij
	7:1	10%	10.43 ± 0.15 a	9.12 ± 0.20 ab	34.33 ± 0.77 a	18.19 ± 1.91 bcd
	7:1	20%	8.11 ± 0.45 b	8.22 ± 0.72 bc	27.82 ± 1.64 de	16.70 ± 1.32 cd
7:1	40%	5.66 ± 0.40 cd	4.05 ± 1.83 ef	21.75 ± 1.13 f	8.25 ± 1.52 fgj	
7:1	60%	3.76 ± 0.22 ef	1.10 ± 0.54 h	15.98 ± 0.36 hi	2.81 ± 1.13 jk	
Formula			ns	ns	ns	ns
Pelleting time			*	**	ns	**
CaO ₂ content			**	**	**	**
Formula × Pelleting time			ns	**	**	ns
Formula × CaO ₂ content			ns	ns	*	ns
CaO ₂ content × Pelleting time			**	**	**	**
Formula × CaO ₂ content × Pelleting time			ns	ns	ns	ns

Table 3. The effect of two seed pelleting formulae with various pelleting times and CaO₂ contents on seedling growth parameters of direct-seeded rice under waterlogging conditions (1.5 cm water layer) at 8 DAS in Experiment 2. Pelleting time is the weight ratio of pelleting agent to rice seeds. Within a column under each pelleting formula, means followed by different letters are significantly different at a 0.05 probability level according to a least significant difference (LSD) test. The naked seeds were not involved in three-way ANOVA. FW: fresh weight. ** and * denote significance at the 0.01 and 0.05 probability level, respectively. ns: non-significant.




Figure 3. Germination dynamics of pelleted rice seeds in two pelleting treatments under waterlogging (1.5 cm water layer) and normal conditions (without water layer) in Experiment 3. P₀: non-pelleted naked seeds; P₁: Seed pelleting time of three with CaO₂ content of 20% in formula 1; P₂: Seed pelleting time of three with CaO₂ content of 20% in formula 2. The seed germination data were recorded from 3 DAS until constant at 8 DAS. Error bars indicate standard error (n = 4).

Water condition	Pelleted Treatment	Shoot length	Root length	Shoot FW	Root FW
		(cm)	(cm)	(mg seedling ⁻¹)	(mg seedling ⁻¹)
Waterlogging (Water depth 1.5 cm)	P ₀	1.99 ± 0.07 c	1.86 ± 0.14 d	4.79 ± 0.30 e	1.70 ± 0.08 c
	P ₁	7.24 ± 0.13 b	9.73 ± 0.24 c	22.09 ± 0.78 d	18.92 ± 1.11 b
	P ₂	7.09 ± 0.30 b	9.84 ± 0.36 c	21.04 ± 0.47 d	19.00 ± 1.67 b
Control (Without standing water)	P ₀	7.99 ± 0.11 a	12.29 ± 0.04 b	24.35 ± 0.49 c	24.11 ± 0.65 a
	P ₁	8.31 ± 0.09 a	13.11 ± 0.57 a	28.71 ± 0.98 a	25.72 ± 1.01 a
	P ₂	8.13 ± 0.14 a	12.80 ± 0.27 ab	26.13 ± 0.49 b	23.76 ± 0.81 a

Table 4. The effect of two seed pelleting treatments on seedling growth parameters of direct-seeded rice under waterlogging (1.5 cm water layer) and normal (without standing water) conditions at 8 DAS in Experiment 3. Within a column under each water condition, means followed by different letters are significantly different at a 0.05 probability level according to a least significant difference (LSD) test. P₀: non-pelleted naked seeds; P₁: Seed pelleting time of three with CaO₂ content of 20% in formula 1; P₂: Seed pelleting time of three with CaO₂ content of 20% in formula 2. FW: fresh weight.

Amylase activity. Waterlogging reduced the α -amylase activity of rice seeds or seedlings compared with normal conditions (Fig. 4). Even under normal conditions, the two pelleting treatments slightly increased the α -amylase activity compared with the naked seed control. Under waterlogging conditions, both of the seed pelleting treatments at the pelleting time of 3:1 with 20% of CaO₂ content significantly improved the starch metabolism of rice seeds or seedlings. In all treatments, α -amylase activities were progressively increased and achieved the highest values at 7 DAS.

ADH and PDC activities. ADH activities in rice seeds or seedlings were significantly higher under waterlogging conditions than under normal conditions (Fig. 5a). Under waterlogging conditions, both seed pelleting treatments significantly decreased the ADH activities in rice seeds or seedlings compared with the naked seeds. There were no variations in ADH activities between two seed pelleting treatments under waterlogging or normal conditions. However, seed pelleting treatments didn't affect the ADH activities in rice seeds or seedlings consistently. The ADH activities in naked seeds under waterlogging conditions continuously increased as seed germination progressed. While in the other five treatments, the ADH activities were progressively decreased as seed germination progressed. Under waterlogging conditions, both seed pelleting treatments significantly decreased the PDC activities in rice seeds or seedlings compared with the naked seeds (Fig. 5b). There were no variations in PDC activities between two seed pelleting treatments under waterlogging or normal conditions.

Field experiment: Performance of pelleting treatments on seed germination and early seedling growth of pelleted rice under waterlogging conditions in field. *Seed germination and Seedling growth.* Seed pelleting treatments at a pelleting time of 3:1 with 20% of CaO₂ significantly ($p \leq 0.05$) enhanced seed germination and early seedling growth compared with naked seeds under waterlogged field conditions, while there were no significant differences between the two seed pelleting treatments (Fig. 6 and Table 5). These




Figure 4. Variations in α -amylase activity of pelleted rice seeds and seedlings in two pelleting treatments under waterlogging (1.5 cm water layer) and normal (without water layer) conditions at 0, 1, 3, 5, and 7 DAS in Experiment 3. P₀: non-pelleted naked seeds; P₁: Seed pelleting time of three with a CaO₂ content of 20% in formula 1; P₂: Seed pelleting time of three with a CaO₂ content of 20% in formula 2. Seeds at 0 DAS were tested under a dry weight basis, while seedlings at 1, 3, 5 and 7 DAS were tested under a fresh weight basis. Error bars above the mean indicate standard error (n = 4).




Figure 5. Variations in alcohol dehydrogenase (ADH) (a) and pyruvate decarboxylase (PDC) (b) activities of pelleted rice seeds and seedlings in two pelleting treatments under waterlogging (1.5 cm water layer) and normal (without water layer) conditions at 0, 1, 3, 5, and 7 DAS in Experiment 3. P₀: non-pelleted naked seeds; P₁: Seed pelleting time of three with a CaO₂ content of 20% in formula 1; P₂: Seed pelleting time of three with a CaO₂ content of 20% in formula 2. Seeds at 0 DAS were tested under a dry weight basis, while seedlings at 1, 3, 5 and 7 DAS were tested under a fresh weight basis. Error bars above the mean indicate standard error (n = 4).

were consistent with the results from the growth chamber experiments. On average, the germination percentages of the two pelleting treatments were 33.75% higher than the naked seeds. The shoot length, root length, shoot fresh weight, and root fresh weight in the pelleting treatments were increased by 65.06%, 33.86%, 172.61%, and 182.53%, respectively, compared with naked seeds control.

Discussion

Waterlogging seriously impeded seed germination and early seedling growth of DSR, pelleting treatments with calcium peroxide significantly increased seed germination and improved seedling growth. Ellis and Setter reported that the presence of O₂ in the floodwater was critical for the survival of seedlings under waterlogging conditions³¹. Miro and Ismail reported that rice seeds failed to develop roots and leaves even though they can germinate in submerged conditions³². Under waterlogging conditions, the elongations of the coleoptile and mesocotyl were stimulated, which facilitated the exposure to air^{33,34}. However, the fast elongation of the coleoptile and mesocotyl consumes much energy. Sometimes, waterlogging results in plant death if energy reserves are depleted before emergence³⁵. In this study, coleoptiles of the pelleted rice seeds with optimal CaO₂ content first expanded in a thin layer, and later, the coleoptile formed a sturdy morphology and rapid expansion of leaf after contact with air, while the naked seeds seldom germinated.




Figure 6. Germination dynamics of pelleted rice seeds in two pelleting treatments under waterlogging conditions in Field. P₀: non-pelleted naked seeds; P₁: Seed pelleting time of three with CaO₂ content of 20% in formula 1; P₂: Seed pelleting time of three with CaO₂ content of 20% in formula 2. The seed germination data were recorded from 3 DAS until constant at 8 DAS. Error bars indicate standard error (n = 4).

Treatment	Shoot length	Root length	Shoot FW	Root FW
	(cm)	(cm)	(mg seedling ⁻¹)	(mg seedling ⁻¹)
P ₀	7.37 ± 0.10 b	5.70 ± 0.27 b	7.76 ± 0.29 b	3.55 ± 0.19 b
P ₁	12.18 ± 0.37 a	7.49 ± 0.19 a	21.16 ± 0.66 a	10.56 ± 0.69 a
P ₂	12.15 ± 0.19 a	7.77 ± 0.26 a	21.15 ± 1.29 a	9.50 ± 0.37 a

Table 5. The effect of two seed pelleting treatments on seedling growth parameters of direct-seeded rice under waterlogging conditions at 9 DAS in Field Experiment. Within a column under each water condition, means followed by different letters are significantly different at a 0.05 probability level according to a least significant difference (LSD) test. P₀: non-pelleted naked seeds; P₁: Seed pelleting time of three with CaO₂ content of 20% in formula 1; P₂: Seed pelleting time of three with CaO₂ content of 20% in formula 2. FW: fresh weight.

Baker and Hatton reported that the available oxygen increased as the content of CaO₂ increased³⁶. In this study, increased content of CaO₂ in the pelleting agents significantly increased the oxygen release to flooding water. In Experiment 2, an increase in the content of CaO₂ from 10% to 60% at a pelleting time of 3:1, resulted in an increase in the amount of oxygen released from CaO₂ by 106% (data not shown). Previous reports suggested that elevated oxygen tension may be highly toxic to embryos, root development and root elongation³⁷. The chemical reactions of CaO₂ and water, which usually occur as $2\text{CaO}_2 + 2\text{H}_2\text{O} \rightarrow 2\text{Ca(OH)}_2 + \text{O}_2$, produce large amounts of calcium hydroxide and could form an alkaline environment for seeds when pelleted with excessive CaO₂. In this study, CaO₂ was added in the pelleting agents increased available oxygen for seed respiration during germination, relieved the anoxia or hypoxia caused by waterlogging conditions. In addition, CaO₂ may inhibit the proliferation of anaerobic microorganisms, which negatively affect the seed germination. A similar research reported that the germination of seeds coated with streptomycin or a mixture of streptomycin and CaO₂ was significantly reduced as compared with that of seeds coated with only CaO₂, which implied that limitation of bacterial pathogens with CaO₂ might not be as important as O₂ supply in crop establishment under waterlogging conditions³⁸. However, we could not exclude the sterilization effect from CaO₂ during seed germination in this study. The importance of the antimicrobial effect from CaO₂ will be evaluated in the future research.

During rice seed germination, oxygen plays an important role on decomposition of reserves in endosperm. While oxygen is limited under waterlogging conditions, CaO₂ produce large amounts of oxygen in water. Under aerobic conditions, α -amylase activity in rice seedlings was maintained at higher level because of higher expression of *Amy1A* gene compared with that in flooded soils³⁹. The α -amylase in rice is likely playing a crucial role for the successful degradation of starch into carbohydrates in the anoxic rice endosperm. Perata *et al.* and Guglielminetti *et al.* indicated that α -amylase activity of the anaerobically germinated rice seeds was markedly lower than that presented in aerobic conditions^{40,41}. Moreover, Saddam *et al.* proclaimed that submergence stress severely reduced α -amylase activity by limiting starch degradation, therefore, rice seed reserves were not well metabolized under submerged conditions⁴². The above findings were consistent with our results that extremely low α -amylase activities were observed for naked rice seeds or seedlings under waterlogging conditions.

Pronounced increases in α -amylase activities were recorded in seed pelleting treatments with CaO_2 under waterlogging conditions.

One of the best characterised plant responses to waterlogging is the metabolic switch from aerobic respiration to anaerobic fermentation. Under submerged conditions, glycolysis and alcoholic fermentation are important processes for the energy production of seeds or seedlings^{31, 42}. The two key enzymes involved in glycolysis and alcoholic fermentation are ADH and PDC^{43–45}. Magneschi and Perata noted that the inhibited aerobic respiration under submergence conditions resulted in reduced energy production and supply of intermediates needed for seed germination and seedling growth⁴⁶. Several reports had revealed the importance of increases in ADH and PDC activities in acceleration of plant growth and survival under anoxia^{47–50}. Tolerance to waterlogging stress during seed germination was expressed as rapid germination, coleoptile elongation, anaerobic respiration to sustain energy supply, and maintenance of the cellular extensibility of the growing embryo^{14, 34}. Energy supply is indeed a key factor for rice tolerance to oxygen deficiency. Plants tolerant of anaerobiosis maintain an active fermentative metabolism for a relatively long time⁴⁶, that accelerates carbohydrate consumption that can be expected to increase the energy production⁵¹, with a 2–3-fold increase in glycolytic flux under anoxia with respect to the aerobic control⁵⁰. Under aerobic conditions, activities of fermentative enzymes, ADH and PDC were decreased to 1/10–1/5 compared to that under anaerobic conditions⁵². In this study, seed pelleting treatments with CaO_2 decreased the ADH and PDC activities in rice seeds or seedlings compared with the naked seeds under waterlogging conditions. This was because CaO_2 increased the solubility of O_2 in water under waterlogging conditions, which may transfer to aerobic respiration in rice seeds or seedlings, thus providing enough energy for the survival or growth of rice seeds or seedlings under submergence conditions.

Conclusively, seed pelleting with CaO_2 can alleviate the negative effects of flooding stress on rice seed emergence and seedling establishment. High seed germination percentage and vigorous seedling growth due to seed pelleting was associated with an increase in α -amylase activity, but decreases in ADH and PDC activities in pelleted seeds. Seed pelleting with CaO_2 could be used as a strategy to improve the crop establishment of direct-seeded rice, when seeds suffered from waterlogging soon after seed sowing. However, more exact mechanisms underlying the waterlogging tolerance induced by seed pelleting with CaO_2 will be addressed in the future. And the effects of seed pelleting treatments need to be wide demonstrated and popularized in practical production.

Materials and Methods

Seed source. Seed of a widely grown Indica rice cultivar viz., Huanghuazhan (HHZ, inbred) were obtained from the Crop Physiology and Production Center, Huazhong Agricultural University, Wuhan, China. The initial germination rate of the seeds was >93%. Healthy and filled seeds selected from the same seed lot were used for all of the experiments. Prior to experimentation, seeds were stored in a refrigerator (-4°C).

Seed pelleting treatments. Based on previous research, four seed pelleting formulae were adopted for present studies (Table 1). In general, the pelleting formula consists of pelleting agents, including clay, talc, attapulgite and bentonite (all were produced by Shijiazhuang Keying Refractory Material Co., LTD, China) and the adhesive (solution of 1.5% (w/w) polyvinyl alcohol, produced by Sinopharm Chemical Reagent Co., LTD, China). The rice seeds were pelleted with a minitype seed coater (produced by Shanghai Huanghai Drug Testing Instrument Distribution CO., LTD, China) using the following procedures: Firstly, the seeds were quickly soaked in the tap water then air-dried without visible moisture on the surface of seeds. Secondly, 1/4 of the total pelleting agents were slowly placed into the rotate pan of the seed coater (200 mm diameter) at the rotating speed of 45 rpm. Then, the pre-treated seeds were placed into the pan, after then, 1/4 of the total adhesive was sprayed onto the seed surface for 2 min while rolling. Thirdly, another 1/4 of the total pelleting agents were added into the rotate pan with 1/4 of the total adhesive sprayed onto the seed surface, this step was repeated twice. Finally, after all of the pelleting agents and adhesive were added, the pan was continuously rotated for 5 min. The pelleted seeds were then air-dried for 2 days at room temperature. Naked seeds were used as a control.

In Experiment 1, four pelleting formulae were used (Table 1). To evaluate the effectiveness of CaO_2 on seed germination and seedling growth under waterlogging conditions, two treatments were set in each formula, with 10% (w/w) CaO_2 (with a purity of 70%) and without CaO_2 , and each treatment included various pelleting times (1, 3, 5 or 7). Because the germination and seedling growth performance of pelleted seeds with CaO_2 was better compared to naked seeds as well as pelleted seeds without CaO_2 , in Experiment 2, formula 1 (best performance) and 2 (previously reported) were selected. In order to examine the effectiveness of different contents of CaO_2 on seed germination and seedling growth, based on the proportions of ingredients unvaried, the CaO_2 content were set as 0%, 10%, 20%, 40% and 60% (w/w) in each pelleting formula with various pelleting times (1, 3, 5 and 7). In Experiment 3, the pelleting formulae were the same as that in Experiment 2, however, only 20% (w/w) of CaO_2 content at a pelleting time of three was adopted in Experiment 3.

Experimentation. Growth chamber experiments were carried out in the Crop Physiology and Production Center, Huazhong Agricultural University, Wuhan, China during October 2015 to July 2016. Plastic trays 31.0 cm \times 20.0 cm \times 12.0 m in size were filled with 7.50 kg of puddled soil which was collected from the field and then soil surface was leveled in trays. In each tray, 30 rice seeds for each treatment were equally sown on the soil surface in experiment 1 and 2 with three replications. In experiment 3, four replications with four trays with 1.5 cm water depth and the other four replications without standing water were used to sample for determinations of α -amylase, alcohol dehydrogenase and pyruvate decarboxylase activity. The experiments were laid out in a completely randomized design. All of the trays were placed in a growth chamber with a 12 h light period and a constant temperature of 25°C . The humidity during the course of the study was maintained at 60% in the growth

chamber. Water was applied to all trays when their moisture content declined. Germination of seeds was recorded on a daily basis according to AOSA until a constant count was achieved⁵³. Seeds were considered to be germinated when the hypocotyl length exceeded 2 mm. Seed germination percentage was taken as the ratio of the number of seeds germinated to the total number of seeds sown and was expressed as a percentage. At 8 DAS, 6 seedlings were randomly sampled from each treatment to record their shoot and root lengths. The seedlings were dissected into roots and shoots and their fresh weights were recorded immediately. For determination of α -amylase activity, 1.0 g of dry seeds (0 DAS) and seedlings sampled at 1, 3, 5 and 7 DAS including shoot and root were ground and mixed with 100 ml distilled water, and left for 24 h at 4 °C and then the mixture was filtered with filter paper (Whatman No. 42). The enzyme activity was determined by the dinitro-salicylic acid (DNS) method⁵⁴. To determine alcohol dehydrogenase activity and pyruvate decarboxylase activity, 0.1 g dry seeds (0 DAS) and seedlings sampled at 1, 3, 5 and 7 DAS were ground and detected by an alcohol dehydrogenase assay kit and a pyruvate decarboxylase test kit (Nanjing Jiancheng Bioengineering Institute).

The field experiments were conducted at the Zhangbang Village, Dajin Town, Wuxue County, Hubei Province, China, in May 2016. In case of seed floating or stress from waterlogging conditions when heavy rains occurred soon after direct sowing, 20% (w/w) of CaO₂ content at a pelleting time of three in formulae 1 and 2, which were consistent with that in Experiment 3. The treatments were randomly arranged using a randomized block design with four replicates. The plot area was 5 m². The water layer in each plot was kept 2–3 cm until the seed germination was stable. The germination of seeds was recorded on daily basis according to AOSA (1990) until a constant count was achieved. Germination percentage was taken as the ratio of the number of seeds germinated to the total number of seeds sown and is expressed as a percentage. At 9 DAS, 10 seedlings were randomly sampled from each plot to record their shoot and root length. The seedlings from each plot were then dissected into roots and shoots, and their fresh weight was recorded immediately.

Statistical Analysis. Data were analyzed to confirm variability following analysis of variance using Statistix 9.0. In the experiment 1 & field experiment, we used one-way ANOVA, pelleting treatments as grouping factors, germination, root length, shoot length, root fresh weight, shoot fresh weight as response variables. In the experiments 2 & 3, we used three-way ANOVAs, with formulae, CaO₂ contents and pelleting times as grouping factors, germination, root length, shoot length, root fresh weight, shoot fresh weight as response variables. But the naked seeds were not involved in three-way ANOVAs. The differences between treatments were separated using a Least Significance Difference (LSD) test at a 0.05 probability level.

References

1. FAOSTAT. FAO Statistical databases. www.fao.org. Food and Agriculture Organization (FAO) of the United Nations, Rome (2014).
2. Peng, S., Tang, Q. & Zou, Y. Current status and challenges of rice production in china. *Plant Prod. Sci.* **12**, 3–8 (2009).
3. FAOSTAT. FAO Statistical databases. www.fao.org. Food and Agriculture Organization (FAO) of the United Nations, Rome (2011).
4. Zhang *et al.* Development and transition of rice planting in China. *Agric. Sci. Technol.* **13**, 1270–1276 (2012).
5. Cai, H. & Chen, Q. Rice research in China in the early 21st century. *Chinese Rice Res. Newsl.* **8**, 14–16 (2000).
6. Li, Y. Water saving irrigation in china. *Irrig. Drain.* **55**, 327–336 (2006).
7. Kumar, V. & Ladha, J. K. Direct Seeding of Rice: Recent Developments and Future Research Needs. *Adv. Agron.* **111**, 297–413 (2011).
8. Tao, Y., Chen, Q., Peng, S., Wang, W. & Nie, L. Lower global warming potential and higher yield of wet direct-seeded rice in central china. *Agron. Sustain Dev.* **36**, 24 (2016).
9. Pandey, S. & Velasco, L. Economics of direct-seeding in Asia: Patterns of adoption and research priorities. In “Direct Seeding: Research Strategies and Opportunities” (Pandey, S., Mortimer, M., Wade, L., Tuong, T. P., Lopez, K. and Hardy, B. Eds.), Los Banos, Philippines: IRRI. pp. 3–14 (2002).
10. Liu *et al.* Dry direct-seeded rice as an alternative to transplanted-flooded rice in central china. *Agron. Sustain Dev.* **35**, 285–294 (2015).
11. Rao, A. N., Johnson, D. E., Sivaprasad, B., Ladha, J. K. & Mortimer, A. M. Weed management in direct-seeded rice. *Adv. Agron.* **93**, 153–255 (2007).
12. Washio, O. Direct seeding rice culture in Japan: its technical outlook. *Farming Japan.* **26**, 11–19 (1992).
13. Du, L. V. & Tuong, T. P. Enhancing performance of dry-seeded rice: effects of seed priming, seeding rate, and time of seeding. In: Pandey, S., Mortimer, M., Wade, L., Tuong, T. P., Lopez, K., Hardy, B. eds. Direct seeding: research strategies and opportunities. Los Banos, Philippines: IRRI, 241–256 (2002).
14. Ismail, A. M., Ella, E. S., Vergara, G. V. & Mackill, D. J. Mechanisms associated with tolerance to flooding during germination and early seedling growth in rice (*oryza sativa*). *Ann. Bot.* **103**, 197–209 (2009).
15. Bhuiyan, S. J., Abedin, M. L., Singh, V. P. & Hardy, B. Rice Research and Development in the Flood-Prone Ecosystem (Eds). International Rice Research Institute, Los Banos, Philippines. 237–249 (2004).
16. Chen, Y., Yin, Y. X., Chen, X. W. & Xu., W. C. Change and future pattern of provincial flood affected areas in China: Possible relationship with climate change. *Disaster Adv.* **5**, 321–326 (2012).
17. Sato, T. & Maruyama, S. Seedling emergence and establishment under drained conditions in rice direct-sown into puddled and leveled soil. -Effect of calcium peroxide seed coating and sowing depth. *Plant Prod. Sci.* **5**, 71–76 (2002).
18. Sharma, A. R. Direct seeding and transplanting for rice production under flood-prone lowland conditions. *Field Crop Res.* **44**, 129–137 (1995).
19. Sasaki, T. Plant breeding: rice in deep water. *Nat.* **442**–635 (2006).
20. Powell, A. A. & Matthews, S. Seed treatments: developments and prospects. *Outlook Agric.* **17**, 93–107 (1988).
21. Butler, R. Coatings, films and treatments. *Seed World.* **1993**, 18–24 (1993).
22. Taylor, A. G. *et al.* Seed enhancement. *J. New Seeds.* **1**, 103–103 (1998).
23. Ryu, C. M., Kim, J., Choi, O., Kim, S. H. & Chang, S. P. Improvement of biological control capacity of *paenibacillus polymyxa*, E681 by seed pelleting on sesame. *Biol. Control.* **39**, 282–289 (2006).
24. Ramesh, K. & Thirumurugan, V. Effect of seed pelleting and foliar nutrition on growth of soybean. *Madras Agric. J.* **88**, 465–468 (2001).
25. Asagi, N., Ueno, H., Sekiya, H. & Ogiwara, H. Establishment of rice seedlings by direct sowing of multiple seed pellets on paddy soil covered with legume living mulch. *Plant Prod. Sci.* **11**, 361–365 (2008).
26. Guan, Y. *et al.* An enhanced drought-tolerant method using SA-loaded PAPMS polymer materials applied on tobacco pelleted seeds. *Sci. world J.* **2014**, 752658–752658 (2014).

27. Yamada, N. Calcium peroxide as an oxygen supplier for crop plants. *Proc. Crop Sci. Soc. Jpn.* **21**, 65–66 (1951). (In Japanese with English summary).
28. Ota, Y. & Nakayama, M. Effect of seed coating with calcium peroxide on germination under submerged conditions in rice plant. *Proc. Crop Sci. Soc. Jpn.* **39**, 535–536 (1970). (In Japanese with English summary).
29. Yamauchi, M. & Chuong, P. V. Rice seedling establishment as affected by cultivar, seed coating with calcium peroxide, sowing depth, and water level. *Field Crop Res.* **41**, 123–134 (1995).
30. Yu, P. Y., Xiong, Y. F., Zou, Y. B., Xu, Q. G. & Xiong, H. R. Effects of seed pelleting agent on the seed establishment and seedling growth in rice. *Crop Res.* **18**, 141–144 (2004). (In Chinese with English abstract).
31. Ellis, M. H. & Setter, T. L. Hypoxia induces anoxia tolerance in completely submerged rice seedlings. *J. Plant Physiol.* **154**, 219–230 (1999).
32. Miro, B. & Ismail, A. M. Tolerance of anaerobic conditions caused by flooding during germination and early growth in rice (*Oryza sativa* L.). *Front. Plant Sci.* **4**, 1466–1478 (2012).
33. Magneschi, L., Kudahettige, R. L., Alpi, A. & Perata, P. Comparative analysis of anoxic coleoptile elongation in rice varieties: relationship between coleoptile length and carbohydrate levels, fermentative metabolism and anaerobic gene expression. *Plant Biology.* **11**, 561–573 (2009).
34. Simon, A. Relationship between coleoptile and mesocotyl elongation of upland rice (*Oryza sativa* L.) seedlings under submergence and soil-sand culture. *Afr. J. Agr. Res.* **6**, 6463–6472 (2011).
35. Bailey-Serres, J. & Voesenek, L. A. Flooding stress: acclimations and genetic diversity. *Annu. Rev. Plant Biol.* **59**, 313–39 (2008).
36. Baker, A. M. & Hatton, W. Calcium peroxide as a seed coating material for padi rice. *Plant Soil.* **99**, 379–386 (1987).
37. Siegel, S. M. & Gerschman, R. A study of the toxic effects of elevated oxygen tension on plants. *Physiol. Plantarum.* **12**, 314–323 (2006).
38. Sarlistyaningsih, L., Sivasithamparam, K. & Setter, T. L. Influence of waterlogging on germination and survival of lupin seeds (*Lupinus angustifolius* L. cv. Gungurru) coated with calcium peroxide and streptomycin. *Aust. J. Exp. Agric.* **35**, 537–541 (1995).
39. Hwang, Y. S., Thomas, B. R. & Rodriguez, R. L. Differential expression of rice alpha-amylase genes during seedling development under anoxia. *Plant Mol. Bio.* **40**, 911–920 (1999).
40. Perata, P., Pozueta-Romero, J., Akazawa, T. & Yamaguchi, J. Effect of anoxia on starch breakdown in rice and wheat seeds. *Planta.* **188**, 611–618 (1992).
41. Guglielminetti, L., Yamaguchi, J., Perata, P. & Alpi, A. Amylolytic activities in cereal seeds under aerobic and anaerobic conditions. *Plant Physiol.* **109**, 1069–1076 (1995).
42. Hussain *et al.* Comparative transcriptional profiling of primed and non-primed rice seedlings under submergence stress. *Front. Plant Sci.* **7**, 1125 (2016).
43. Reggiani, R., Brambilla, I. & Bertani, A. Effect of exogenous nitrate on anaerobic metabolism in excised rice roots. *J. Exp. Bot.* **36**, 1193–1199 (1985).
44. Rivoal, J., Ricard, B. & Pradet, A. Glycolytic and fermentative enzyme induction during anaerobiosis in rice seedlings. *Plant Physiol. Bioch.* **1**, 43–52 (1989).
45. Ito, O., Ella, E. & Kawano, N. Physiological basis of submergence tolerance in rainfed lowland rice ecosystem. *Field Crop Res.* **64**, 75–90 (1999).
46. Magneschi, L. & Perata, P. Rice germination and seedling growth in the absence of oxygen. *Ann. Bot.* **103**, 181–196 (2008).
47. Avadhani, P. N., Greenway, H., Lefroy, R. & Prior, L. Alcoholic fermentation and malate metabolism in rice germinating at low oxygen concentrations. *Funct. Plant Biol.* **5**, 15–25 (1978).
48. Alpi, A. & Beevers, H. Effects of O₂ concentration on rice seedlings. *Plant Physiol.* **71**, 30–4 (1983).
49. Setter, T. L. & Ella, E. S. Relationship between coleoptile elongation and alcoholic fermentation in rice exposed to anoxia. I. importance of treatment conditions and different tissues. *Ann. Bot.* **74**, 265–271 (1994).
50. Gibbs, J., Morrell, S., Valdez, A., Setter, T. L. & Greenway, H. Regulation of alcoholic fermentation in coleoptiles of two rice cultivars differing in tolerance to anoxia. *J. Exp. Bot.* **51**, 785–96 (2000).
51. Waters, I., Morrell, S., Greenway, H. & Colmer, T. D. Effects of anoxia on wheat seedlings: II. Influence of O₂ supply prior to anoxia on tolerance to anoxia, alcoholic fermentation, and sugar levels. *J. Exp. Bot.* **42**, 1437–1447 (1991).
52. Mustrup, A. *et al.* Organ specific analysis of the anaerobic primary metabolism in rice and wheat seedlings II: Light exposure reduces needs for fermentation and extends survival during anaerobiosis. *Planta.* **225**, 139–152 (2006).
53. AOSA. Rules for testing seeds. Association of Official Seed Analysts (AOSA). *J. Seed. Technol.* **12**, 1–112 (1990).
54. Bernfeld, P. *Methods in enzymology* [149–158] (Academic Press, New York, USA, 1955).

Acknowledgements

This work is supported by the National Natural Science Foundation of China (Project No. 31371571), the National High Technology Research and Development Program of China (863 Program) (Project No. 2014AA10A605), and Special Fund for Agro-scientific Research in the Public Interest of China (Project No. 201203096).

Author Contributions

J.M. and L.N. initiated and designed the research, J.M. and Q.W. performed the experiments, J.M. and L.N. analyzed the data and wrote the manuscript, S.P. revised and edited the manuscript and also provided advice on the experiments.

Additional Information

Competing Interests: The authors declare that they have no competing interests.

Publisher's note: Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons license and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>.

© The Author(s) 2017