



OPEN

## Seasonal variations of the relationships between measures of training monotony and strain in professional soccer players

Hadi Nobari<sup>1,2,3,4</sup>✉, Alexandre Duarte Martins<sup>5,6,7</sup>, Rafael Oliveira<sup>5,6,8</sup> & Luca Paolo Ardigo<sup>9</sup>

The purposes of this study were (a) to determine the variations in internal and external measures of training monotony (TM) and strain (TS) in professional soccer players according to periods of the season and playing positions, and (b) to analyze the relationships between internal and external measures of TM and TS. Twenty male professional players (age = 29.4 ± 4.4 years) were followed for 20 weeks through session rating of perceived exertion (s-RPE), total distance (TD), high-speed running distance (HSRD) and sprint distance (SpD). Regardless of measure, highest mean TM and TS scores were observed in mid-season and end-season. In general, wingers and strikers tended to have greater values in TM. Midfielders exhibited greater TS of TD and SpD. Correlation results for TM revealed that s-RPE was positively associated with SpD in early-season ( $r = 0.608$ ) and negatively associated in mid-season ( $r = -0.506$ ). Regarding the TS, result demonstrated that s-RPE is negatively associated with HSRD in early-season ( $r = -0.464$ ) and positively associated in mid-season ( $r = 0.476$ ). In general, there different meanings in correlations between internal and external measures across the season. On the one hand, our findings highlighted that TM and TS of professional soccer players is sensitive to period of the season and player's position, but on other hand, correlation analyses proved that changes in one external/internal measure does not cause changes in another external/internal measure which support the constant monitoring of these values across the season.

The knowledge of load dynamics in soccer would help coaches, their staff, and practitioners to improve performance and at the same time to avoid fatigue, injury and illness<sup>1</sup>. In addition, it is well known that load measures may vary from session to session<sup>2-4</sup> week to week<sup>5-8</sup>, mesocycle to mesocycle<sup>2,5,9</sup> and/or period to period<sup>4,6,10-13</sup>.

Some indexes that allow to interpret load variations are known as training monotony (TM) and training strain (TS)<sup>14</sup>. While TM is calculated through the daily mean load divided by the week standard deviation, TS is based on TM multiplied by the accumulated load of the week<sup>15</sup>.

Another major factor that influences load dynamic is the positions of the players. Previous studies reported that player positions have different physical roles and consequently different load during matches<sup>16-20</sup>. Such differences were also revealed in training<sup>2,4,21</sup> and recently it was shown that external defenders and wingers presented greater TS for high-speed running and number of sprints during the season compared to the remaining positions. However, another study found no significant differences between positions for TM and TS calculated through

<sup>1</sup>Department of Exercise Physiology, Faculty of Educational Sciences and Psychology, University of Mohagheh Ardabili, Ardabil, Iran. <sup>2</sup>Department of Motor Performance, Faculty of Physical Education and Mountain Sports, Transilvania University of Braşov, 500068 Brasov, Romania. <sup>3</sup>Faculty of Sport Sciences, University of Extremadura, 10003 Cáceres, Spain. <sup>4</sup>Sepahan Football Club, Isfahan 81887-78473, Iran. <sup>5</sup>Sports Science School of Rio Maior-Polytechnic Institute of Santarém, 2040-413 Rio Maior, Portugal. <sup>6</sup>Life Quality Research Centre, 2040-413 Rio Maior, Portugal. <sup>7</sup>Departamento de Desporto e Saúde, Comprehensive Health Research Centre (CHRC), Escola de Saúde e Desenvolvimento Humano, Universidade de Évora, Largo dos Colegiais, 7000-727 Évora, Portugal. <sup>8</sup>Research Center in Sport Sciences, Health Sciences and Human Development, 5001-801 Vila Real, Portugal. <sup>9</sup>Department of Neurosciences, Biomedicine and Movement Sciences, School of Exercise and Sport Science, University of Verona, 37131 Verona, Italy. ✉email: hadi.nobari1@gmail.com

decelerations, accelerations, impacts and high metabolic load distance<sup>10</sup>. Thus, more research on those metrics and between player positions is needed to confirm or not the results of the previous studies.

Beyond the information given before about TM and TS, some studies calculated both indexes through running distance variables<sup>5–7</sup> and through s-RPE variable<sup>5,7,9,22</sup>. The study of Oliveira et al.<sup>5</sup> included both external and internal workload measures in simultaneously but failed to analyze them taking into account the player positions while the other study of Oliveira et al.<sup>7</sup> seems to be the only one that **analyzed** both external and internal workload measures considering player positions. Specifically, this study found significant differences between player positions with moderate to very large effect across 10 mesocycles of the in-season. However, the previous study had small sample size and recommended more research on this topic. Moreover, the differences in the periods analyzed (10 mesocycles) reinforced that more analysis could be performed considering different periods of the season (e.g. pre-season and in-season). Furthermore, none of the previous studies<sup>5–7,9</sup> showed the relationships between TM and TS calculated through internal and external measures.

The relationship between internal and external load measures have been analyzed in previous studies<sup>23–25</sup> although without considering TM and TS indexes. Specifically, a study with professional soccer players showed that rating perceived exertion (RPE) correlate with distances covered between 14.4 and 19.8 and between 19.9 and 25.1 km/h<sup>23</sup>. Another study with professional soccer players also found a relationship between session-RPE (s-RPE) and total distance and between s-RPE and distances covered at > 19.8 km/h<sup>24</sup>. Such findings were also confirmed in young soccer players<sup>25</sup>.

Therefore, the aim of this study was (a) to describe and compare the in-season variations of TM and TS through s-RPE, total distance (TD), high-speed running distance (HSRD) and sprint distance (SpD) across different periods of a professional soccer season (early-season, mid-season, and end-season) and according to player positions (defenders, midfielders, wingers and strikers), and (b) to analyze the relationship of the aforementioned internal with external workload indexes measures across different periods of the season, respectively.

## Methods

**Participants.** Twenty professional players from an Asian First League (29.4 ± 4.4 years old; 75.0 ± 3.9 kg; 1.8 ± 0.1 cm; BMI: 23.4 ± 1.8 kg/m<sup>2</sup>) participated in this study. Five players from each position were selected from the entire number of participants, including defenders (DF), midfielders (MF), wingers (WG), and strikers (SF). It were included only players, who (1) were part of the team from week 1 to week 20 and (2) participated in 80% of weekly training sessions. It were not included players (1) with prolonged injury or a lack of participation in training for at least two consecutive weeks, (2) who showed the initial physical fitness test scores two standard deviations below the squad mean and (3) whose position was goal keeper due to differences in training activities and workload in training and matches. At the very beginning of the research, the players were informed about the study design and procedures. Thereafter, the players signed a free consent about their participation in the study. They did it even if this research's methods were already part of their club daily routine. This research fulfilled the requirements of the Declaration of Helsinki under the approval of the Ardabil University of Medical Sciences research ethics committee.

**Experimental design.** This research makes use of a descriptive-longitudinal approach. Players' monitoring occurred over 20 consecutive in-season weeks. All team's main training sessions were part of this research. Rehabilitation and recuperation sessions were not taken into account. Training sessions were made of warm-up, main and slow-down phases in addition to stretching. Coaching staff designed all training sessions, while researchers standardized only first and final 30 min (i.e., start and end of each session). Research took place from October 30, 2017 (early-season) until March 18, 2018 (end-season). Whole season was made of early-season (weeks 1–7), mid-season (weeks 8–13) and end-season (weeks 14–20; Table 1). Table 1 shows training sessions and matches numbers over the three season's periods, as well.

**External load monitoring.** During each session, players were monitored by a GPS (GPSports systems Pty Ltd, Model: SPI High-Performance Unit (HPU); Australian) and the study measures were collected daily during the in-season (i.e., all training sessions and matches). This study aimed to describe and compare the in-season variations of acute: TM and TS through s-RPE, total distance (TD), high-speed running distance (HSRD) and SpD across different periods of a professional soccer season (early-season, mid-season, and end-season) according to players' positions.

Global navigation satellite systems for professional athletes, such as the SPI HPU, include a 15 Hz GPS sensor in addition to a tri-axial accelerometer. As already shown in the literature, used device shows high validity and reliability (Cohen's *d* of differences between gold standard and device from trivial to small and intraclass correlation coefficients > 0.95)<sup>26</sup>. Throughout season, temperature and humidity resulted from 10 and 26 °C and from 22 to 48%, respectively. Special vests for the devices were placed on players' shoulders before trainings

Periods of the in-season	Early-season	Mid-season	End-season
Number of weeks	7	7	6
Training sessions (N)	15	14	18
Number of matches (N)	7	8	5

**Table 1.** Weeks and training sessions and number of competitive matches.

and matches starts. After activities, devices were removed from the players and checked by the team's match analyst before downloading recorded data to a computer equipped with the Team AMS software. Then, devices' memories were "cleaned" from old data and devices were put on an electric re-charge station. Devices' software was used according to manufacturer's instructions including putting into it players' anthropometric information and personal vest's assignment.

**Internal load monitoring.** Players were daily monitored for their RPE using the CR-10 Borg's scale<sup>27</sup>, adapted by Foster et al.<sup>28</sup>. Previous study demonstrated the validity and reliability of this scale to estimate the session intensity<sup>29</sup>. Thirty minutes after the end of each training session, players rated their RPE value using an app on a tablet. The scores provided by the players were also multiplied by the training duration, to obtain the s-RPE<sup>28,30</sup>. The players were previously familiarized with the scale, and all the answers were provided individually to avoid non-valid scores.

**Calculations of training indexes.** Through s-RPE, TD, HSRD and SpD, the following measures were calculated: (1) TM (mean of training load during the 7 days of the week divided by the standard deviation of the training load of the 7 days<sup>5-7,31</sup>) and (2) TS (sum of the training load for all training sessions during a week multiplied by training monotony<sup>5-7,31</sup>).

**Statistical analysis.** Descriptive statistics were used to characterize the sample. Shapiro–Wilk was used to test normality of results. Results were presented as mean  $\pm$  standard deviation (SD). The relationship between all variables at the different periods was verified using bivariate correlations<sup>32</sup> (Pearson product-moment correlation coefficient ( $r$ )). The effect size of the correlations was determined by considering the following thresholds:  $< 0.1$  = trivial;  $0.1-0.3$  = small;  $> 0.3-0.5$  = moderate;  $> 0.5-0.7$  = large;  $> 0.7-0.9$  = very large; and  $> 0.9$  = nearly perfect<sup>33,34</sup>.

All measures obtained a normal distribution (Shapiro–Wilk  $> 0.05$ ), it was used a repeated measures ANOVA test and the Bonferroni post-hoc test to compare measures for periods of the in-season and groups. The results are significant for a  $p \leq 0.05$ . Hedge's  $g$  effect size (ES) was also calculated to determine the magnitude of pairwise comparisons. The Hopkins threshold was utilized as follows:  $g \leq 0.2$ , trivial;  $0.2 < g \leq 0.6$ , small;  $0.6 < g \leq 1.2$ , moderate;  $1.2 < g \leq 2.0$ , large;  $2.0 < g \leq 4.0$ , very large; and  $g > 4.0$ , nearly perfect<sup>33</sup>. All data were analysed using IBM SPSS Statistics (version 22, IBM Corporation (SPSS Inc., Chicago, IL).

**Ethics approval and consent to participants.** To engage in this study, both the players and their staff coach signed an informed consent form. The study has approved by the Ardabil university of medical sciences Ethics Committee prior to its start, and the Helsinki Declaration was used to follow the recommendations of Human Ethics in Research.

## Results

Figures 1, 2, 3 and 4 show an overall view of the weekly average for TM and TS calculated through s-RPE, TD, HSRD, and SpD across different periods of a professional soccer season (early-season, mid-season, and end-season) between players' positions.

The weekly changes of TM and TS for s-RPE can be found in Fig. 1. The highest  $TM_{s-RPE}$  occurred in week 9 in mid-season (MF =  $10.02 \pm 3.00$  arbitrary units (AU)) and the lowest values happened in week 1 in early-season (DF =  $1.00 \pm 0.01$  AU and MF =  $1.00 \pm 0.01$  AU), week 4 (early season) (MF =  $1.00 \pm 2.00$  AU and ST =  $1.00 \pm 0.01$  AU), week 10 (mid-season) (DF =  $1.00 \pm 0.01$  AU) and week 20 in end-season (MF =  $1.00 \pm 0.01$  AU). The  $TS_{s-RPE}$  was the highest in week 8 in mid-season (MF =  $10,996.00 \pm 6968.00$  AU) and the lowest in in week 4 in early-season (MF =  $10,996.00 \pm 6968.00$  AU).

The weekly changes of TM and TS for TD can be seen in Fig. 2. The highest  $TM_{TD}$  occurred in week 12 in mid-season (DF =  $13.00 \pm 11.00$  AU) and the lowest values happened in week 1 in early-season (MF =  $1.00 \pm 0.02$  AU, WG =  $1.00 \pm 0.01$  AU and ST =  $1.00 \pm 0.01$  AU), week 4 (early season) (MF =  $1.00 \pm 0.01$  AU), week 5 (early-season) (MF =  $1.00 \pm 0.01$  AU and ST =  $1.00 \pm 0.01$  AU), week 7 (early-season) (MF =  $1.00 \pm 0.01$  AU), week 8 (mid-season) (MF =  $1.00 \pm 0.01$  AU and ST =  $1.00 \pm 0.01$  AU) and week 16 in end-season (DF =  $1.00 \pm 0.01$  AU and MF =  $1.00 \pm 0.01$  AU).  $TS_{TD}$  was the highest in week 18 in end-season (ST =  $250,402.00 \pm 346,684.00$  AU) and the lowest in week 16 in end-season (DF =  $18,423.00 \pm 5765.00$  AU).


The weekly changes of TM and TS for HSRD can be seen in Fig. 3. The highest  $TM_{HSRD}$  occurred in week 14 in end-season (DF =  $14.00 \pm 17.00$  AU) and the lowest happened in week 8 in mid-season (MF =  $0.55 \pm 0.16$  AU). The  $TS_{HSRD}$  was the highest in week 14 in end-season (WG =  $111,872.00 \pm 117,710.00$  AU) and the lowest in week 7 in early-season (DF =  $966.00 \pm 647.00$  AU).

The weekly changes of TM and TS SpD can be found in Fig. 4. The highest  $TM_{SpD}$  occurred in week 18 in end-season (ST =  $9.00 \pm 6.00$  AU) and the lowest happened in week 4 in early-season (MF =  $0.78 \pm 0.18$  AU). The  $TS_{SpD}$  was the highest in week 14 in end-season (ST =  $16,580.00 \pm 19,639.00$  AU) and the lowest in week 7 in early-season (DF =  $1444.00 \pm 1580.00$  AU).

Table 2 presents the differences between the early-season, mid-season, and end-season for TM and TS calculated through s-RPE, TD, HSRD, and SpD. To simplify the description, only large to nearly perfect effect sizes will be described here. There was no significant difference for  $TM_{s-RPE}$ .

The  $TS_{s-RPE}$  presents a significant higher value in mid-season than early-season [large effect] and shows a significant higher value in end-season than early-season [very large effect].

The  $TM_{HSRD}$  presents a significant higher value in end-season than early-season [large effect] and shows a significant higher value in end-season than mid-season [very large effect]. The  $TS_{HSRD}$  shows a significant



**Figure 1.** TM (A) and TS (B) variations calculated through the s-RPE across 20 weeks between players' positions.

higher value in mid-season than early-season [large effect], shows a significant higher value in end-season than early-season [very large effect], and presents a significant higher value in end-season than mid-season [very large effect].

Finally, the  $TS_{SpD}$  presents a significant higher value in mid-season than early-season [large effect] and shows a significant higher value in end-season than early-season [large effect].

Table 3 presents the differences between player positions for TM and TS calculated through s-RPE, TD, HSRD, and SpD during in-season. There were no meaningful differences for  $TS_{HSRD}$ . To simplify the description, only large to nearly perfect effect sizes will be described here.


The  $TM_{s-RPE}$  shows a significant higher value in WG than MF [large effect] and shows a significant higher value in ST than MF [very large effect].

The  $TM_{TD}$  shows a significant higher value in WG than DF [very large effect], shows a significant higher value in ST than DF [nearly perfect effect], and presents a significant higher value in WG than MF [large effect]. The  $TS_{TD}$  shows a significant higher value in MF than DF [large effect].

The  $TM_{HSRD}$  shows a significant higher value in WG than DF [large effect], presents a significant higher value in ST than DF [large effect], shows a significant higher value in WG than MF [very large effect], and shows a significant higher value in ST than MF [very large effect].

Finally, the  $TM_{SpD}$  presents a significant higher value in WG than DF [nearly perfect effect].

Table 4 shows the correlation coefficient of all measures in the study for the team. In early-season, two positive correlations were denoted between:  $TM_{TD}$  and  $TM_{s-RPE}$ ;  $TM_{SpD}$  and  $TM_{s-RPE}$ . Two negative correlations were also denoted between:  $TS_{HSRD}$  and  $TM_{s-RPE}$ ;  $TS_{HSRD}$  and  $TS_{s-RPE}$ . In mid-season, one positive correlation (between  $TS_{HSRD}$  and  $TS_{s-RPE}$ ) and three negative correlations were denoted between:  $TM_{SpD}$  and  $TM_{s-RPE}$ ;  $TM_{TD}$  and  $TS_{s-RPE}$ ;  $TM_{SpD}$  and  $TS_{s-RPE}$ . The correlations with large effects are presented in Fig. 5.



**Figure 2.** TM (A) and TS (B) variations calculated through the TD across 20 weeks between players' positions.


## Discussion

The aims of this study were to investigate the variations in internal and external workload measures of TM and TS in professional male soccer players according to periods of the season and playing positions, and to determine the associations between the same internal and external workload measures of TM and TS. The results revealed that regardless of measure, the highest mean TM and TS scores were observed in mid-season or end-season. This result contrasts with the findings of Fessi et al.<sup>35</sup>. They analyze the weekly variations of training-related monotony and strain in professional soccer players and found significantly higher scores in TM and TS during pre-season when compared to in-season. On the other side, results of the present study seem to be in line with the findings of Clemente et al.<sup>22</sup>. Those authors monitored the training load variables of professional soccer players across a 10-week period and found highest values of TM in pre-season and highest values of TS in early competitive season. Lastly, the results found by Oliveira et al.<sup>7</sup> did not find significant differences across 10 mesocycles of the in-season period. As stated in the experimental design, coaches were responsible for training plan during the full-season which may help to explain the inconsistent results when analyzing other studies<sup>22,35</sup> since different coaches may have different philosophies for training. Nonetheless, our study highlights the importance of quantifying load through the full-season to better understand the intensity variations of all players.

The results on the position-related differences in TM showed greater values for wingers and strikers which is not consistent with a previous study<sup>7</sup> that showed a w-shape variation across 10 mesocycles from the in-season for all positions considering TM of HSRD (> 19 km/h) while the remaining TM values calculated through total distance or session rated perceived exertion were similar for all positions.

When TD and SpD were considered, midfielders exhibited greater TS scores. Contrasting results were obtained in a recent study<sup>6</sup> that examine the differences between playing positions for TM and TS in professional players. They found no significant difference for both measures between positions. In another recent similar study<sup>10</sup> weekly variations of external training loads throughout a professional soccer season were studied. While significantly greater TS values were reported for wingers and central defenders, no significant differences were found for TM between positions. Additionally, a recent study<sup>7</sup> did not find such results. They found a tendency of higher values of TS of TD and HSRD (> 19 km/h) for wide defenders than central midfielders over the 10 mesocycles of the in-season.






**Figure 3.** TM (A) and TS (B) variations calculated through the HSRD across 20 weeks between players' positions.

Overall and considering TM, it shows the uniformity of exercises during microcycles as well. These results warn the coaches that due to the training content of the game positions, they should pay more attention to the midfield positions (e.g., when compared to defenders and strikers), which can keep them from the uniformity of the training. Whilst, if not observed, can lead to a decline in player performance and possibly detraining. While wingers and strikers, due to the nature of their playing positions in training and competitions, this problem is often not faced.

Correlation analysis for TM revealed that s-RPE is significantly associated with SpD in both early and mid-season. Results also showed a significant association between s-RPE and TD in early season. Regarding the TS, the results demonstrated that s-RPE is significantly associated with HSRD in early and mid-season. Current literature provides limited evidence on the relationships between different load measures to estimate TM and TS. Nevertheless, this result is in agreement with the findings of a previous study<sup>23</sup> that investigated the association between s-RPE and external training load measures. Supportively, a significant association was noted between s-RPE and HSRD for a group of soccer players competing in the English Premier League<sup>36</sup>. According to Nobari et al.<sup>8</sup>, increasing internal intensity (e.g., HR and RPE) is linked to higher TM and TS, implying that increasing external intensity raises rating perceived. According to a previous studies<sup>9,37,38</sup>, an increase in TM can lead to



**Figure 4.** TM (A) and TS (B) variations calculated through the sprint distance across 20 weeks between players' positions.

overtraining, which is one of the consequences of a not well-adjusted training plan and, as a result, it raises the internal intensity during training and competition.

The present study has several limitations that should be taken into account. Firstly, the study data were obtained from one soccer team and thus it was conducted on a small sample. Secondly, generalizability of the results is limited to male professional soccer players. Lastly, the study lacks information about the injury records of players during training and match play across the different periods of the season. Therefore, further examinations are warranted to analyze the relationships between training load indices to estimate monotony and strain and injury in larger group of male and female soccer players from different age categories and competitive levels. The final limitation of this study was the lack of internal and external load monitoring in resistance training and competition sessions which should be considered in future studies.

## Conclusion

This study is original in the sense that it provides information regarding the variations in various internal and external training load measures of TM and TS with respect to the period of the season and the positions of the players. Our findings highlighted that TM and TS of professional soccer players is sensitive to period of the season, player's position, and the measure used to estimate training workloads. Therefore, coaching staff should take into account these variabilities in order to identify the training requirements of players.

Measures	EarlyS (Mean ± SD)	MidS (Mean ± SD)	EndS (Mean ± SD)	<i>p</i>	Hedges' <i>g</i> (95% CI)
TM <sub>s-RPE</sub> (AU)	3.00 ± 0.10	5.00 ± 2.00	4.09 ± 0.01	EarS versus MidS: 0.059	–
				EarS versus EndS: 1.000	–
				MidS versus EndS: 0.077	–
TS <sub>s-RPE</sub> (AU)	2007.09 ± 418.00	4119.00 ± 1884.00	3700.00 ± 806.00	<b>EarS versus MidS: 0.001</b>	–1.52 [–2.25, –0.83]#
				<b>EarS versus EndS: EndS: &lt; 0.01</b>	–2.58 [–3.48, –1.77]§
				MidS versus EndS: 1.000	–
TM <sub>TD</sub> (AU)	3.00 ± 2.00	4.00 ± 3.09	5.00 ± 2.00	EarS versus MidS: 1.000	–
				<b>EarS versus EndS: 0.010</b>	–0.98 [–1.65, –0.33]*
				<b>MidS versus EndS: EndS: &lt; 0.01</b>	–0.38 [–1.01, 0.24]&
TS <sub>TD</sub> (AU)	73,645.00 ± 38,650.00	64,607.00 ± 22,224.00	85,703.00 ± 36,003.00	<b>EarS versus MidS: MidS: &lt; 0.01</b>	0.28 [–0.34, 0.91]&
				<b>EarS versus EndS: EndS: &lt; 0.01</b>	–0.32 [–0.94, 0.30]&
				<b>MidS versus EndS: 0.032</b>	–0.69 [–1.34, –0.06]&
TM <sub>HSRD</sub> (AU)	2.01 ± 0.02	1.00 ± 0.10	4.02 ± 2.00	EarS versus MidS: 1.000	–
				<b>EarS versus EndS: EndS: &lt; 0.01</b>	–1.39 [–2.11, –0.72]#
				<b>MidS versus EndS: EndS: &lt; 0.01</b>	–2.09 [–2.91, –1.34]§
TS <sub>HSRD</sub> (AU)	3402.00 ± 980.00	8866.00 ± 5909.00	33,956.00 ± 13,164.00	<b>EarS versus MidS: 0.002</b>	–1.26 [–1.97, –0.59]#
				<b>EarS versus EndS: EndS: &lt; 0.01</b>	–3.21 [–4.22, –2.30]§
				<b>MidS versus EndS: EndS: &lt; 0.01</b>	–2.41 [–3.28, –1.62]§
TM <sub>SpD</sub> (AU)	2.00 ± 1.03	3.00 ± 1.02	3.00 ± 1.00	<b>EarS versus MidS: 0.011</b>	–0.96 [–1.63, –0.31]*
				<b>EarS versus EndS: 0.003</b>	–0.97 [–1.64, –0.32]*
				MidS versus EndS: 1.000	–
TS <sub>SpD</sub> (AU)	3092.00 ± 609.00	5339.00 ± 1800.00	6201.00 ± 2498.00	<b>EarS versus MidS: MidS: &lt; 0.01</b>	–1.64 [–2.39, –0.94]#
				<b>EarS versus EndS: EndS: &lt; 0.01</b>	–1.68 [–2.43, –0.97]#
				<b>MidS versus EndS: EndS: &lt; 0.01</b>	–0.38 [–1.02, 0.23]&

**Table 2.** Descriptive statistics (mean ± SD) of all measures in early-season, mid-season and end-season. Significant differences between periods are highlighted in bold ( $p \leq 0.05$ ). AU arbitrary units, *EarlyS* early-season, *MidS* mid-season, *EndS* end-season, *SD* standard deviation, *TM* training monotony, *TS* train strain, *s-RPE* session rate of perceived exertion, *TD* total distance, *HSRD* high-speed running distance, *SpD*, sprint distance. &small effect; \*, moderate effect; #, large effect; §, very large effect; £, nearly perfect effect.




Measures	DF (Mean ± SD)	MF (Mean ± SD)	WG (Mean ± SD)	ST (Mean ± SD)	<i>p</i>	Hedges' <i>g</i> (95% CI)
TM <sub>s-RPE</sub> (AU)	4.00 ± 0.02	3.00 ± 0.01	4.00 ± 1.02	4.00 ± 0.58	DF versus MF: 1.000	–
					DF versus WG: 1.000	–
					DF versus ST: 1.000	–
					<b>MF versus WG: &lt;0.01</b>	– 1.25 [– 2.76, 0.05]#
					<b>MF versus ST: &lt;0.01</b>	– 2.20 [– 4.11, – 0.69]§
					WG versus ST: 1.000	–
TS <sub>s-RPE</sub> (AU)	3943.00 ± 712.00	2990.00 ± 65.00	3062.00 ± 777.00	2937.00 ± 270.00	DF versus MF: 0.083	–
					<b>DF versus WG: &lt;0.01</b>	1.07 [– 0.21, 2.52]*
					DF versus ST: 0.061	–
					MF versus WG: 1.000	–
					MF versus ST: 1.000	–
					WG versus ST: 1.000	–
TM <sub>TD</sub> (AU)	2.00 ± 0.01	3.00 ± 1.00	5.00 ± 1.00	6.00 ± 1.00	DF versus MF: 1.000	–
					<b>DF versus WG: 0.049</b>	– 3.83 [– 6.59, – 1.83]§
					<b>DF versus ST: 0.010</b>	– 5.11 [– 8.62, – 2.65]£
					<b>MF versus WG: &lt;0.01</b>	– 1.81 [– 3.53, – 0.39]#
					<b>MF versus ST: &lt;0.01</b>	– 0.90 [– 2.30, 0.36]&
					WG versus ST: 1.000	–
TS <sub>TD</sub> (AU)	68,306.00 ± 6446.00	91,790.00 ± 28,738.06	77,714.00 ± 36,359.00	62,806.00 ± 14,552.00	<b>DF versus MF: &lt;0.01</b>	– 1.02 [– 2.45, 0.25]#
					DF versus WG: 1.000	–
					DF versus ST: 1.000	–
					MF versus WG: 1.000	–
					<b>MF versus ST: &lt;0.01</b>	1.15 [– 0.14, 2.62]*
					WG versus ST: 1.000	–
TM <sub>HSRD</sub> (AU)	2.03 ± 0.01	1.00 ± 0.01	3.00 ± 1.00	3.00 ± 1.00	DF versus MF: 1.000	–
					<b>DF versus WG: &lt;0.01</b>	– 1.24 [– 2.74, 0.06]#
					<b>DF versus ST: &lt;0.01</b>	– 1.24 [– 2.74, 0.06]#
					<b>MF versus WG: &lt;0.01</b>	– 2.55 [– 4.63, – 0.95]§
					<b>MF versus ST: &lt;0.01</b>	– 2.55 [– 4.63, – 0.95]§
					WG versus ST: 1.000	–
Continued						

Measures	DF (Mean ± SD)	MF (Mean ± SD)	WG (Mean ± SD)	ST (Mean ± SD)	<i>p</i>	Hedges' <i>g</i> (95% CI)
TS <sub>HSRD</sub> (AU)	17,612.00 ± 3265.00	14,354.00 ± 672.00	16,391.00 ± 6809.00	14,583.00 ± 6603.00	DF versus MF: 1.000	–
					DF versus WG: 1.000	–
					DF versus ST: 1.000	–
					MF versus WG: 1.000	–
					MF versus ST: 1.000	–
					WG versus ST: 1.000	–
TM <sub>SpD</sub> (AU)	2.07 ± 0.08	3.00 ± 1.00	3.00 ± 0.22	3.00 ± 1.00	<b>DF versus MF: &lt;0.01</b>	–1.18 [–2.67, 0.11]*
					<b>DF versus WG: &lt;0.01</b>	–5.07 [–8.56, –2.63]£
					<b>DF versus ST: 0.014</b>	–1.18 [–2.67, 0.11]*
					MF versus WG: 1.000	–
					MF versus ST: 1.000	–
					WG versus ST: 1.000	–
TS <sub>SpD</sub> (AU)	5009.00 ± 390.00	6070.00 ± 1938.00	4190.00 ± 1014.00	4149.00 ± 705.00	DF versus MF: 1.000	–
					DF versus WG: 1.000	–
					DF versus ST: 1.000	–
					<b>MF versus WG: &lt;0.01</b>	–1.09 [0.19, 2.56]*
					<b>MF versus ST: &lt;0.01</b>	1.16 [–0.13, 2.64]*
					WG versus ST: 1.000	–

**Table 3.** Descriptive statistics (mean ± SD) of all measures between players' positions. Significant differences between player positions are highlighted in bold ( $p \leq 0.05$ ). AU arbitrary units, DF defenders, MF midfielders, WG wingers, ST strikers, SD standard deviation, ACWR acute: chronic workload ratio, EWMA exponentially weighted moving averages, CP coupled, UCP uncoupled, *s-RPE* session rate of perceived exertion, TD total distance, HSRD high-speed running distance, SpD sprint distance. &small effect; \*, moderate effect; #, large effect; §, very large effect; £, nearly perfect effect.

Measures	TM <sub>s-RPE</sub> (AU)	TS <sub>s-RPE</sub> (AU)
<b>Early-season</b>		
TM <sub>TD</sub> (AU)	<b>0.474*</b>	0.243
TS <sub>TD</sub> (AU)	-0.035	0.071
TM <sub>HSRD</sub> (AU)	0.403	0.032
TS <sub>HSRD</sub> (AU)	<b>-0.588#</b>	<b>-0.464*</b>
TM <sub>SpD</sub> (AU)	<b>0.608#</b>	0.342
TS <sub>SpD</sub> (AU)	-0.047	-0.028
<b>Mid-season</b>		
TM <sub>TD</sub> (AU)	-0.338	<b>-0.463*</b>
TS <sub>TD</sub> (AU)	-0.441	-0.405
TM <sub>HSRD</sub> (AU)	-0.013	-0.031
TS <sub>HSRD</sub> (AU)	0.282	<b>0.476*</b>
TM <sub>SpD</sub> (AU)	<b>-0.506#</b>	<b>-0.486*</b>
TS <sub>SpD</sub> (AU)	-0.295	-0.044
<b>End-season</b>		
TM <sub>TD</sub> (AU)	0.069	-0.313
TS <sub>TD</sub> (AU)	0.333	0.190
TM <sub>HSRD</sub> (AU)	-0.030	-0.334
TS <sub>HSRD</sub> (AU)	0.084	0.027
TM <sub>SpD</sub> (AU)	0.103	-0.275
TS <sub>SpD</sub> (AU)	0.115	0.040

**Table 4.** Correlation analysis between external and internal load measures during the three periods of the in-season by the overall team. Significant differences are highlighted in bold ( $p \leq 0.05$ ). AU arbitrary units, TM training monotony, TS training strain, s-RPE session rated perceived exertion, TD total distance, HSRD high-speed running distance, SpD sprint distance. \*moderate effect; #, large effect.



**Figure 5.** Pearson correlations in Early-season between TS HSRD and TM S-RPE (A); between TS SpD and TM s-RPE (B) and in Mid-season between TM HSRD and TM s-RPE (C).

**Data availability**

The datasets generated during and analyzed during the current study are available from the corresponding author on reasonable request.

Received: 21 September 2021; Accepted: 21 June 2022

Published online: 29 June 2022

## References

- Gabbett, T. J. *et al.* The athlete monitoring cycle: A practical guide to interpreting and applying training monitoring data. *Br. J. Sports Med.* **51**, 1451–1452 (2017).
- Oliveira, R. *et al.* In-season internal and external training load quantification of an elite European soccer team. *PLoS ONE* **14**, 1–18 (2019).
- Oliveira, R. *et al.* In-season training load quantification of one-, two- and three-game week schedules in a top European professional soccer team. *Physiol. Behav.* **201**, 146–156 (2019).
- Malone, J. J. *et al.* Seasonal training-load quantification in elite English premier league soccer players. *Int. J. Sports Physiol. Perform.* **10**, 489–497 (2015).
- Oliveira, R. *et al.* In-season internal and external workload variations between starters and non-starters: A case study of a top elite European soccer team. *Med.* **57**, 1–15 (2021).
- Clemente, F. M. *et al.* Weekly load variations of distance-based variables in professional soccer players: A full-season study. *Int. J. Environ. Res. Pub. Health* **17**, 3300 (2020).
- Oliveira, R. *et al.* In: season monotony, strain and acute/chronic workload of perceived exertion, global positioning system running based variables between player positions of a top elite soccer team. *BMC Sports Sci. Med. Rehabil* <https://doi.org/10.1186/s13102-021-00356-3> (2021).
- Nobari, H., Gholizadeh, R., Martins, A. D., Badicu, G. & Oliveira, R. In-season quantification and relationship of external and internal intensity, sleep quality, and psychological or physical stressors of semi-professional soccer players. *Biology (Basel)* **11**, 467 (2022).
- Martins, A. D. *et al.* Intra-season variations in workload parameters in europe's elite young soccer players: A comparative pilot study between starters and non-starters. *Healthcare* **9**, 977 (2021).
- Clemente, F. M. *et al.* Accelerometry-based variables in professional soccer players: Comparisons between periods of the season and playing positions. *Biol. Sport* **37**, 389–403 (2020).
- Nobari, H. *et al.* Comparisons of accelerometer variables training monotony and strain of starters and non-starters: A full-season study in professional soccer players. *Int. J. Environ. Res. Public Heal.* **17**, 6547 (2020).
- Nobari, H. *et al.* Wearable inertial measurement unit to accelerometer-based training monotony and strain during a soccer season: A within-group study for starters and non-starters. *Int. J. Environ. Res. Pub. Health* **18**, 1–12 (2021).
- Nobari, H. *et al.* Variations of accelerometer and metabolic power global positioning system variables across a soccer season: A within-group study for starters and non-starters. *Appl. Sci.* **11**, 6747 (2021).
- Jaspers, A., Brink, M. S., Probst, S. G. M., Frencken, W. G. P. & Helsen, W. F. Relationships between training load indicators and training outcomes in professional soccer. *Sport. Med.* **47**, 533–544 (2017).
- Foster, C. Monitoring training in athletes with reference to overtraining syndrome. *Med. Sci. Sports Exerc.*, Vol. 30, No.7, pp. 1164–1168, 1998. *Med. Sci. Sports Exerc.* **30**, 1164–1168 (1998).
- Di Salvo, V. *et al.* Performance characteristics according to playing position in elite soccer. *Int. J. Sports Med.* **28**, 222–227 (2007).
- Barrett, S., McLaren, S., Spears, L., Ward, P. & Weston, M. The influence of playing position and contextual factors on soccer players' match differential ratings of perceived exertion: A preliminary investigation. *Sports* **6**, 13 (2018).
- Brito, A. *et al.* Effects of pitch surface and playing position on external load activity profiles and technical demands of young soccer players in match play. *Int. J. Perform. Anal. Sport* **17**, 902–918 (2017).
- Barron, D. J., Atkins, S., Edmundson, C. & Fewtrell, D. Accelerometer derived load according to playing position in competitive youth soccer accelerometer derived load according. *Int. J. Perform. Anal. Sport* **14**, 734–743 (2014).
- Owen, A. L. *et al.* A contemporary positional multi modal assessment approach to training monitoring in elite professional soccer. *J. Complement. Med. Altern. Healthc.* **10**, 1–10 (2019).
- Clemente, F. M. *et al.* Internal training load and its longitudinal relationship with seasonal player wellness in elite professional soccer. *Physiol. Behav.* **179**, 262–267 (2017).
- Clemente, F. M. *et al.* Variations of training load, monotony, and strain and dose-response relationships with maximal aerobic speed, maximal oxygen uptake, and isokinetic strength in professional soccer players. *PLoS ONE* **14**, e0225522 (2019).
- Rago, V., Brito, J., Figueiredo, P., Krstrup, P. & Rebelo, A. Relationship between external load and perceptual responses to training in professional football: Effects of quantification method. *Sports* **7**, 68 (2019).
- Delaney, J. A., Duthie, G. M., Thornton, H. R. & Pyne, D. B. Quantifying the relationship between internal and external work in team sports: Development of a novel training efficiency index. *Sci. Med. Footb.* **2**, 149–156 (2018).
- de Dios-Álvarez, V. *et al.* Relationships between RPE-derived internal training load parameters and GPS-based external training load variables in elite young soccer players. *Res. Sport. Med* <https://doi.org/10.1080/15438627.2021.1937165> (2021).
- Tessaro, E. & Williams, J. H. Validity and reliability of a 15 Hz GPS device for court-based sports movements. *Sport Perform. Sci. Rep.* **29**, 1–4 (2018).
- Borg, G. Perceived exertion as an indicator of somatic stress. *Scand. J. Rehabil. Med.* **2**, 92–98 (1970).
- Foster, C. *et al.* A new approach to monitoring exercise training. *J. Strength Cond. Res.* **15**, 109–115 (2001).
- Impellizzeri, F. M., Rampinini, E., Coutts, A. J., Sassi, A. & Marcora, S. M. Use of RPE-based training load in soccer. *Med. Sci. Sports Exerc.* **36**, 1042–1047 (2004).
- Foster, C. *et al.* Effects of specific versus cross-training on running performance. *Eur. J. Appl. Physiol. Occup. Physiol.* **70**, 367–372 (1995).
- Nobari, H. *et al.* Comparisons of new body load and metabolic power average workload indices between starters and non-starters A full-season study in professional soccer players. *Proc. Inst. Mech. Eng. Part P J. Sport. Eng. Technol.* **10**, 1–10 (2020).
- Cohen, J., Cohen, P., West, S. G. & Aiken, L. S. *Applied Multiple Regression/Correlation Analysis for the Behavioral Sciences* (Routledge, 2002). <https://doi.org/10.4324/9780203774441>.
- Hopkins, W. G., Marshall, S. W., Batterham, A. M. & Hanin, J. Progressive statistics for studies in sports medicine and exercise science. *Med. Sci. Sports Exerc.* **41**, 3–13 (2009).
- Batterham, A. M. & Hopkins, W. G. Making meaningful inferences about magnitudes. *Int. J. Sports Physiol. Perform.* **1**, 50–57 (2006).
- Fessi, M. S. *et al.* Changes of the psychophysical state and feeling of wellness of professional soccer players during pre-season and in-season periods. *Res. Sport. Med.* **24**, 375–386 (2016).
- Gaudino, P. *et al.* Factors influencing perception of effort (session rating of perceived exertion) during elite soccer training. *Int. J. Sports Physiol. Perform.* **10**, 860–864 (2015).
- Nobari, H. *et al.* Weekly variations in the workload of Turkish national youth wrestlers: A season of complete preparation. *Int. J. Environ. Res. Pub. Health* **18**, 3832 (2021).
- Nobari, H., Kharatzadeh, M., Khalili, S. M., Jorge, P. & Paolo, L. Fluctuations of training load variables in elite soccer players U-14 throughout the competition season. *Healthc.* **9**, 1418 (2021).

## Acknowledgements

The authors thank Sepahan Football Club for your full cooperation in this study.

### Author contributions

Conceptualization, H.N., A.D.M. and R.O., methodology, H.N., A.D.M. and R.O., data collection, H.N., and A.D.M. analysis, H.N., and R.O., writing—original draft preparation, A.D.M., and R.O., writing—review and editing, H.N., A.D.M., L.P.A., and R.O. All authors have read and agreed to the published version of the manuscript.

### Funding

This research did not receive any specific grant from any funding agency in the public, commercial or not-for-profit sector.

### Competing interests

The authors declare no competing interests.

### Additional information

**Correspondence** and requests for materials should be addressed to H.N.

**Reprints and permissions information** is available at [www.nature.com/reprints](http://www.nature.com/reprints).

**Publisher's note** Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.



**Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

© The Author(s) 2022, corrected publication 2023