



OPEN Irreversible and quantum thermodynamic considerations on the quantum zeno effect

Umberto Lucia

The quantum zeno effect slows down the quantum system's time evolution under frequent measurements. This paper aims to study this quantum effect by introducing the definition of time based on an irreversible thermodynamic analysis of quantum systems. Consequently, the quantum zeno effect requires (i) high values of the electromagnetic entropy generation rate related to the spontaneously down-converted light and (ii) a decrease in the quantum system's entropy value. So, the quantum zeno effect is a quantum process related to the interaction between a quantum system and the electromagnetic waves of the measurement device, causing a quantum thermodynamic stationary state. Last, the fundamental role of irreversibility emerges.

The quantum zeno effect, also known as Turing's paradox, is a feature of quantum mechanical systems that allows the time evolution of a particle to be slowed down by measuring it frequently enough against a chosen measurement setting¹.

Alan Turing was the first to mention this phenomenon in his private correspondence with his colleague Robin Grandy dated 1954². Turing pointed out this paradoxical result. Frequent measurements can slow the evolution of a quantum system, hindering transitions to states different from the initial one³. This phenomenon is considered related to the quadratic behaviour of the survival probability at short times for the Schrödinger equation⁴, as pointed out by John von Neumann⁵. Degasperis, Fonda and Ghirardi⁶ carried out the first approach to the phenomenon, while Baidyanath Misra and George Sudarshan¹ developed its rigorous description.

In 1988, Cook⁷ proposed an experimental approach, based on oscillating systems, to verify the quantum zeno effect, then carried out by Itano et al.⁸. Then, other experimental confirmations of the phenomenon were obtained in different physical contexts, such as photon polarization⁹, nuclear spin isomers¹⁰, optical pumping¹¹, etc. In the analysis of the experiment carried out by Itano et al.⁸, Venugopalan and Ghosh¹² showed that the experimental results are related to the environment-induced decoherence theory¹³, for which the environment of a quantum system can monitor some of the system's observables, with the result that the eigenstates of those observables are continuously subjected to decoherence, showing classical-like states. Indeed, they highlighted that, during the measurement process, the quantum system is coupled to its external environment, which leads to decoherence over a characteristic time scale, named *decoherence time*. In the context of quantum zeno effect, Facchi and Pascazio¹⁴ highlight that a down-conversion process in a non-linear crystal can be studied as the decay of a pump photon into a pair of photons of lower frequency, and the energy of the spontaneously down-converted light monotonously increases. Moreover, in 2006, Streed et al. observed experimentally the dependence of the quantum zeno effect on measurement pulse characteristics¹⁵.

Today, the growing interest in the quantum zeno effect is related to its theoretical implication in the fundamentals of quantum mechanics, and also in its applications to spin polarization in gases¹⁶, control of decoherence in quantum computing¹⁷, etc. The physical meaning of decoherence has been clarified by means of quantum and classical irreversibility¹⁸, highlighting that decoherence is an irreversible process¹⁹, due to interaction between the quantum system and its environment.

The Second Law of Thermodynamics states that the entropy of an isolated system increases with time (for irreversible processes) or remains constant²⁰. Remembering that measurement on a quantum system is an irreversible process^{5,21}, the entropy is expected to increase, while on the contrary, the information obtained by frequent observation on a quantum system has been proven to decrease the entropy of the system itself²².

Recently, a thermodynamic approach^{23–25} to irreversibility in quantum systems has been developed based on the continuous interaction between the environmental electromagnetic waves and the matter, analysing the absorption-emission of a photon by an atomic electron, obtaining a thermophysical model of quantum

¹Dipartimento Energia "Galileo Ferraris", Politecnico di Torino, Corso Duca degli Abruzzi 24, 10129 Torin, Italy. ²INFN-Sezione di Torino, Turin, Italy. email: umberto.lucia@polito.it

thermodynamics in agreement with the experimental results^{26–29}. In this context, quantum system is considered as an open system, due to the photon inflow and outflow.

This approach can be extended to the quantum zeno effect, considering the coupled quantum system-experimental device as the isolated system while the quantum system and the device are independently two open subsystems³⁰.

This paper aims to develop this thermophysical analysis of the quantum zeno effect, pointing out the fundamental role of irreversibility and its relation with the definition of time.

Results

The result of this paper are Eqs. (7) and (11), with respect to which we can highlight:

- The interaction between a quantum system and a measurement device can be studied as the interaction between a quantum system and a photon. Following Einstein, this interaction is irreversible, and concerning it, we have introduced the definition of the time interval for which the quantum system evolves. So, the unitary evolution operator results related to this time interval definition;
- The quantum zeno effects requires that the eigenvalue of the unitary operator results in 1. This can be obtained by high values of electromagnetic entropy generation rate, Σ , which is related to the spontaneously down-converted light, following the physical evidence summarised in Ref.¹⁴;
- The quantum zeno effect requires small values of a time interval. This implies a high electromagnetic entropy generation rate, Σ , but also small values of entropy generation, σ , which means that the quantum system decreases its entropy value, in agreement with the experimental evidence as reported in Ref.²²
- When the rate σ/Σ is not small, the eigenvalue of the unitary evolution operator does not result in 1, so the evolution inhibition does not occur, and the Quantum anti-Zeno Effect emerges³¹.

These considerations lead to the state that the quantum zeno effect is a quantum process related to the interaction between a quantum system and the electromagnetic waves of the measurement device, such that the result is a quantum thermodynamic stationary state ($dS/dt = 0$, with S entropy of the control volume composed by the quantum system and the measurement device). Indeed, considering the thermodynamic universe composed by the quantum system and the measurement device, this effect results in a dynamical process in which the system exchanges photons inflow and outflow such that the entropy generation rate Σ emerges from the entropy balance for the quantum system, as follows:

$$\frac{dS}{dt} = \sum_{in} \dot{n}_{ph,in} S_{ph,in} - \sum_{out} \dot{n}_{ph,out} S_{ph,out} + \Sigma \xrightarrow{dS/dt=0} \Sigma = \sum_{out} \dot{n}_{ph,out} S_{ph,out} - \sum_{in} \dot{n}_{ph,in} S_{ph,in} \quad (1)$$

where \dot{n}_{ph} is the photon flux, s is the entropy generated by the interaction with each photon, *in* and *out* mean inflow and outflow respectively. Consequently, the total entropy of the thermodynamic universe results constant ($dS = 0$), such that:

- The device (environment) increases its entropy due to the entropy generation rate, Σ , produced by the electromagnetic waves interaction;
- The quantum system reduces its entropy; indeed, its entropy variation results $\Delta S_{sys} = \Delta S - \int_0^\tau \Sigma dt = -\sigma$, where $S = -\text{Tr}(\hat{\rho} \ln(\hat{\rho})) = -\sum_n w_n \ln(w_n)$, where $\hat{\rho}$ is the density operator, Tr is the trace over a complete set of states and w_n denotes the probability of such a state $|n_i\rangle$ ^{32,33}.

All these considerations allow us to point out that the thermodynamic approach suggested can describe and explain the quantum systems' behaviour concerning the quantum zeno effect.

Discussion

The quantum zeno effect is a quantum-mechanical process that slows down a quantum system's time evolution measuring it frequently enough for some chosen measurement setting¹. The effect consists in freezing the evolution of a quantum system by measuring it frequently enough in its known initial state. Recently, the quantum zeno effect definition has been extended considering it as the suppression of unitary time evolution in quantum systems provided by a variety of interactions^{16,34}. It appears only in systems with distinguishable quantum states, so it cannot be applied in classical system and macroscopic bodies³⁵.

The study of this effect has pointed out that application of a series of strong and fast perturbations of a quantum system can decouple it from its *decohering* environment³⁶. Thus, frequent measurements can inhibit decay of a system, while measurements applied more slowly can enhance decay rates (Quantum anti-Zeno Effect)³¹.

In this context, the transitions from the subspace without decoherent loss of a qubit to a state with a qubit lost in a quantum computer are particularly interesting for application to quantum computer and physics of computation³⁷; indeed, for the qubit correction, it is sufficient to determine whether the decoherence has already occurred or not³⁷.

During periodical measurements, with a finite interval of time τ , at each measurement, the wave function collapses to an eigenstate of the measurement operator. Then, among the measurements, the system evolves into a superposition of states. When this superposition state is measured, it collapses with a probability proportional to τ^2 , so for very short time intervals, the probability of collapse back to the initial state becomes around 1³⁸. Following the decoherence theory, the collapsing time is related to the decoherence time of the system coupled to the environment with thermal noise. The stronger the coupling is, the shorter the decoherence time is, and the faster it will collapse³⁹.

The thermodynamic approach allows us to address answers to the experimental evidences by involving time definition based on the evaluation of the irreversibility. So, the fundamental role of irreversibility³⁰ emerges also in quantum zeno effect. Particularly interesting are the thermodynamic considerations of quantum computing. Indeed, it is accepted that quantum computing is in principle reversible: an ideal quantum algorithm consists in initializing the data register in a state of the computational basis, followed by a unitary operation, and a final measurement in the computational basis. There is no back-action associated with this final measurement, so there is no heat dissipation associated with the reset. But, any single gate activation requires 10^{-21} J of energy, which is the same order of magnitude as the heat dissipated by the erasure of the single bit. These quantum processes generate entropy production, so quantum computing can be considered reversible, but quantum circuits, i.e., quantum computers, are irreversible. Their irreversibility leads to decoherence, and the quantum zeno effect has been shown to play a fundamental role in the control of the decoherence in the analysis of the Josephson junction in superconducting qubits⁴.

This new frontier of thermodynamics could represent a topic of investigation for the optimisation of quantum circuits. Indeed, the quantum zeno effect has attracted interest in the thermodynamic analysis of quantum information. In the context of information, entropy fluctuations represent a very interesting topic: Ahmadi et al.⁴⁰ have analysed the stochastic quantum processes due to quantum correlations concerning also memory effects concerning the apparent violation of the second law of thermodynamics. They extend the second law to quantum processes by incorporating information explicitly into this thermodynamic law, by providing the meaning of flows and backflows of information, proving that quantum thermodynamic force is responsible for encoding and decoding information even when a feedback controller outside the system is involved in the process.

Methods

The effect of a measure on a quantum system⁴¹, from a mathematical viewpoint, consists in the collapse of its full wave function $|\psi\rangle$ into an eigenstate, $|\psi_i\rangle$, of the state bases of Hilbert space \mathcal{H} , i.e.,⁴²:

$$\begin{aligned}\langle\psi_i|\psi_i\rangle &= 1 \\ \langle\psi_i|\psi_j\rangle &= 0, \forall j \neq i\end{aligned}\quad (2)$$

where $|\psi_j\rangle$ are the states in the Hilbert space \mathcal{H} and $|\psi\rangle$ is the quantum pure state such that⁴³:

$$|\psi\rangle \neq |\psi_1\rangle \otimes |\psi_2\rangle \otimes \dots \otimes |\psi_n\rangle \quad (3)$$

where \otimes is the tensorial product.

The time evolution of the interaction is described by the Schrödinger equation⁴²:

$$H|\psi(t)\rangle = -i\hbar \frac{\partial\psi}{\partial t} \quad (4)$$

where H is the Hamiltonian operator, $\hbar = h/2\pi$ with $h = 6.62607004 \times 10^{-34}$ J s is the Planck constant, t is the time and $i = \sqrt{-1}$. If the Hamiltonian is independent of time, then the unitary time evolution operator $U(t)$ can be introduced, obtaining⁴²:

$$|\phi(\tau)\rangle = U(\tau)|\psi(0)\rangle = e^{-iH\tau/\hbar}|\psi(0)\rangle \quad (5)$$

where $|\psi(0)\rangle$ is the state function evaluated in $t = 0$, the initial state, and $|\phi(\tau)\rangle$ is the state function evaluated in $t = \tau$, with τ time interval.

Consequently, the quantum zeno effect can be analysed by considering the state function evolution as follows:

$$|\phi(\tau)\rangle = e^{-iH\tau/\hbar}|\psi(0)\rangle = |\psi(0)\rangle \quad (6)$$

which allows us to focus our study on the definition of time interval τ .

Indeed, recently, a thermodynamic approach to time interval definition has been developed^{44–46} about the analysis of irreversibility^{23,23,24} in photon-atomic-electron interaction^{26–30}.

In this approach, time is conjectured to be related both to the entropy production (also called entropy generation in engineering thermodynamics) and to the entropy production rate (also called entropy generation rate in engineering thermodynamics), in agreement with the approach of Planck and Einstein, who highlighted that the law of system evolution consists of the law of entropy evolution⁴⁷.

The physic bases of the thermodynamic approach are the followings:

- The atom, without interaction, can be considered an isolated system, and any process inside it is completely reversible;
- The atom, in interaction with a photon, is an open system where inflow and outflow of photons can occur;
- The atom in interaction is subjected to an irreversible process due to the perturbation of its center of mass: the irreversibility emerges as interaction with the environment;
- During the absorption-emission phenomena, electrons seem to follow a reversible energetic pathway^{48–50} (Franck-Condon approximation), because considering a single atom or molecule, the energy perturbation of the center of mass results of the order of 10^{-13} J, small compared to the electron transition energy which is of the order of 10^{-8} J, with a related excited state lifetime of the order of 10^{-8} s⁵⁰. But, the reversible atom is only an approximation^{48–50}, that cannot be introduced in the study of irreversibility, due to the need to

- consider all the phenomena of the system photon-atom-environment⁵¹, during the photon-atomic electron interaction, following some recent experimental results⁵²;
- The consequence of the interaction between the bound electron and the photon is an entropic footprint in the quantum system.

Similarly to rational mechanics, where position and velocity can be used as independent variables for the state space, we use the entropy production σ and the entropy production rate Σ as independent variables of the state space $\Omega = \{(\sigma, \Sigma)\}$ ^{44,46}. In this context, some comments on entropy must be included. Indeed, different analytical forms of entropy have been introduced in thermodynamics, statistical physics and information sciences^{53–56}. The entropy variation ΔS is a footprint of the change of the state of the system, during its evolution between its initial and final states^{54,55}. Different approaches have been developed to derive an irreversible description of processes^{57–59}. An interesting approach to irreversible processes^{53,60,61} is to evaluate $\sigma = \sum_i J_i X_i > 0$, where J_i are flows of the matter, heat, etc., and X_i are generalized driving forces for vector transport processes or for chemical reactions, etc. In this paper, this approach is interesting in relation to the study of Ahmadi et al. in relation to information and entropy fluctuation, because it is based on the definition of thermodynamic forces. The J_i and X_i are linearly related when the system is not too far from equilibrium. Consequently, $J_i = \sum_j L_{ij} X_j$, where the L_{ij} are called phenomenological transport coefficients. The aim of the nonequilibrium statistical mechanics is to obtain the macroscopic thermodynamic laws from microscopic time reversible dynamics^{53–55,60–64}, still under intense studies from various viewpoint⁶⁵. Irreversibility emerges from the interaction between systems and their environment. In the analysis of the quantum zero effect, the interconnection between macroscopic approach to irreversibility and microscopic behaviour of the systems must be considered in the context of its temporal evolution. This analysis is based on the recent thorough study^{54,55} of the concept of time, and on the definition of time⁶⁶. The fundamental assumption is that time may be defined by means of the entropy production and the entropy rate, both due to irreversibility, and considered as independent variables as usually done for the generalised coordinates and the generalised velocities in Rational Mechanics⁶⁷. Consequently, a possibility to design a kind of *thermodynamic clock*⁶⁶, by using certain properties of a black body, has been introduced. Some example of applications of this approach are shown in Ref.⁴⁶.

Moreover, radiative processes in matter and its link to entropy variation are well known processes^{68–74} in non-equilibrium thermodynamics. A recent analysis of irreversibility concerning the electromagnetic interaction with atoms and molecules, has highlighted that the continuous photon–electron interaction due to the environmental electromagnetic fields, due to the continuous thermal nonequilibrium, represents an energy footprint in the environments^{23,66}, in agreement with the experimental results on the irreversibility of the electromagnetic interaction with atoms and molecules, developed in the late sixties. Here, we use these results. A deep analysis of this thermodynamic approach is developed in Ref.⁴⁶, while a deep discussion on entropy and electromagnetic fields is developed in Ref.⁷⁵.

Consequently, the thermodynamic definition of time interval, τ , has been introduced as^{25,44,46,46}:

$$\tau = \frac{\sigma}{\Sigma} \quad (7)$$

where the entropy production rate Σ results⁷⁵:

$$T_0 \Sigma = \int_V \left(\frac{1}{2} \varepsilon_0 c E_{el}^2 + \frac{1}{2\mu_0} c B_m^2 \right) dV \approx \frac{V}{2} \varepsilon_0 c E_{el}^2 + \frac{V}{2\mu_0} c B_m^2 \quad (8)$$

where E_{el} is the electric field, B_m is the magnetic field, $c = 299792458 \text{ m s}^{-1}$ is the speed of light, $\varepsilon_0 = 8.8541878128(13) \times 10^{-12} \text{ F m}^{-1}$ is the electric permittivity in vacuum and $\mu_0 = 4\pi \times 10^{-7} \text{ H m}^{-1}$ is the magnetic permeability in vacuum, A is the area of the border of the thermodynamic control volume, and T_0 is the environmental temperature, while the entropy production has been evaluated with respect to the semi-classical analysis of the photon-bound electron interaction²³:

$$T_0 \sigma = \frac{m_e}{M} E_\gamma \quad (9)$$

where T_0 is the environmental temperature, and E_γ is the energy of the photon. So, the time interval has been evaluated by means of measurable physical quantities as follows:

$$\tau = \frac{2 m_e}{V M c} \frac{E_\gamma}{\varepsilon_0 E_{el}^2 + \mu_0^{-1} B_m^2} \quad (10)$$

As a consequence of this definition the unitary operator evaluated in the time $t = \tau$ results:

$$U(\tau) = \exp \left(- \frac{i}{\hbar} H \frac{\sigma}{\Sigma} \right) \quad (11)$$

Data availability

All data generated or analysed during this study are included in this published article.

Received: 16 April 2023; Accepted: 1 July 2023

Published online: 04 July 2023

References

- Misra, B. B. & Sudarshan, E. C. G. The zeno's paradox in quantum theory. *J. Math. Phys.* **18**, 756 (1977).
- Teuscher, C. & Hofstadter, D. *Alan Turing: Life and Legacy of a Great Thinker* (Springer, 2004).
- Facchi, P. & Pascazio, S. Quantum zeno dynamics: Mathematical and physical aspects. *J. Phys. A Math. Theor.* **41**, 493001 (2008).
- Facchi, P., Fazio, R., Florio, G., Pascazio, S. & Yoneda, T. Zeno subspaces for coupled superconducting qubits. *Found. Phys.* **36**, 500–511 (2006).
- von Neumann, J. *Mathematical Foundations of Quantum Mechanics*. (Princeton University Press, 2018).
- Degasperis, A., Fonda, L. & Ghirardi, G. C. Does the lifetime of an unstable system depend on the measuring apparatus? *Il Nuovo Cimento A* **21**, 471 (1974).
- Cook, R. J. What are quantum jumps?. *Phys. Scr.* **1988**, 49 (1988).
- Itano, W. M., Heinzen, D. J., Bollinger, J. J. & Wineland, D. J. Quantum zeno effect. *Phys. Rev. A* **41**, 2295 (1990).
- Kwiat, P., Weinfurter, H., Herzog, T., Zeilinger, A. & Kasevich, M. A. Interaction-free measurement. *Phys. Rev. Lett.* **74**, 4763 (1995).
- Nagels, B., Hermans, L. J. F. & Chapovsky, P. L. Quantum zeno effect induced by collisions. *Phys. Rev. Lett.* **79**, 3097 (1997).
- Mølhave, K. & Drewsen, M. Demonstration of the continuous quantum zeno effect in optical pumping. *Phys. Lett. A* **45–49**, 3097 (2000).
- Venugopalan, A. & Ghosh, R. Decoherence and the quantum zeno effect. *Phys. Lett. A* **204**, 11–15 (1995).
- Zurek, W. H. Decoherence and the transition from quantum to classical. *Phys. Today* **44**, 36 (1991).
- Facchi, P. & Pascazio, S. Chapter 3—Quantum zeno and inverse quantum zeno effects. In Wolf, E. (ed.) *Progress in Optics*, vol. 42 of *Progress in Optics*, 147–217 (Elsevier, 2001).
- Streed, E. W. *et al.* Continuous and pulsed quantum zeno effect. *Phys. Rev. Lett.* **97**, 260402 (2006).
- Nakanishi, T., Yamane, K. & Kitano, M. Absorption-free optical control of spin systems: The quantum zeno effect in optical pumping. *Phys. Rev. A* **65**, 013404 (2001).
- Facchi, P. *et al.* Control of decoherence: Analysis and comparison of three different strategies. *Phys. Rev. A* **71**, 022302 (2005).
- Fortin, S. & Lombardi, O. Understanding decoherence as an irreversible process. *Int. J. Quantum Found.* **4**, 247–267 (2018).
- Omnès, R. Decoherence as an irreversible process. In Blanchard, P., Joos, E., Giulini, D., Kiefer, C. & Stamatescu, I.-O. (eds.) *Decoherence: Theoretical, Experimental, and Conceptual Problems*, 291–298 (Springer, 2000).
- Bejan, A. *Advanced Engineering Thermodynamics* 2nd edn. (Wiley, 1997).
- Brillouin, L. *Science and Information Theory* (Academic Press Inc., 1963).
- Pati, A. K. Understanding decoherence as an irreversible process. *arXiv arXiv:quant-ph/0006089*, 1–6 (2000).
- Lucia, U. Macroscopic irreversibility and microscopic paradox: A constructal law analysis of atoms as open systems. *Sci. Rep.* **6**, 35792 (2016).
- Lucia, U. Unreal perpetual motion machine, Rydberg constant and Carnot non-unitary efficiency as a consequence of the atomic irreversibility. *Phys. A* **492**, 962–968 (2018).
- Lucia, U. Some considerations on molecular machines and loschmidt paradox. *Chem. Phys. Lett.* **623**, 98–100 (2015).
- ao, T. B. *et al.* Irreversibility and the arrow of time in a quenched quantum system. *Phys. Rev. Lett.* **115**, 190601 (2015).
- Doyle, R. O. *Great Problems in Philosophy & Physics—Solved?* (Information Philosopher, 2016).
- Doyle, R. O. *The Origin of Irreversibility* (Information Philosopher, www.informationphilosopher.org, 2019 (Last access)).
- Doyle, R. O. The continuous spectrum of the hydrogen quasi-molecule. *J. Quant. Spectrosc. Radiati. Transf.* **492**, 1555–1569 (1968).
- Lucia, U. Considerations on non equilibrium thermodynamics of interactions. *Phys. A* **447**, 314–319 (2016).
- Wilkinson, S. *et al.* Experimental evidence for non-exponential decay in quantum tunnelling. *Nature* **384**, 575–577 (1997).
- H.-T. Elze. Open quantum systems, entropy and chaos. In T. Kodama *et al.* (eds.) *Relativistic Aspects of Nuclear Physics*, vol. 42 of *Proceedings of the Fifth Rio de Janeiro International Workshop*, 264–287 (World Scientific, 1998).
- Gemmer, J., Michel, M. & Mahler, G. *Quantum Thermodynamics* (Springer, 2009).
- Ghasemi, F. & Shafiee, A. A new approach to study the zeno effect for a macroscopic quantum system under frequent interactions with a harmonic environment. *Sci. Rep.* **9**, 15265 (2019).
- Bedingham, D. & Halliwell, J. J. Classical limit of the quantum zeno effect by environmental decoherence. *Phys. Rev. A* **89**, 042116 (2014).
- Facchi, P., Lidar, D. A. & Pascazio, S. Unification of dynamical decoupling and the quantum zeno effect. *Phys. Rev. A* **69**, 032314 (2004).
- Stolze, J. & Suter, D. *Quantum Computing: A Short Course from Theory to Experiment* (Wiley, 2008).
- Engelhardt, G. & Schaller, G. Maxwell's demon in the quantum-zeno regime and beyond. *New J. Phys.* **20**, 023011 (2018).
- Mensky, M. B. *Quantum Measurements and Decoherence* (Springer, 2000).
- Ahmadi, B., Salimi, S. & Khorashad, A. S. Irreversible work and Maxwell demon in terms of quantum thermodynamic force. *Sci. Rep.* **11**, 2301 (2021).
- Bohr, N. The quantum postulate and the recent development of atomic theory. *Nature* **121**, 580–590 (1928).
- Sakurai, J. J. & Napolitano, J. *Modern Quantum Mechanics* (Cambridge University Press, 2020).
- Schrödinger, E. Die gegenwärtige situation in der quantenmechanik. *Naturwissenschaften* **23**, 807–812 (1935).
- Lucia, U. & Grisolia, G. Time: A constructal viewpoint & its consequences. *Sci. Rep.* **9**, 10454 (2019).
- Lucia, U. & Grisolia, G. Time & clocks: A thermodynamic approach. *Results Phys.* **16**, 102977 (2020).
- Lucia, U., Grisolia, G. & Kuzemsky, A. L. Irreversibility and entropy production in nonequilibrium systems. *Entropy* **22**, 887 (2020).
- Einstein, A. *Autobiographical Notes (Translated by Schilpp, P. A.)* (Open Court Publishing Company, 1982).
- Franck, J. Elementary processes of photochemical reactions. *Trans. Faraday Soc.* **21**, 536–542 (1926).
- Condon, E. A theory of intensity distribution in band systems. *Phys. Rev.* **28**, 1182–1201 (1926).
- Alonso, M. & Finn, E. J. *Fundamental University Physics. Vol. III. Quantum and Statistical Physics* (Addison Wesley, 1968).
- Condon, E. Nuclear motions associated with electron transitions in diatomic molecules. *Phys. Rev.* **32**, 858–872 (1928).
- Kukk, E. *et al.* Violation of the franck-condon principle due to recoil effects in high energy molecular core-level photoionization. *Phys. Rev. Lett.* **95**, 133001 (2005).
- Kuzemsky, A. L. *Statistical Mechanics and the Physics of Many-Particle Model Systems* (World Scientific, 2017).
- Kuzemsky, A. L. Temporal evolution, directionality of time and irreversibility. *Rivista del Nuovo Cimento* **41**, 513–574 (2018).
- Kuzemsky, A. L. In search of time lost: Asymmetry of time and irreversibility in natural processes. *Found. Sci.* **25** (2020).
- Lin, S. K. Diversity and entropy. *Entropy* **1**, 1–3 (1999).
- Beattie, J. A. & Oppenheim, I. *Principles of Thermodynamics* (Elsevier, 1979).
- Guggenheim, E. A. *Thermodynamics. An Advanced Treatment for Chemists and Physicists* (Elsevier, 1985).
- Landsberg, P. T. *Thermodynamics and Statistical Mechanics* (Dover Publications, 1990).
- Zubarev, D. N. *Nonequilibrium Statistical Thermodynamics* (Consultant Bureau, 1974).
- Kuzemsky, A. L. Theory of transport processes and the method of the nonequilibrium statistical operator. *Int. J. Mod. Phys.* **21**, 2821–2949 (2007).
- Mackey, M. C. The dynamic origin of increasing entropy. *Rev. Mod. Phys.* **61**, 981–1015 (1989).
- Mackey, M. C. *Time's Arrow: The Origin of Thermodynamic Behavior* (Springer, 1992).

64. Hoover, W. G. & Hoover, C. G. Time-irreversibility is hidden within Newtonian mechanics. *Mol. Phys.* **116**(21–22), 3085–3096 (2018).
65. Lighthill, J. The recently recognized failure of predictability in Newtonian dynamics. *Proc. R. Soc.* **407**, 35–50 (1986).
66. Lucia, U. & Grisolia, G. Time: A footprint of irreversibility. *Atti dell'Accademia Peloritana dei Pericolanti* **97**, SC1–SC4 (2019).
67. Landau, L. D. & Lifshitz, E. M. *Mechanics* (Butterworth-Heinemann, 1976).
68. Planck, M. *The Theory of Heat Radiation* (Dover Publications, 1959).
69. Heitler, W. *The Quantum Theory of Radiation* (Dover Publications, 2010).
70. Jammer, M. *The Conceptual Development of Quantum Mechanics* (McGraw Hill, 1966).
71. Surdin, M., Braffort, P. & Taroni, F. Black-body Radiation Law deduced from Stochastic Electrodynamics. *Nature* **210**, 405–406 (1966).
72. Rueda, A. On the irreversible thermodynamics of radiative processes. *Found. Phys.* **4**, 215–226 (1974).
73. Fonseca, J. M., Gomes, A. H. & Moura-Melo, W. A. Emission and absorption of photons and the black-body spectrum in Lorentz-odd electrodynamics. *Phys. Lett.* **671**, 280–283 (2009).
74. Boyer, T. H. Blackbody radiation in classical physics: A historical perspective. *Am. J. Phys.* **86**, 495–509 (2018).
75. Beretta, G. P. & Gyftopoulos, E. P. Electromagnetic radiation: A carrier of energy and entropy. *J. Energy Resour. Technol.* **137**, 021005 (2015).

Author contributions

The sole author All data generated or analysed during this study are included in this published article.

Competing interests

The author declares no competing interests

Additional information

Correspondence and requests for materials should be addressed to U.L.

Reprints and permissions information is available at www.nature.com/reprints.

Publisher's note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

© The Author(s) 2023