



OPEN

Investigation of the dynamical structures of double-chain deoxyribonucleic acid model in biological sciences

Muhammad Zain Yousaf¹, Muhammad Abbas¹, Tahir Nazir¹, Farah Aini Abdullah², Asnake Birhanu³✉ & Homan Emadifar^{4,5}

The present research investigates the double-chain deoxyribonucleic acid model, which is important for the transfer and retention of genetic material in biological domains. This model is composed of two lengthy uniformly elastic filaments, that stand in for a pair of polynucleotide chains of the deoxyribonucleic acid molecule joined by hydrogen bonds among the bottom combination, demonstrating the hydrogen bonds formed within the chain's base pairs. The modified extended Fan sub equation method effectively used to explain the exact travelling wave solutions for the double-chain deoxyribonucleic acid model. Compared to the earlier, now in use methods, the previously described modified extended Fan sub equation method provide more innovative, comprehensive solutions and are relatively straightforward to implement. This method transforms a non-linear partial differential equation into an ODE by using a travelling wave transformation. Additionally, the study yields both single and mixed non-degenerate Jacobi elliptic function type solutions. The complexiton, kink wave, dark or anti-bell, V, anti-Z and singular wave shapes soliton solutions are a few of the creative solutions that have been constructed utilizing modified extended Fan sub equation method that can offer details on the transversal and longitudinal moves inside the DNA helix by freely chosen parameters. Solitons propagate at a consistent rate and retain their original shape. They are widely used in nonlinear models and can be found everywhere in nature. To help in understanding the physical significance of the double-chain deoxyribonucleic acid model, several solutions are shown with graphics in the form of contour, 2D and 3D graphs using computer software Mathematica 13.2. All of the requisite constraint factors that are required for the completed solutions to exist appear to be met. Therefore, our method of strengthening symbolic computations offers a powerful and effective mathematical tool for resolving various moderate nonlinear wave problems. The findings demonstrate the system's potentially very rich precise wave forms with biological significance. The fundamentals of double-chain deoxyribonucleic acid model diffusion and processing are demonstrated by this work, which marks a substantial development in our knowledge of double-chain deoxyribonucleic acid model movements.

Keywords Modified extended Fan sub equation approach, Double-chain deoxyribonucleic acid model, Exact traveling wave solutions, 3D graph, 2D line graph, Contour plot

Each cell's primary means of storing data is deoxyribonucleic acid. It resembles a very significant straight molecule that holds the inherited instructions that distinguish one species from the others in the sort of structure of a nucleotide chain. Because its significance with the origins of existence, the structure and function of DNA molecules have become one of the foremost fascinating issues in the discipline of contemporary biophysics. Because it consists of two complimentary chains or strands, DNA is dual-stranded. Since both of these chains

¹Department of Mathematics, University of Sargodha, 40100 Sargodha, Pakistan. ²School of Mathematical Sciences, Universiti Sains Malaysia, 11800 Penang, Malaysia. ³Department of Mathematics, College of Science, Hawassa University, Hawassa, Ethiopia. ⁴Department of Mathematics, Saveetha School of Engineering, Saveetha Institute of Medical and Technical Sciences, Saveetha University, Chennai, Tamil Nadu 602 105, India. ⁵MEU Research Unit, Middle East University, Amman, Jordan. ✉email: asnakeb@hu.edu.et

are joined by bonds of hydrogen and operate counter to one another, they are referred to as oppositely strands. In DNA, the term “nucleic acid” speaks to the phosphate particles, whereas “deoxyribo” defines the sugar.

The field of genetics evolved in 1953 when Watson and Crick identified the framework of the double-chain deoxyribonucleic acid (D-CDNA) model. More significantly, the double helix concept contributes to the understanding of the recurrence DNA process in addition to indicating the known layout of DNA. Nonlinear (NL) science is being applied to analyzing D-CDNA frameworks in order to understand further about their characteristics. This is why the characteristics of D-CDNA can be examined using NL models that combine biological tools and scientific methodologies. When attempting to investigate the configuration of D-CDNA from a NL science perspective, the appropriate NL mathematical models must be identified. The intricate layout of the D-CDNA mechanism and the existence of distinct motions make it challenging to imitate at first using a particular mathematical model.

Since strands of DNA are the building blocks of life, their structure and function represent one of the most interesting biophysical mysteries of our day. Over the years, researchers have examined the makeup of DNA. Predicting the emergence of significant NL structures has proved effective in the study of DNA dynamics. It has been demonstrated that nonlinearity causes confined waves. The capacity of these confined waves to transfer energy with no dissipating makes them intriguing. Numerous scientists have attempted to create mathematical representations of the mathematical biology. Indeed, this is the one of the main reasons to take this model into consideration. Numerous scholars have created and employed a wide range of approaches in the literature for resolving these mathematical structures. As an illustration, F-expansion approach¹, Hirota bilinear method^{2,3}, Kudryashov technique⁴, translation method⁵, inverse scattering scheme⁶, homogeneous balance approach⁷, $\frac{G}{G}$ -expansion approach⁸, tanh-function approach⁹, fractional dual-function scheme¹⁰, exp-function approach¹¹, sine cosine approach¹², Φ^6 -model expansion approach¹³, improved sardar sub-equation method¹⁴, generalized exponential rational function method¹⁵, direct algebraic technique¹⁶.

Numerous mathematicians and scientists have examined the dynamics of the D-CDNA model in^{17–22}. ϕ^6 -model expansion technique utilized in²³ to analyze mathematical dynamics in the D-CDNA model. A helicoidal D-CDNA model's semi-discrete breather is examined in²⁴. A DNA structure is supplied in²⁵ using both the wormlike-chain and bead models. The D-CDNA dynamical structure within the context of a pandemic is analyzed in²⁶. The DNA model's precise closed and solitonic structures can be reviewed in²⁷. In order to calculate the circuitry of a D-CDNA, a tight-binding framework has been developed and provided in²⁸. For the development of D-CDNA loops, a self-avoiding wormlike chain model is provided in²⁹. In³⁰, an unpredictable genome evolution in a D-CDNA system has been investigated using lie symmetry. The exact travelling wave (TW) solutions for NLDEs in a D-CDNA Model is demonstrated in³¹. In³², the behavior of soliton solutions in the D-CDNA model can be observed via two analytical techniques. The D-CDNA model has many accurate solutions that are closed-form utilising GERF provided in³³. The exact analytical solution for PDE that can be achieved through the bilinear residual network and bilinear neural network methods^{34–37} is considerably better compared to the traditional neural network numerical method.

The research presented here used modified extended Fan sub equation method (MEFSEM) to construct the D-CDNA model's exact TW solutions, which are crucial for biologically linked research. After reading through the literature, it is discovered that MEFSEM has not yet been used in the construction of exact TW solutions of the D-CDNA model. Consequently, with this navigate, we're going to effectively employ the suggested technique and generate exact TW solutions for the D-CDNA model in this work. The come upon solutions are innovative and have prospective uses in NL sciences. Solitons have been utilized to investigate several significant real-world problems across domains including fluid dynamics, plasma, nonlinear optics, astrophysics, and molecular biology.

As an illustration, the conceptual framework of solitons has been specifically used in fiber optics to the long-range transfer of digital information. In addition to their implementation in communication, solitons are employed in optical converters. Despite its significant potential applications in technological innovation, optical solitons continue to be among the foremost study areas in soliton theory today. Numerous types of solitary waves have been employed in biology to explain low frequency collective motion in proteins and DNA, the nervous system, and the transmission of signals and energy in biomembranes. Additionally, as plasmas are made up of considerably highly charged particles, soliton research is important in this field as well. For instance, NL oscillator chains that permit a variety of solitary wave solutions have been utilized to describe dusty plasmas, which are composed of micro charged dust particles.

Analytical techniques and numerical methods are the primary areas of investigation for soliton solutions of various types for tackling NLPDEs. The MEFSEM, an inexpensive and effective analytical technique, is employed in this work. The foundation of this technique is a widely recognized approximation. By using MEFSEM, one can acquire well-formed solutions that provide immediate insight. Heavy computing, which naturally requires a lot of time and resources, is not necessary with this technique. But there are also drawbacks to adopting MEFSEM in biotechnology, such as time and expense, particularly when working with complicated products, massive manufacturing, and rapidly evolving markets. The first-order derivatives problems arise in many technical domains, including fluid dynamics, biological sciences, physics, and mechanics are also extremely challenging to solve employing MEFSEM.

The subsequent format demonstrates how this study is designed: A description of the suggested MEFSEM is provided in section “Algorithm of the MEFSEM”. Section “Mathematical formulation and implementation of the MEFSEM” explains the construction of the DNA model mathematically and the implementation of MEFSEM to it. Section “Graphical Findings and discussion” analyzes the visual behaviors of the exact TW solutions of the D-CDNA model using MEFSEM. A few endure observations are presented in section “Conclusion”.

Algorithm of the MEFSEM

The procedure for the MEFSEM utilizing the generalized elliptic equation is provided in this section. Below is an outline of the key steps of our methodology.

Examine the subsequent non-linear partial differential equation (NLPDE), which has two independent spatial and temporal variables: s and t .

$$A(\Theta, \Theta_s, \Theta_t, \Theta_{ss}, \Theta_{tt}, \Theta_{st}, \dots) = 0, \quad (1)$$

where in above Eq. (1) $\Theta(s, t)$ is unknown function, A is the polynomial with $\Theta(s, t)$ and its partial derivatives indicated by the subscripts, which comprise NL factors and derivatives of the greatest order. In the following steps the contexts of the MEFSEM are described.

Step 1 The creation of a single variable ϱ from the separate spatial and temporal variables such that:

$$\Theta(s, t) = \Upsilon(\varrho), \quad \varrho = \xi s + \zeta t. \quad (2)$$

To determine the TW solutions, interpret Eq. (1) to the succeeding ODE by employing the wave transformation provided in Eq. (2)

$$B(\Upsilon, \Upsilon', \Upsilon'', \Upsilon''', \dots) = 0, \quad (3)$$

where B is a polynomial in $\Upsilon(\varrho)$ combine with derivatives of $\Upsilon(\varrho)$ and prime signifies the derivative in terms of ϱ which means $\Upsilon'(\varrho) = \frac{d\Upsilon}{d\varrho}$, $\Upsilon''(\varrho) = \frac{d^2\Upsilon}{d\varrho^2}$ and so on.

Step 2 In the subsequent finite series format, the solution to ODE (3) is presumed

$$\Upsilon(\varrho) = \sum_{l=0}^M \eta_l \chi^l(\varrho), \quad (4)$$

where M is a numerical value which needs to be determined, and η_l 's are real constants with $\eta_M \neq 0$ to be identified. The function $\Upsilon(\varrho)$ is the following elliptic equation

$$\chi'^2(\varrho) = c_0 + c_1\chi(\varrho) + c_2\chi^2(\varrho) + c_3\chi^3(\varrho) + c_4\chi^4(\varrho), \quad (5)$$

where c_0, c_1, c_2, c_3 and c_4 are constants that need to be established. Under certain situations, there can be three parameters σ, μ, ρ if c_0, c_1, c_2, c_3 and $c_4 \neq 0$ such that

$$\chi'^2(\varrho) = c_0 + c_1\chi(\varrho) + c_2\chi^2(\varrho) + c_3\chi^3(\varrho) + c_4\chi^4(\varrho) = (\sigma + \mu\chi(\varrho) + \rho\chi^2(\varrho))^2. \quad (6)$$

Equation (6) is solely satisfied in the scenario that subsequent relations hold.

$$c_0 = \sigma^3, \quad c_1 = 2\sigma\mu, \quad c_2 = 2\rho\sigma + \mu^2, \quad c_3 = 2\mu\rho, \quad c_4 = \rho^2. \quad (7)$$

In certain instances, if c_0, c_1, c_3 and $c_4 \neq 0$ and $c_2 = 0$, three factors σ, μ, ρ could possibly be present such that

$$\chi'^2(\varrho) = c_0 + c_1\chi(\varrho) + c_3\chi^3(\varrho) + c_4\chi^4(\varrho) = (\sigma + \mu\chi(\varrho) + \rho\chi^2(\varrho))^2. \quad (8)$$

Equation (8) is solely satisfied in the scenario that subsequent relations hold.

$$c_0 = \sigma^3, \quad c_1 = 2\sigma\mu, \quad c_3 = 2\mu\rho, \quad c_4 = \rho^2. \quad (9)$$

Among σ, μ and ρ parameters, the following constraint is needed to be present

$$\mu^2 = -2\sigma\rho, \quad \sigma\rho < 0. \quad (10)$$

Therefore the generalized Riccati equation (GRE) is obtained by simplifying the general elliptic equation (GEE) using Eqs. (6) and (8). This auxiliary equation is produced from the GEE for $c_0 = c_1 = 0$

$$\chi'^2(\varrho) = c_2\chi^2(\varrho) + c_3\chi^3(\varrho) + c_4\chi^4(\varrho). \quad (11)$$

The GEE generates the elliptic equation when $c_1 = c_3 = 0$

$$\chi'^2(\varrho) = c_0 + c_2\chi^2(\varrho) + c_4\chi^4(\varrho), \quad (12)$$

where in above Eq. (12) there is the Riccati equation.

$$\chi'^2(\varrho) = (D + \chi^2(\varrho))^2. \quad (13)$$

The GEE takes on its subsequent form when $c_2 = c_4 = 0$

$$\chi'^2(\varrho) = c_0 + c_1\chi(\varrho) + c_3\chi^3(\varrho). \quad (14)$$

Step 3 Through considering the homogeneous balance method between the linear and NL factors of the highest order given in Eq. (3), the positive integer M observe in solution (4) can easily be obtained.

Step 4 A polynomial in $\chi(\varrho)$ can be initiated through swapping the solution (4) into Eq. (3) with the support of Eq. (5). Next, combine each term that have the same $\chi(\varrho)$ powers. Thus, through applying the same powers

of $\chi(\varrho)$ equal to zero, a system of algebraic equations can be generated. Mathematica 13.2 is used to solve these equations consequently, it is feasible to discover the crucial constant values η_i 's. The exact TW solutions of Eq. (1) can be retrieved by inserting all constants and Eqs. (4)–(5) into Eq. (3).

Special cases

- *Case 1*

Type 1 $\mu^2 - 4\rho\sigma > 0, \mu\rho \neq 0$ and $\rho\sigma \neq 0$:

$$\begin{cases} \chi_1^I(\varrho) = -\frac{\mu + \sqrt{\mu^2 - 4\rho\sigma} \tanh\left(\frac{\sqrt{\mu^2 - 4\rho\sigma}\varrho}{2}\right)}{2\rho}, \\ \chi_2^I(\varrho) = -\frac{\mu + \sqrt{\mu^2 - 4\rho\sigma} \coth\left(\frac{\sqrt{\mu^2 - 4\rho\sigma}\varrho}{2}\right)}{2\rho}; \end{cases} \tag{15}$$

$$\begin{cases} \chi_3^I(\varrho) = -\frac{\mu + \sqrt{\mu^2 - 4\rho\sigma} \left(\tanh\left(\frac{\sqrt{\mu^2 - 4\rho\sigma}\varrho}{2}\right) \pm \operatorname{tsech}\left(\frac{\sqrt{\mu^2 - 4\rho\sigma}\varrho}{2}\right) \right)}{2\rho}, \\ \chi_4^I(\varrho) = -\frac{\mu + \sqrt{\mu^2 - 4\rho\sigma} \left(\coth\left(\frac{\sqrt{\mu^2 - 4\rho\sigma}\varrho}{2}\right) \pm \operatorname{csch}\left(\frac{\sqrt{\mu^2 - 4\rho\sigma}\varrho}{2}\right) \right)}{2\rho}, \\ \chi_5^I(\varrho) = -\frac{2\mu + \sqrt{\mu^2 - 4\rho\sigma} \left(\tanh\left(\frac{\sqrt{\mu^2 - 4\rho\sigma}\varrho}{4}\right) + \coth\left(\frac{\sqrt{\mu^2 - 4\rho\sigma}\varrho}{4}\right) \right)}{4\rho}; \end{cases} \tag{16}$$

$$\begin{cases} \chi_6^I(\varrho) = \frac{1}{2\rho} \left(-\mu + \frac{\sqrt{(E^2 + F^2)(\mu^2 - 4\rho\sigma)} - E(\sqrt{\mu^2 - 4\rho\sigma}) \cosh(\sqrt{\mu^2 - 4\rho\sigma}\varrho)}{E \sinh(\sqrt{\mu^2 - 4\rho\sigma}\varrho) + F} \right), \\ \chi_7^I(\varrho) = \frac{1}{2\rho} \left(-\mu - \frac{\sqrt{(F^2 - E^2)(\mu^2 - 4\rho\sigma)} + E(\sqrt{\mu^2 - 4\rho\sigma}) \sinh(\sqrt{\mu^2 - 4\rho\sigma}\varrho)}{E \cosh(\sqrt{\mu^2 - 4\rho\sigma}\varrho) + F} \right); \end{cases} \tag{17}$$

where E and F are non-zero constants that fulfill $F^2 - E^2 > 0$.

$$\begin{cases} \chi_8^I(\varrho) = \frac{2\sigma \cosh\left(\frac{\sqrt{\mu^2 - 4\rho\sigma}\varrho}{2}\right)}{\sqrt{\mu^2 - 4\rho\sigma} \sinh\left(\frac{\sqrt{\mu^2 - 4\rho\sigma}\varrho}{2}\right) - \mu \cosh\left(\frac{\sqrt{\mu^2 - 4\rho\sigma}\varrho}{2}\right)}, \\ \chi_9^I(\varrho) = \frac{-2\sigma \sinh\left(\frac{\sqrt{\mu^2 - 4\rho\sigma}\varrho}{2}\right)}{\mu \sinh\left(\frac{\sqrt{\mu^2 - 4\rho\sigma}\varrho}{2}\right) - \sqrt{\mu^2 - 4\rho\sigma} \cosh\left(\frac{\sqrt{\mu^2 - 4\rho\sigma}\varrho}{2}\right)}; \end{cases} \tag{18}$$

$$\begin{cases} \chi_{10}^I(\varrho) = \frac{2\sigma \cosh\left(\frac{\sqrt{\mu^2 - 4\rho\sigma}\varrho}{2}\right)}{\sqrt{\mu^2 - 4\rho\sigma} \sinh\left(\frac{\sqrt{\mu^2 - 4\rho\sigma}\varrho}{2}\right) - (\mu \cosh\left(\frac{\sqrt{\mu^2 - 4\rho\sigma}\varrho}{2}\right) \pm \sqrt{\mu^2 - 4\rho\sigma})}, \\ \chi_{11}^I(\varrho) = \frac{2\sigma \sinh\left(\frac{\sqrt{\mu^2 - 4\rho\sigma}\varrho}{2}\right)}{(\sqrt{\mu^2 - 4\rho\sigma} \cosh\left(\frac{\sqrt{\mu^2 - 4\rho\sigma}\varrho}{2}\right) \pm (\sqrt{\mu^2 - 4\rho\sigma})) - \mu \sinh\left(\frac{\sqrt{\mu^2 - 4\rho\sigma}\varrho}{2}\right)}; \end{cases} \tag{19}$$

$$\chi_{12}^I(\varrho) = \frac{4\sigma \cosh\left(\frac{\sqrt{\mu^2 - 4\rho\sigma}\varrho}{4}\right) \sinh\left(\frac{\sqrt{\mu^2 - 4\rho\sigma}\varrho}{4}\right)}{-2\mu \cosh\left(\frac{\sqrt{\mu^2 - 4\rho\sigma}\varrho}{4}\right) \sinh\left(\frac{\sqrt{\mu^2 - 4\rho\sigma}\varrho}{4}\right) + 2\sqrt{\mu^2 - 4\rho\sigma} \cosh^2\left(\frac{\sqrt{\mu^2 - 4\rho\sigma}\varrho}{4}\right) - (\sqrt{\mu^2 - 4\rho\sigma})}, \tag{20}$$

Type 2 $\mu^2 - 4\rho\sigma < 0, \mu\rho \neq 0$ and $\rho\sigma \neq 0$:

$$\begin{cases} \chi_{13}^I(\varrho) = \frac{-\mu + \sqrt{4\rho\sigma - \mu^2} \tan\left(\frac{\sqrt{4\rho\sigma - \mu^2}\varrho}{2}\right)}{2\rho}, \\ \chi_{14}^I(\varrho) = \frac{-\mu - \sqrt{4\rho\sigma - \mu^2} \cot\left(\frac{\sqrt{4\rho\sigma - \mu^2}\varrho}{2}\right)}{2\rho}; \end{cases} \tag{21}$$

$$\begin{cases} \chi_{15}^I(\varrho) = \frac{-\mu + \sqrt{4\rho\sigma - \mu^2} \left(\tan\left(\frac{\sqrt{4\rho\sigma - \mu^2}\varrho}{2}\right) \pm \sec\left(\frac{\sqrt{4\rho\sigma - \mu^2}\varrho}{2}\right) \right)}{2\rho}, \\ \chi_{16}^I(\varrho) = \frac{-\mu - \sqrt{4\rho\sigma - \mu^2} \left(\cot\left(\frac{\sqrt{4\rho\sigma - \mu^2}\varrho}{2}\right) \pm \csc\left(\frac{\sqrt{4\rho\sigma - \mu^2}\varrho}{2}\right) \right)}{2\rho}, \\ \chi_{17}^I(\varrho) = \frac{-2\mu + \sqrt{4\rho\sigma - \mu^2} \left(\tan\left(\frac{\sqrt{4\rho\sigma - \mu^2}\varrho}{4}\right) - \cot\left(\frac{\sqrt{4\rho\sigma - \mu^2}\varrho}{4}\right) \right)}{4\rho}; \end{cases} \tag{22}$$

$$\begin{cases} \chi_{18}^I(\varrho) = \frac{1}{2\rho}(-\mu + \frac{\pm\sqrt{(E^2-F^2)(4\rho\sigma-\mu^2)}-E(\sqrt{4\rho\sigma-\mu^2})\cos(\sqrt{4\rho\sigma-\mu^2}\varrho)}{E\sin(\sqrt{4\rho\sigma-\mu^2}\varrho)+F}), \\ \chi_{19}^I(\varrho) = \frac{1}{2\rho}(-\mu - \frac{\pm\sqrt{(E^2-F^2)(4\rho\sigma-\mu^2)}-E(\sqrt{4\rho\sigma-\mu^2})\sin(\sqrt{4\rho\sigma-\mu^2}\varrho)}{E\cos(\sqrt{4\rho\sigma-\mu^2}\varrho)+F}); \end{cases} \tag{23}$$

where E and F are non-zero constants that fulfill $E^2 - F^2 > 0$.

$$\begin{cases} \chi_{20}^I(\varrho) = \frac{2\sigma\cos(\frac{\sqrt{4\rho\sigma-\mu^2}}{2}\varrho)}{\sqrt{4\rho\sigma-\mu^2}\sin(\frac{\sqrt{4\rho\sigma-\mu^2}}{2}\varrho)+\mu\cos(\frac{\sqrt{4\rho\sigma-\mu^2}}{2}\varrho)}, \\ \chi_{21}^I(\varrho) = \frac{2\sigma\sin(\frac{\sqrt{4\rho\sigma-\mu^2}}{2}\varrho)}{-\mu\sin(\frac{\sqrt{4\rho\sigma-\mu^2}}{2}\varrho)+\sqrt{4\rho\sigma-\mu^2}\cos(\frac{\sqrt{4\rho\sigma-\mu^2}}{2}\varrho)}; \end{cases} \tag{24}$$

$$\begin{cases} \chi_{22}^I(\varrho) = -\frac{2\sigma\cos(\sqrt{4\rho\sigma-\mu^2}\varrho)}{\sqrt{4\rho\sigma-\mu^2}\sin(\sqrt{4\rho\sigma-\mu^2}\varrho)+\mu\cos(\sqrt{4\rho\sigma-\mu^2}\varrho)\pm\sqrt{4\rho\sigma-\mu^2}}, \\ \chi_{23}^I(\varrho) = \frac{2\sigma\sin(\sqrt{4\rho\sigma-\mu^2}\varrho)}{(\sqrt{4\rho\sigma-\mu^2}\cos(\sqrt{4\rho\sigma-\mu^2}\varrho)\pm(\sqrt{4\rho\sigma-\mu^2}))-\mu\sin(\sqrt{4\rho\sigma-\mu^2}\varrho)}; \end{cases} \tag{25}$$

$$\chi_{24}^I(\varrho) = \frac{4\sigma\cos(\frac{\sqrt{4\rho\sigma-\mu^2}}{4}\varrho)\sin(\frac{\sqrt{4\rho\sigma-\mu^2}}{4}\varrho)}{-2\mu\cos(\frac{\sqrt{4\rho\sigma-\mu^2}}{4}\varrho)\sin(\frac{\sqrt{4\rho\sigma-\mu^2}}{4}\varrho)+2\sqrt{\mu^2-4\rho\sigma}\cos^2(\frac{\sqrt{4\rho\sigma-\mu^2}}{4}\varrho)-(\sqrt{4\rho\sigma-\mu^2})}. \tag{26}$$

• Case 2

Type 1 $\rho\sigma < 0$ and $\rho\sigma \neq 0$:

$$\begin{cases} \chi_1^{II}(\varrho) = -\frac{\pm\sqrt{-2\rho\sigma}+\sqrt{-6\rho\sigma}\tanh(\frac{\sqrt{-6\rho\sigma}\varrho}{2})}{2\rho}, \\ \chi_2^{II}(\varrho) = -\frac{\pm\sqrt{-2\rho\sigma}+\sqrt{-6\rho\sigma}\coth(\frac{\sqrt{-6\rho\sigma}\varrho}{2})}{2\rho}; \end{cases} \tag{27}$$

$$\begin{cases} \chi_3^{II}(\varrho) = -\frac{\pm\sqrt{-2\rho\sigma}+\sqrt{-6\rho\sigma}(\tanh(\sqrt{-6\rho\sigma}\varrho)\pm\text{tsech}(\sqrt{-6\rho\sigma}\varrho))}{2\rho}, \\ \chi_4^{II}(\varrho) = -\frac{\pm\sqrt{-2\rho\sigma}+\sqrt{-6\rho\sigma}(\coth(\sqrt{-6\rho\sigma}\varrho)\pm\text{tsch}(\sqrt{-6\rho\sigma}\varrho))}{2\rho}, \\ \chi_5^{II}(\varrho) = -\frac{\pm\sqrt{-2\rho\sigma}+\sqrt{-6\rho\sigma}(\tanh(\frac{\sqrt{-6\rho\sigma}\varrho}{4})+\coth(\frac{\sqrt{-6\rho\sigma}\varrho}{4}))}{4\rho}; \end{cases} \tag{28}$$

$$\begin{cases} \chi_6^{II}(\varrho) = \frac{1}{2\rho}\left(\mp\sqrt{-2\rho\sigma} + \frac{\sqrt{(E^2+F^2)(-6\rho\sigma)}-E(\sqrt{-6\rho\sigma})\cosh(\sqrt{-6\rho\sigma}\varrho)}{E\sinh(\sqrt{-6\rho\sigma}\varrho)+F}\right), \\ \chi_7^{II}(\varrho) = \frac{1}{2\rho}\left(\mp\sqrt{-2\rho\sigma} - \frac{\sqrt{(F^2-E^2)(-6\rho\sigma)}+E(\sqrt{-6\rho\sigma})\sinh(\sqrt{-6\rho\sigma}\varrho)}{E\cosh(\sqrt{-6\rho\sigma}\varrho)+F}\right); \end{cases} \tag{29}$$

where E and F are non-zero constants that fulfill $F^2 - E^2 > 0$.

$$\begin{cases} \chi_8^{II}(\varrho) = \frac{2\sigma\cosh(\frac{\sqrt{-6\rho\sigma}\varrho}{2})}{\sqrt{-6\rho\sigma}\sinh(\frac{\sqrt{-6\rho\sigma}\varrho}{2})\mp\sqrt{-2\rho\sigma}\cosh(\frac{\sqrt{-6\rho\sigma}\varrho}{2})}, \\ \chi_9^{II}(\varrho) = \frac{-2\sigma\sinh(\frac{\sqrt{-6\rho\sigma}\varrho}{2})}{\pm\sqrt{-2\rho\sigma}\sinh(\frac{\sqrt{-6\rho\sigma}\varrho}{2})-\sqrt{-6\rho\sigma}\cosh(\frac{\sqrt{-6\rho\sigma}\varrho}{2})}; \end{cases} \tag{30}$$

$$\begin{cases} \chi_{10}^{II}(\varrho) = \frac{2\sigma\cosh(\sqrt{-6\rho\sigma}\varrho)}{\sqrt{-6\rho\sigma}\sinh(\sqrt{-6\rho\sigma}\varrho)\mp\sqrt{-2\rho\sigma}\cosh(\sqrt{-6\rho\sigma}\varrho)\pm\sqrt{-6\rho\sigma}}, \\ \chi_{11}^{II}(\varrho) = \frac{2\sigma\sinh(\sqrt{-6\rho\sigma}\varrho)}{\mp\sqrt{-2\rho\sigma}\sinh(\sqrt{-6\rho\sigma}\varrho)+\sqrt{-6\rho\sigma}\cosh(\sqrt{-6\rho\sigma}\varrho)\pm\sqrt{-6\rho\sigma}}; \end{cases} \tag{31}$$

$$\chi_{12}^{II}(\varrho) = \frac{4\sigma\cosh(\frac{\sqrt{-6\rho\sigma}\varrho}{4})\sinh(\frac{\sqrt{-6\rho\sigma}\varrho}{4})}{\pm 2\sqrt{-2\rho\sigma}\cosh(\frac{\sqrt{-6\rho\sigma}\varrho}{4})\sinh(\frac{\sqrt{-6\rho\sigma}\varrho}{4})+2\sqrt{-6\rho\sigma}\cosh^2(\frac{\sqrt{-6\rho\sigma}\varrho}{4})-(\sqrt{-6\rho\sigma})}. \tag{32}$$

• Case 3

Type 1 When $c_2 = 1, c_3 = -\frac{2\alpha_3}{\alpha_1}, c_4 = \frac{\alpha_3^2-\alpha_2^2}{\alpha_1^2}$, listed below is the solution to Eq. (11)

$$\chi_1^{III}(\varrho) = \frac{\alpha_1\text{sech}(\varrho)}{\alpha_2+\alpha_3\text{sech}(\varrho)}. \tag{33}$$

Type 2 When $c_2 = 1, c_3 = -\frac{2\alpha_3}{\alpha_1}, c_4 = \frac{\alpha_3^2 + \alpha_2^2}{\alpha_1^2}$, listed below is the solution to Eq. (11)

$$\left\{ \chi_2^{III}(\varrho) = \frac{\alpha_1 \operatorname{csch}(\varrho)}{\alpha_2 + \alpha_3 \operatorname{csch}(\varrho)}. \right. \tag{34}$$

Type 3 When $c_2 = 4, c_3 = -\frac{4(2\alpha_2 + \alpha_4)}{\alpha_1}, c_4 = \frac{\alpha_3^2 + 4\alpha_2^2 + 4\alpha_2\alpha_4}{\alpha_1^2}$, listed below is the solution to Eq. (11)

$$\left\{ \chi_3^{III}(\varrho) = \frac{\alpha_1 \operatorname{sech}^2(\varrho)}{\alpha_2 \operatorname{sech}^2(\varrho) + \alpha_3 \tanh(\varrho) + \alpha_4}. \right. \tag{35}$$

Type 4 When $c_2 = 4, c_3 = -\frac{4(\alpha_4 - 2\alpha_2)}{\alpha_1}, c_4 = \frac{\alpha_3^2 + 4\alpha_2^2 - 4\alpha_2\alpha_4}{\alpha_1^2}$, listed below is the solution to Eq. (11)

$$\left\{ \chi_4^{III}(\varrho) = \frac{\alpha_1 \operatorname{csch}^2(\varrho)}{\alpha_2 \operatorname{coth}^2(\varrho) + \alpha_3 \tanh(\varrho) + \alpha_4}. \right. \tag{36}$$

Type 5 When $c_2 = \alpha_1^2, c_3 = 2\alpha_1\alpha_2, c_4 = \alpha_2^2$, listed below is the solution to Eq. (11)

$$\begin{cases} \chi_5^{III}(\varrho) = -\frac{\alpha_1\alpha_3}{\alpha_2(\cosh(\alpha_1\varrho) - \sinh(\alpha_1\varrho) + \alpha_3)}, \\ \chi_6^{III}(\varrho) = -\frac{\alpha_1(\sinh(\alpha_1\varrho) + \cosh(\alpha_1\varrho))}{\alpha_2(\sinh(\alpha_1\varrho) + \cosh(\alpha_1\varrho) + \alpha_3)}. \end{cases} \tag{37}$$

Type 6 When $c_2 = -1, c_3 = \frac{2\alpha_3}{\alpha_1}, c_4 = -\frac{\alpha_3^2 - \alpha_2^2}{\alpha_1^2}$, listed below is the solution to Eq. (11)

$$\left\{ \begin{aligned} \chi_7^{III}(\varrho) &= \frac{\alpha_1 \sec(\varrho)}{\alpha_2 + \alpha_3 \sec(\varrho)}, \\ \chi_8^{III}(\varrho) &= \frac{\alpha_1 \csc(\varrho)}{\alpha_2 + \alpha_3 \csc(\varrho)}. \end{aligned} \right. \tag{38}$$

Type 7 When $c_2 = -4, c_3 = \frac{4(2\alpha_2 + \alpha_4)}{\alpha_1}, c_4 = -\frac{-\alpha_3^2 + 4\alpha_2^2 + 4\alpha_2\alpha_4}{\alpha_1^2}$, listed below is the solution to Eq. (11)

$$\left\{ \begin{aligned} \chi_9^{III}(\varrho) &= \frac{\alpha_1 \sec^2(\varrho)}{\alpha_2 \sec^2(\varrho) + \alpha_3 \tan(\varrho) + \alpha_4}, \\ \chi_{10}^{III}(\varrho) &= \frac{\alpha_1 \csc^2(\varrho)}{\alpha_2 \csc^2(\varrho) + \alpha_3 \cot(\varrho) + \alpha_4}. \end{aligned} \right. \tag{39}$$

where the constants $\alpha_1, \alpha_2, \alpha_3$ and α_4 are arbitrary.

- Case 4 In the present scenario, the common solutions to Eq. (12) are the single and combination non-degenerative Jacobi elliptic functions (JEFs). The association among the parameters of c_0, c_2 and c_4 in regard of solution of JEF given in NLODE Eq. (12), are depicted in a subsequent Table 1. Tables 1, 2, and 3 listed below, accordingly, describe the categories.
- Case 5 By way of illustration, in the case where Eq. (14) is grunted the answer is Weierstrass elliptic doubly periodic type solution.

$$\chi_1^V(\varrho) = F\left(\frac{\sqrt{c_3}}{2}\varrho, \frac{-4c_1}{c_3}, \frac{-4c_0}{c_3}\right) \quad c_3 > 0. \tag{40}$$

c_0	c_2	c_4	$\chi_i^{IV}(\varrho)$
1	$-(1 - q^2)$	q^2	$\chi_1^{IV}(\varrho) = cn\varrho, \chi_2^{IV}(\varrho) = cd\varrho = \frac{cn\varrho}{cd\varrho}$
$1 - q^2$	$2q^2 - 1$	$-q^2$	$\chi_3^{IV}(\varrho) = cn\varrho$
$q^2 - 1$	$2 - q^2$	-1	$\chi_4^{IV}(\varrho) = dn\varrho$
q^2	$-(1 + q^2)$	1	$\chi_5^{IV}(\varrho) = nr\varrho = (rn\varrho)^{-1}, \chi_6^{IV}(\varrho) = dc\varrho = \frac{dn\varrho}{cn\varrho}$
$-q^2$	$2q^2 - 1$	$1 - q^2$	$\chi_7^{IV}(\varrho) = nc\varrho = (cn\varrho)^{-1}$
-1	$2 - q^2$	$q^2 - 1$	$\chi_8^{IV}(\varrho) = nd\varrho = (dn\varrho)^{-1}$
1	$2 - q^2$	$1 - q^2$	$\chi_9^{IV}(\varrho) = rc\varrho = \frac{rn\varrho}{cn\varrho}$
1	$2q^2 - 1$	$-q^2(1 - q^2)$	$\chi_{10}^{IV}(\varrho) = rd\varrho = \frac{rn\varrho}{dn\varrho}$
$1 - q^2$	$2 - q^2$	1	$\chi_{11}^{IV}(\varrho) = cr\varrho = \frac{cn\varrho}{rn\varrho}$
$-q^2(1 - q^2)$	$2q^2 - 1$	1	$\chi_{12}^{IV}(\varrho) = dr\varrho = \frac{dn\varrho}{rn\varrho}$
$\frac{1}{4}$	$\frac{1-2q^2}{2}$	$\frac{1}{4}$	$\chi_{13}^{IV}(\varrho) = nr\varrho \pm cr\varrho$
$\frac{1-q^2}{4}$	$\frac{1+q^2}{2}$	$\frac{1-q^2}{4}$	$\chi_{14}^{IV}(\varrho) = nc\varrho \pm rc\varrho$
$\frac{q^2}{4}$	$\frac{q^2-2}{2}$	$\frac{1}{4}$	$\chi_{15}^{IV}(\varrho) = nr\varrho \pm dr\varrho$
$\frac{q^2}{4}$	$\frac{q^2-2}{2}$	$\frac{q^2}{4}$	$\chi_{16}^{IV}(\varrho) = rn\varrho \pm cr\varrho$

Table 1. The association among the parameters of c_0, c_2 and c_4 in regard of solution of JEF given in Eq. (12) where $0 \leq q \leq 1$.

JEF	HF	JEF	HF
$m \varrho$	$\tanh \varrho$	$nr \varrho$	$\coth \varrho$
$cn \varrho$	$\operatorname{sech} \varrho$	$nc \varrho$	$\cosh \varrho$
$dn \varrho$	$\operatorname{sech} \varrho$	$nd \varrho$	$\cosh \varrho$
$rc \varrho$	$\sinh \varrho$	$cr \varrho$	$\operatorname{csch} \varrho$
$rd \varrho$	$\sinh \varrho$	$dr \varrho$	$\operatorname{csch} \varrho$
$cd \varrho$	1	$dc \varrho$	1

Table 2. The JEFs become hyperbolic functions if $q \rightarrow 1$.

JEF	HF	JEF	HF
$m \varrho$	$\sin \varrho$	$nr \varrho$	$\csc \varrho$
$cn \varrho$	$\cos \varrho$	$nc \varrho$	$\sec \varrho$
$dn \varrho$	1	$nd \varrho$	1
$rc \varrho$	$\tan \varrho$	$cr \varrho$	$\cot \varrho$
$rd \varrho$	$\sin \varrho$	$dr \varrho$	$\csc \varrho$
$cd \varrho$	$\cos \varrho$	$dc \varrho$	$\sec \varrho$

Table 3. The JEFs become trigonometric functions if $q \rightarrow 0$.

Mathematical formulation and implementation of the MEFSEM

In current portion, A D-CDNA model fabricated from elastic rods is laid out. NL equations demonstrate the dynamics of this model, that can be expressed as

$$\Theta_{tt} - \Lambda_1^2 \Theta_{ss} = \Lambda_2 \Theta + \Lambda_3 \Theta \Xi + \Lambda_4 \Theta^3 + \Lambda_5 \Theta \Xi^2, \tag{41}$$

$$\Xi_{tt} - \Lambda_6^2 \Xi_{ss} = \Lambda_7 \Xi + \Lambda_8 \Theta^2 + \Lambda_9 \Theta^2 \Xi + \Lambda_{10} \Xi^3 + \Lambda_{11}, \tag{42}$$

where the function Θ , reflects the disparity between the longitudinal displacements of the foremost and bottom conductors in Eqs. (41) and (42), whereas the variation between the highest and lowest strands' transverse displacements is represented by Ξ . The constants $\Lambda_1, \Lambda_2, \dots, \Lambda_{11}$ are written as

$$\begin{aligned} \Lambda_1 &= \pm \frac{Z}{\delta}, \Lambda_2 = \pm \frac{-2\omega(y - a_0)}{\delta \Gamma h}, \Lambda_3 = 2\Lambda_8 = \frac{2\sqrt{2}\omega a_0}{\Delta \Gamma h^2}, \\ \Lambda_4 &= \Lambda_9 = \frac{-2\omega a_0}{\Delta \Gamma h^2}, \Lambda_5 = \Lambda_{10} = \frac{4\omega a_0}{\Delta \Gamma h^3}, \Lambda_6 = \pm \frac{g}{\Delta}, \Lambda_{11} = \frac{\sqrt{2}\omega(h - a_0)}{\Delta \Gamma}, \end{aligned} \tag{43}$$

where, $a_0, h, \omega, g, Z, \Gamma$ and Δ demonstrate the membrane's height, separation of the two strands, the elastic membrane's rigidity, density of stress in every thread, Young modulus, cross-sectional area and mass density respectively.

Assume the subsequent transformation

$$\Xi = b_0 \Theta + b_1, \tag{44}$$

where b_0 and b_1 are constants. Equations (41) and (42) are streamlined into the two subsequent forms because of (44).

$$\Theta_{tt} - \Lambda_1^2 \Theta_{ss} = \Theta^3 (\Lambda_4 + \Lambda_5 b_0^2) + \Theta^2 (2\Lambda_5 b_0 b_1 + \Lambda_3 b_0) + \Theta (\Lambda_2 + b_1 \Lambda_3 + \Lambda_5 b_1^2), \tag{45}$$

$$\Theta_{tt} - \Lambda_6^2 \Theta_{ss} = \Theta^3 (\Lambda_9 + \Lambda_{10} b_0^2) + \Theta^2 (3\Lambda_{10} b_0 b_1 + \frac{\Lambda_8}{b_0} + \frac{\Lambda_9 b_1}{b_0}) + \Theta (\Lambda_7 + 3\Lambda_{10} b_1^2) + \frac{\Lambda_7 b_1}{b_0} + \frac{\Lambda_{10} b_1^3}{b_0} + \frac{\Lambda_{11}}{b_0}. \tag{46}$$

Equations (45) and (46) can be compared with Eq. (44) to determine that $b_1 = \frac{h}{\sqrt{2}}$ and $Z = g$. At this point, Eqs. (45) and (46) can be expressed as

$$\Theta_{tt} - \Lambda_1^2 \Theta_{ss} = \Phi_1 \Theta^3 + \Phi_2 \Theta^2 + \Phi_3 \Theta, \tag{47}$$

where $\Phi_1 = \frac{\kappa(-2+4b_0^2)}{h^3}$, $\Phi_2 = \frac{6\sqrt{2}b_0\kappa}{h^2}$, $\Phi_3 = \frac{-2\kappa}{a_0} + \frac{6\kappa}{h}$, $\kappa = \frac{\omega a_0}{\Delta \Gamma}$ and $\Lambda_1 = \pm \frac{Z}{\delta}$.

Applying the wave transformations specified below

$$\Theta(s, t) = \Upsilon(\varrho), \quad \varrho = \xi s + \zeta t. \tag{48}$$




Figure 1. Algorithm flowchart of the MEFSEM.

Equation (47) reduce to the listed below system of ODE as a result of replacing Eq. (48) into Eq. (47)

$$(\zeta^2 - \xi^2 \Lambda_1^2) \Upsilon'' - \Phi_1 \Upsilon^3 - \Phi_2 \Upsilon^2 - \Phi_3 \Upsilon = 0, \quad \zeta^2 - \xi^2 \Lambda_1^2 \neq 0, \quad (49)$$

where prime stands for derivative with respect to ϱ . Employing the homogeneous balancing technique on the highest order derivative linear term Υ'' and the highest power non-linear term Υ^3 given in Eq. (49) provides $3M = M + 2$ which gives $M = 1$.

Accordingly, Eq. (4) translates to

$$\Upsilon(\varrho) = \eta_0 + \eta_1 \chi(\varrho). \tag{50}$$

The polynomial equation in the shape of $\chi(\varrho)$ for Eq. (49) can be uncovered in the following manner with the help of the Eqs. (50) and (5)

$$\begin{aligned} & (4\zeta^2 c_4 \eta_1 - 4\xi^2 c_4 \eta_1 \Lambda_1^2 - 2\eta_1^3 \Phi_1)(\chi(\varrho))^3 + (3\zeta^2 c_3 \eta_1 - 3\xi^2 c_3 \eta_1 \Lambda_1^2 - 6\eta_0 \eta_1^2 \Phi_1 - 2\eta_1^2 \Phi_2)(\chi(\varrho))^2 \\ & + (2\zeta^2 c_2 \eta_1 - 2\xi^2 c_2 \eta_1 \Lambda_1^2 - 2\eta_1 \Phi_3 - 6\eta_0^2 \eta_1 \Phi_1 - 4\eta_0 \eta_1 \Phi_2)(\chi(\varrho)) \\ & + \zeta^2 c_1 \eta_1 - \xi^2 c_1 \eta_1 \Lambda_1^2 - 2\eta_0^3 \Phi_1 - 2\eta_0^2 \Phi_2 - 2\eta_0 \Phi_3 = 0. \end{aligned} \tag{51}$$

In order to get the set of algebraic equations, the coefficient of comparable power of $\chi(\varrho)$ in Eq. (51) equal to zero.

$$\begin{cases} (\chi(\varrho))^3 : 4\zeta^2 c_4 \eta_1 - 4\xi^2 c_4 \eta_1 \Lambda_1^2 - 2\eta_1^3 \Phi_1 = 0, \\ (\chi(\varrho))^2 : 3\zeta^2 c_3 \eta_1 - 3\xi^2 c_3 \eta_1 \Lambda_1^2 - 6\eta_0 \eta_1^2 \Phi_1 - 2\eta_1^2 \Phi_2 = 0, \\ (\chi(\varrho))^1 : 2\zeta^2 c_2 \eta_1 - 2\xi^2 c_2 \eta_1 \Lambda_1^2 - 2\eta_1 \Phi_3 - 6\eta_0^2 \eta_1 \Phi_1 - 4\eta_0 \eta_1 \Phi_2 = 0, \\ (\chi(\varrho))^0 : \zeta^2 c_1 \eta_1 - \xi^2 c_1 \eta_1 \Lambda_1^2 - 2\eta_0^3 \Phi_1 - 2\eta_0^2 \Phi_2 - 2\eta_0 \Phi_3 = 0. \end{cases}$$

Solve above system of algebraic equations with the aid of Mathematica 13.2 gives

$$\eta_0 = \frac{3\zeta^2 c_3 - 3\xi^2 c_3 \Lambda_1^2 - 2\eta_1 \Phi_2}{6\eta_1 \Phi_1}, \eta_1 = \mp \sqrt{\frac{2(\zeta^2 c_4 - \xi^2 c_4 \Lambda_1^2)}{\Phi_1}}. \tag{52}$$

The TW solutions of Eqs. (41)–(42) that result from replacing Eqs. (52) and (15)–(40) into Eq. (50) are formulated as follows:

Special cases

Case 1

Type 1: $\mu^2 - 4\rho\sigma > 0, \mu\rho \neq 0, \rho\sigma \neq 0$

$$\Theta_{1,1,1}(s, t) = \frac{3\zeta^2 c_3 - 3\xi^2 c_3 \Lambda_1^2 - 2\eta_1 \Phi_2}{6\eta_1 \Phi_1} \mp \sqrt{\frac{2(\zeta^2 c_4 - \xi^2 c_4 \Lambda_1^2)}{\Phi_1}} \left(-\frac{\mu + \sqrt{\mu^2 - 4\rho\sigma} \left(\tanh\left(\sqrt{\mu^2 - 4\rho\sigma}\varrho\right) \pm \text{sech}\left(\sqrt{\mu^2 - 4\rho\sigma}\varrho\right) \right)}{2\rho} \right), \tag{53}$$

$$\Theta_{1,1,2}(s, t) = \frac{3\zeta^2 c_3 - 3\xi^2 c_3 \Lambda_1^2 - 2\eta_1 \Phi_2}{6\eta_1 \Phi_1} \mp \sqrt{\frac{2(\zeta^2 c_4 - \xi^2 c_4 \Lambda_1^2)}{\Phi_1}} \left(-\frac{\mu + \sqrt{\mu^2 - 4\rho\sigma} \coth\left(\frac{\sqrt{\mu^2 - 4\rho\sigma}\varrho}{2}\right)}{2\rho} \right), \tag{54}$$

$$\Theta_{1,1,3}(s, t) = \frac{3\zeta^2 c_3 - 3\xi^2 c_3 \Lambda_1^2 - 2\eta_1 \Phi_2}{6\eta_1 \Phi_1} \mp \sqrt{\frac{2(\zeta^2 c_4 - \xi^2 c_4 \Lambda_1^2)}{\Phi_1}} \left(-\frac{\mu + \sqrt{\mu^2 - 4\rho\sigma} \left(\tanh\left(\sqrt{\mu^2 - 4\rho\sigma}\varrho\right) \pm \text{sech}\left(\sqrt{\mu^2 - 4\rho\sigma}\varrho\right) \right)}{2\rho} \right), \tag{55}$$

$$\Theta_{1,1,4}(s, t) = \frac{3\zeta^2 c_3 - 3\xi^2 c_3 \Lambda_1^2 - 2\eta_1 \Phi_2}{6\eta_1 \Phi_1} \mp \sqrt{\frac{2(\zeta^2 c_4 - \xi^2 c_4 \Lambda_1^2)}{\Phi_1}} \left(-\frac{\mu + \sqrt{\mu^2 - 4\rho\sigma} \left(\coth\left(\sqrt{\mu^2 - 4\rho\sigma}\varrho\right) \pm \text{csch}\left(\sqrt{\mu^2 - 4\rho\sigma}\varrho\right) \right)}{2\rho} \right), \tag{56}$$

$$\Theta_{1,1,5}(s, t) = \frac{3\zeta^2 c_3 - 3\xi^2 c_3 \Lambda_1^2 - 2\eta_1 \Phi_2}{6\eta_1 \Phi_1} \mp \sqrt{\frac{2(\zeta^2 c_4 - \xi^2 c_4 \Lambda_1^2)}{\Phi_1}} \left(\frac{2\mu + \sqrt{\mu^2 - 4\rho\sigma} \left(\tanh\left(\frac{\sqrt{\mu^2 - 4\rho\sigma}}{4} \varrho\right) + \coth\left(\frac{\sqrt{\mu^2 - 4\rho\sigma}}{4} \varrho\right) \right)}{4\rho} \right), \quad (57)$$

$$\Theta_{1,1,6}(s, t) = \frac{3\zeta^2 c_3 - 3\xi^2 c_3 \Lambda_1^2 - 2\eta_1 \Phi_2}{6\eta_1 \Phi_1} \mp \sqrt{\frac{2(\zeta^2 c_4 - \xi^2 c_4 \Lambda_1^2)}{\Phi_1}} \left(\frac{1}{2\rho} \left(-\mu + \frac{\sqrt{(E^2 + F^2)(\mu^2 - 4\rho\sigma)} - E(\sqrt{\mu^2 - 4\rho\sigma}) \cosh(\sqrt{\mu^2 - 4\rho\sigma} \varrho)}{E \sinh(\sqrt{\mu^2 - 4\rho\sigma} \varrho) + F} \right) \right), \quad (58)$$

$$\Theta_{1,1,7}(s, t) = \frac{3\zeta^2 c_3 - 3\xi^2 c_3 \Lambda_1^2 - 2\eta_1 \Phi_2}{6\eta_1 \Phi_1} \mp \sqrt{\frac{2(\zeta^2 c_4 - \xi^2 c_4 \Lambda_1^2)}{\Phi_1}} \left(\frac{1}{2\rho} \left(-\mu - \frac{\sqrt{(F^2 - E^2)(\mu^2 - 4\rho\sigma)} + E(\sqrt{\mu^2 - 4\rho\sigma}) \sinh(\sqrt{\mu^2 - 4\rho\sigma} \varrho)}{E \cosh(\sqrt{\mu^2 - 4\rho\sigma} \varrho) + F} \right) \right), \quad (59)$$

$$\Theta_{1,1,8}(s, t) = \frac{3\zeta^2 c_3 - 3\xi^2 c_3 \Lambda_1^2 - 2\eta_1 \Phi_2}{6\eta_1 \Phi_1} \mp \sqrt{\frac{2(\zeta^2 c_4 - \xi^2 c_4 \Lambda_1^2)}{\Phi_1}} \left(\frac{2\sigma \cosh\left(\frac{\sqrt{\mu^2 - 4\rho\sigma}}{2} \varrho\right)}{\sqrt{\mu^2 - 4\rho\sigma} \sinh\left(\frac{\sqrt{\mu^2 - 4\rho\sigma}}{2} \varrho\right) - \mu \cosh\left(\frac{\sqrt{\mu^2 - 4\rho\sigma}}{2} \varrho\right)} \right), \quad (60)$$

$$\Theta_{1,1,9}(s, t) = \frac{3\zeta^2 c_3 - 3\xi^2 c_3 \Lambda_1^2 - 2\eta_1 \Phi_2}{6\eta_1 \Phi_1} \mp \sqrt{\frac{2(\zeta^2 c_4 - \xi^2 c_4 \Lambda_1^2)}{\Phi_1}} \left(\frac{-2\sigma \sinh\left(\frac{\sqrt{\mu^2 - 4\rho\sigma}}{2} \varrho\right)}{\mu \sinh\left(\frac{\sqrt{\mu^2 - 4\rho\sigma}}{2} \varrho\right) - \sqrt{\mu^2 - 4\rho\sigma} \cosh\left(\frac{\sqrt{\mu^2 - 4\rho\sigma}}{2} \varrho\right)} \right), \quad (61)$$

$$\Theta_{1,1,10}(s, t) = \frac{3\zeta^2 c_3 - 3\xi^2 c_3 \Lambda_1^2 - 2\eta_1 \Phi_2}{6\eta_1 \Phi_1} \mp \sqrt{\frac{2(\zeta^2 c_4 - \xi^2 c_4 \Lambda_1^2)}{\Phi_1}} \left(\frac{2\sigma \cosh(\sqrt{\mu^2 - 4\rho\sigma} \varrho)}{\sqrt{\mu^2 - 4\rho\sigma} \sinh(\sqrt{\mu^2 - 4\rho\sigma} \varrho) - (\mu \cosh(\sqrt{\mu^2 - 4\rho\sigma} \varrho) \pm i\sqrt{\mu^2 - 4\rho\sigma})} \right), \quad (62)$$

$$\Theta_{1,1,11}(s, t) = \frac{3\zeta^2 c_3 - 3\xi^2 c_3 \Lambda_1^2 - 2\eta_1 \Phi_2}{6\eta_1 \Phi_1} \mp \sqrt{\frac{2(\zeta^2 c_4 - \xi^2 c_4 \Lambda_1^2)}{\Phi_1}} \left(\frac{2\sigma \sinh(\sqrt{\mu^2 - 4\rho\sigma} \varrho)}{(\sqrt{\mu^2 - 4\rho\sigma} \cosh(\sqrt{\mu^2 - 4\rho\sigma} \varrho) \pm (\sqrt{\mu^2 - 4\rho\sigma})) - \mu \sinh(\sqrt{\mu^2 - 4\rho\sigma} \varrho)} \right), \quad (63)$$

$$\Theta_{1,1,12}(s, t) = \frac{3\zeta^2 c_3 - 3\xi^2 c_3 \Lambda_1^2 - 2\eta_1 \Phi_2}{6\eta_1 \Phi_1} \mp \sqrt{\frac{2(\zeta^2 c_4 - \xi^2 c_4 \Lambda_1^2)}{\Phi_1}} \left(\frac{4\sigma \cosh\left(\frac{\sqrt{\mu^2 - 4\rho\sigma}}{4} \varrho\right) \sinh\left(\frac{\sqrt{\mu^2 - 4\rho\sigma}}{4} \varrho\right)}{-2\mu \cosh\left(\frac{\sqrt{\mu^2 - 4\rho\sigma}}{4} \varrho\right) \sinh\left(\frac{\sqrt{\mu^2 - 4\rho\sigma}}{4} \varrho\right) + 2\sqrt{\mu^2 - 4\rho\sigma} \cosh^2\left(\frac{\sqrt{\mu^2 - 4\rho\sigma}}{4} \varrho\right) - (\sqrt{\mu^2 - 4\rho\sigma})} \right). \quad (64)$$

Type 2: $\mu^2 - 4\rho\sigma < 0, \mu\rho \neq 0, \rho\sigma \neq 0$

$$\Theta_{1,2,13}(s, t) = \frac{3\zeta^2 c_3 - 3\xi^2 c_3 \Lambda_1^2 - 2\eta_1 \Phi_2}{6\eta_1 \Phi_1} \mp \sqrt{\frac{2(\zeta^2 c_4 - \xi^2 c_4 \Lambda_1^2)}{\Phi_1}} \left(\frac{-\mu + \sqrt{4\rho\sigma - \mu^2} \tan\left(\frac{\sqrt{4\rho\sigma - \mu^2} \varrho}{2}\right)}{2\rho} \right), \quad (65)$$

$$\Theta_{1,2,14}(s, t) = \frac{3\zeta^2 c_3 - 3\xi^2 c_3 \Lambda_1^2 - 2\eta_1 \Phi_2}{6\eta_1 \Phi_1} \mp \sqrt{\frac{2(\zeta^2 c_4 - \xi^2 c_4 \Lambda_1^2)}{\Phi_1}} \left(\frac{-\mu - \sqrt{4\rho\sigma - \mu^2} \cot\left(\frac{\sqrt{4\rho\sigma - \mu^2} \varrho}{2}\right)}{2\rho} \right), \quad (66)$$

$$\Theta_{1,2,15}(s, t) = \frac{3\zeta^2 c_3 - 3\xi^2 c_3 \Lambda_1^2 - 2\eta_1 \Phi_2}{6\eta_1 \Phi_1} \mp \sqrt{\frac{2(\zeta^2 c_4 - \xi^2 c_4 \Lambda_1^2)}{\Phi_1}} \left(\frac{-\mu + \sqrt{4\rho\sigma - \mu^2} \left(\tan\left(\frac{\sqrt{4\rho\sigma - \mu^2} \varrho}{2}\right) \pm \sec\left(\frac{\sqrt{4\rho\sigma - \mu^2} \varrho}{2}\right) \right)}{2\rho} \right), \quad (67)$$

$$\Theta_{1,2,16}(s, t) = \frac{3\zeta^2 c_3 - 3\xi^2 c_3 \Lambda_1^2 - 2\eta_1 \Phi_2}{6\eta_1 \Phi_1} \mp \sqrt{\frac{2(\zeta^2 c_4 - \xi^2 c_4 \Lambda_1^2)}{\Phi_1}} \left(\frac{-\mu - \sqrt{4\rho\sigma - \mu^2} \left(\cot\left(\frac{\sqrt{4\rho\sigma - \mu^2} \varrho}{2}\right) \pm \csc\left(\frac{\sqrt{4\rho\sigma - \mu^2} \varrho}{2}\right) \right)}{2\rho} \right), \quad (68)$$

$$\Theta_{1,2,17}(s, t) = \frac{3\zeta^2 c_3 - 3\xi^2 c_3 \Lambda_1^2 - 2\eta_1 \Phi_2}{6\eta_1 \Phi_1} \mp \sqrt{\frac{2(\zeta^2 c_4 - \xi^2 c_4 \Lambda_1^2)}{\Phi_1}} \left(\frac{-2\mu + \sqrt{4\rho\sigma - \mu^2} \left(\tan\left(\frac{\sqrt{4\rho\sigma - \mu^2} \varrho}{4}\right) - \cot\left(\frac{\sqrt{4\rho\sigma - \mu^2} \varrho}{4}\right) \right)}{4\rho} \right), \quad (69)$$

$$\Theta_{1,2,18}(s, t) = \frac{3\zeta^2 c_3 - 3\xi^2 c_3 \Lambda_1^2 - 2\eta_1 \Phi_2}{6\eta_1 \Phi_1} \mp \sqrt{\frac{2(\zeta^2 c_4 - \xi^2 c_4 \Lambda_1^2)}{\Phi_1}} \left(\frac{1}{2\rho} \left(-\mu + \frac{\pm \sqrt{(E^2 - F^2)(4\rho\sigma - \mu^2)} - E(\sqrt{4\rho\sigma - \mu^2}) \cos\left(\frac{\sqrt{4\rho\sigma - \mu^2} \varrho}{2}\right)}{E \sin\left(\frac{\sqrt{4\rho\sigma - \mu^2} \varrho}{2}\right) + F} \right) \right), \quad (70)$$

$$\Theta_{1,2,19}(s, t) = \frac{3\zeta^2 c_3 - 3\xi^2 c_3 \Lambda_1^2 - 2\eta_1 \Phi_2}{6\eta_1 \Phi_1} \mp \sqrt{\frac{2(\zeta^2 c_4 - \xi^2 c_4 \Lambda_1^2)}{\Phi_1}} \left(\frac{1}{2\rho} \left(-\mu - \frac{\pm \sqrt{(E^2 - F^2)(4\rho\sigma - \mu^2)} - E(\sqrt{4\rho\sigma - \mu^2}) \sin\left(\frac{\sqrt{4\rho\sigma - \mu^2} \varrho}{2}\right)}{E \cos\left(\frac{\sqrt{4\rho\sigma - \mu^2} \varrho}{2}\right) + F} \right) \right), \quad (71)$$

$$\Theta_{1,2,20}(s, t) = \frac{3\zeta^2 c_3 - 3\xi^2 c_3 \Lambda_1^2 - 2\eta_1 \Phi_2}{6\eta_1 \Phi_1} \mp \sqrt{\frac{2(\zeta^2 c_4 - \xi^2 c_4 \Lambda_1^2)}{\Phi_1}} \left(\frac{2\sigma \cos\left(\frac{\sqrt{4\rho\sigma - \mu^2} \varrho}{2}\right)}{\sqrt{4\rho\sigma - \mu^2} \sin\left(\frac{\sqrt{4\rho\sigma - \mu^2} \varrho}{2}\right) + \mu \cos\left(\frac{\sqrt{4\rho\sigma - \mu^2} \varrho}{2}\right)} \right), \quad (72)$$

$$\Theta_{1,2,21}(s, t) = \frac{3\xi^2 c_3 - 3\xi^2 c_3 \Lambda_1^2 - 2\eta_1 \Phi_2}{6\eta_1 \Phi_1} \mp \sqrt{\frac{2(\xi^2 c_4 - \xi^2 c_4 \Lambda_1^2)}{\Phi_1}} \left(\frac{2\sigma \sin\left(\frac{\sqrt{4\rho\sigma - \mu^2}}{2} \varrho\right)}{-\mu \sin\left(\frac{\sqrt{4\rho\sigma - \mu^2}}{2} \varrho\right) + \sqrt{4\rho\sigma - \mu^2} \cos\left(\frac{\sqrt{4\rho\sigma - \mu^2}}{2} \varrho\right)} \right), \tag{73}$$

$$\Theta_{1,2,22}(s, t) = \frac{3\xi^2 c_3 - 3\xi^2 c_3 \Lambda_1^2 - 2\eta_1 \Phi_2}{6\eta_1 \Phi_1} \mp \sqrt{\frac{2(\xi^2 c_4 - \xi^2 c_4 \Lambda_1^2)}{\Phi_1}} \left(\frac{2\sigma \cos\left(\sqrt{4\rho\sigma - \mu^2} \varrho\right)}{\sqrt{4\rho\sigma - \mu^2} \sin\left(\sqrt{4\rho\sigma - \mu^2} \varrho\right) + \mu \cos\left(\sqrt{4\rho\sigma - \mu^2} \varrho\right) \pm \sqrt{4\rho\sigma - \mu^2}} \right), \tag{74}$$

$$\Theta_{1,2,23}(s, t) = \frac{3\xi^2 c_3 - 3\xi^2 c_3 \Lambda_1^2 - 2\eta_1 \Phi_2}{6\eta_1 \Phi_1} \mp \sqrt{\frac{2(\xi^2 c_4 - \xi^2 c_4 \Lambda_1^2)}{\Phi_1}} \left(\frac{2\sigma \sin\left(\sqrt{4\rho\sigma - \mu^2} \varrho\right)}{\left(\sqrt{4\rho\sigma - \mu^2} \cos\left(\sqrt{4\rho\sigma - \mu^2} \varrho\right) \pm \left(\sqrt{4\rho\sigma - \mu^2}\right) - \mu \sin\left(\sqrt{4\rho\sigma - \mu^2} \varrho\right)\right)} \right), \tag{75}$$

$$\Theta_{1,2,24}(s, t) = \frac{3\xi^2 c_3 - 3\xi^2 c_3 \Lambda_1^2 - 2\eta_1 \Phi_2}{6\eta_1 \Phi_1} \mp \sqrt{\frac{2(\xi^2 c_4 - \xi^2 c_4 \Lambda_1^2)}{\Phi_1}} \left(\chi_{24}^I(\varrho) = \frac{4\sigma \cos\left(\frac{\sqrt{4\rho\sigma - \mu^2}}{4} \varrho\right) \sin\left(\frac{\sqrt{4\rho\sigma - \mu^2}}{4} \varrho\right)}{-2\mu \cos\left(\frac{\sqrt{4\rho\sigma - \mu^2}}{4} \varrho\right) \sin\left(\frac{\sqrt{4\rho\sigma - \mu^2}}{4} \varrho\right) + 2\sqrt{\mu^2 - 4\rho\sigma} \cos^2\left(\frac{\sqrt{4\rho\sigma - \mu^2}}{4} \varrho\right) - \left(\sqrt{4\rho\sigma - \mu^2}\right)} \right). \tag{76}$$

Case 2

Type 1: $\rho\sigma < 0$ and $\rho\sigma \neq 0$

$$\Theta_{2,1,25}(s, t) = \frac{3\xi^2 c_3 - 3\xi^2 c_3 \Lambda_1^2 - 2\eta_1 \Phi_2}{6\eta_1 \Phi_1} \mp \sqrt{\frac{2(\xi^2 c_4 - \xi^2 c_4 \Lambda_1^2)}{\Phi_1}} \left(\frac{\pm\sqrt{-2\rho\sigma} + \sqrt{-6\rho\sigma} \tanh\left(\frac{\sqrt{-6\rho\sigma}\varrho}{2}\right)}{2\rho} \right), \tag{77}$$

$$\Theta_{2,1,26}(s, t) = \frac{3\xi^2 c_3 - 3\xi^2 c_3 \Lambda_1^2 - 2\eta_1 \Phi_2}{6\eta_1 \Phi_1} \mp \sqrt{\frac{2(\xi^2 c_4 - \xi^2 c_4 \Lambda_1^2)}{\Phi_1}} \left(\frac{\pm\sqrt{-2\rho\sigma} + \sqrt{-6\rho\sigma} \coth\left(\frac{\sqrt{-6\rho\sigma}\varrho}{2}\right)}{2\rho} \right), \tag{78}$$

$$\Theta_{2,1,27}(s, t) = \frac{3\xi^2 c_3 - 3\xi^2 c_3 \Lambda_1^2 - 2\eta_1 \Phi_2}{6\eta_1 \Phi_1} \mp \sqrt{\frac{2(\xi^2 c_4 - \xi^2 c_4 \Lambda_1^2)}{\Phi_1}} \left(\frac{\pm\sqrt{-2\rho\sigma} + \sqrt{-6\rho\sigma} (\tanh(\sqrt{-6\rho\sigma}\varrho) \pm \operatorname{sech}(\sqrt{-6\rho\sigma}\varrho))}{2\rho} \right), \tag{79}$$

$$\Theta_{2,1,28}(s, t) = \frac{3\zeta^2 c_3 - 3\xi^2 c_3 \Lambda_1^2 - 2\eta_1 \Phi_2}{6\eta_1 \Phi_1} \mp \sqrt{\frac{2(\zeta^2 c_4 - \xi^2 c_4 \Lambda_1^2)}{\Phi_1}} \left(-\frac{\pm \sqrt{-2\rho\sigma} + \sqrt{-6\rho\sigma} (\coth(\sqrt{-6\rho\sigma}\varrho) \pm \iota \operatorname{csch}(\sqrt{-6\rho\sigma}\varrho))}{2\rho} \right), \tag{80}$$

$$\Theta_{2,1,29}(s, t) = \frac{3\zeta^2 c_3 - 3\xi^2 c_3 \Lambda_1^2 - 2\eta_1 \Phi_2}{6\eta_1 \Phi_1} \mp \sqrt{\frac{2(\zeta^2 c_4 - \xi^2 c_4 \Lambda_1^2)}{\Phi_1}} \left(-\frac{\pm \sqrt{-2\rho\sigma} + \sqrt{-6\rho\sigma} \left(\tanh\left(\frac{\sqrt{-6\rho\sigma}\varrho}{4}\right) + \coth\left(\frac{\sqrt{-6\rho\sigma}\varrho}{4}\right) \right)}{4\rho} \right), \tag{81}$$

$$\Theta_{2,1,30}(s, t) = \frac{3\zeta^2 c_3 - 3\xi^2 c_3 \Lambda_1^2 - 2\eta_1 \Phi_2}{6\eta_1 \Phi_1} \mp \sqrt{\frac{2(\zeta^2 c_4 - \xi^2 c_4 \Lambda_1^2)}{\Phi_1}} \left(\frac{1}{2\rho} \left(\mp \sqrt{-2\rho\sigma} + \frac{\sqrt{(E^2 + F^2)(-6\rho\sigma)} - E(\sqrt{-6\rho\sigma}) \cosh(\sqrt{-6\rho\sigma}\varrho)}{E \sinh(\sqrt{-6\rho\sigma}\varrho) + F} \right) \right), \tag{82}$$

$$\Theta_{2,1,31}(s, t) = \frac{3\zeta^2 c_3 - 3\xi^2 c_3 \Lambda_1^2 - 2\eta_1 \Phi_2}{6\eta_1 \Phi_1} \mp \sqrt{\frac{2(\zeta^2 c_4 - \xi^2 c_4 \Lambda_1^2)}{\Phi_1}} \left(\frac{1}{2\rho} \left(\mp \sqrt{-2\rho\sigma} - \frac{\sqrt{(F^2 - E^2)(-6\rho\sigma)} + E(\sqrt{-6\rho\sigma}) \sinh(\sqrt{-6\rho\sigma}\varrho)}{E \cosh(\sqrt{-6\rho\sigma}\varrho) + F} \right) \right), \tag{83}$$

where E and F are non-zero constants that fulfill $F^2 - E^2 > 0$.

$$\Theta_{2,1,32}(s, t) = \frac{3\zeta^2 c_3 - 3\xi^2 c_3 \Lambda_1^2 - 2\eta_1 \Phi_2}{6\eta_1 \Phi_1} \mp \sqrt{\frac{2(\zeta^2 c_4 - \xi^2 c_4 \Lambda_1^2)}{\Phi_1}} \left(\frac{2\sigma \cosh\left(\frac{\sqrt{-6\rho\sigma}\varrho}{2}\right)}{\sqrt{-6\rho\sigma} \sinh\left(\frac{\sqrt{-6\rho\sigma}\varrho}{2}\right) \mp \sqrt{-2\rho\sigma} \cosh\left(\frac{\sqrt{-6\rho\sigma}\varrho}{2}\right)} \right), \tag{84}$$

$$\Theta_{2,1,33}(s, t) = \frac{3\zeta^2 c_3 - 3\xi^2 c_3 \Lambda_1^2 - 2\eta_1 \Phi_2}{6\eta_1 \Phi_1} \mp \sqrt{\frac{2(\zeta^2 c_4 - \xi^2 c_4 \Lambda_1^2)}{\Phi_1}} \left(\frac{-2\sigma \sinh\left(\frac{\sqrt{-6\rho\sigma}\varrho}{2}\right)}{\pm \sqrt{-2\rho\sigma} \sinh\left(\frac{\sqrt{-6\rho\sigma}\varrho}{2}\right) - \sqrt{-6\rho\sigma} \cosh\left(\frac{\sqrt{-6\rho\sigma}\varrho}{2}\right)} \right), \tag{85}$$

$$\Theta_{2,1,34}(s, t) = \frac{3\zeta^2 c_3 - 3\xi^2 c_3 \Lambda_1^2 - 2\eta_1 \Phi_2}{6\eta_1 \Phi_1} \mp \sqrt{\frac{2(\zeta^2 c_4 - \xi^2 c_4 \Lambda_1^2)}{\Phi_1}} \left(\frac{2\sigma \cosh(\sqrt{-6\rho\sigma}\varrho)}{\sqrt{-6\rho\sigma} \sinh(\sqrt{-6\rho\sigma}\varrho) \mp \sqrt{-2\rho\sigma} \cosh(\sqrt{-6\rho\sigma}\varrho) \pm \iota(\sqrt{-6\rho\sigma})} \right), \tag{86}$$

$$\Theta_{2,1,35}(s, t) = \frac{3\zeta^2 c_3 - 3\xi^2 c_3 \Lambda_1^2 - 2\eta_1 \Phi_2}{6\eta_1 \Phi_1} \mp \sqrt{\frac{2(\zeta^2 c_4 - \xi^2 c_4 \Lambda_1^2)}{\Phi_1}} \left(\frac{2\sigma \sinh(\sqrt{-6\rho\sigma}\varrho)}{\mp \sqrt{-2\rho\sigma} \sinh(\sqrt{-6\rho\sigma}\varrho) + \sqrt{-6\rho\sigma} \cosh(\sqrt{-6\rho\sigma}\varrho) \pm \iota(\sqrt{-6\rho\sigma})} \right), \tag{87}$$

$$\Theta_{2,1,36}(s, t) = \frac{3\zeta^2 c_3 - 3\xi^2 c_3 \Lambda_1^2 - 2\eta_1 \Phi_2}{6\eta_1 \Phi_1} \mp \sqrt{\frac{2(\zeta^2 c_4 - \xi^2 c_4 \Lambda_1^2)}{\Phi_1}} \left(\frac{4\sigma \cosh\left(\frac{\sqrt{-6\rho\sigma}\varrho}{4}\right) \sinh\left(\frac{\sqrt{-6\rho\sigma}\varrho}{4}\right)}{\pm 2\sqrt{-2\rho\sigma} \cosh\left(\frac{\sqrt{-6\rho\sigma}\varrho}{4}\right) \sinh\left(\frac{\sqrt{-6\rho\sigma}\varrho}{4}\right) + 2\sqrt{-6\rho\sigma} \cosh^2\left(\frac{\sqrt{-6\rho\sigma}\varrho}{4}\right) - (\sqrt{-6\rho\sigma})} \right). \tag{88}$$

Case 3

Type 1: When $c_2 = 1, c_3 = -\frac{2\alpha_3}{\alpha_1}, c_4 = \frac{\alpha_3^2 - \alpha_2^2}{\alpha_1^2}$

$$\Theta_{3,1,37}(s, t) = \frac{3\zeta^2 c_3 - 3\xi^2 c_3 \Lambda_1^2 - 2\eta_1 \Phi_2}{6\eta_1 \Phi_1} \mp \sqrt{\frac{2(\zeta^2 c_4 - \xi^2 c_4 \Lambda_1^2)}{\Phi_1}} \left(\frac{\alpha_1 \operatorname{sech}(\varrho)}{\alpha_2 + \alpha_3 \operatorname{sech}(\varrho)} \right). \tag{89}$$

Type 2: When $c_2 = 1, c_3 = -\frac{2\alpha_3}{\alpha_1}, c_4 = \frac{\alpha_3^2 + \alpha_2^2}{\alpha_1^2}$

$$\Theta_{3,2,38}(s, t) = \frac{3\zeta^2 c_3 - 3\xi^2 c_3 \Lambda_1^2 - 2\eta_1 \Phi_2}{6\eta_1 \Phi_1} \mp \sqrt{\frac{2(\zeta^2 c_4 - \xi^2 c_4 \Lambda_1^2)}{\Phi_1}} \left(\frac{\alpha_1 \operatorname{csch}(\varrho)}{\alpha_2 + \alpha_3 \operatorname{csch}(\varrho)} \right). \tag{90}$$

Type 3: When $c_2 = 4, c_3 = -\frac{4(2\alpha_2 + \alpha_4)}{\alpha_1}, c_4 = \frac{\alpha_3^2 + 4\alpha_2^2 + 4\alpha_2\alpha_4}{\alpha_1^2}$

$$\Theta_{3,3,39}(s, t) = \frac{3\zeta^2 c_3 - 3\xi^2 c_3 \Lambda_1^2 - 2\eta_1 \Phi_2}{6\eta_1 \Phi_1} \mp \sqrt{\frac{2(\zeta^2 c_4 - \xi^2 c_4 \Lambda_1^2)}{\Phi_1}} \left(\frac{\alpha_1 \operatorname{sech}^2(\varrho)}{\alpha_2 \operatorname{sech}^2(\varrho) + \alpha_3 \tanh(\varrho) + \alpha_4} \right). \tag{91}$$

Type 4: When $c_2 = 4, c_3 = -\frac{4(\alpha_4 - 2\alpha_2)}{\alpha_1}, c_4 = \frac{\alpha_3^2 + 4\alpha_2^2 - 4\alpha_2\alpha_4}{\alpha_1^2}$

$$\Theta_{3,4,40}(s, t) = \frac{3\zeta^2 c_3 - 3\xi^2 c_3 \Lambda_1^2 - 2\eta_1 \Phi_2}{6\eta_1 \Phi_1} \mp \sqrt{\frac{2(\zeta^2 c_4 - \xi^2 c_4 \Lambda_1^2)}{\Phi_1}} \left(\frac{\alpha_1 \operatorname{csch}^2(\varrho)}{\alpha_2 \coth^2(\varrho) + \alpha_3 \tanh(\varrho) + \alpha_4} \right). \tag{92}$$

Type 5: When $c_2 = \alpha_1^2, c_3 = 2\alpha_1\alpha_2, c_4 = \alpha_2^2$

$$\Theta_{3,5,41}(s, t) = \frac{3\zeta^2 c_3 - 3\xi^2 c_3 \Lambda_1^2 - 2\eta_1 \Phi_2}{6\eta_1 \Phi_1} \mp \sqrt{\frac{2(\zeta^2 c_4 - \xi^2 c_4 \Lambda_1^2)}{\Phi_1}} \left(-\frac{\alpha_1 \alpha_3}{\alpha_2 (\cosh(\alpha_1 \varrho) - \sinh(\alpha_1 \varrho) + \alpha_3)} \right), \tag{93}$$

$$\Theta_{3,5,42}(s, t) = \frac{3\zeta^2 c_3 - 3\xi^2 c_3 \Lambda_1^2 - 2\eta_1 \Phi_2}{6\eta_1 \Phi_1} \mp \sqrt{\frac{2(\zeta^2 c_4 - \xi^2 c_4 \Lambda_1^2)}{\Phi_1}} \left(-\frac{\alpha_1 (\sinh(\alpha_1 \varrho) + \cosh(\alpha_1 \varrho))}{\alpha_2 (\sinh(\alpha_1 \varrho) + \cosh(\alpha_1 \varrho) + \alpha_3)} \right). \tag{94}$$

Type 6: When $c_2 = -1, c_3 = \frac{2\alpha_3}{\alpha_1}, c_4 = -\frac{\alpha_3^2 - \alpha_2^2}{\alpha_1^2}$

$$\Theta_{3,6,43}(s, t) = \frac{3\zeta^2 c_3 - 3\xi^2 c_3 \Lambda_1^2 - 2\eta_1 \Phi_2}{6\eta_1 \Phi_1} \mp \sqrt{\frac{2(\zeta^2 c_4 - \xi^2 c_4 \Lambda_1^2)}{\Phi_1}} \left(\frac{\alpha_1 \sec(\varrho)}{\alpha_2 + \alpha_3 \sec(\varrho)} \right), \tag{95}$$

$$\Theta_{3,6,44}(s, t) = \frac{3\zeta^2 c_3 - 3\xi^2 c_3 \Lambda_1^2 - 2\eta_1 \Phi_2}{6\eta_1 \Phi_1} \mp \sqrt{\frac{2(\zeta^2 c_4 - \xi^2 c_4 \Lambda_1^2)}{\Phi_1}} \left(\frac{\alpha_1 \csc(\varrho)}{\alpha_2 + \alpha_3 \csc(\varrho)} \right). \tag{96}$$

Type 7: When $c_2 = -4$, $c_3 = \frac{4(2\alpha_2 + \alpha_4)}{\alpha_1}$, $c_4 = -\frac{-\alpha_3^2 + 4\alpha_2^2 + 4\alpha_2\alpha_4}{\alpha_1^2}$

$$\Theta_{3,7,45}(s, t) = \frac{3\zeta^2 c_3 - 3\xi^2 c_3 \Lambda_1^2 - 2\eta_1 \Phi_2}{6\eta_1 \Phi_1} \mp \sqrt{\frac{2(\zeta^2 c_4 - \xi^2 c_4 \Lambda_1^2)}{\Phi_1}} \left(\frac{\alpha_1 \sec^2(\varrho)}{\alpha_2 \sec^2(\varrho) + \alpha_3 \tan(\varrho) + \alpha_4} \frac{\alpha_1 \sec^2(\varrho)}{\alpha_2 \sec^2(\varrho) + \alpha_3 \tan(\varrho) + \alpha_4} \right), \quad (97)$$

$$\Theta_{3,7,46}(s, t) = \frac{3\zeta^2 c_3 - 3\xi^2 c_3 \Lambda_1^2 - 2\eta_1 \Phi_2}{6\eta_1 \Phi_1} \mp \sqrt{\frac{2(\zeta^2 c_4 - \xi^2 c_4 \Lambda_1^2)}{\Phi_1}} \left(\frac{\alpha_1 \csc^2(\varrho)}{\alpha_2 \csc^2(\varrho) + \alpha_3 \cot(\varrho) + \alpha_4} \right), \quad (98)$$

where the constants $\alpha_1, \alpha_2, \alpha_3$ and α_4 are arbitrary.

Case 4

Type 1: When $c_0 = 1$, $c_2 = -(1 - q^2)$, $c_4 = q^2$

$$\Theta_{4,1,47}(s, t) = \frac{3\zeta^2 c_3 - 3\xi^2 c_3 \Lambda_1^2 - 2\eta_1 \Phi_2}{6\eta_1 \Phi_1} \mp \sqrt{\frac{2(\zeta^2 c_4 - \xi^2 c_4 \Lambda_1^2)}{\Phi_1}} (cn\varrho), \quad (99)$$

$$\Theta_{4,1,48}(s, t) = \frac{3\zeta^2 c_3 - 3\xi^2 c_3 \Lambda_1^2 - 2\eta_1 \Phi_2}{6\eta_1 \Phi_1} \mp \sqrt{\frac{2(\zeta^2 c_4 - \xi^2 c_4 \Lambda_1^2)}{\Phi_1}} \left(\frac{cn\varrho}{cd\varrho} \right). \quad (100)$$

Type 2: When $c_0 = 1 - q^2$, $c_2 = 2q^2 - 1$, $c_4 = -q^2$

$$\Theta_{4,2,49}(s, t) = \frac{3\zeta^2 c_3 - 3\xi^2 c_3 \Lambda_1^2 - 2\eta_1 \Phi_2}{6\eta_1 \Phi_1} \mp \sqrt{\frac{2(\zeta^2 c_4 - \xi^2 c_4 \Lambda_1^2)}{\Phi_1}} (cn\varrho). \quad (101)$$

Type 3: When $c_0 = q^2 - 1$, $c_2 = 2 - q^2$, $c_4 = -1$

$$\Theta_{4,3,50}(s, t) = \frac{3\zeta^2 c_3 - 3\xi^2 c_3 \Lambda_1^2 - 2\eta_1 \Phi_2}{6\eta_1 \Phi_1} \mp \sqrt{\frac{2(\zeta^2 c_4 - \xi^2 c_4 \Lambda_1^2)}{\Phi_1}} (dn\varrho). \quad (102)$$

Type 4: When $c_0 = q^2$, $c_2 = -(1 + q^2)$, $c_4 = 1$

$$\Theta_{4,4,51}(s, t) = \frac{3\zeta^2 c_3 - 3\xi^2 c_3 \Lambda_1^2 - 2\eta_1 \Phi_2}{6\eta_1 \Phi_1} \mp \sqrt{\frac{2(\zeta^2 c_4 - \xi^2 c_4 \Lambda_1^2)}{\Phi_1}} (rn\varrho)^{-1}, \quad (103)$$

$$\Theta_{4,4,52}(s, t) = \frac{3\zeta^2 c_3 - 3\xi^2 c_3 \Lambda_1^2 - 2\eta_1 \Phi_2}{6\eta_1 \Phi_1} \mp \sqrt{\frac{2(\zeta^2 c_4 - \xi^2 c_4 \Lambda_1^2)}{\Phi_1}} \left(\frac{dn\varrho}{cn\varrho} \right). \quad (104)$$

Type 5: When $c_0 = -q^2$, $c_2 = 2q^2 - 1$, $c_4 = 1 - q^2$

$$\Theta_{4,5,53}(s, t) = \frac{3\zeta^2 c_3 - 3\xi^2 c_3 \Lambda_1^2 - 2\eta_1 \Phi_2}{6\eta_1 \Phi_1} \mp \sqrt{\frac{2(\zeta^2 c_4 - \xi^2 c_4 \Lambda_1^2)}{\Phi_1}} (cn\varrho)^{-1}. \quad (105)$$

Type 6: When $c_0 = -1$, $c_2 = 2 - q^2$, $c_4 = q^2 - 1$

$$\Theta_{4,6,54}(s, t) = \frac{3\zeta^2 c_3 - 3\xi^2 c_3 \Lambda_1^2 - 2\eta_1 \Phi_2}{6\eta_1 \Phi_1} \mp \sqrt{\frac{2(\zeta^2 c_4 - \xi^2 c_4 \Lambda_1^2)}{\Phi_1}} (dn\varrho)^{-1}. \quad (106)$$

Type 7: When $c_0 = 1, c_2 = 2 - q^2, c_4 = 1 - q^2$

$$\Theta_{4,7,55}(s, t) = \frac{3\zeta^2 c_3 - 3\xi^2 c_3 \Lambda_1^2 - 2\eta_1 \Phi_2}{6\eta_1 \Phi_1} \mp \sqrt{\frac{2(\zeta^2 c_4 - \xi^2 c_4 \Lambda_1^2)}{\Phi_1}} \left(\frac{rn\varrho}{cn\varrho} \right). \quad (107)$$

Type 8: When $c_0 = 1, c_2 = 2q^2 - 1, c_4 = -q^2(1 - q^2)$

$$\Theta_{4,8,56}(s, t) = \frac{3\zeta^2 c_3 - 3\xi^2 c_3 \Lambda_1^2 - 2\eta_1 \Phi_2}{6\eta_1 \Phi_1} \mp \sqrt{\frac{2(\zeta^2 c_4 - \xi^2 c_4 \Lambda_1^2)}{\Phi_1}} \left(\frac{rn\varrho}{dn\varrho} \right). \quad (108)$$

Type 9: When $c_0 = 1 - q^2, c_2 = 2 - q^2, c_4 = 1$

$$\Theta_{4,9,57}(s, t) = \frac{3\zeta^2 c_3 - 3\xi^2 c_3 \Lambda_1^2 - 2\eta_1 \Phi_2}{6\eta_1 \Phi_1} \mp \sqrt{\frac{2(\zeta^2 c_4 - \xi^2 c_4 \Lambda_1^2)}{\Phi_1}} \left(\frac{cn\varrho}{rn\varrho} \right). \quad (109)$$

Type 10: When $c_0 = -q^2(1 - q^2), c_2 = 2q^2 - 1, c_4 = 1$

$$\Theta_{4,10,58}(s, t) = \frac{3\zeta^2 c_3 - 3\xi^2 c_3 \Lambda_1^2 - 2\eta_1 \Phi_2}{6\eta_1 \Phi_1} \mp \sqrt{\frac{2(\zeta^2 c_4 - \xi^2 c_4 \Lambda_1^2)}{\Phi_1}} \left(\frac{dn\varrho}{rn\varrho} \right). \quad (110)$$

Type 11: When $c_0 = \frac{1}{4}, c_2 = \frac{1-2q^2}{2}, c_4 = \text{frac}14$

$$\Theta_{4,11,59}(s, t) = \frac{3\zeta^2 c_3 - 3\xi^2 c_3 \Lambda_1^2 - 2\eta_1 \Phi_2}{6\eta_1 \Phi_1} \mp \sqrt{\frac{2(\zeta^2 c_4 - \xi^2 c_4 \Lambda_1^2)}{\Phi_1}} (nr\varrho \pm cr\varrho). \quad (111)$$

Type 12: When $c_0 = \frac{1-q^2}{4}, c_2 = \frac{1+q^2}{2}, c_4 = \frac{1-q^2}{4}$

$$\Theta_{4,12,60}(s, t) = \frac{3\zeta^2 c_3 - 3\xi^2 c_3 \Lambda_1^2 - 2\eta_1 \Phi_2}{6\eta_1 \Phi_1} \mp \sqrt{\frac{2(\zeta^2 c_4 - \xi^2 c_4 \Lambda_1^2)}{\Phi_1}} (nc\varrho \pm rc\varrho). \quad (112)$$

Type 13: When $c_0 = \frac{q^2}{4}, c_2 = \frac{q^2-2}{2}, c_4 = \frac{1}{4}$

$$\Theta_{4,13,61}(s, t) = \frac{3\zeta^2 c_3 - 3\xi^2 c_3 \Lambda_1^2 - 2\eta_1 \Phi_2}{6\eta_1 \Phi_1} \mp \sqrt{\frac{2(\zeta^2 c_4 - \xi^2 c_4 \Lambda_1^2)}{\Phi_1}} (nr\varrho \pm dr\varrho). \quad (113)$$

Type 14: When $c_0 = \frac{q^2}{4}, c_2 = \frac{q^2-2}{2}, c_4 = \frac{q^2}{4}$

$$\Theta_{4,14,62}(s, t) = \frac{3\zeta^2 c_3 - 3\xi^2 c_3 \Lambda_1^2 - 2\eta_1 \Phi_2}{6\eta_1 \Phi_1} \mp \sqrt{\frac{2(\zeta^2 c_4 - \xi^2 c_4 \Lambda_1^2)}{\Phi_1}} (rn\varrho \pm icr\varrho), \quad (114)$$

where q is the JEF satisfying $0 \leq q \leq 1$. Equations (99)–(114) can be stated as follows when $q \rightarrow 1$, JEFs degenerate into hyperbolic functions, which is illustrated in Table 2.

$$\Theta_{4,1,63}(s, t) = \frac{3\zeta^2 c_3 - 3\xi^2 c_3 \Lambda_1^2 - 2\eta_1 \Phi_2}{6\eta_1 \Phi_1} \mp \sqrt{\frac{2(\zeta^2 c_4 - \xi^2 c_4 \Lambda_1^2)}{\Phi_1}} (\text{sech}\varrho), \quad (115)$$

$$\Theta_{4,1,64}(s, t) = \frac{3\zeta^2 c_3 - 3\xi^2 c_3 \Lambda_1^2 - 2\eta_1 \Phi_2}{6\eta_1 \Phi_1} \mp \sqrt{\frac{2(\zeta^2 c_4 - \xi^2 c_4 \Lambda_1^2)}{\Phi_1}} (\text{sech}\varrho), \quad (116)$$

$$\Theta_{4,2,65}(s, t) = \frac{3\zeta^2 c_3 - 3\xi^2 c_3 \Lambda_1^2 - 2\eta_1 \Phi_2}{6\eta_1 \Phi_1} \mp \sqrt{\frac{2(\zeta^2 c_4 - \xi^2 c_4 \Lambda_1^2)}{\Phi_1}} (\operatorname{sech} \varrho), \quad (117)$$

$$\Theta_{4,3,66}(s, t) = \frac{3\zeta^2 c_3 - 3\xi^2 c_3 \Lambda_1^2 - 2\eta_1 \Phi_2}{6\eta_1 \Phi_1} \mp \sqrt{\frac{2(\zeta^2 c_4 - \xi^2 c_4 \Lambda_1^2)}{\Phi_1}} (\operatorname{sech} \varrho), \quad (118)$$

$$\Theta_{4,4,67}(s, t) = \frac{3\zeta^2 c_3 - 3\xi^2 c_3 \Lambda_1^2 - 2\eta_1 \Phi_2}{6\eta_1 \Phi_1} \mp \sqrt{\frac{2(\zeta^2 c_4 - \xi^2 c_4 \Lambda_1^2)}{\Phi_1}} (\tanh \varrho)^{-1}, \quad (119)$$

$$\Theta_{4,4,68}(s, t) = \frac{3\zeta^2 c_3 - 3\xi^2 c_3 \Lambda_1^2 - 2\eta_1 \Phi_2}{6\eta_1 \Phi_1} \mp \sqrt{\frac{2(\zeta^2 c_4 - \xi^2 c_4 \Lambda_1^2)}{\Phi_1}} \left(\frac{\operatorname{sech} \varrho}{\operatorname{sech} \varrho} \right), \quad (120)$$

$$\Theta_{4,5,69}(s, t) = \frac{3\zeta^2 c_3 - 3\xi^2 c_3 \Lambda_1^2 - 2\eta_1 \Phi_2}{6\eta_1 \Phi_1} \mp \sqrt{\frac{2(\zeta^2 c_4 - \xi^2 c_4 \Lambda_1^2)}{\Phi_1}} (\operatorname{sech} \varrho)^{-1}, \quad (121)$$

$$\Theta_{4,6,70}(s, t) = \frac{3\zeta^2 c_3 - 3\xi^2 c_3 \Lambda_1^2 - 2\eta_1 \Phi_2}{6\eta_1 \Phi_1} \mp \sqrt{\frac{2(\zeta^2 c_4 - \xi^2 c_4 \Lambda_1^2)}{\Phi_1}} (\operatorname{sech} \varrho)^{-1}, \quad (122)$$

$$\Theta_{4,7,71}(s, t) = \frac{3\zeta^2 c_3 - 3\xi^2 c_3 \Lambda_1^2 - 2\eta_1 \Phi_2}{6\eta_1 \Phi_1} \mp \sqrt{\frac{2(\zeta^2 c_4 - \xi^2 c_4 \Lambda_1^2)}{\Phi_1}} \left(\frac{\tanh \varrho}{\operatorname{sech} \varrho} \right), \quad (123)$$

$$\Theta_{4,8,72}(s, t) = \frac{3\zeta^2 c_3 - 3\xi^2 c_3 \Lambda_1^2 - 2\eta_1 \Phi_2}{6\eta_1 \Phi_1} \mp \sqrt{\frac{2(\zeta^2 c_4 - \xi^2 c_4 \Lambda_1^2)}{\Phi_1}} \left(\frac{\sinh \varrho}{\operatorname{sech} \varrho} \right), \quad (124)$$

$$\Theta_{4,9,73}(s, t) = \frac{3\zeta^2 c_3 - 3\xi^2 c_3 \Lambda_1^2 - 2\eta_1 \Phi_2}{6\eta_1 \Phi_1} \mp \sqrt{\frac{2(\zeta^2 c_4 - \xi^2 c_4 \Lambda_1^2)}{\Phi_1}} \left(\frac{\operatorname{sech} \varrho}{\tanh \varrho} \right), \quad (125)$$

$$\Theta_{4,10,74}(s, t) = \frac{3\zeta^2 c_3 - 3\xi^2 c_3 \Lambda_1^2 - 2\eta_1 \Phi_2}{6\eta_1 \Phi_1} \mp \sqrt{\frac{2(\zeta^2 c_4 - \xi^2 c_4 \Lambda_1^2)}{\Phi_1}} \left(\frac{\operatorname{sech} \varrho}{\tanh \varrho} \right), \quad (126)$$

$$\Theta_{4,11,75}(s, t) = \frac{3\zeta^2 c_3 - 3\xi^2 c_3 \Lambda_1^2 - 2\eta_1 \Phi_2}{6\eta_1 \Phi_1} \mp \sqrt{\frac{2(\zeta^2 c_4 - \xi^2 c_4 \Lambda_1^2)}{\Phi_1}} (\coth \varrho \pm \operatorname{csch} \varrho), \quad (127)$$

$$\Theta_{4,12,76}(s, t) = \frac{3\zeta^2 c_3 - 3\xi^2 c_3 \Lambda_1^2 - 2\eta_1 \Phi_2}{6\eta_1 \Phi_1} \mp \sqrt{\frac{2(\zeta^2 c_4 - \xi^2 c_4 \Lambda_1^2)}{\Phi_1}} (\cosh \varrho \pm \sinh \varrho), \quad (128)$$

$$\Theta_{4,13,77}(s, t) = \frac{3\zeta^2 c_3 - 3\xi^2 c_3 \Lambda_1^2 - 2\eta_1 \Phi_2}{6\eta_1 \Phi_1} \mp \sqrt{\frac{2(\zeta^2 c_4 - \xi^2 c_4 \Lambda_1^2)}{\Phi_1}} (\coth \varrho \pm \operatorname{csch} \varrho), \quad (129)$$

$$\Theta_{4,14,78}(s, t) = \frac{3\zeta^2 c_3 - 3\xi^2 c_3 \Lambda_1^2 - 2\eta_1 \Phi_2}{6\eta_1 \Phi_1} \mp \sqrt{\frac{2(\zeta^2 c_4 - \xi^2 c_4 \Lambda_1^2)}{\Phi_1}} (\tanh \varrho \pm \operatorname{csch} \varrho), \quad (130)$$

likewise, when $q \rightarrow 0$, JEFs degenerate into trigonometric functions, as shown in Table 3, Eqs. (99)–(114) can be expressed as subsequently

$$\Theta_{4,1,79}(s, t) = \frac{3\zeta^2 c_3 - 3\xi^2 c_3 \Lambda_1^2 - 2\eta_1 \Phi_2}{6\eta_1 \Phi_1} \mp \sqrt{\frac{2(\zeta^2 c_4 - \xi^2 c_4 \Lambda_1^2)}{\Phi_1}} (\cos \varrho), \quad (131)$$

$$\Theta_{4,1,80}(s, t) = \frac{3\zeta^2 c_3 - 3\xi^2 c_3 \Lambda_1^2 - 2\eta_1 \Phi_2}{6\eta_1 \Phi_1} \mp \sqrt{\frac{2(\zeta^2 c_4 - \xi^2 c_4 \Lambda_1^2)}{\Phi_1}} \quad (132)$$

$$\Theta_{4,2,81}(s, t) = \frac{3\zeta^2 c_3 - 3\xi^2 c_3 \Lambda_1^2 - 2\eta_1 \Phi_2}{6\eta_1 \Phi_1} \mp \sqrt{\frac{2(\zeta^2 c_4 - \xi^2 c_4 \Lambda_1^2)}{\Phi_1}} (\cos \varrho), \quad (133)$$

$$\Theta_{4,3,82}(s, t) = \frac{3\zeta^2 c_3 - 3\xi^2 c_3 \Lambda_1^2 - 2\eta_1 \Phi_2}{6\eta_1 \Phi_1} \mp \sqrt{\frac{2(\zeta^2 c_4 - \xi^2 c_4 \Lambda_1^2)}{\Phi_1}}, \quad (134)$$

$$\Theta_{4,4,83}(s, t) = \frac{3\zeta^2 c_3 - 3\xi^2 c_3 \Lambda_1^2 - 2\eta_1 \Phi_2}{6\eta_1 \Phi_1} \mp \sqrt{\frac{2(\zeta^2 c_4 - \xi^2 c_4 \Lambda_1^2)}{\Phi_1}} (\sin \varrho)^{-1}, \quad (135)$$

$$\Theta_{4,4,84}(s, t) = \frac{3\zeta^2 c_3 - 3\xi^2 c_3 \Lambda_1^2 - 2\eta_1 \Phi_2}{6\eta_1 \Phi_1} \mp \sqrt{\frac{2(\zeta^2 c_4 - \xi^2 c_4 \Lambda_1^2)}{\Phi_1}} \left(\frac{d\eta \varrho}{\cos \varrho} \right), \quad (136)$$

$$\Theta_{4,5,85}(s, t) = \frac{3\zeta^2 c_3 - 3\xi^2 c_3 \Lambda_1^2 - 2\eta_1 \Phi_2}{6\eta_1 \Phi_1} \mp \sqrt{\frac{2(\zeta^2 c_4 - \xi^2 c_4 \Lambda_1^2)}{\Phi_1}} (\cos \varrho)^{-1}, \quad (137)$$

$$\Theta_{4,6,86}(s, t) = \frac{3\zeta^2 c_3 - 3\xi^2 c_3 \Lambda_1^2 - 2\eta_1 \Phi_2}{6\eta_1 \Phi_1} \mp \sqrt{\frac{2(\zeta^2 c_4 - \xi^2 c_4 \Lambda_1^2)}{\Phi_1}}, \quad (138)$$

$$\Theta_{4,7,87}(s, t) = \frac{3\zeta^2 c_3 - 3\xi^2 c_3 \Lambda_1^2 - 2\eta_1 \Phi_2}{6\eta_1 \Phi_1} \mp \sqrt{\frac{2(\zeta^2 c_4 - \xi^2 c_4 \Lambda_1^2)}{\Phi_1}} \left(\frac{\sin \varrho}{\cos \varrho} \right), \quad (139)$$

$$\Theta_{4,8,88}(s, t) = \frac{3\zeta^2 c_3 - 3\xi^2 c_3 \Lambda_1^2 - 2\eta_1 \Phi_2}{6\eta_1 \Phi_1} \mp \sqrt{\frac{2(\zeta^2 c_4 - \xi^2 c_4 \Lambda_1^2)}{\Phi_1}} (\sin \varrho), \quad (140)$$

$$\Theta_{4,9,89}(s, t) = \frac{3\zeta^2 c_3 - 3\xi^2 c_3 \Lambda_1^2 - 2\eta_1 \Phi_2}{6\eta_1 \Phi_1} \mp \sqrt{\frac{2(\zeta^2 c_4 - \xi^2 c_4 \Lambda_1^2)}{\Phi_1}} \left(\frac{\cos \varrho}{\sin \varrho} \right), \quad (141)$$

$$\Theta_{4,10,90}(s, t) = \frac{3\zeta^2 c_3 - 3\xi^2 c_3 \Lambda_1^2 - 2\eta_1 \Phi_2}{6\eta_1 \Phi_1} \mp \sqrt{\frac{2(\zeta^2 c_4 - \xi^2 c_4 \Lambda_1^2)}{\Phi_1}} \left(\frac{1}{\sin \varrho} \right), \quad (142)$$

$$\Theta_{4,11,91}(s, t) = \frac{3\zeta^2 c_3 - 3\xi^2 c_3 \Lambda_1^2 - 2\eta_1 \Phi_2}{6\eta_1 \Phi_1} \mp \sqrt{\frac{2(\zeta^2 c_4 - \xi^2 c_4 \Lambda_1^2)}{\Phi_1}} (\csc \varrho \pm \cot \varrho), \quad (143)$$

$$\Theta_{4,12,92}(s, t) = \frac{3\zeta^2 c_3 - 3\xi^2 c_3 \Lambda_1^2 - 2\eta_1 \Phi_2}{6\eta_1 \Phi_1} \mp \sqrt{\frac{2(\zeta^2 c_4 - \xi^2 c_4 \Lambda_1^2)}{\Phi_1}} (\sec \varrho \pm \tan \varrho), \quad (144)$$

$$\Theta_{4,13,93}(s, t) = \frac{3\zeta^2 c_3 - 3\xi^2 c_3 \Lambda_1^2 - 2\eta_1 \Phi_2}{6\eta_1 \Phi_1} \mp \sqrt{\frac{2(\zeta^2 c_4 - \xi^2 c_4 \Lambda_1^2)}{\Phi_1}} (\csc \varrho \pm \csc \varrho), \quad (145)$$

$$\Theta_{4,14,94}(s, t) = \frac{3\zeta^2 c_3 - 3\xi^2 c_3 \Lambda_1^2 - 2\eta_1 \Phi_2}{6\eta_1 \Phi_1} \mp \sqrt{\frac{2(\zeta^2 c_4 - \xi^2 c_4 \Lambda_1^2)}{\Phi_1}} (\sin \varrho \pm \iota \cot \varrho), \quad (146)$$

where $\varrho = \xi s + \zeta t$.

Graphical findings and discussion

The visual representations of the D-CDNA model is examined in this section. When trying to construct the exact TW solutions, the visual behavior related to the D-CDNA model is developed using the MEFSEM for various parametric assumptions. To evaluate the structural design of D-CDNA model with manipulating the parameters set, the Mathematica 13.2 computer software is utilized. By adjusting the settings of the parameter, the graphs appearance of D-CDNA can be modified. Along with adding to the 3D plots and related 2D line graphs, we additionally provided contour plots for straightforward understanding. Distinct wave patterns can be developed via assigning distinct values for the parameters. By implementing MEFSEM into practice, numerous solutions,

such as complexiton, kink wave, dark or anti-bell, V, anti-Z and singular wave shapes soliton solutions can be encountered.

Figure 2 exhibits the 3D, 2D at multiple values of t and contour plots of the solution $\Theta_{1,1,1}(s, t)$ given in Eq. (53) while considering parametric values $\zeta = 4, \rho = 0.01, \Lambda_1 = 1, \xi = 1, \Phi_2 = 1, \Phi_1 = 1, \sigma = 1$ and $\mu = 1$, which displays a kink wave soliton. Figure 3 exhibits the 3D, 2D at multiple values of t and contour plots of the solution $\Theta_{1,1,2}(s, t)$ given in Eq. (54) while considering parametric values $\zeta = 0.5, \rho = 0.1, \Lambda_1 = 1, \xi = 0.01, \Phi_2 = 10, \Phi_1 = 10, \sigma = 0$ and $\mu = 1.5$, which displays a singular shape soliton. Figure 4 exhibits the 3D, 2D at multiple values of t and contour plots of the solution $\Theta_{1,1,8}(s, t)$ given in Eq. (60) while considering parametric values $\zeta = 1.5, \rho = 2, \Lambda_1 = 1, \xi = 1, \Phi_2 = 1, \Phi_1 = 5, \sigma = 1$ and $\mu = 3.5$, which displays a complexitons shape soliton. Figure 5 exhibits the 3D, 2D at multiple values of t and contour plots of the solution $\Theta_{1,1,10}(s, t)$ given in Eq. (62) while considering parametric values $\zeta = 3, \rho = 2, \Lambda_1 = 1, \xi = 1, \Phi_2 = 1, \Phi_1 = 5, \sigma = 1$ and $\mu = 3$, which displays a anti-Z shape soliton. Figure 6 exhibits the 3D, 2D at multiple values of t and contour plots of the solution $\Theta_{1,2,13}(s, t)$ given in Eq. (65) while considering parametric values $\zeta = 4, \rho = 10, \Lambda_1 = 2, \xi = 1, \Phi_2 = 1, \Phi_1 = 1, \sigma = 1$ and $\mu = 1$, which displays a anti-bell shape or dark soliton soliton. Figure 7 exhibits the 3D, 2D at multiple values of t and contour plots of the solution $\Theta_{1,2,15}(s, t)$ given in Eq. (67) while considering parametric values $\zeta = 3.5, \rho = 0.01, \Lambda_1 = 5, \xi = 1, \Phi_2 = 1, \Phi_1 = 1, \sigma = 1$ and $\mu = 3.5$, which displays a kink wave soliton. Figure 8 exhibits the 3D, 2D at multiple values of t and contour plots of the solution $\Theta_{2,1,25}(s, t)$ given in Eq. (77) while considering parametric values $\zeta = 5, \rho = 0.1, \Lambda_1 = 1, \xi = 2, \Phi_2 = 1, \Phi_1 = 10, \sigma = 1$ and $\mu = 10$, which displays a kink wave soliton. Figure 9 exhibits the 3D, 2D at multiple values of t and contour plots of the solution $\Theta_{2,1,32}(s, t)$ given in Eq. (84) while considering parametric values $\zeta = 5, \rho = 0.01, \Lambda_1 = 3.5, \xi = 2.5, \Phi_2 = 1, \Phi_1 = 5, \sigma = 1$ and $\mu = 3$, which displays a V-shape soliton.

Conclusion

In the present research, by utilizing the MEFSEM to find exact TW solutions to D-CDNA model, that, due to its biological basis, constitutes one of the many intriguing issues in contemporary biophysics. This technique frequently produces deeper, original, general solutions and accurate findings than rival methods, which is its main advantage. Several soliton solutions have been developed for a range of parametric factors. Using MEFSEM, a vast array of novel solutions have been created, such as the complexiton, kink wave, dark or anti-bell, V, anti-Z and




Figure 2. (a) A 3D plot of $\Theta_{1,1,1}(s, t)$ given in Eq. (53) is kink wave soliton, (b) analogous 2D line graphs for numerous values of t and (c) associated contour plot when $\zeta = 4, \rho = 0.01, \Lambda_1 = 1, \xi = 1, \Phi_2 = 1, \Phi_1 = 1, \sigma = 1$ and $\mu = 1$.




Figure 3. (a) A 3D plot of $\Theta_{1,1,2}(s, t)$ given in Eq. (54) is singular wave soliton, (b) analogous 2D line graphs for numerous values of t and (c) associated contour plot when $\zeta = 0.5, \rho = 0.1, \Lambda_1 = 1, \xi = 0.01, \Phi_2 = 10, \Phi_1 = 10, \sigma = 0$ and $\mu = 1.5$.




Figure 4. (a) A 3D plot of $\Theta_{1,1,8}(s, t)$ given in Eq. (60) is complexiton shape soliton, (b) analogous 2D line graphs for numerous values of t and (c) associated contour plot when $\zeta = 1.5, \rho = 2, \Lambda_1 = 1, \xi = 1, \Phi_2 = 1, \Phi_1 = 5, \sigma = 1$ and $\mu = 3.5$.




Figure 5. (a) A 3D plot of $\Theta_{1,1,10}(s, t)$ given in Eq. (62) is anti-Z shape soliton, (b) analogous 2D line graphs for numerous values of t and (c) associated contour plot when $\zeta = 3, \rho = 2, \Lambda_1 = 1, \xi = 1, \Phi_2 = 1, \Phi_1 = 5, \sigma = 1$ and $\mu = 3$.




Figure 6. (a) A 3D plot of $\Theta_{1,2,13}(s, t)$ given in Eq. (65) is anti-bell shape or dark soliton soliton, (b) analogous 2D line graphs for numerous values of t and (c) associated contour plot when $\zeta = 4, \rho = 10, \Lambda_1 = 2, \xi = 1, \Phi_2 = 1, \Phi_1 = 1, \sigma = 1$ and $\mu = 1$.

singular wave shapes soliton solutions. Numerous examples of soliton solutions can be seen visually. In pure and applied mathematics, soliton solution are widely used especially in disciplines like differential equations, algebraic and differential geometry, Lie groups, and Lie algebras. Soliton- admitting models possess extensive mathematical architecture and characteristics. An infinite number of conservation laws and related symmetries especially are closely linked to the integrability of the these models, became one of their fundamental characteristics. An further characteristic is the presence of a Hamiltonian or bi-Hamiltonian form, which enables the analysis and description of a system without adequately the need for explicit solution of the related models. In contrast to bright or bell shape solitons, dark or anti-bell shape solitons in fiber lasers exhibit greater stability and resistance




Figure 7. (a) A 3D plot of $\Theta_{1,2,15}(s, t)$ given in Eq. (67) is kink wave soliton, (b) analogous 2D line graphs for numerous values of t and (c) associated contour plot when $\zeta = 3.5, \rho = 0.01, \Lambda_1 = 5, \xi = 1, \Phi_2 = 1, \Phi_1 = 1, \sigma = 1$ and $\mu = 3.5$.




Figure 8. (a) A 3D plot of $\Theta_{2,1,25}(s, t)$ given in Eq. (77) is kink wave soliton, (b) analogous 2D line graphs for numerous values of t and (c) associated contour plot when $\zeta = 5, \rho = 0.1, \Lambda_1 = 1, \xi = 2, \Phi_2 = 1, \Phi_1 = 10, \sigma = 1$ and $\mu = 10$.




Figure 9. (a) A 3D plot of $\Theta_{2,1,32}(s, t)$ given in Eq. (84) is V-shape soliton, (b) analogous 2D line graphs for numerous values of t and (c) associated contour plot when $\zeta = 5, \rho = 0.01, \Lambda_1 = 3.5, \xi = 2.5, \Phi_2 = 1, \Phi_1 = 5, \sigma = 1$ and $\mu = 3$.

to distortion in noise-filled environments. Therefore, dark or anti-bell shape solitons are frequently employed in the disciplines of nonlinear optics, optic exchanges, and optic sensor. The kink soliton is a semi-local NL pattern that shows a steep curve in the spectrum across an unchanged bottom height and numerous fading oscillation tails, comparable to other soliton forms. Intensive short-pulse characteristics are influenced by the waveform structure because kink solitons cause self-steepness or NL impacts in NL fibers. Kink solitons have real-world applications as optical logic devices or as polarized switches among two distinct domains. Mathematica 13.2, a straightforward mathematical software, is employed to validate the reliability of the aforementioned soliton solutions findings. Additionally, there are 2D and 3D visualizations that show how the noticed soliton solutions

behave dynamically. Because the contour plots enable in comprehending the dynamical characteristics and trends presented by these soliton solutions, they also feature in this paper. The findings rendered it obvious that some of the soliton solutions listed are distinct and had never been seen before. Furthermore, the MEFSEM could potentially be utilized for NLPDEs in diverse areas, hence boosting its practicality as an instrument for further research endeavors. The results of the present research will provide ideas and impetus for future discussions in the NL sciences, particularly those with biological applications. The method's ease of use render it a significant addition for the study of NLPDEs and could unlock the door to further advances in this discipline. Since, there exists no single method for resolving the analytical solution of the NLPDE in the domain of integrable systems. A general symbolic computing approach for the analytical solution of a nonlinear partial differential equation is made possible by the implied neural network-based symbol calculation method, which also establishes the groundwork for a universal symbolic calculation method for analytical expressions. This might deliver some novel insights into how the model examined in this publication could be resolved in further research employing this method.

Data availability

All data generated or analyzed during this study are included in this manuscript.

Received: 15 December 2023; Accepted: 27 February 2024

Published online: 17 March 2024

References

- Zhang, J. L., Wang, M. L., Wang, Y. M. & Fang, Z. D. The improved F-expansion method and its applications. *Phys. Lett. A* **350**(1–2), 103–109 (2006).
- Younas, U. *et al.* A diversity of patterns to new $(3 + 1)$ -dimensional Hirota bilinear equation that models dynamics of waves in fluids. *Results Phys.* **54**, 107124 (2023).
- Younas, U., Sulaiman, T. A., Ismael, H. F., Shah, N. A. & Eldin, S. M. On the lump interaction phenomena to the conformable fractional $(2 + 1)$ -dimensional KdV equation. *Results Phys.* **52**, 106863 (2023).
- Mirzazadeh, M., Alqahtani, R. T. & Biswas, A. Optical soliton perturbation with quadratic-cubic nonlinearity by Riccati-Bernoulli sub-ODE method and Kudryashov's scheme. *Optik* **145**, 74–78 (2017).
- Drouiche, I., Harrouni, S. & Arab, A. H. A new approach for modelling the aging PV module upon experimental I-V curves by combining translation method and five-parameters model. *Electric Power Syst. Res.* **163**, 231–241 (2018).
- Ali, M. R., Khattab, M. A. & Mabrouk, S. M. Optical soliton solutions for the integrable Lakshmanan-Porsezian-Daniel equation via the inverse scattering transformation method with applications. *Optik* **272**, 170256 (2023).
- Zayed, E. M. E., Zedan, H. A. & Gepreel, K. A. On the solitary wave solutions for nonlinear Hirota-Satsuma coupled KdV of equations. *Chaos Solitons Fractals* **22**(2), 285–303 (2004).
- Wang, M., Li, X. & Zhang, J. The $(G'G)$ -expansion method and travelling wave solutions of nonlinear evolution equations in mathematical physics. *Phys. Lett. A* **372**(4), 417–423 (2008).
- Wazwaz, A. M. The tan h method: Solitons and periodic solutions for the Dodd-Bullough-Mikhailov and the Tzitzeica-Dodd-Bullough equations. *Chaos Solitons Fractals* **25**(1), 55–63 (2005).
- Wang, B. H., Wang, Y. Y., Dai, C. Q. & Chen, Y. X. Dynamical characteristic of analytical fractional solitons for the space-time fractional Fokas-Lenells equation. *Alex. Eng. J.* **59**(6), 4699–4707 (2020).
- Bekir, A. & Boz, A. Exact solutions for nonlinear evolution equations using Exp-function method. *Phys. Lett. A* **372**(10), 1619–1625 (2008).
- Wazwaz, A. M. A sine-cosine method for handling nonlinear wave equations. *Math. Comput. Model.* **40**(5–6), 499–508 (2004).
- Tipu, G. H. *et al.* The optical exact soliton solutions of Shynaray-IIA equation with Φ^6 -model expansion approach. *Opt. Quan. Electron.* **56**(2), 226 (2024).
- Faridi, W. A., Tipu, G. H., Myrzakulova, Z., Myrzakulov, R. & Akinyemi, L. Formation of optical soliton wave profiles of Shynaray-IIA equation via two improved techniques: A comparative study. *Opt. Q. Electron.* **56**(1), 132 (2024).
- Younas, U. & Ren, J. On the study of optical soliton molecules of Manakov model and stability analysis. *Int. Modern Phys. B* **36**(26), 2250180 (2022).
- Younas, U., Seadawy, A. R., Younis, M., Rizvi, S. T. & Althobaiti, S. Diverse wave propagation in shallow water waves with the Kadomtsev-Petviashvili-Benjamin-Bona-Mahony and Benney-Luke integrable models. *Open Phys.* **19**(1), 808–818 (2021).
- Shi, D. *et al.* Analytical study of the dynamics in the double-chain model of DNA. *Results Phys.* **52**, 106787 (2023).
- De-Xing, K., Sen-Yue, L. & Jin, Z. Nonlinear dynamics in a new double chain-model of DNA. *Commun. Theor. Phys.* **36**(6), 737 (2001).
- Abdou, M. A., Ouahid, L., Al Shahrani, J. S., Alanazi, M. M. & Kumar, S. New analytical solutions and efficient methodologies for DNA (Double-Chain Model) in mathematical biology. *Modern Phys. Lett. B* **36**(24), 2250124 (2022).
- Bilal, M., Younas, U. & Ren, J. Dynamics of exact soliton solutions in the double-chain model of deoxyribonucleic acid. *Math. Methods Appl. Sci.* **44**(17), 13357–13375 (2021).
- Forinash, K., Bishop, A. R. & Lomdahl, P. S. Nonlinear dynamics in a double-chain model of DNA. *Phys. Rev. B* **43**(13), 10743 (1991).
- Xian-Min, Q. I. A. N. & Sen-Yue, L. O. U. Exact solutions of nonlinear dynamics equation in a new double-chain model of DNA. *Commun. Theor. Phys.* **39**(4), 501 (2003).
- Seadawy, A. R. *et al.* Analytical mathematical approaches for the double-chain model of DNA by a novel computational technique. *Chaos Solitons Fractals* **144**, 110669 (2021).
- Okaly, J. B., Mvogo, A., Woulaché, R. L. & Kofané, T. C. Semi-discrete breather in a helicoidal DNA double chain-model. *Wave Motion* **82**, 1–15 (2018).
- Jian, H., Vologodskii, A. V. & Schlick, T. A combined wormlike-chain and bead model for dynamic simulations of long linear DNA. *J. Comput. Phys.* **136**(1), 168–179 (1997).
- Yao, S. W., Mabrouk, S. M., Inc, M. & Rashed, A. S. Analysis of double-chain deoxyribonucleic acid dynamical system in pandemic confrontation. *Results Phys.* **42**, 105966 (2022).
- Kumar, S., Kumar, A. & Kharbanda, H. Abundant exact closed-form solutions and solitonic structures for the double-chain deoxyribonucleic acid (DNA) model. *Braz. J. Phys.* **51**(4), 1043–1068 (2021).
- Iguchi, K. Tight-binding model for DNA double chains: Metal-insulator transition due to the formation of a double strand of DNA. *Int. J. Modern Phys. B* **11**(20), 2405–2423 (1997).
- Pollak, Y., Goldberg, S. & Amit, R. Self-avoiding wormlike chain model for double-stranded-DNA loop formation. *Phys. Rev. E* **90**(5), 052602 (2014).

30. Saleh, R., Mabrouk, S. M. & Wazwaz, A. M. Lie symmetry analysis of a stochastic gene evolution in double-chain deoxyribonucleic acid system. *Waves Random Complex Media* **32**(6), 2903–2917 (2022).
31. De-Xing, K., Sen-Yue, L. & Jin, Z. Nonlinear dynamics in a new double chain-model of DNA. *Commun. Theor. Phys.* **36**(6), 737 (2001).
32. Bilal, M., Younas, U. & Ren, J. Dynamics of exact soliton solutions in the double-chain model of deoxyribonucleic acid. *Math. Methods Appl. Sci.* **44**(17), 13357–13375 (2021).
33. Kumar, S., Kumar, A. & Kharbanda, H. Abundant exact closed-form solutions and solitonic structures for the double-chain deoxyribonucleic acid (DNA) model. *Braz. J. Phy.* **51**(4), 1043–1068 (2021).
34. Zhang, R. F. & Li, M. C. Bilinear residual network method for solving the exactly explicit solutions of nonlinear evolution equations. *Nonlinear Dyn.* **108**(1), 521–531 (2022).
35. Zhang, R. F. & Bilige, S. Bilinear neural network method to obtain the exact analytical solutions of nonlinear partial differential equations and its application to p-gBKP equation. *Nonlinear Dyn.* **95**, 3041–3048 (2019).
36. Zhang, R. F., Li, M. C. & Yin, H. M. Rogue wave solutions and the bright and dark solitons of the (3 + 1)-dimensional Jimbo-Miwa equation. *Nonlinear Dyn.* **103**, 1071–1079 (2021).
37. Zhang, R., Bilige, S. & Chaolu, T. Fractal solitons, arbitrary function solutions, exact periodic wave and breathers for a nonlinear partial differential equation by using bilinear neural network method. *J. Syst. Sci. Complex.* **34**, 122–139 (2021).

Acknowledgements

The authors are grateful to anonymous referees for their valuable suggestions, which significantly improved this manuscript.

Author contributions

Conceptualization, M. Z. Y., M. A. and A. B.; Formal analysis, M.A. and T. N.; Investigation, M. Z. Y., M. A., A. B., and H. E.; Methodology, M. Z. Y., M.A. and T. N.; Project administration, M.A., F.A.A. and T. N.; Software, M. Z. Y., M.A., F.A.A., and H. E.; Supervision, M.A. and T. N.; Validation, M.A.; Visualization, M. Z. Y., M. A., T. N., F. A. A., A. B., and H. E.; Writing—original draft, M. Z. Y., M. A., T. N., F. A. A., and A. B.; Writing—review and editing, M. Z. Y., M. A., T. N., F. A. A., A. B., and H. E.; All authors have read and agreed to the published version of the manuscript.

Competing interests

The authors declare no competing interests.

Additional information

Correspondence and requests for materials should be addressed to A.B.

Reprints and permissions information is available at www.nature.com/reprints.

Publisher's note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

© The Author(s) 2024