



OPEN

SUBJECT AREAS:
ANTIBIOTICS
BACTERIAL GENETICSReceived
30 November 2014Accepted
3 February 2015Published
4 March 2015Correspondence and
requests for materials
should be addressed to
H.T. (tanhr@im.ac.cn)
or G.N. (niugq@im.ac.
cn)* These authors
contributed equally to
this work.

Genome engineering and direct cloning of antibiotic gene clusters via phage ϕ BT1 integrase-mediated site-specific recombination in *Streptomyces*

Deyao Du^{1,2*}, Lu Wang^{1,3*}, Yuqing Tian¹, Hao Liu³, Huarong Tan¹ & Guoqing Niu¹¹State Key Laboratory of Microbial Resources, Institute of Microbiology, Chinese Academy of Sciences, Beijing, China, ²University of Chinese Academy of Sciences, Beijing, China, ³Key Laboratory of Industrial Fermentation Microbiology, Ministry of Education, College of Biotechnology, Tianjin University of Science and Technology, Tianjin, China.

Several strategies have been used to clone large DNA fragments directly from bacterial genome. Most of these approaches are based on different site-specific recombination systems consisting of a specialized recombinase and its target sites. In this study, a novel strategy based on phage ϕ BT1 integrase-mediated site-specific recombination was developed, and used for simultaneous *Streptomyces* genome engineering and cloning of antibiotic gene clusters. This method has been proved successful for the cloning of actinorhodin gene cluster from *Streptomyces coelicolor* M145, napsamycin gene cluster and daptomycin gene cluster from *Streptomyces roseosporus* NRRL 15998 at a frequency higher than 80%. Furthermore, the system could be used to increase the titer of antibiotics as we demonstrated with actinorhodin and daptomycin, and it will be broadly applicable in many *Streptomyces*.

Streptomyces are high-GC Gram-positive bacteria well known for their ability to produce a wide variety of medically and agriculturally useful antibiotics and related compounds¹. Genes responsible for the biosynthesis of a specific secondary metabolite are usually arranged in clusters that vary in size from a few to over 100 kb². To gain insight into the biosynthesis and regulation of antibiotics in *Streptomyces*, it is of great importance to clone their gene clusters. Recently, various approaches have been developed to clone gene clusters directly from bacterial genomic DNA. These methods include RecET-mediated linear-plus-linear homologous recombination (LLHR)³, *oriT*-directed capture system⁴ and transformation-associated recombination (TAR)⁵. The RecET-mediated LLHR was successful in cloning gene clusters (10 to 52 kb in length) from the genome of *Photorehabdus luminescens* into expression vectors in *Escherichia coli*³. The *oriT*-directed capture system has been used to clone regions up to 140 kb from the genome of *Burkholderia pseudomallei*⁶ and 200 kb from megaplasmid of *Sinorhizobium meliloti*⁴. However, the use of this system was limited to Gram-negative bacteria that can be established as conjugation donors⁶. Taking advantage of the natural *in vivo* homologous recombination of *Saccharomyces cerevisiae*, TAR cloning strategy was used to capture a 21.3 kb enterocin gene cluster from *Salinispora pacifica* CNT-150⁷ and a 67 kb taromycin A biosynthetic gene cluster from *Saccharomonospora* sp. CNQ-490⁸.

The ability to delete large genomic fragments within *Streptomyces* genome is of great interest for genetic manipulations of *Streptomyces*. Several strategies have been developed for a number of bacteria. Some methods are based on the meganuclease I-SceI system which involves the meganuclease I-SceI of *Saccharomyces cerevisiae* and its 18 bp recognition sequence^{9,10}. Many of them are based on site-specific recombination systems consisting of a specialized recombinase and its target sites. Nearly all site-specific recombinases fall into two families, the tyrosine recombinases and the serine recombinases¹¹. The recombination systems of the tyrosine recombinase family include Cre/loxP from the P1 phage¹², Dre/rox from the P1-like transducing phage D6¹³ and the Flp/FRT from yeast¹⁴. The Cre, Dre and Flp proteins are the tyrosine recombinases which catalyze reciprocal site-specific recombination of DNA at loxP, rox and FRT sites, respectively. Integrases (Int) from *Streptomyces* temperate phage ϕ C31 and ϕ BT1 belong to serine recombinase family. They catalyze site-specific recombination of the phage attachment site (*attP*) with the bacterial attachment site (*attB*), resulting in the formation of two hybrid sites (*attL* and *attR*)^{15,16}. Both ϕ C31 and ϕ BT1 *attP-int* loci have been used to construct versatile vectors which can




Figure 1 | Schematic diagram of antibiotic gene cluster cloning from *Streptomyces* chromosome. Initially, a pUC119-based suicide plasmid (*pSV::attB₆Up*) carrying *attB₆* and a region homologous to 5' end of the cluster is introduced into the chromosome by a single crossover. A second plasmid *pKC1139::attP₆Dn* is based on *pKC1139* carrying *attP₆* and a region homologous to 3' end of the cluster. When the incubation temperature is higher than 34°C, *pKC1139::attP₆Dn* turns into a non-replicating plasmid and then is integrated into the chromosome by a single crossover. Expression of ϕ BT1 integrase (encoded in the plasmid pIJ10500) leads to excision of the *pKC1139* backbone with gene cluster of interest, leaving behind the suicide vector *pUC119::neo* and 42 bp *attL₆* site. *aac(3)IV*: apramycin resistance gene; *neo*: kanamycin resistance gene; *ori*: temperature-sensitive origin of replication from pSG5; *rep*: *rep* encoding a replication initiator protein from pSG5.

integrate into different *attB* sites in *Streptomyces*^{15,16}. To increase the diversity of *attP-attB* pair of ϕ BT1, 15 mutated *attP-attB* pairs (*attP₀₁-attB₀₁* → *attP₁₅-attB₁₅*) were generated by PCR mutagenesis of the central dinucleotide sequence of *attB* and *attP*¹⁷. The Cre/loxP system was successfully used for the deletion of large fragments in *Magnetospirillum gryphiswaldense* and several *Streptomyces* species^{18–20}. However, the use of ϕ C31 and ϕ BT1 integrase in this aspect has not been exploited.

We devised a novel strategy for *Streptomyces* genome engineering and cloning of antibiotic gene clusters. This method is based on phage ϕ BT1 *attP-attB-int* system and requires two single crossovers for targeted integration of mutated *attB* and *attP* into the recipient chromosome. Using the system, we easily cloned 25 kb fragment containing actinorhodin (*act*) gene cluster from *S. coelicolor* M145, 45 kb fragment containing napsamycin (*nap*) gene cluster and 157 kb fragment containing daptomycin (*dap*) gene cluster from *S. roseosporus* NRRL 15998. In addition, this method could be used to improve the titer of antibiotics by increasing copy numbers of antibiotic gene clusters.

Results

Construction of pUC119- and pKC1139-based plasmids. Our strategy used in this study requires both homologous and site-specific recombinations. The homologous recombinations were used for targeted integration of the mutated *attB* and *attP* into *Streptomyces* chromosome, while the ϕ BT1 integrase-mediated site-specific recombination was employed to excise targeted region of interest from the chromosome (Fig. 1). The mutated *attB* and *attP* sites were chosen to avoid site specific recombination with the

endogenous *attB* site in *Streptomyces* genome and consequently undesirable DNA rearrangements. Sites of *attB₆* and *attP₆* were randomly chosen from the 15 mutated *attP-attB* pairs. For the integration of *attB₆* into *Streptomyces* chromosome, pUC119-based suicide plasmids (*pSV::attB₆-act*, *pSV::attB₆-nap* and *pSV::attB₆-dap*) were constructed. These plasmids are derivatives of pUC119 containing the kanamycin-resistance gene (*neo*), the origin of transfer (*oriT*) from plasmid RK2 (for the intergeneric conjugation between *E. coli* and *Streptomyces*), *attB₆* and a 2.0 kb homologous region flanking 5' end of the targeted regions (Fig. S1a). We also constructed pKC1139-based plasmids (*pKC1139::attP₆-act*, *pKC1139::attP₆-nap* and *pKC1139::attP₆-dap*) for the integration of *attP₆* into *Streptomyces* chromosome. These plasmids are derivatives of pKC1139 containing *attP₆* and a 2.0 kb homologous region flanking 3' end of the targeted regions (Fig. S1b).

Cloning of *act* gene cluster from *S. coelicolor* M145. To test this strategy, we first chose to clone the well-studied *act* gene cluster (SCO5070-SCO5092) from *S. coelicolor* M145. For this purpose, *pSV::attB₆-act* and *pKC1139::attP₆-act* were introduced into the recipient chromosomes via single-crossover homologous recombination to obtain double-co-integrate strain Sco-actB₆P₆ (Fig. 1). Further introduction of pIJ10500 (an integrative plasmid containing the ϕ BT1 integrase gene) into Sco-actB₆P₆ allowed subsequent excision of 23 kb *act* gene cluster from *S. coelicolor* M145, leaving behind the suicide vector pUC119::*neo*, a scar of 42 bp *attL* site and pIJ10500 integrated within SCO4848. Excision of the gene cluster was confirmed by PCR analysis using both genomic and plasmid DNA as templates (Fig. 2). For 9 out of 10




Figure 2 | Confirmation of the excision events by PCR amplifications. (A) The schematic diagram showing the position of primers in the chromosome of double-cointegrate strains. (B) Agarose gel electrophoresis showing PCR amplified fragments. PCR templates in the upper panels are genomic DNAs from *S. coelicolor* M145 or *S. roseosporus* NRRL 15998 (G) and ten randomly selected double-cointegrate strains with pIJ10500 (M145-MCact, Sro-MCnap or Sro-MCdap), while PCR templates in the lower panels are plasmid DNAs including pKC1139 (P) and ten different clones of pKC1139::act, pKC1139::nap and pKC1139::dap. The primers used and the expected size of amplification fragments were indicated.

exconjugates tested, the ϕ BT1 integrase-mediated excision of *act* gene cluster occurred at a frequency of 90%. Furthermore, the presence of *attL* and *attR* in the amplified fragments was confirmed by DNA sequencing (Fig. 3). To recover the plasmid containing the entire *act* gene cluster (pKC1139::act) from *Streptomyces*, the DNA extract containing pKC1139::act from M145-MCact was used to transform *E. coli* Top10. Plasmid DNA from four apramycin resistant *E. coli* colonies was confirmed by BamHI and NotI digestion, respectively. The restriction fragments showed correct band patterns (Fig. S2a).

To verify the cluster is complete, the recombinant plasmid pKC1139::act was introduced into *S. coelicolor* M1146 (M1146) to obtain M1146-MCact. Unlike M1146 and M1146-pKC1139 (*S. coelicolor* M1146 containing empty vector pKC1139), M1146-MCact regained the ability to produce the blue pigment actinorhodin (Fig. 4a). These results showed that the cloned *act* gene cluster was complete and functional.

Deletion of *act* gene cluster from *S. coelicolor* M145. To delete the *act* gene cluster from *S. coelicolor* M145, a single colony of M145-MCact was randomly chosen for the removal of pKC1139::act. After three rounds of nonselective growth at 28°C and subsequent cultivation at 40°C, approximately 5% of M145-MCact colonies lost pKC1139::act. Strains lacking the *act* gene cluster (M145-Dact) were first confirmed by PCR (data not shown), and then patched on R5MS solid agar plate for visual comparison of actinorhodin production. Unlike *S. coelicolor* M145 that could produce both blue pigment actinorhodin and red pigment undecylprodigiosins, M145-Dact could only produce the red pigment undecylprodigiosins (Fig. 4a). This was further validated by no actinorhodin production of M145-Dact in R5MS liquid culture (Fig. 4b).

Cloning and deletion of *nap* and *dap* gene cluster from *S. roseosporus* NRRL 15998. To clone gene cluster of medium and large sizes, we used the same strategy to clone *nap* and *dap* gene cluster from *S. roseosporus* NRRL 15998. Excision of *nap* gene cluster from *S. roseosporus* NRRL 15998 occurred in 9 out of 10 exconjugates, and excision of *dap* gene cluster from *S. roseosporus* NRRL 15998 occurred in 8 out of 10 exconjugates (Fig. 2). Like pKC1139::act, plasmid containing *nap* gene cluster (pKC1139::nap) was passed through *E. coli* Top10 and isolated plasmid DNA was confirmed by BglII and EcoRI digestion, respectively (Fig. S2b). The cloned fragment covers a contiguous DNA region of 45 kb from SSGG02973 to SSGG03009. For pKC1139::dap, the plasmid was isolated directly from *Streptomyces* and confirmed with restriction digestion (Fig. S2c). The 157 kb fragment covering SSGG00215-SSGG00287 contains the complete *dap* gene cluster. Similar to that of *act* gene cluster, the removal of pKC1139::nap and pKC1139::dap from Sro-MCnap and Sro-MCdap generated strains lacking *nap* and *dap* gene clusters (Sro-Dnap and Sro-Ddap).

Improvement of antibiotic titers. The pKC1139 contains a temperature-sensitive origin of replication from pSG5, which is a medium copy plasmid with an approximate 20–50 copy numbers per chromosome²¹. When cultured at 28°C, pKC1139 exists as autonomous plasmid in *Streptomyces*. In *S. coelicolor* M145, there is only one copy of *act* gene cluster in the chromosome. After the ϕ BT1 integrase-mediated excision, the *act* gene cluster was transferred into pKC1139. An increase in copy number of *act* gene cluster will improve actinorhodin production. This was confirmed both on R5MS agar plate (Fig. 4a) and in R5MS liquid culture (Fig. 4b). It should be noted that the titer of actinorhodin in M1146-MCact was even higher than that of M145-MCact. Similarly, daptomycin titer




Figure 3 | Representative excision of the *act* gene cluster from *S. coelicolor* M145. (A) Nucleotide sequence of $attB_6$, $attP_6$, $attL_6$ and $attR_6$. The mutated core dinucleotide (CT) at which the crossover occurs is in bold. (B) Verification of $attL_6$ and $attR_6$ by DNA sequencing. Sequences of $attL_6$ and $attR_6$ from DNA sequencing are underlined.

could also increase after the excision of *dap* gene cluster from its chromosome location in *S. roseosporus* NRRL 15998. Cultures of Sro-MCdap and *S. roseosporus* NRRL 15998 were subjected to bioassay against *S. aureus*, the results showed that Sro-MCdap exhibited bigger inhibition zones against *S. aureus* than *S. roseosporus* NRRL 15998 at time intervals from 2–5 days (Fig. 5a). This was further verified by comparison of daptomycin from fermentation broth of *S. roseosporus* NRRL 15998 and Sro-MCdap by high-performance liquid chromatography (HPLC) analysis (Fig. 5b). In addition, we noticed that existence of extra copy numbers of antibiotic gene clusters caused a slowdown in growth of *Streptomyces*. When cultured on AS-1 agar medium, growth of Sro-MCnap and Sro-MCdap are severely impaired, especially at earlier stages of cultivation (Fig. S3). This phenotype was most likely attributed to the metabolic burden of extra copy numbers of antibiotic gene clusters. This assumption is based on the observation that growth of Sro-Dnap (devoid of *nap* gene cluster) and Sro-Ddap (devoid of *dap* gene cluster) are converted back to that of *S. roseosporus* NRRL 15998 (Fig. S3).

To examine the stability of multiple copy plasmids in *Streptomyces*, two randomly chosen strains of Sro-MCdap were passed consecutively for five or ten times on AS-1 plates supplemented with or without apramycin. Biological activities of these stains (G_5 and G_{10}) were compared with that of the original Sro-MCdap (G_0). All Sro-MCdap strains exhibited similar inhibitory activity against *S. aureus* (Fig. S4), suggesting that pKC1139-derived large plasmids are stable in the engineered *Streptomyces* in the presence or absence of selective pressure.

Discussion

We have established an efficient method for genome engineering and direct cloning of gene clusters in *Streptomyces*. The strategy is based on phage ϕ BT1 *attP-attB-int* system and provides several advantages over similar methods. First, it can be used for the deletion of large fragment (up to 157 kb) from *Streptomyces* genome. In the meantime, the large fragment containing gene cluster of interest was cloned into pKC1139 simultaneously. Another advantage of our strategy is that it could clone gene cluster in size up to 157 kb. This is the largest size ever reported in Gram positive bacteria and should be good enough for most antibiotic gene clusters. Last, our strategy can be used to improve the titer of industrial important antibiotics by creating strains with extra copy numbers of antibiotic biosynthetic gene clusters.

The ϕ BT1 *attP-attB-int* system is helpful for genetic modifications of *Streptomyces* genome at multiple sites. In addition to the intact *attP-attB* pair, there are 15 mutated *attP-attB* pairs which can be recognized by ϕ BT1 integrase¹⁷. Multiple rounds of large fragment deletion can be achieved with the following modifications. (1) Relocation of the $attB_6$ sequence (or any other mutated *attB*) to the downstream of the 2.0 kb homologous fragment in pSV:: $attB_6$ Up. This change will allow the excision of the pUC119::neo backbone from *Streptomyces* genome together with pKC1139. (2) Construction of an autonomous helper plasmid containing a temperature-sensitive origin of replication from pSG5, the origin of transfer (*oriT*) from plasmid RK2 and ϕ BT1 integrase gene. This plasmid can ensure the high efficient excision of large fragment from *Streptomyces* genome and subsequent removal of ϕ BT1 inte-




Figure 4 | Comparison of actinorhodin production in *S. coelicolor* M145 and its derivatives. (A) Comparison of actinorhodin production (blue pigment) of *S. coelicolor* M145 and its derivatives. Photograph was taken from the bottom of the plate after grown on R5MS agar medium for 4 days at 28°C. Representative image of three independent experiments with similar results was shown. (B) Actinorhodin titers of *S. coelicolor* M145 and its derivatives grown in 50 ml of R5MS at 28°C. Error bars show standard deviations.

grase. With these modifications, there is only 42 bp *attL* site left in the chromosome of *Streptomyces*.

Genome analysis suggested that *S. roseosporus* NRRL 15998 has potential capacity to produce napsamycins^{22,23}. However, the production of napsamycins in *S. roseosporus* NRRL 15998 has not been reported. With this strategy, we cloned *nap* gene cluster in pKC1139 to generate pKC1139::*nap*. It can be manipulated extensively in *E. coli*. These manipulations include replacement of vector backbone with integrative plasmid and deletion or constitutive expression of regulatory gene by PCR targeting²⁴. The modified gene cluster can be transferred into heterologous hosts for expression. It can also be transferred back into the mutant devoid of *nap* gene cluster after removal of pKC1139::*nap*. Detection of napsamycin in these strains will shed light on the activation of cryptic gene clusters in *Streptomyces*.

In some industrial overproducing strains generated by traditional mutagenesis, amplification of biosynthetic gene cluster has been observed^{25–27}. Based on these observations, controlled amplification of gene cluster was used to increase the productivity of commercially




Figure 5 | Analysis of daptomycin production in *S. roseosporus* NRRL 15998 (WT) and Sro-MCdap. (A) Bioassay of daptomycin against *S. aureus*. After grown on AS-1 agar for 2–5 days at 28°C, the patches of WT and Sro-MCdap were overlaid with cultures of *S. aureus* and the zone of inhibition was assessed after overnight incubation at 37°C. Representative images of three independent experiments with similar results are shown. (B) HPLC analysis of fermentation filtrates from WT and Sro-MCdap after incubation for 4 days. Components of daptomycin were indicated by comparison with standards.

important antibiotics. Integration of an additional copy of gene cluster for nikkomycin and gougerotin biosynthesis led to an increased production of nikkomycin and gougerotin by *Streptomyces ansochromogene*²⁸ and *Streptomyces gramineus*²⁹, respectively. The *zouA*-mediated gene amplification of *act* gene cluster in *S. coelicolor* M145 led to a 20-fold increase in actinorhodin production³⁰. The *zouA* encodes a site-specific relaxase similar to TraA protein which catalyzes RecA-independent site-specific recombination. The recombination sites of ZouA are *oriT*-like RsA and RsB³⁰. In this study, we reported the amplification of gene clusters mediated by phage ϕ BT1 integrase and improved antibiotic titers in the engineered *Streptomyces* strains. We believe that the system described here could be used readily to increase antibiotic titers in many *Streptomyces* and possible other actinomycetes.

Methods

Bacterial strains, plasmids, primers and growth conditions. Bacterial strains and plasmids used in this study are listed in Table 1, and primers are listed in Table S1. *S. coelicolor* M145 and *S. roseosporus* NRRL 15998 were used for cloning of *act*, *nap* and *dap* gene clusters. *S. coelicolor* M1146 is an engineered derivative of *S. coelicolor* M145 that lacks gene clusters for actinorhodin (ACT), undecylprodigiosins (RED), cryptic polyketide (CPK) and calcium-dependent antibiotic (CDA) biosynthesis³¹. *Staphylococcus aureus* was used as an indicator strain for daptomycin bioassay. *E. coli* Top10 was used as a general host for propagating plasmids. *E. coli* ET12567 (pUZ8002) was used as a host for transferring DNA from *E. coli* to *Streptomyces* by intergeneric conjugation³².

For general purpose, *S. coelicolor* M145 and its derivatives were grown on mannitol soya flour medium (MS) agar or in yeast extract-malt extract (YEME) liquid med-



Table 1 | Strains and plasmids used in this study

Strains/plasmids	Genotype/description	Reference/source
<i>E. coli</i>		
Top10	F- <i>mcrA</i> Δ(<i>mrr-hsdRMS-mcrBC</i>) φ80(<i>lacZ</i> ΔM15 Δ <i>lacX74</i> <i>recA1 araD139</i> Δ(<i>ara</i> <i>leu</i>)7697 <i>galU galK rpsL</i> (StrR) <i>endA1 nupG</i>	Invitrogen
ET12567	F- <i>dam-13::Tn9 dcm-6 hsdM hsdR zjj-202::Tn10 recF143 galK2 galT22 ara-14 lacY1 xyl-5 leuB6 thi-1 tonA31 rpsL136 hisG4 tsx-78 mtl-1 glnV44</i>	40
<i>Staphylococcus</i>		
<i>S. aureus</i>	A indicator strain	41
<i>Streptomyces</i>		
<i>S. coelicolor</i> M1145	Prototrophic; SCP1 - SCP2 - Pgl ⁺	32
<i>S. coelicolor</i> M1146	Δ <i>act</i> Δ <i>red</i> Δ <i>cpk</i> Δ <i>acdA</i>	31
<i>Sco-actB₆P₆</i>	A derivative of <i>S. coelicolor</i> M1145 with <i>attB₆</i> and <i>attP₆</i> flanking <i>act</i> gene cluster	This study
<i>S. roseosporus</i> NRRL 15998	A daptomycin-producing strain	Broad Institute
Sro-pKC1139	A derivative of <i>S. roseosporus</i> NRRL 15998 containing pKC1139	This study
Sro-napB ₆ P ₆	A derivative of <i>S. roseosporus</i> NRRL 15998 with <i>attB₆</i> and <i>attP₆</i> flanking <i>nap</i> gene cluster	This study
Sro-dapB ₆ P ₆	A derivative of <i>S. roseosporus</i> NRRL 15998 with <i>attB₆</i> and <i>attP₆</i> flanking <i>dap</i> gene cluster	This study
M1145-Dact	Δ <i>act</i>	This study
M145-MCact	A derivative of <i>S. coelicolor</i> M1145 containing multicopy of <i>act</i> gene cluster	This study
M1146-pKC1139	<i>S. coelicolor</i> M1146 containing pKC1139	This study
M1146-MCact	<i>S. coelicolor</i> M1146 containing pKC1139:: <i>act</i>	This study
Sro-MCnap	A derivative of <i>S. roseosporus</i> NRRL 15998 containing multicopy of <i>nap</i> gene cluster	This study
Sro-MCdap	A derivative of <i>S. roseosporus</i> NRRL 15998 containing multicopy of <i>dap</i> gene cluster	This study
Plasmids		
pUZ8002	<i>Ira neo</i> RP4	42
pUC119::neo	pUC119 containing kanamycin resistance gene (<i>neo</i>)	43
pKC1139	<i>E. coli</i> : <i>Streptomyces</i> shuttle plasmid contains a <i>Streptomyces</i> temperature-sensitive origin of replication	15
pJ10500	A derivative of pMS82 containing φBT1 integrase gene	36
pUC119::neo-attB ₆ -act	A derivation of pUC119::neo containing <i>attB₆</i> and 2.0 kb homologous region flanking the 5' end of <i>act</i> gene cluster	This study
pSV::attB ₆ -act	A derivation of pUC119::neo containing the origin of transfer (<i>oriT</i>) from plasmid RK2, <i>attB₆</i> and 2.0 kb homologous region flanking the 5' end of <i>act</i> gene cluster	This study
pSV::attB ₆ -nap	A derivation of pUC119::neo containing the origin of transfer (<i>oriT</i>) from plasmid RK2, <i>attB₆</i> and 2.0 kb homologous region flanking the 5' end of <i>nap</i> gene cluster	This study
pSV::attB ₆ -dap	A derivation of pUC119::neo containing the origin of transfer (<i>oriT</i>) from plasmid RK2, <i>attB₆</i> and 2.0 kb homologous region flanking the 5' end of <i>dap</i> gene cluster	This study
pKC1139::attP ₆ -act	A derivation of pKC1139 containing <i>attP₆</i> and 2.0 kb homologous region flanking the 3' end of <i>act</i> gene cluster	This study
pKC1139::attP ₆ -nap	A derivation of pKC1139 containing <i>attP₆</i> and 2.0 kb homologous region flanking the 3' end of <i>nap</i> gene cluster	This study
pKC1139::attP ₆ -dap	A derivation of pKC1139 containing <i>attP₆</i> and 2.0 kb homologous region flanking the 3' end of <i>dap</i> gene cluster	This study
pKC1139::act	A derivation of pKC1139 containing 25 kb fragment including actinorhodin gene cluster and its flanking sequences	This study
pKC1139::nap	A derivation of pKC1139 containing 45 kb fragment including napsamycin gene cluster and its flanking sequences	This study
pKC1139::dap	A derivation of pKC1139 containing 157 kb fragment including daptomycin gene cluster and its flanking sequences	This study



ium³². For actinorhodin production, *S. coelicolor* M145 and its derivatives were grown on R5MS agar or in R5MS liquid medium³³. *S. roseosporus* NRRL 15998 was cultured on AS-1 agar medium or in tryptic soy broth (TSB) liquid medium³⁴. All *Streptomyces* strains were maintained at 28 °C unless specified otherwise. General approaches for *E. coli* or *Streptomyces* manipulations were performed according to standard protocols^{32,35}. When necessary, the final antibiotic concentrations used for selection of *E. coli* transformants were as follows: ampicillin, 100 µg ml⁻¹; apramycin, 100 µg ml⁻¹; kanamycin, 100 µg ml⁻¹; chloramphenicol, 12.5 µg ml⁻¹. For selection of *Streptomyces* transformants, the final antibiotic concentrations were, kanamycin, 50 µg ml⁻¹ in MS for *S. coelicolor* and 20 µg ml⁻¹ in AS-1 for *S. roseosporus*; apramycin, 50 µg ml⁻¹ in MS for *S. coelicolor* and 10 µg ml⁻¹ in AS-1 for *S. roseosporus*; hygromycin, 50 µg ml⁻¹ in MS or AS-1 for *Streptomyces*; nalidixic acid, 25 µg ml⁻¹ in MS or AS-1 for *Streptomyces*.

Construction of plasmids. Of the 15 mutated *attP-attB* pairs¹⁷, *attP₆-attB₆* was randomly chosen for this experiment. The sequences of *attB₆* and *attP₆* were obtained by overlapping PCR. For construction of pSV::*attB₆-act*, a 2.0 kb fragment flanking 5' end of the *act* gene cluster was amplified from genomic DNA of *S. coelicolor* M145 with primer pair act-Up F/act-Up R. The amplicon was diluted 1:100 and used as templates for the second round of PCR with primer pair attB₆-in F/act-Up R. The product from the second amplification reaction was diluted again and underwent a third run with primer pair attB₆-out F/act-Up R. The final product was digested with HindIII/BamHI and then inserted into the corresponding sites of pUC119::*neo* to generate pUC119::*neo-attB₆-act*. The origin of transfer (*oriT*) from plasmid RK2 was amplified from pKC1139 with primer pair oriT F and oriT R, subsequently digested with EcoRI and inserted into the EcoRI site of pUC119::*neo-attB₆-act* to generate pSV::*attB₆-act* (Fig. S1a). For construction of pKC1139::*attP₆-act*, a 2.0 kb fragment flanking 3' end of the *act* gene cluster was amplified from genomic DNA of *S. coelicolor* M145 with primer pair act-Dn F/act-Dn R. The amplicon with 1:100 dilution served as templates for the second round of PCR with primer pair attP₆ F/act-Dn R. The final product was digested with HindIII/EcoRI and then inserted into the corresponding sites of pKC1139 to generate pKC1139::*attP₆-act* (Fig. S1b).

For construction of pSV::*attB₆-nap* and pSV::*attB₆-dap*, a 2.0 kb fragment flanking 5' end of the *nap* and *dap* gene cluster was amplified from genomic DNA of *S. roseosporus* NRRL 15998 with primer pairs nap-Up F/nap-Up R and dap-UpF/dap-Up R, respectively. The product was digested with XbaI/BamHI and used to replace the 2.0 kb fragment upstream of the *act* gene cluster in pSV::*attB₆-act*. For construction of pKC1139::*attP₆-nap* and pKC1139::*attP₆-dap*, a 2.0 kb fragment flanking 3' end of the *nap* and *dap* gene cluster was amplified from genomic DNA of *S. roseosporus* NRRL 15998 using primer pairs nap-Dn F/nap-Dn R and dap-Dn F/dap-Dn R, respectively. The product was digested with BamHI/EcoRI and used to replace the 2.0 kb fragment downstream of the *act* gene cluster in pKC1139::*attP₆-act*. To ensure the authenticity of DNA sequences, all PCR products were verified by sequencing.

Construction of double-cointegrate strains. To insert *attB₆* and *attP₆* at sites flanking the *act* gene cluster of *S. coelicolor*, pSV::*attB₆-act* and pKC1139::*attP₆-act* were conjugated into *S. coelicolor* M145. The pSV::*attB₆-act* is unable to replicate alone in *Streptomyces* and selection with kanamycin allows to select exconjugants in which pSV::*attB₆-act* is inserted into the *S. coelicolor* genome. The pKC1139::*attP₆-act* is a derivative of the *E. coli-Streptomyces* shuttle vector pKC1139 that contains a *Streptomyces* temperature-sensitive origin of replication from pSG5¹⁵. When the incubation temperature is higher than 34 °C, pKC1139::*attP₆-act* turns into non-replicating plasmid and *attP₆* was then inserted into *S. coelicolor* genome with selection of apramycin to obtain double-cointegrate strain Sco-actB₆P₆. Similar strategy was used for the construction of double-cointegrate strains Sro-*nap*B₆P₆ and Sro-*dap*B₆P₆.

Excision of targeted regions. The integrative plasmid pIJ10500³⁶ is a derivative of pMS82 which contains the phage φBT1 integrase gene and integrates intragenically into SCO4848 encoding a putative integral membrane protein¹⁶. It was conjugated into a randomly selected strain Sco-actB₆P₆, Sro-*nap*B₆P₆ and Sro-*dap*B₆P₆, respectively. The exconjugants were initially selected with hygromycin and ten randomly chosen exconjugants were passed twice on MS or AS-1 plates supplemented with kanamycin and apramycin, and subject to genomic and plasmid extraction. Excision of targeted region from *Streptomyces* genome was analyzed by PCR amplifications using genomic DNA templates and primer pairs B₆-VF/actDn-VR, B₆-VF/napDn-VR and B₆-VF/dapDn-VR. In the meantime, PCR amplifications were performed with plasmid DNA template by using primer pairs P₆-VF/actUp-VR, B₆-VF/napUp-VR and B₆-VF/dapUp-VR. Strains with excision of the targeted regions were designated as M145-MCact, Sro-MCnap and Sro-MCdap, respectively.

Deletion of the antibiotic gene clusters from *Streptomyces*. A single colony of M145-MCact was randomly chosen and passed three times on nonselective MS plates at 28 °C. Spores were harvested, serially diluted, and then spread on MS agar. After growing for 4 days at 40 °C, colonies were replicated on MS agar plates containing kanamycin or apramycin. Strains lacking the *act* gene cluster (M145-Dact) are apramycin sensitive (Apr^r) and kanamycin resistant (Kan^r). Apr^r and Kan^r strains were further verified by PCR. In a similar way, strains lacking the *nap* and *dap* gene clusters (Sro-Dnap and Sro-Ddap) were obtained by the removal of pKC1139::*nap* and pKC1139::*dap* from Sro-MCnap and Sro-MCdap.

Actinorhodin quantification. To quantitate actinorhodin production, *S. coelicolor* M145 and its derivatives were grown in 50 ml of R5MS at 28 °C. 1 ml culture was harvested in a time-course and treated with KOH (1 N final concentration), and titer was calculated by measuring the absorbance at 640 nm³⁷.

Production and analysis of daptomycin. Small-scale fermentation of daptomycin was carried out by following the procedures described previously^{38,39} with minor modifications. In brief, starter culture was grown in TSB for 48 h, 1 ml of starter culture was transferred to A355 (1% [wt/vol] glucose, 1.5% [vol/vol] glycerol, 1.5% [wt/vol] soya peptone, 0.3% [wt/vol] NaCl, 0.5% [wt/vol] malt extract, 0.5% [wt/vol] yeast extract, 0.1% [vol/vol] Tween 80 and 2% [wt/vol] MOPS, pH 7.0) and grown for 36 h as seed culture, and 1 ml of seed culture was transferred into a shake flask containing 50 ml A346 (1% [wt/vol] glucose, 2% [wt/vol] soluble starch, 0.5% [wt/vol] yeast extract, 0.5% [wt/vol] casein and 4.6% [wt/vol] MOPS, pH 7.0). The cultures were incubated for different time points at 28 °C before fermentation broths were collected by centrifugation.

For daptomycin analysis, culture broths were centrifuged at 13,000 × g for 10 min to remove the mycelia. The supernatants were filtered through a Millipore membrane (pore diameter, 0.22 µm) and 50 µl of sample was used for HPLC analysis. Separation of daptomycin was achieved with an Agilent 1100 HPLC system and a ZORBAX SB-Aq column (5 µm pore size, 4.6 by 250 mm). HPLC conditions were described as follows: gradient elution with buffer A (0.01% [vol/vol] trifluoroacetic acid in acetonitrile) and buffer B (0.01% [vol/vol] trifluoroacetic acid in ddH₂O), flow rate at 1.0 ml/min, ultraviolet detection at wavelength of 224 nm. The elution profile was a linear gradient of 10%–100% buffer A over 22 min, a hold at 100% buffer A over 3 min, a linear gradient of 100%–10% buffer A over 2 min and a final hold at 10% buffer A over 3 min.

Bioassay against *S. aureus* was performed as previously described with modifications³⁸. In brief, *S. roseosporus* and its derivatives were patched on AS-1 agar. After incubation for 2–5 days at 28 °C, agar plugs were prepared from the patches, placed on the surface of an empty Petri dish, and overlaid with culture of indicator strain in soft nutrient agar containing 5 mM CaCl₂. The zone of inhibition was assessed after overnight incubation at 37 °C.

- Niu, G. & Tan, H. Biosynthesis and regulation of secondary metabolites in microorganisms. *Sci. China Life Sci.* **56**, 581–583 (2013).
- Bibb, M. J. Regulation of secondary metabolism in streptomycetes. *Curr. Opin. Microbiol.* **8**, 208–215 (2005).
- Fu, J. *et al.* Full-length RecE enhances linear-linear homologous recombination and facilitates direct cloning for bioprospecting. *Nat. Biotechnol.* **30**, 440–446 (2012).
- Chain, P. S., Hernandez-Lucas, I., Golding, B. & Finan, T. M. *oriT*-directed cloning of defined large regions from bacterial genomes: identification of the *Sinorhizobium meliloti* pExo megaplasmid replicator region. *J. Bacteriol.* **182**, 5486–5494 (2000).
- Shao, Z., Luo, Y. & Zhao, H. Rapid characterization and engineering of natural product biosynthetic pathways via DNA assembler. *Mol. Biosyst.* **7**, 1056–1059 (2011).
- Kvitko, B. H., McMillan, I. A. & Schweizer, H. P. An improved method for *oriT*-directed cloning and functionalization of large bacterial genomic regions. *Appl. Environ. Microbiol.* **79**, 4869–4878 (2013).
- Bonet, B., Teufel, R., Crusemann, M., Ziemert, N. & Moore, B. S. Direct capture and heterologous expression of *Salinispora* natural product genes for the biosynthesis of enterocin. *J. Nat. Prod.* doi:10.1021/np500664q (2014).
- Yamanaka, K. *et al.* Direct cloning and refactoring of a silent lipopeptide biosynthetic gene cluster yields the antibiotic taromycin A. *Proc. Natl. Acad. Sci. U.S.A.* **111**, 1957–1962 (2014).
- Fernandez-Martinez, L. T. & Bibb, M. J. Use of the meganuclease I-SceI of *Saccharomyces cerevisiae* to select for gene deletions in actinomycetes. *Sci. Rep.* **4**, 7100, doi:10.1038/srep07100 (2014).
- Lu, Z., Xie, P. & Qin, Z. Promotion of markerless deletion of the actinorhodin biosynthetic gene cluster in *Streptomyces coelicolor*. *Acta Biochim. Biophys. Sin. (Shanghai)* **42**, 717–721 (2010).
- Grindley, N. D., Whiteson, K. L. & Rice, P. A. Mechanisms of site-specific recombination. *Annu. Rev. Biochem.* **75**, 567–605 (2006).
- Sternberg, N. & Hamilton, D. Bacteriophage P1 site-specific recombination. I. Recombination between loxP sites. *J. Mol. Biol.* **150**, 467–486 (1981).
- Sauer, B. & McDermott, J. DNA recombination with a heterospecific Cre homolog identified from comparison of the pac-c1 regions of P1-related phages. *Nucleic Acids Res.* **32**, 6086–6095 (2004).
- Schweizer, H. P. Applications of the *Saccharomyces cerevisiae* FLP-FRT system in bacterial genetics. *J. Mol. Microbiol. Biotechnol.* **5**, 67–77 (2003).
- Bierman, M. *et al.* Plasmid cloning vectors for the conjugal transfer of DNA from *Escherichia coli* to *Streptomyces* spp. *Gene* **116**, 43–49 (1992).
- Gregory, M. A., Till, R. & Smith, M. C. Integration site for *Streptomyces* phage φBT1 and development of site-specific integrating vectors. *J. Bacteriol.* **185**, 5320–5323 (2003).
- Zhang, L. *et al.* DNA cleavage is independent of synapsis during *Streptomyces* phage φBT1 integrase-mediated site-specific recombination. *J. Mol. Cell. Biol.* **2**, 264–275 (2010).



18. Ullrich, S. & Schuler, D. Cre-lox-based method for generation of large deletions within the genomic magnetosome island of *Magnetospirillum gryphiswaldense*. *Appl. Environ. Microbiol.* **76**, 2439–2444 (2010).
19. Komatsu, M., Uchiyama, T., Omura, S., Cane, D. E. & Ikeda, H. Genome-minimized *Streptomyces* host for the heterologous expression of secondary metabolism. *Proc. Natl. Acad. Sci. U.S.A.* **107**, 2646–2651 (2010).
20. Herrmann, S. *et al.* Site-specific recombination strategies for engineering actinomycete genomes. *Appl. Environ. Microbiol.* **78**, 1804–1812 (2012).
21. Muth, G., Wohlleben, W. & Puhler, A. The minimal replicon of the *Streptomyces ghanaensis* plasmid pSG5 identified by subcloning and Tn5 mutagenesis. *Mol. Gen. Genet.* **211**, 424–429 (1988).
22. Kaysser, L. *et al.* Identification of a napsamycin biosynthesis gene cluster by genome mining. *Chembiochem* **12**, 477–487 (2011).
23. Niu, G. & Tan, H. Nucleoside antibiotics: biosynthesis, regulation, and biotechnology. *Trends Microbiol.* **23**, 110–119 (2015).
24. Gust, B., Challis, G. L., Fowler, K., Kieser, T. & Chater, K. F. PCR-targeted *Streptomyces* gene replacement identifies a protein domain needed for biosynthesis of the sesquiterpene soil odor geosmin. *Proc. Natl. Acad. Sci. U.S.A.* **100**, 1541–1546 (2003).
25. Peschke, U., Schmidt, H., Zhang, H. Z. & Piepersberg, W. Molecular characterization of the lincomycin-production gene cluster of *Streptomyces lincolnensis* 78-11. *Mol. Microbiol.* **16**, 1137–1156 (1995).
26. Fierro, F. *et al.* The penicillin gene cluster is amplified in tandem repeats linked by conserved hexanucleotide sequences. *Proc. Natl. Acad. Sci. U.S.A.* **92**, 6200–6204 (1995).
27. Yanai, K., Murakami, T. & Bibb, M. Amplification of the entire kanamycin biosynthetic gene cluster during empirical strain improvement of *Streptomyces kanamyceticus*. *Proc. Natl. Acad. Sci. U.S.A.* **103**, 9661–9666 (2006).
28. Liao, G. *et al.* Cloning, reassembling and integration of the entire nikkomycin biosynthetic gene cluster into *Streptomyces ansochromogenes* lead to an improved nikkomycin production. *Microb. Cell Fact.* **9**, 6 (2010).
29. Jiang, L., Wei, J., Li, L., Niu, G. & Tan, H. Combined gene cluster engineering and precursor feeding to improve gougerotin production in *Streptomyces graminearus*. *Appl. Microbiol. Biotechnol.* **97**, 10469–10477 (2013).
30. Murakami, T., Burian, J., Yanai, K., Bibb, M. J. & Thompson, C. J. A system for the targeted amplification of bacterial gene clusters multiplies antibiotic yield in *Streptomyces coelicolor*. *Proc. Natl. Acad. Sci. U.S.A.* **108**, 16020–16025 (2011).
31. Gomez-Escribano, J. P. & Bibb, M. J. Engineering *Streptomyces coelicolor* for heterologous expression of secondary metabolite gene clusters. *Microb. Biotechnol.* **4**, 207–215 (2011).
32. Kieser, T., Bibb, M. J., Buttner, M. J., Chater, K. F. & Hopwood, D. A. *Practical Streptomyces genetics*. (John Innes Foundation, Norwich, U.K., 2000).
33. Okamoto, S., Taguchi, T., Ochi, K. & Ichinose, K. Biosynthesis of actinorhodin and related antibiotics: discovery of alternative routes for quinone formation encoded in the act gene cluster. *Chem. Biol.* **16**, 226–236 (2009).
34. Nguyen, K. T. *et al.* A glutamic acid 3-methyltransferase encoded by an accessory gene locus important for daptomycin biosynthesis in *Streptomyces roseosporus*. *Mol. Microbiol.* **61**, 1294–1307 (2006).
35. Sambrook, J. & Russell, D. W. *Molecular cloning: a laboratory manual*, 3rd ed. (Cold Spring Harbor Laboratory Press, NY, 2001).
36. Pullan, S. T., Chandra, G., Bibb, M. J. & Merrick, M. Genome-wide analysis of the role of GlnR in *Streptomyces venezuelae* provides new insights into global nitrogen regulation in actinomycetes. *BMC Genomics* **12**, 175 (2011).
37. Bystrykh, L. V. *et al.* Production of actinorhodin-related “blue pigments” by *Streptomyces coelicolor* A3(2). *J. Bacteriol.* **178**, 2238–2244 (1996).
38. Miao, V. *et al.* Genetic engineering in *Streptomyces roseosporus* to produce hybrid lipopeptide antibiotics. *Chem. Biol.* **13**, 269–276 (2006).
39. Miao, V. *et al.* Daptomycin biosynthesis in *Streptomyces roseosporus*: cloning and analysis of the gene cluster and revision of peptide stereochemistry. *Microbiology* **151**, 1507–1523 (2005).
40. MacNeil, D. J. *et al.* Analysis of *Streptomyces avermitilis* genes required for avermectin biosynthesis utilizing a novel integration vector. *Gene* **111**, 61–68 (1992).
41. Yang, H. *et al.* Autoregulation of *hpdR* and its effect on CDA biosynthesis in *Streptomyces coelicolor*. *Microbiology* **156**, 2641–2648 (2010).
42. Paget, M. S., Chamberlin, L., Atrih, A., Foster, S. J. & Buttner, M. J. Evidence that the extracytoplasmic function sigma factor σ^E is required for normal cell wall structure in *Streptomyces coelicolor* A3(2). *J. Bacteriol.* **181**, 204–211 (1999).
43. Li, R. *et al.* *polR*, a pathway-specific transcriptional regulatory gene, positively controls polyoxin biosynthesis in *Streptomyces cacaoi* subsp. *asoensis*. *Microbiology* **155**, 1819–1831 (2009).

Acknowledgments

This work was supported by grants from the Ministry of Science and Technology of China (grant nos. 2012CB721103 and 2013CB734001) and the National Natural Science Foundation of China (grant nos. 31270110 and 31171202). We would like to thank Professor Mervyn Bibb and Dr. Chris D. Den Hengst (John Innes Centre, Norwich, UK) for providing *S. coelicolor* M1146 and pJJ10500, respectively. We also thank Dr. Guojian Liao (Southwest University, Chongqing, China) for the gift of daptomycin standard.

Author contributions

D.D. and W.L. performed the experiments. T.Y. assisted with design of the project. L.H. assisted with the primary data analysis. N.G. conceived and designed the project, and wrote the manuscript. T.H. supervised the project and revised the manuscript.

Additional information

Supplementary information accompanies this paper at <http://www.nature.com/scientificreports>

Competing financial interests: The authors declare no competing financial interests.

How to cite this article: Du, D. *et al.* Genome engineering and direct cloning of antibiotic gene clusters via phage ϕ BT1 integrase-mediated site-specific recombination in *Streptomyces*. *Sci. Rep.* **5**, 8740; DOI:10.1038/srep08740 (2015).



This work is licensed under a Creative Commons Attribution 4.0 International License. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in the credit line; if the material is not included under the Creative Commons license, users will need to obtain permission from the license holder in order to reproduce the material. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>