

SCIENTIFIC REPORTS



OPEN

NF- κ B signaling is essential for resistance to heat stress-induced early stage apoptosis in human umbilical vein endothelial cells

Received: 04 February 2015

Accepted: 30 July 2015

Published: 04 September 2015

Yanan Liu¹, Gengbiao Zhou², Zhenglian Wang², Xiaohua Guo³, Qiulin Xu^{4,5}, Qiaobing Huang³ & Lei Su^{1,4}

Cell apoptosis induced by heat stress is regulated by a complex signaling network. We previously reported that a p53-dependent pathway is involved. Here, we present evidence that NF- κ B signaling plays a crucial role in preventing heat stress-induced early apoptosis. Human umbilical vein endothelial cells (HUVECs) were examined and increased phosphorylation of p65 and I κ B α were detected, without I κ B α degradation. When NF- κ B signaling was inhibited by BAY11-7082, or a small interference RNA (siRNA) targeting p65, a significant increase in cell apoptosis and caspase-3 activity was observed, as well as reduced expression and translocation of HSP27 into the nucleus, an accumulation of reactive oxygen species, and prolonged phosphorylation of mitogen-activated protein kinases (MAPKs). In addition, an association between HSP27 and p65 was identified which may enhance NF- κ B activation. When HSP27 was overexpressed, pretreatment of HUVECs with the antioxidant, apocynin, or N-acetyl cysteine, suppressed apoptosis. Similarly, inhibition of JNK and p38 with SP600125 and SB203580, respectively, also suppressed apoptosis, whereas siRNA-mediated HSP27 knockdown and treatment with the ERK 1/2 inhibitor PD98059 did otherwise. In conclusion, these findings suggest a novel role for an NF- κ B signaling pathway involving HSP27, ROS, and MAPKs that confers a protective effect against heat stress-induced cell apoptosis.

Heatstroke is a life-threatening condition that typically develops following exposure to extended periods of high temperatures. It is characterized by a rapid increase in core temperature to more than 40°C and multiple organ dysfunction syndrome (MODS)^{1–3}. The critical maximum temperature for the human body is between 41.6°C and 42.0°C. Previous studies have suggested that apoptosis is a major cause of cell death in heatstroke, and that it can be induced within a few hours^{4,5}. It is hypothesized that endothelial cell activation/injury contributes to the pathophysiology of heat stroke⁶, and endothelial damage has been detected in heatstroke patients^{7,8}. In addition, recent studies have reported that the acute phase of heat stress induces significant apoptosis in endothelial cells⁹, and we recently reported that intense heat stress induces early apoptosis via a transcription-independent mitochondrial p53 pathway¹⁰. However, the mechanisms mediating cell death in the late phase of heat stress remain unclear.

NF- κ B is an important intracellular signaling protein that controls the transcription of several genes involved in cell growth, inflammatory responses, cell survival, and cell apoptosis¹¹. When NF- κ B is associated with inhibitory molecules of the I κ B family in the cytosol, it is inactive. Correspondingly, most of

¹Southern Medical University, Guangzhou, China. ²Guangzhou University of Chinese Medicine, Guangzhou, China. ³Department of Pathophysiology, Southern Medical University, Guangzhou, China. ⁴Department of ICU, General Hospital of Guangzhou Military Command, Key Laboratory of Tropical Zone Trauma Care and Tissue Repair of PLA, Guangzhou, China. ⁵Postdoctoral Workstation, Huabo Bio-pharmaceutical Research Institute, Guangzhou, China. Correspondence and requests for materials should be addressed to Q.X. (email: qiul1978@gmail.com) or Q.H. (email: bing@smu.edu.cn) or L.S. (email: sulei_ICU@163.com)

the inducers that activate NF- κ B use a common pathway that involves phosphorylation-induced degradation of I κ B proteins. The latter includes the major protein, I κ B α , which was the first protein described for this family and is also the most extensively studied I κ B protein to date¹². Phosphorylation and degradation of I κ B α requires phosphorylation of the upstream target, I κ B kinase (IKK), which contains two catalytic subunits, IKK α and IKK β ¹³. Upon release from the NF- κ B/I κ B α dimer, NF- κ B translocates from the cytoplasm into the nucleus to bind DNA and regulate transcription.

The NF- κ B signaling pathway has a critical role in regulating various aspects of the apoptotic program¹⁴. For example, NF- κ B activation has been shown to down-regulate pro-apoptotic JNK signaling in many cell types, thereby preventing apoptosis^{15,16}. However, in certain pathological conditions, such as ischemia, the excessive accumulation of reactive oxygen species (ROS) can induce apoptosis or necrosis by activating mitogen-activated protein kinase (MAPK) and caspase signaling cascades, and/or by disrupting mitochondrial membrane potential in Jurkat and in HeLa cells¹⁷. NF- κ B has also been shown to exert pro-survival functions by inhibiting TNF- α -induced ROS accumulation-mediated prolongation of MAPK activation and necrotic cell death in murine embryonic fibroblasts¹⁸. Despite these insights, however, it remains unknown whether ROS play a critical role in heat stress-induced MAPK activation, and whether NF- κ B has a role in mediating oxidative stress and MAPK signaling pathways under physiological conditions in HUVECs.

Heat shock proteins (HSPs) are an evolutionarily conserved set of proteins that mediate a cell's response to heat stress, and a subset of HSPs protects cells against an induction of cell death (including apoptosis and necrosis) in response to a variety of stresses¹⁹. In particular, HSP27 and HSP70 have been shown to contribute to the regulation of NF- κ B activation in many different cell types^{20–24}, with a direct link observed between HSP27 and regulation of the NF- κ B signaling pathway in cell apoptosis. For example, in macrophage-conditioned intestinal epithelial cells stimulated with interleukin-1 β (IL-1 β), HSP27 was shown to bind and suppress IKK to regulate NF- κ B activation²⁵. Similar mechanisms have been found in keratinocytes stimulated with tumor necrosis factor- α (TNF- α) and UV irradiation²⁶, and in HeLa cells stimulated with TNF- α ²⁰. Moreover, when HSP27 was overexpressed in response to various stimuli, it facilitated proteasome-mediated proteolysis via phosphorylated I κ B α and enhanced NF- κ B activity²⁷. The latter observation is consistent with the antiapoptotic properties reported for HSP27²⁷. To investigate whether regulation of NF- κ B activation by HSP27 affects heat stress-induced cell apoptosis, various experiments were performed using HUVECs as a model. As a result, a novel NF- κ B signaling pathway was identified that includes HSP27 protein expression and translocation into the nucleus, the accumulation of ROS, and subsequent MAPK activation.

Results

Heat stress activates NF- κ B during the recovery period following heat stress. The transcription factor, NF- κ B, has been shown to be activated during the recovery period following heat stress in HeLa cells²⁸. Therefore, in this study, it was investigated whether heat stress activates NF- κ B in human umbilical vein endothelial cells (HUVECs). After HUVEC cells were grown in culture media for 48 h, the culture dishes were sealed with parafilm and immersed in a circulating water bath maintained at 43 °C to induce heat stress²⁹. After 90 min, the culture media was replaced with fresh media and the cells were further incubated at 37 °C for various periods of time (e.g., 0, 2, 6, and 12 h) before being assayed. Indirect immunofluorescence studies demonstrated that the distribution of p65 in nucleus was obviously increased after 6–12 h of heat stress recovery at 37 °C (Fig. 1a). Then nuclear and cytoplasmic extracts of the same timepoints were collected and analyzed by western blot. The result revealed that, unexpectedly, the amount of p65 in nuclear and cytoplasm were both increased (Fig. 1b). It is possible that this increase of nucleus p65 is due to the increase of p65 synthesis, rather than the shuttling of p65 from cytoplasm to nucleus.

To address this issue, 0.5 μ g/ml actinomycin D was added to HUVECs 5 min prior to the heat stress treatment in order to block transcription and translation²⁸. An ELISA-based TransAM NF- κ B Activation kit was then used to quantify NF- κ B binding to DNA^{30,31}. Pretreatment with actinomycin D did not modify the DNA-binding kinetics of NF- κ B during the heat stress recovery period (Fig. 1c), thereby suggesting that no new translational events are required during heat stress recovery for NF- κ B activation. Taken together, these data suggest that heat stress induces the translocation and activation of NF- κ B during the recovery period in HUVECs.

NF- κ B activation during heat stress recovery occurs with prior phosphorylation, and without degradation of the I κ B subunits. To determine whether activation of NF- κ B is associated with the phosphorylation of I κ B α , IKK- α/β , and/or p65, as well as the degradation of I κ B α , Western blot assays were performed for HUVEC extracts collected during heat stress treatment (Fig. 2a) and during the recovery period following the heat stress treatment (Fig. 2b). Phosphorylation levels of IKK- α/β were high at 37 °C, and were unchanged by heat stress. In contrast, levels of phosphorylated I κ B α and p65 were low at 37 °C, but increased after 15 min and 90 min of heat treatment (Fig. 2a,b). The levels of phosphorylated I κ B α and p65 then further increased 24 h after the heat stress treatment was completed.

There were no signs of I κ B α degradation observed during the heat stress treatment or during the recovery periods assayed. Thus, it appears that NF- κ B activation is accompanied by phosphorylation of p65 and I κ B α , and not by degradation of I κ B α in HUVECs.




Figure 1. Relocalization of p65 from the cytosol into the nucleus of heat stressed HUVECs. Cells were incubated at 37°C (CONT) or were subjected to a heat stress (HS) treatment at 43°C for 90 min, followed by a recovery period at 37°C for 0 h (R0), 2 h (R2), 6 h (R6), or 12 h (R12). (a) The cells were then fixed and processed for indirect immunofluorescence analysis using an antibody raised against the p65 subunit of NF- κ B. Representative images are shown. (b) Expression of p65 was detected in cytoplasmic (CE) and nuclear (NC) fractions of HUVECs by Western blotting. The cropped images represent blotting experiments that were performed under the same experimental conditions. (c) HUVECs were pretreated with or without 0.5 μ g/ml actinomycin D 5 min before being incubated at 37°C (CONT) or being subjected a HS treatment. Whole cell extracts were prepared and NF- κ B binding to DNA was quantified with the Trans-AMTMp65 transcription factor assay kit. Each value represents the mean \pm SD of three independent experiments. ** $P < 0.01$ and *** $P < 0.001$ versus control.




Figure 2. NF- κ B activation during the heat stress recovery period involves the phosphorylation of I κ B α and p65, without degradation of I κ B α . HUVECs were either maintained at 37°C (CONT) or were subjected to a heat stress treatment (HS) at 43°C for 15, 30, or 90 min (a), or a HS at 43°C for 90 min, followed by a recovery period at 37°C for 0 h (R0), 2 h (R2), 6 h (R6), 12 h (R12), or 24 h (R24) (b). Protein expression levels of I κ B α , p-I κ B α , p65, p-p65, p-IKK α/β , IKK α , and GAPDH for each set of samples were determined by Western blotting. The cropped images represent blotting experiments that were performed under the same experimental conditions. Each value represents the mean \pm SD of three independent experiments. * $P < 0.05$, ** $P < 0.01$, *** $P < 0.001$ versus control.

NF- κ B activation decreases heat stress-induced apoptosis. To investigate whether NF- κ B activation affects heat stress-induced apoptosis, Annexin V-FITC/PI staining was used to differentiate early apoptosis from late apoptosis and necrosis. HUVECs that were pretreated with the NF- κ B inhibitor, BAY11-7082, prior to heat stress treatment were more susceptible to high levels of early apoptosis

compared with the HUVECs that were pretreated with DMSO (Fig. 3a). When HUVECs were transfected with a p65-targeted siRNA, lower levels of p65 protein were detected (Fig. 3b), and these cells were more susceptible to heat stress-induced early apoptosis (Fig. 3c).

Caspase-3 activity was also assayed for HUVECs that were pretreated with DMSO or the NF- κ B inhibitor, BAY11-7082 (5 μ M), for 1 h, and for HUVECs that were transfected with p65-targeted siRNA for 48 h, then underwent a heat stress treatment (Fig. 3d,e). Higher levels of caspase-3 activity were detected when NF- κ B was inhibited and when levels of p65 were knocked down. Taken together, these results indicate that activation of NF- κ B is essential for HUVECs to resist heat stress-induced apoptosis.

Role of HSP27 in activating heat stress-induced apoptosis in HUVECs. To investigate the role of HSP27 in heat stress-induced cell apoptosis, HUVECs were transfected with an HSP27-targeted siRNA or an adenovirus expressing HSP27, and then were subjected to a heat stress treatment followed by a 24 h recovery period. The knockdown and overexpression of HSP27 that was initially achieved was detected in Western blot assays (Fig. 4a). Levels of apoptosis were subsequently analyzed by flow cytometry using Annexin V-FITC/PI staining. Higher levels of heat stress-induced apoptosis were detected in HUVECs following the knockdown of HSP27, while an increase in apoptosis levels were detected in cells overexpressing HSP27 (Fig. 4b). A similar profile was obtained when the corresponding cell lysates were analyzed for caspase-3 activity (Fig. 4c). In combination, these data suggest that HSP27 protects HUVECs from heat stress-induced apoptosis.

Co-localization and functional correlation of HSP27 with NF- κ B in HUVECs during the recovery period following heat stress. Based on the anti-apoptotic properties of HSP27 during heat stress recovery, and previous observations that HSP27 and NF- κ B translocate into the nucleus in response to heat stress^{28,32}, we hypothesized that NF- κ B activation after heat stress may be linked to HSP27. Therefore, immunofluorescence studies of heat-stressed HUVECs were performed to detect the expression and localization of HSP27 and p65. Heat stress was found to stimulate the translocation of both HSP27 and NF- κ B from the cytoplasm into the nucleus (Fig. 5a). To investigate the potential for interactions between NF- κ B and HSP27, HUVECs were subjected to a heat stress treatment, they recovered at 37 °C for 6 h, and then whole cell lysates were prepared. Subsequent co-immunoprecipitation assays were performed with an anti-NF- κ B p65 antibody and Western blotting revealed that NF- κ B p65 co-immunoprecipitated with HSP27 both at 37 °C and 6 h after a heat stress treatment (Fig. 5b).

To further assess the interactions between NF- κ B and HSP27, HUVECs were transfected with an HSP27-targeted siRNA or an adenovirus expressing HSP27, and then were subjected to a heat stress treatment followed by a 6 h recovery period. Following the knockdown of HSP27 expression, DNA binding by NF- κ B in the nucleus decreased (Fig. 5c). Although the NF- κ B DNA-binding capacity did not notably increase after HSP27 overexpression, it may be that the levels of HSP27 themselves were increased after heat stress treatment. These results suggest that HSP27 may facilitate the nuclear import of NF- κ B.

Meanwhile, blocking NF- κ B activation by the inhibitor BAY11-7082 also prevents Hsp27 from going to the nucleus (Fig. 5d). However, when the HUVECs were pretreated with BAY11-7082 before heat treatment, the nuclear and cytoplasmic levels of p65 decreased. It is possible that a feedback loop exists whereby NF- κ B regulates p65 expression. Moreover, our data showed that heat stress significantly increased HSP27 protein expression during heat recovery periods, which were decreased by inactivation of NF- κ B with BAY11-7082 or p65siRNA (Fig. 5e). These results suggested that NF- κ B signaling regulates the translocation and expression of HSP27.

Influence of NF- κ B signaling on MAPK activation. Extracellular signal-regulated kinase (ERK), c-Jun NH2-terminal kinase (JNK), and p38 are members of well-characterized subfamilies of MAPK, and these enzymes have been implicated in the increased sensitivity to heat stress-induced cell apoptosis exhibited by IEC-6 cells³³. To examine the role of these three kinases in HUVECs during heat stress and after various periods of recovery following heat stress, HUVEC extracts were analyzed by Western blot. Phosphorylation of ERK and phosphorylation of p38 were significantly induced after 15 min at 43 °C, while phosphorylation of JNK occurred after 60 min at 43 °C (Fig. 6a). During the recovery period, MAPK phosphorylation was rapidly induced and was maintained at high levels for more than 4 h before decreasing to baseline levels after a 6 h recovery period (Fig. 6b). Since the decrease in MAPK levels temporally coincided with NF- κ B activation, it was hypothesized that activation of NF- κ B inhibited MAPK phosphorylation. Therefore, HUVECs were pretreated with BAY11-7082, SN50, or p65-targeted siRNA, then were subjected to a heat stress treatment and a 6 h recovery period (Fig. 6c,d). Decreasing levels of MAPK phosphorylation were observed except when NF- κ B was inactivated. These data suggest that heat stress-induced phosphorylation of MAPK proteins is inhibited by NF- κ B activation 6 h after a heat stress event.

ROS mediates the inhibition of MAPK phosphorylation by NF- κ B activation. As shown in Fig. 7a, the antioxidant apocynin (APO) inhibited ROS accumulation in the heat stress group, but the effects were not complete in untreated HUVECs. We utilized the fluorescent dye DCFH-DA, which produces enhanced fluorescence when cells generate ROS, and analyzed fluorescent signals by flow





Figure 3. Influence of NF- κ B on endothelial cell apoptosis induced by heat stress. (a) Cells were pretreated with DMSO or 5 μ M BAY11-7082 (BAY) for 1 h, then were incubated at 37 $^{\circ}$ C (CONT) or 43 $^{\circ}$ C for 90 min (HS), followed by a recovery period at 37 $^{\circ}$ C for 24 h. Levels of apoptosis were detected with Annexin V-FITC/PI staining and were analyzed by flow cytometry. Levels of early apoptosis are shown in the bar graph at the far right. (b) HUVECs were transfected with a negative control siRNA (NCsiRNA) or a p65-targeted siRNA (p65siRNA) for 48 h. Protein expression of p65 was detected by Western blot and band density was quantified relative to GAPDH and is graphed at the right. (c) Apoptosis was detected for the samples characterized in (b) following staining with Annexin V-FITC/PI. Levels of early apoptosis were analyzed and are quantitated in the bar graph. (d) Enzymatic activity of caspase-3 was measured for the cell lysates indicated using the fluorogenic substrate, Ac-DEVD-AMC. Data are expressed as a relative increase to the control (CONT) and represent the mean \pm SD of three independent experiments. * P < 0.05, ** P < 0.01.






Figure 5. HSP27 associates and interacts with NF- κ B in heat stressed HUVECs. HUVECs were incubated at 37°C (CONT) or were subjected to a heat stress (HS) treatment at 43°C for 90 min, then were further incubated for 6 h at 37°C. **(a)** The cells were subsequently stained with an anti-HSP27 antibody (shown in green) and an anti-p65 antibody (shown in red). Co-staining with DAPI was used to visualize the nuclei (shown in blue). Merged images of these three stainings are shown in the right panels. **(b)** Whole cell lysates were prepared and probed in co-immunoprecipitation assays with an anti-p65 antibody, while a normal IgG antibody was used as a negative control. Western blot assays were then performed for the immunoprecipitated samples with an anti-HSP27 antibody. **(c)** HUVECs were transfected with a negative control siRNA (NCsiRNA) or an *Hsp27*-targeted siRNA (HSP27siRNA), or with an empty adenovirus vector (Ad-empty) and a adenovirus vector expressing HSP27 (Ad-HSP27), respectively. These transfected cells were then incubated at 37°C (CONT) or were subjected to a heat stress (HS) treatment at 43°C for 90 min, and recovery period at 37°C for 6 h. Binding of NF- κ B to DNA was quantified by using a Trans-AMTMp65 transcription factor assay kit. The values represent the relative binding that was measured at 450 nm. **(d–e)** HUVECs were pretreated with DMSO or 5 μ M BAY11-7082 for 1 h **(d)**, or were transfected with a negative control siRNA (NCsiRNA) or a p65-targeted siRNA (p65siRNA) for 48 h **(e)**. Then both sets of cells were incubated at 37°C (CONT) or 43°C for 90 min (HS), followed by recovery period at 37°C for 2 h (R2), 6 h (R6), or 12 h (R12). Expression levels of p65 and HSP27 were detected in Western blot assays performed. The cropped images represent blotting experiments that were performed under the same experimental conditions. Each value represents the mean \pm SD of three independent experiments. * $P < 0.05$, ** $P < 0.01$, ns: no significance. CE: cytoplasmic; NC: nuclear.




Figure 6. Prolonged MAPK activation detected in heat stressed HUVECs. (a) HUVECs were incubated at 37°C (CONT) or were subjected to heat stress (HS) treatments at 43°C for the indicated times. (b) HUVECs were incubated at 37°C (CONT) or were subjected to a heat stress treatment at 43°C for 90 min, and then were further incubated at 37°C for the indicated times. (c) HUVECs were treated with BAY11-7082 or SN50 for 1 h prior to a heat stress treatment at 43°C for 90 min. The cells were then further incubated at 37°C for 6 h. (d) HUVECs were transfected with a negative control siRNA (NCsiRNA) or a p65-targeted siRNA (p65siRNA) and were incubated at 37°C (CONT) or were subjected to a heat stress (HS) treatment at 43°C for 90 min, followed by a recovery period at 37°C for the indicated times. For a-d, the lysates were blotted with antibodies specific for activated forms of JNK (phospho-JNK), p38 (phospho-p38), and ERK (phospho-ERK), with detection of GAPDH used as a loading control. The membranes were reblotted with antibodies to detect levels of total JNK, p38, and ERK. All of the cropped images represent blotting experiments that were performed under the same experimental conditions.

cytometry. H_2O_2 was used as a positive control. Consistent with the partial inhibitory effect of APO on heat stress-induced ROS accumulation, APO only partially inhibited heat stress-induced MAPK activation (Fig. 7b). These results demonstrate that accumulation of ROS perfectly coincides with prolonged MAPK activation.

The observed inhibition of MAPK activation by APO treatment in HUVECs prompted us to examine whether heat stress stimulation induces ROS accumulation in BAY11-7082- and SN50-pretreated




Figure 7. Inactivation of NF- κ B induced the accumulation of ROS in HUVECs during the heat stress recovery period. (a) HUVECs were pretreated with DMSO or APO and then were subjected to 37°C (CONT) or heat stress (HS) treatment for 90 min, followed by a second incubation at 37°C for 6 h. HUVECs were treated with 1 mM H₂O₂ as a positive control for ROS. Treated cells were labeled with DCFH-DA for the last 30 min. Fluorescence was detected by flow cytometry. (b) HUVECs were untreated or pretreated with APO (250 μ M) for 30 min, and then were incubated at 37°C (CONT) or were subjected to a heat stress (HS) treatment at 43°C for the indicated time periods. The corresponding cell lysates were analyzed as described in Fig. 6, and the cropped gels were run under the same experimental conditions. (c) HUVECs were pretreated with DMSO or the NF- κ B inhibitors, BAY11-7082 and SN50, for 1 h, or were transfected with a negative control siRNA (NCsiRNA) or a p65-targeted siRNA (p65siRNA) for 48 h. These cells were subsequently incubated at 37°C (CONT) or 43°C (HS) for 90 min, followed by a recovery period of 6 h at 37°C. The cells were labeled with DCFH-DA for the last 30 min and fluorescence was detected by flow cytometry. (d) HUVECs were transfected with NCsiRNA or p65siRNA for 48 h, then were incubated at 37°C (CONT) or 43°C (HS) for 90 min, followed by a recovery period at 37°C for 6 h. Mitochondrial membrane potential ($\Delta\Psi$ m) was assessed with JC-1 staining: red cells represent JC-1 aggregates due to an intact $\Delta\Psi$ m. Conversely, the green cells represent JC-1 monomers due to a disrupted $\Delta\Psi$ m. Each value represents the mean \pm SD from three independent experiments. *P < 0.05, **P < 0.01, ***P < 0.001.

cells and p65-depleted HUVEC cells. A substantial increase in the fluorescent signals of DCFH-DA was observed in BAY11-7082- or SN50-pretreated cells during the heat stress recovery period compared to the untreated group. The same results were obtained with p65-depleted HUVECs (Fig. 7c). In addition, p65-depleted HUVEC cells significantly increased the heat stress-induced the loss of mitochondrial membrane potential ($\Delta\Psi_m$) (Fig. 7d), thereby suggesting that mitochondria may provide a permanent source of ROS in heat stressed HUVECs. Overall, the results of these experiments suggest that ROS contribute to the signaling pathway involving NF- κ B activation-induced phosphorylation of MAPKs in response to heat stress.

Contribution of ROS and MAPKs to heat stress-induced cell apoptosis in HUVECs. To examine whether accumulated ROS or MAPK activation participates in heat stress-induced cell apoptosis, HUVECs were stimulated with heat stress in the presence or absence of inhibitors for ROS or MAPKs. As shown in Fig. 8a, treatment with APO or N-acetyl-L-cysteine (NAC) alone substantially decreased the number of cells undergoing early apoptosis. Furthermore, while HUVECs pretreated with PD98059, a specific inhibitor of ERK, exhibited an increase in cell apoptosis, HUVECs pretreated respectively with specific inhibitors of JNK and p38, SP600125 and SB203580, exhibited a significant decrease in cell early apoptosis (Fig. 8b). Caspase-3 activity was also assayed (Fig. 8c,d), and these results are consistent with the apoptosis data. Finally, We verified that these inhibitors actually inhibited MAP kinase activities by using antibodies specific for phosphorylated form of JNK, ERK, or p38 (Fig. 8e). The three MAPK inhibitors examined also did not affect the heat stress-induced accumulation of ROS in HUVECs (data not shown), thereby indicating that MAPK activation is a downstream event of ROS accumulation.

Discussion

In this study, we demonstrate that activation of the NF- κ B signaling pathway is essential for resistance to heat stress-induced apoptosis in HUVECs, and this pathway includes roles for NF- κ B in mediating an increase in the expression and translocation of HSP27 into the nucleus, as well as an inhibition of ROS accumulation and subsequent MAPK activation. Together, these actions by NF- κ B lead to its anti-apoptotic role in HUVECs. Moreover, the association between HSP27 and p65 was found to regulate the heat stress-induced activation of NF- κ B that affects HUVEC survival.

In HeLa cells, it has been demonstrated that heat shock is a very powerful inducer of NF- κ B-dependent transcription²⁸. Accordingly, it has been hypothesized that hyperthermic stress as a result of heat stroke can lead to recruitment of NF- κ B. However, in a previous study, binding of DNA by NF- κ B was not detected in cells that underwent heat shock^{34,35}. It has been proposed a mild heat treatment, or one of short duration, without a recovery period does not lead to NF- κ B activation²⁸. In the present study, NF- κ B activation in HUVECs was not observed until 6 h after administration of a heat stress treatment. In addition, NF- κ B was found to only bind DNA during the recovery period.

To date, there have been several exceptions to the universal pathway of NF- κ B activation depending on the cell type and extracellular stimuli studied¹⁴. For example, activation of NF- κ B in response to UV radiation or amino acid analog treatment requires I κ B α degradation without prior phosphorylation^{36,37}, while anoxia stimulates I κ B α phosphorylation at Tyr42, leading to the release of p65/p50 from the NF- κ B-I κ B α complex³⁸. NF- κ B can also be activated by Ang II in vascular smooth muscle cells via phosphorylation of p65 at Ser536, and not via phosphorylation of I κ B α ³⁹. In HeLa cells, NF- κ B activation during the heat stress recovery period has been associated with the thermostability of the NF- κ B-I κ B α complex, independent of prior phosphorylation or degradation of the I κ B subunits²⁸. In the present study, a novel mechanism for heat stress-induced NF- κ B activation is characterized in endothelial cells, and this mechanism involves phosphorylation of p65 and I κ B α , without I κ B α degradation, and this differs from the NF- κ B activation pathway that has been characterized in HeLa cells.

In diverse cell types, NF- κ B signaling has been shown to have a critical role in regulating the apoptotic program. Moreover, whether NF- κ B promotes or inhibits apoptosis appears to depend on the cell type or inducer studied¹⁴. In the experiments performed in the present study to determine whether NF- κ B activation plays a role in heat stress-induced apoptosis in HUVECs, a significant increase in the levels of early cell apoptosis were detected 24 h after the heat stress treatment. Furthermore, when NF- κ B activation was inhibited by BAY11-7082 or p65-targeting siRNA, the levels of cell apoptosis markedly increased. Nivon *et al.*⁴⁰ previously described a significant increase in late apoptosis and necrosis in HeLa cells during the recovery period following heat stress, and these levels were markedly reduced by NF- κ B-activated autophagy. Interestingly, early apoptosis was the predominant event observed in the present study following heat treatment, while only a few late apoptosis and necrosis events were detected. This difference is attributed to the different cell types that were examined³³.

Here, the anti-apoptotic effect of NF- κ B was linked with HSP27, an important anti-apoptosis protein that translocates into the nucleus in response to heat stress. The findings of the present study also support the hypothesis that HSP27 facilitates the nuclear transport of NF- κ B to regulate cell apoptosis. For example, HSP27 and NF- κ B p65 exhibited similar cytoplasm-nucleus translocation events, HSP27 and NF- κ B p65 were co-immunoprecipitated from heat stressed HUVEC lysates, and HSP27 and NF- κ B p65 appeared to colocalize at 37°C and 43°C in HUVECs. In addition, knockdown of HSP27 abolished NF- κ B transactivation. Thus, attenuation of cell apoptosis by HSP27 is likely due, at least in part, to its enhancement of NF- κ B activation.




Figure 8. The role of ROS and MAPKs in heat stress induced cell apoptosis in HUVECs. (a) HUVECs were pretreated with DMSO or APO or NAC prior to being incubated at 37°C (CONT) or 43°C (HS) for 90 min, followed by a recovery period at 37°C for 6 h. Apoptosis was analyzed by flow cytometry using Annexin V-FITC/PI staining. (b) HUVECs were pretreated with DMSO, PD98059, SP600125, and SB203580 (specific inhibitors of ERK, JNK, and p38, respectively), prior to being incubated at 37°C (CONT) or 43°C (HS) for 90 min, followed by a recovery period at 37°C for 6 h. Apoptosis was analyzed by flow cytometry using Annexin V-FITC/PI staining and was quantitated. (c,d) Enzymatic activity of caspase-3 was detected for the cell lysates of the cells analyzed in panels A and B. Data are expressed as the fluorescence increase relative to the control at 37°C. (e) HUVECs were pretreated with or without MAPK inhibitors (PD98059, SP600125, and SB203580, respectively) for 30 min prior to a heat stress (HS) treatment at 43°C for 90 min. Cell lysates were subsequently collected and analyzed by Western blot as described in Fig. 6. The cropped gels were run under the same experimental conditions. Each value represents the mean \pm SD of three independent experiments. * $P < 0.05$, ** $P < 0.01$, ns: no significance.

There are many potential mechanisms by which HSP27 may regulate NF- κ B activation given the broad range of cellular functions that HSP27 participates in. However, we hypothesize that HSP27 stabilizes and protects stress-labile NF- κ B p65, thereby allowing for efficient translocation of NF- κ B into the nucleus to modulate gene transcription and apoptosis. To our knowledge, the present study is the first to identify HSP27 as a positive regulator of the translocation of p65-associated NF- κ B into the nucleus in response to heat stress. Furthermore, these results are in agreement with the data collected following the treatment of HUVECs with TNF- α , where HSP27 appeared to be essential for sustained NF- κ B activation⁴¹. Meanwhile, NF- κ B was found to increase HSP27 expression and its translocation into the nucleus since inhibition of NF- κ B with BAY11-7082, or following depletion of p65 with siRNA, led to a decrease in the expression and translocation of HSP27 during heat stress recovery. While additional studies are needed to determine the mechanism by which NF- κ B modulates HSP27 expression, the results of the present study suggest that HSP27 is involved in regulating the NF- κ B signaling pathway, and NF- κ B functions upstream of HSP27 to regulate its transcription and nuclear accumulation. It is also possible that a positive feedback loop exists between HSP27 and the activation of NF- κ B. Regardless, the data of the present study suggest that NF- κ B/HSP27 signaling protects endothelial cells from heat stress-induced apoptosis.

The results of the present study are consistent with those from IEC-6 cells where heat stress was found to induce the accumulation of ROS and to reduce anti-oxidase activity and MAPK signaling to affect apoptosis^{33,42}. Under physiological conditions, ROS are rapidly eliminated by antioxidant enzymes. However, the excessive accumulation of ROS can lead to the activation of MAPK and caspase signaling cascades, as well as the loss of $\Delta\Psi_m$, thereby leading to apoptotic or necrotic cell death¹⁷. In the present study, the accumulation of ROS significantly increased cell apoptosis and led to enhanced phosphorylation of ERK1/2, JNK, and p38. Moreover, these phosphorylation events were inhibited by activation of NF- κ B, or the antioxidant, APO. These results suggest that NF- κ B can inhibit heat-induced ROS accumulation that mediates MAPK activation. It is generally believed that ERKs are important for cell survival, while JNKs and p38 MAPKs have been characterized as stress-responsive factors, and thus, have a role in apoptosis⁴³. Therefore, the dynamic balance between heat stress-activated ERK and JNK-p38 signaling pathways are hypothesized to be important for determining whether a cell survives or undergoes apoptosis. Here, heat stress-induced apoptosis was attenuated by the antioxidant, APO, or NAC, as well as by inhibitors of JNK (SP600125) and p38 (SB203580). In contrast, heat stress-induced apoptosis was exacerbated by inhibition of ERK1/2 by PD98059. Taken together, these data suggest that heat stress can induce both pro- and anti-apoptotic pathways via an NF- κ B/ROS/MAPK signaling pathway in HUVECs. The most apparent effects under the conditions examined were anti-apoptotic.

In conclusion, the present study provides evidence that apoptotic cell death occurs in the late phase of heat stress recovery and it is regulated by a complex signaling network that is stimulated by NF- κ B activation. This network also involves the expression and translocation of HSP27, ROS accumulation, and MAPK activation. Furthermore, the association of HSP27 with p65 appears to regulate heat stress-induced NF- κ B activation (Fig. 9). Taken together, these results suggest that NF- κ B signaling plays a novel role in the tolerance of cells to intense heat stress and it represents a potential therapeutic target for heat stroke.

Methods

Cell culture and treatments. HUVECs were purchased from the Shanghai Institute of Cell Biology, Chinese Academy of Sciences. Cells were grown in DMEM/F12 supplemented with 10% (v/v) fetal bovine serum (FBS), 100 U/ml of penicillin, and 100 μ g/ml of streptomycin (Invitrogen Life Technology, USA) at 37°C in a humidified atmosphere of 5% CO₂ and 95% air. For the heat stress treatment, the bottom of each culture dish was placed into a circulating water bath that was maintained at 43 \pm 0.5°C for various periods of time as indicated. The cell culture media for each plate was then replaced with fresh media and the cells were further incubated at 37°C as indicated. As a control, cell culture dishes were placed in a circulating water bath maintained at 37 \pm 0.5°C.

Immunofluorescence analysis. Cell culture media was removed and cells were rinsed twice with PBS (2 min each) before being fixed in 3.5% paraformaldehyde and permeabilized in 0.1% Triton X-100 for 10 min at room temperature. After the cells were rinsed four times in PBS, the cells were incubated with PBS containing 5% FBS and 5% glycerol at 37°C. After 45 min, the cells were incubated with a rabbit anti-human p65 antibody (1:50) or a mouse anti-HSP27 antibody (1:100) overnight at 4°C. The cells were then washed with PBS twice (5 min each) and were incubated with an appropriate secondary antibody (1:100) at 37°C. After 1 h, the cells were incubated with mounting solution containing 0.2 μ g/ml DAPI for another 10 min. Images were acquired using a confocal fluorescence microscope.

Western blot analysis. HUVECs were subjected to a heat stress treatment and then were lysed in NE-PER extraction reagent (Pierce) according to the manufacturer's protocol to obtain cytoplasmic and nuclear protein extracts. Total protein concentrations were determined by the Bradford assay. Western blotting was performed as described previously⁴⁴. The primary antibodies used included: p65 and HSP27 from Abcam (USA), and I κ B α , p-I κ B α , p-p65, p-IKK α / β , IKK α , p-JNK, p-p38, p-ERK, JNK, p38, ERK, and GAPDH from Cell Signaling Technology (USA). All of the antibodies were diluted 1:2000, and a horseradish peroxidase (HRP)-conjugated anti-rabbit IgG antibody was diluted 1:5000 as the secondary




Figure 9. NF- κ B activation during heat stress recovery follows a non-canonical signal transduction pathway in HUVECs with increased phosphorylation of p65 and I κ B α , without I κ B α degradation.

Initially, heat stress stimulation induces NF- κ B activation, and this leads to an anti-apoptosis effect involving inhibition of heat stress-induced ROS accumulation that normally mediates the activation of MAPKs. In heat-stressed HUVECs, ERK1/2 activation has an anti-apoptotic role, while activation of JNK and p38 is pro-apoptotic. The results of the present study confirm that HSP27 is an important anti-apoptotic protein, and NF- κ B may modulate HSP27 expression levels and its translocation into the nucleus. In addition, HSP27 acting as a molecular chaperone may facilitate activation of this NF- κ B pathway in HUVECs during heat stress recovery. Taken together, our findings suggest that this NF- κ B signaling pathway involving HSP27, ROS, and MAPK is activated in response to heat stress and it confers a protective effect against heat stress-induced cell apoptosis.

antibody (Zhongshan Inc., China). Signals were detected using enhanced chemiluminescence reagents (Pierce, USA).

Measurement of NF- κ B DNA-binding capacity by enzyme-linked immunosorbent assay (ELISA). Nuclear extracts were prepared from treated and control HUVECs using a nuclear extraction kit (Active Motif, Inc). These extracts were then assayed for the ability of nuclear NF- κ B p65 to bind a DNA consensus sequence provided by an ELISA-based TransAM NF- κ B p65 kit (Active Motif), according to the manufacturer's protocol.

Analysis of cell apoptosis using Annexin V-FITC/PI staining and flow cytometry. Cell apoptosis was analyzed with an Annexin V-FITC apoptosis detection kit and flow cytometry according to the manufacturer's protocol (Invitrogen). Briefly, $\sim 1 \times 10^6$ cells were collected, washed in ice-cold PBS, and resuspended in binding buffer containing a suitable amount of Annexin V-FITC. After a 10 min incubation at room temperature, the buffer was removed by centrifugation and the cells were resuspended in a reaction buffer containing propidium iodide (PI). After 10 min, a flow cytometric analysis was performed for each sample to detect apoptosis.

Caspase-3 activity assay. After a heat stress treatment at 43°C for 90 min, HUVECs were harvested and cell lysates were prepared at -80°C for 30 min. The cells were then incubated with caspase-3

Target gene		Sequence (5'–3')
p65	Sense	GCCCUAUCCCUUUACGUCATT
	Antisense	UGACGAAAAGGGAUAGGGCTT
HSP27	Sense	GUCUCAUCGGAUUUUGCAGC
	Antisense	GCUGCAAAAUCCGAUGAGACTT
Negative Control	Sense	UUCUCCGAACGUGUCACGUTT
	Antisense	ACGUGACACGUUCGGAGAATT

Table 1. Oligonucleotide sequences used.

substrates at 37°C and caspase-3 activity was measured based on cleavage of the fluorogenic peptide substrate⁴⁵, Ac-DEVD-AMC, that was detected with a Quadruple Monochromator Microplate Reader (Infinite M1000, USA). Levels of caspase-3 activity are reported as relative cumulative fluorescence of the kinetic reaction compared to untreated controls.

Transfection of siRNA. siRNA targeting *p65* and *Hsp27* were designed and synthesized by GenePharma (Shanghai, China). The sequence of each gene and their corresponding negative controls are shown in Table 1. For transfection, HUVECs were plated onto 6-well plates (Nest, Biotech, China) at 30–50% confluence. Twenty-four hours later, HUVECs were incubated with siRNAMate Transfection Reagent (GenePharma Company, Shanghai, China) and siRNAs according to the manufacturer's protocol. The transfected cells were collected 48–72 h later.

Adenoviral infection. Adenoviruses (Ad) constitutively overexpressing HSP27 versus an empty control vector (Ad-empty) were constructed by ViGenen Biosciences (China). Cells were infected with the adenoviruses in serum-free DMEM for 6 h before the media was replaced with DMEM supplemented with 10% FBS.

Co-immunoprecipitation (co-IP) assays. HUVEC cells were subjected to a heat treatment at 43°C for 90 min, followed by an additional incubation period at 37°C for the times indicated. Cell lysates (containing ~1.5 mg total protein) were subsequently harvested, and co-IP assays were performed by using the Thermo Scientific Pierce co-IP kit according to the manufacturer's protocol.

Measurement of ROS levels. Levels of intracellular ROS were assessed with a kit (Beyotime, Nanjing, China) that employs dichlorofluorescein diacetate (DCFH-DA, Molecular Probes, Beyotime) as a reagent that enters cells and reacts with ROS, thereby producing the fluorophore, dichlorofluorescein (DCF). Briefly, cells were maintained at 37°C or were heated to 43°C for 90 min, followed by a recovery period of various lengths at 37°C. Approximately 3×10^5 cells were then harvested, washed with serum-free DMEM culture medium, and stained with 1 μ M DCFH-DA for 30 min at 37°C in the dark. After the cells were centrifuged, washed, and resuspended in serum-free DMEM culture medium three times, the fluorescence intensity of each sample was determined with a flow cytometer.

Mitochondrial membrane potential ($\Delta\Psi_m$) assay. The fluorescent dye, 5,5',6,6'-tetrachloro-1,1',3,3'-tetraethylbenzimidazolcarbocyanine iodide (JC-1; Molecular Probes, Eugene, OR, USA), was used to detect $\Delta\Psi_m$. Briefly, after a heat stress treatment, HUVECs were further incubated at 37°C for various periods of time as indicated. After the cells were washed three times with PBS, a JC-1 kit and a fluorescence microscope were used to detect the mitochondrial cross membrane potential.

Statistical analysis. All data were analyzed for statistical significance using SPSS 13.0 software (SPSS, USA). Data are expressed as the mean \pm standard deviation (SD) from at least three independent experiments that were performed in duplicate. Statistical comparisons of the results were made using analysis of one-way ANOVA. * $P < 0.05$, ** $P < 0.01$, and *** $P < 0.001$ were considered statistically significant.

References

- Bouchama, A. & Knochel, J. P. Heat stroke. *N Engl J Med* **346**, 1978–1988 (2002).
- Hemon, D. & Jouglu, E. [The heat wave in France in August 2003]. *Rev Epidemiol Sante Publique* **52**, 3–5 (2004).
- Robine, J. M. *et al.* Death toll exceeded 70,000 in Europe during the summer of 2003. *C R Biol* **331**, 171–178 (2008).
- Roberts, G. T. *et al.* Microvascular injury, thrombosis, inflammation, and apoptosis in the pathogenesis of heatstroke: a study in baboon model. *Arterioscler Thromb Vasc Biol* **28**, 1130–1136 (2008).
- Sakaguchi, Y. *et al.* Apoptosis in tumors and normal tissues induced by whole body hyperthermia in rats. *Cancer Res* **55**, 5459–5464 (1995).
- Bouchama, A., Hammami, M. M., Haq, A., Jackson, J. & al-Sedairy, S. Evidence for endothelial cell activation/injury in heatstroke. *Crit Care Med* **24**, 1173–1178 (1996).
- Chao, T. C., Sinniah, R. & Pakiam, J. E. Acute heat stroke deaths. *Pathology* **13**, 145–156 (1981).

8. Sohal, R. S., Sun, S. C., Colcolough, H. L. & Burch, G. E. Heat stroke. An electron microscopic study of endothelial cell damage and disseminated intravascular coagulation. *Arch Intern Med* **122**, 43–47 (1968).
9. Brinton, M. R. *et al.* Thermal sensitivity of endothelial cells on synthetic vascular graft material. International journal of hyperthermia : the official journal of European Society for Hyperthermic Oncology, *Int J Hyperthermia* **28**, 163–174 (2012).
10. Gu, Z. T. *et al.* Heat stress induces apoptosis through transcription-independent p53-mediated mitochondrial pathways in human umbilical vein endothelial cell. *Sci Rep* **4**, 4469 (2014).
11. Baldwin, A. S. Control of oncogenesis and cancer therapy resistance by the transcription factor NF-kappaB. *J Clin Invest* **107**, 241–246 (2001).
12. Pahl, H. L. Activators and target genes of Rel/NF-kappaB transcription factors. *Oncogene* **18**, 6853–6866 (1999).
13. Israel, A. The IKK complex, a central regulator of NF-kappaB activation. *Cold Spring Harb Perspect Biol* **2**, a000158 (2010).
14. Monks, N. R., Biswas, D. K. & Pardee, A. B. Blocking anti-apoptosis as a strategy for cancer chemotherapy: NF-kappaB as a target. *J Cell Biochem* **92**, 646–650 (2004).
15. Tang, G. *et al.* Inhibition of JNK activation through NF-kappaB target genes. *Nature* **414**, 313–317 (2001).
16. De Smaele, E. *et al.* Induction of gadd45beta by NF-kappaB downregulates pro-apoptotic JNK signalling. *Nature* **414**, 308–313 (2001).
17. Fiers, W., Beyaert, R., Declercq, W. & Vandenabeele, P. More than one way to die: apoptosis, necrosis and reactive oxygen damage. *Oncogene* **18**, 7719–7730 (1999).
18. Sakon, S. *et al.* NF-kappaB inhibits TNF-induced accumulation of ROS that mediate prolonged MAPK activation and necrotic cell death. *EMBO J* **22**, 3898–3909 (2003).
19. Fulda, S., Gorman, A. M., Hori, O. & Samali, A. Cellular stress responses: cell survival and cell death. *Int J Cell Biol* **2010**, 214074 (2010).
20. Park, K. J., Gaynor, R. B. & Kwak, Y. T. Heat shock protein 27 association with the I kappa B kinase complex regulates tumor necrosis factor alpha-induced NF-kappa B activation. *The Journal of biological chemistry* **278**, 35272–35278 (2003).
21. Shi, Y. *et al.* The inhibition of LPS-induced production of inflammatory cytokines by HSP70 involves inactivation of the NF-kappaB pathway but not the MAPK pathways. *Shock (Augusta, Ga.)* **26**, 277–284 (2006).
22. Chen, Y., Voegeli, T. S., Liu, P. P., Noble, E. G. & Currie, R. W. Heat shock paradox and a new role of heat shock proteins and their receptors as anti-inflammation targets. *Inflammation & allergy drug targets* **6**, 91–100 (2007).
23. Weiss, Y. G. *et al.* Enhanced heat shock protein 70 expression alters proteasomal degradation of IkappaB kinase in experimental acute respiratory distress syndrome. *Critical care medicine* **35**, 2128–2138 (2007).
24. Chen, Y. & Currie, R. W. Small interfering RNA knocks down heat shock factor-1 (HSF-1) and exacerbates pro-inflammatory activation of NF-kappaB and AP-1 in vascular smooth muscle cells. *Cardiovascular research* **69**, 66–75 (2006).
25. Kammanadiminti, S. J. & Chadee, K. Suppression of NF-kappaB activation by Entamoeba histolytica in intestinal epithelial cells is mediated by heat shock protein 27. *The Journal of biological chemistry* **281**, 26112–26120 (2006).
26. Sur, R., Lyte, P. A. & Southall, M. D. Hsp27 regulates pro-inflammatory mediator release in keratinocytes by modulating NF-kappaB signaling. *The Journal of investigative dermatology* **128**, 1116–1122 (2008).
27. Parcellier, A. *et al.* HSP27 is a ubiquitin-binding protein involved in I-kappaBalpha proteasomal degradation. *Mol Cell Biol* **23**, 5790–5802 (2003).
28. Kretz-Remy, C., Munsch, B. & Arrigo, A. P. NFkappa B-dependent transcriptional activation during heat shock recovery. Thermolability of the NF-kappaB.Ikappa B complex. *J Biol Chem* **276**, 43723–43733 (2001).
29. Park, C. H. *et al.* Heat shock-induced matrix metalloproteinase (MMP)-1 and MMP-3 are mediated through ERK and JNK activation and via an autocrine interleukin-6 loop. *The Journal of investigative dermatology* **123**, 1012–1019 (2004).
30. Steinbrecher, K. A., Wilson, W., 3rd, Cogswell, P. C. & Baldwin, A. S. Glycogen synthase kinase 3beta functions to specify gene-specific, NF-kappaB-dependent transcription. *Molecular and cellular biology* **25**, 8444–8455 (2005).
31. Chen, G. *et al.* Phosphorylated FADD induces NF-kappaB, perturbs cell cycle, and is associated with poor outcome in lung adenocarcinomas. *Proceedings of the National Academy of Sciences of the United States of America* **102**, 12507–12512 (2005).
32. Loktionova, S. A., Ilyinskaya, O. P., Gabai, V. L. & Kabakov, A. E. Distinct effects of heat shock and ATP depletion on distribution and isoform patterns of human Hsp27 in endothelial cells. *FEBS letters* **392**, 100–104 (1996).
33. Yu, J. *et al.* Involvement of oxidative stress and mitogen-activated protein kinase signaling pathways in heat stress-induced injury in the rat small intestine. *Stress* **16**, 99–113 (2013).
34. Kretz-Remy, C. & Arrigo, A. P. The kinetics of HIV-1 long terminal repeat transcriptional activation resemble those of hsp70 promoter in heat-shock treated HeLa cells. *FEBS letters* **351**, 191–196 (1994).
35. Schreck, R., Rieber, P. & Baeuerle, P. A. Reactive oxygen intermediates as apparently widely used messengers in the activation of the NF-kappa B transcription factor and HIV-1. *The EMBO journal* **10**, 2247–2258 (1991).
36. Bender, K., Gottlicher, M., Whiteside, S., Rahmsdorf, H. J. & Herrlich, P. Sequential DNA damage-independent and -dependent activation of NF-kappaB by UV. *EMBO J* **17**, 5170–5181 (1998).
37. Li, N. & Karin, M. Ionizing radiation and short wavelength UV activate NF-kappaB through two distinct mechanisms. *Proc Natl Acad Sci USA* **95**, 13012–13017 (1998).
38. Imbert, V. *et al.* Tyrosine phosphorylation of I kappa B-alpha activates NF-kappa B without proteolytic degradation of I kappa B-alpha. *Cell* **86**, 787–798 (1996).
39. Douillette, A. *et al.* The proinflammatory actions of angiotensin II are dependent on p65 phosphorylation by the IkappaB kinase complex. *J Biol Chem* **281**, 13275–13284 (2006).
40. Nivon, M., Richet, E., Codogno, P., Arrigo, A. P. & Kretz-Remy, C. Autophagy activation by NFkappaB is essential for cell survival after heat shock. *Autophagy* **5**, 766–783 (2009).
41. Gorska, M. M. *et al.* MK2 controls the level of negative feedback in the NF-kappaB pathway and is essential for vascular permeability and airway inflammation. *The Journal of experimental medicine* **204**, 1637–1652 (2007).
42. Lee, S. J., Yang, E. S., Kim, S. Y., Shin, S. W. & Park, J. W. Regulation of heat shock-induced apoptosis by sensitive to apoptosis gene protein. *Free Radic Biol Med* **45**, 167–176 (2008).
43. Xia, Z., Dickens, M., Raingeaud, J., Davis, R. J. & Greenberg, M. E. Opposing effects of ERK and JNK-p38 MAP kinases on apoptosis. *Science* **270**, 1326–1331 (1995).
44. Gomes, D. A. *et al.* c-Met must translocate to the nucleus to initiate calcium signals. *J Biol Chem* **283**, 4344–4351 (2008).
45. Pallepati, P. & Averill-Bates, D. Mild thermotolerance induced at 40 degrees C increases antioxidants and protects HeLa cells against mitochondrial apoptosis induced by hydrogen peroxide: Role of p53. *Arch Biochem Biophys* **495**, 97–111 (2010).

Acknowledgments

This study was supported by National Natural Science Foundation of China (NO. 81471839), the project team of Natural Science Foundation of Guangdong Province (s2013030013217).

Author Contributions

Y.N.L., Q.B.H. and Q.L.X. performed the study and composed this manuscript. Y.N.L. was responsible for primary data generation and analysis. X.H.G., G.B.Z. and Z.L.W. participated in measurements of ROS and caspase activity. Y.N.L. performed apoptosis test and Weston blots. L.S., Q.B.H. and Q.L.X. are the principal investigator and corresponding author for these studies. All authors reviewed the manuscript.

Additional Information

Competing financial interests: The authors declare no competing financial interests.

How to cite this article: Liu, Y. *et al.* NF- κ B signaling is essential for resistance to heat stress-induced early stage apoptosis in human umbilical vein endothelial cells. *Sci. Rep.* **5**, 13547; doi: 10.1038/srep13547 (2015).



This work is licensed under a Creative Commons Attribution 4.0 International License. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in the credit line; if the material is not included under the Creative Commons license, users will need to obtain permission from the license holder to reproduce the material. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>