

# Computable Embeddings of Classes of Structures Under Enumeration and Turing Operators

I. Sh. Kalimullin\*

(Submitted by M. M. Arslanov)

*N.I. Lobachevskii Institute of Mathematics and Mechanics,  
Kazan (Volga Region) Federal University, ul. Kremlevskaya 18, Kazan, Tatarstan, 420008 Russia*

Received April 4, 2017

**Abstract**—In the paper we study the differences and partial characterizations of the Turing and enumeration computable embeddings of classes of structures

**DOI:** 10.1134/S1995080218010146

Keywords and phrases: *Erhsov's hierarchy, Turing degrees, enumeration degrees, elementary theories, structural properties.*

## 1. INTRODUCTION

In the papers [1] and [2] the following two notions were introduced as a computability analog of Borel embedding.

**Definition 1.** Let  $K_0, K_1$  be classes of structures in finite languages (for each class the language is the same).

1) We say that  $K_0$  is *computably embeddable via an e-operator* into  $K_1$  (and write  $K_0 \leq_c K_1$ ) iff there are a function  $f : K_0 \rightarrow K_1$  and an e-operator  $\Phi$  such that  $D(f(\mathcal{A})) = \Phi(D(\mathcal{A}))$  for any  $\mathcal{A} \in K_0$  and and for any  $\mathcal{A}_1, \mathcal{A}_2 \in K_0$

$$\mathcal{A}_0 \cong \mathcal{A}_1 \Leftrightarrow f(\mathcal{A}_0) \cong f(\mathcal{A}_1).$$

2) We say that  $K_0$  is *computably embeddable via an Turing operator* into  $K_1$  (and write  $K_0 \leq_{tc} K_1$ ) iff there are a function  $f : K_0 \rightarrow K_1$  and a Turing operator  $\varphi_e$  such that  $\chi_{D(f(\mathcal{A}))} = \varphi_e^{D(\mathcal{A})}$  for any  $\mathcal{A} \in K_0$  and and for any  $\mathcal{A}_1, \mathcal{A}_2 \in K_0$

$$\mathcal{A}_0 \cong \mathcal{A}_1 \Leftrightarrow f(\mathcal{A}_0) \cong f(\mathcal{A}_1).$$

It follows from the next proposition then  $K_0 \leq_c K_1$  implies  $K_0 \leq_{tc} K_1$ .

**Proposition 2.**  $K_0 \leq_{tc} K_1$  iff there are a function  $f : K_0 \rightarrow K_1$  and an integer  $e \in \omega$  such that  $D(f(\mathcal{A})) = W_e^{D(\mathcal{A})}$  for any  $\mathcal{A} \in K_0$  and for any  $\mathcal{A}_1, \mathcal{A}_2 \in K_0$

$$\mathcal{A}_0 \cong \mathcal{A}_1 \Leftrightarrow f(\mathcal{A}_0) \cong f(\mathcal{A}_1).$$

*Proof.* ( $\implies$ ) Obvious.

( $\impliedby$ ) Without loss of generality we can assume that  $\text{card}(W_{e,s+1}^X - W_{e,s}^X) \leq 1$  for all  $s$  and  $X$ . We denote via  $T(a)$  the atomic sentence  $a = a$  for each  $a \in \omega$ .

Suppose that  $\mathcal{A} \in K_0$  is given. Define

$$S = \{s \in \omega : (\exists a)[T(a) \in W_{e,s+1}^{D(\mathcal{A})} - W_{e,s}^{D(\mathcal{A})}]\}$$

\*E-mail: ikalimul@gmail.com

and  $T(a_s) \in W_{e,s+1}^{D(\mathcal{A})} - W_{e,s}^{D(\mathcal{A})}$  for each  $s \in S$ . Let  $\mathcal{S}_{\mathcal{A}}$  be the structure with universe  $S$  such that  $\mathcal{S}_{\mathcal{A}} \cong \Phi(\mathcal{A})$  via the isomorphism  $s \mapsto a_s$ . It is easy to see that there is an index  $i$  such that  $D(\mathcal{S}_{\mathcal{A}}) = \varphi_i^{D(\mathcal{A})}$  for each  $\mathcal{A} \in K_0$ .  $\square$

**Corollary 3.** *If  $K_0 \leq_c K_1$  then  $K_0 \leq_{tc} K_1$ .*

To see that the reverse implication is not true we can note that for the case when  $K_0 = \{\mathcal{X} : \mathcal{X} \cong \mathcal{A}\}$  and  $K_1 = \{\mathcal{X} : \mathcal{X} \cong \mathcal{B}\}$  the embedding  $K_0 \leq_{tc} K_1$  is equivalent to the Medvedev reducibility (the uniform Turing reducibility)  $\mathcal{B} \leq_{uT} \mathcal{A}$ , and the embedding  $K_0 \leq_c K_1$  is equivalent to  $\mathcal{B} \leq_{ue} \mathcal{A}$  (the uniform enumeration reducibility). It follows from [3] that there are structures  $\mathcal{A}$  and  $\mathcal{B}$  such that  $\mathcal{B} \leq_{uT} \mathcal{A}$  and  $\mathcal{B} \not\leq_{ue} \mathcal{A}$ . But such structures  $\mathcal{A}$  and  $\mathcal{B}$  should not have a computable presentation, while the embeddings  $\leq_c$  and  $\leq_{tc}$  are more interesting for the classes of computable and even finite structures.

## 2. EMBEDDING OF CLASSES WITH FINITELY MANY ISOMORPHIC TYPES

The following two theorem give a full descriptions of  $c$ - and  $tc$ -embeddings of classes with finitely many isomorphic types. These will give an easy example of classes of computable structures for which the  $c$ - and  $tc$ -embeddings differ.

**Theorem 4.** *Let a class of finite structures  $K_0$  contains only finitely many different isomorphic types. Then  $K_0 \leq_c K_1$  iff there is a function  $f$  from  $K_0$  into a subclass of  $K_1$  containing only computable structures such that for any  $\mathcal{A}, \mathcal{B} \in K_0$   $f(\mathcal{A}) \cong f(\mathcal{B}) \implies \mathcal{A} \cong \mathcal{B}$ , and*

$$\mathcal{A} \text{ is embeddable into } \mathcal{B} \implies f(\mathcal{A}) \subseteq f(\mathcal{B}).$$

**Theorem 5.** *Let a class of finite structures  $K_0$  contains only finitely many different isomorphic types. Then  $K_0 \leq_{tc} K_1$  iff there is a function  $f$  from  $K_0$  into a subclass of  $K_1$  containing only computable structures such that for any  $\mathcal{A}, \mathcal{B} \in K_0$*

$$\mathcal{A} \cong \mathcal{B} \Leftrightarrow f(\mathcal{A}) \cong f(\mathcal{B}) \Leftrightarrow f(\mathcal{A}) = f(\mathcal{B}), \text{ and}$$

$$\mathcal{A} \text{ is embeddable into } \mathcal{B} \implies Th_{\exists}(f(\mathcal{A})) \subseteq Th_{\exists}(f(\mathcal{B})).$$

**Corollary 6.** *Let a class of finite structures  $K_0$  contains only finitely many different isomorphic types. Then for all classes of finite structures  $K_1$  we have  $K_0 \leq_c K_1 \Leftrightarrow K_0 \leq_{tc} K_1$ .*

**Corollary 7.** *There are classes  $K_0$  and  $K_1$  such that  $K_0 \leq_{tc} K_1$  and  $K_0 \not\leq_c K_1$ .*

*Proof of Corollary 7.* Let  $K_0$  consists from the empty linear ordering and all one-element orderings, and let  $K_1$  consists from all linear orderings isomorphic either to  $\omega$ , or to  $\omega^*$ . By Theorem 4  $K_0 \not\leq_c K_1$ . Since all infinite linear orderings have one existensial theory we have  $K_0 \leq_{tc} K_1$  by Theorem 5.  $\square$

*Proof of Theorem 4.* ( $\implies$ ) Let  $K_0 \leq_c K_1$  via an e-operator  $\Phi$ . Since  $K_0$  contains only finitely many different isomorphic types there is a finite collection  $I_0$  of structures from  $K_0$  such that any structure from  $K_0$  has an isomorphic copy in  $I_0$  and such that from any  $\mathcal{A}_0, \mathcal{A}_1 \in K_0$  and any  $\mathcal{C}_0, \mathcal{C}_1 \in I_0$

$$\mathcal{A}_0 \cong \mathcal{C}_0 \ \& \ \mathcal{A}_1 \cong \mathcal{C}_1 \ \& \ \mathcal{A}_0 \text{ is embeddable into } \mathcal{A}_1 \implies \mathcal{C}_0 \subseteq \mathcal{C}_1.$$

For any  $\mathcal{A} \in K_0$  we define  $f(\mathcal{A})$  as the structure from  $K_1$  such that  $D(f(\mathcal{A})) = \Phi(D(\mathcal{C}_{\mathcal{A}}))$ , where  $\mathcal{C}_{\mathcal{A}} \in I_0$  and  $\mathcal{C}_{\mathcal{A}} \cong \mathcal{A}$ .

( $\impliedby$ ) Suppose that such function  $f : K_0 \rightarrow K_1$  exists. We define an e-operator  $\Phi$  via the c.e. set of all axioms  $\langle \varphi, D(\mathcal{A}) \rangle$ , where  $\mathcal{A} \in K_0$  and  $\varphi \in D(f(\mathcal{A}))$ . Then  $K_0 \leq_c K_1$  via the e-operator  $\Phi$ .  $\square$

*Proof of Theorem 5.* ( $\implies$ ) Let  $K_0 \leq_{tc} K_1$  via a Turing operator  $\varphi_e$  and let  $I_0$  be as in the proof of Theorem 4. For any  $\mathcal{A} \in K_0$  we define  $f(\mathcal{A})$  as the structure from  $K_1$  such that  $D(f(\mathcal{A})) = \varphi_e^{D(\mathcal{C}_{\mathcal{A}})}$ , where  $\mathcal{C}_{\mathcal{A}} \in I_0$  and  $\mathcal{C}_{\mathcal{A}} \cong \mathcal{A}$ . Then the implications  $\mathcal{A} \cong \mathcal{B} \Leftrightarrow f(\mathcal{A}) \cong f(\mathcal{B}) \Leftrightarrow f(\mathcal{A}) = f(\mathcal{B})$  are obvious. Suppose that  $\mathcal{A} \in K_0$  is embeddable into  $\mathcal{B} \in K_0$  and  $f(\mathcal{A}) \models \theta$  for some existensial sentence  $\theta$ . Let  $\mathcal{F}$  be a finite substructure of  $f(\mathcal{A})$  such that  $\mathcal{F} \models \theta$  and let  $n$  be such integer that  $\varphi_e^{D(\mathcal{C}_{\mathcal{A}}) \upharpoonright n}(\psi) \downarrow = 1$  for all  $\psi \in D(\mathcal{F})$ . Since  $\mathcal{A}$  is embeddable into  $\mathcal{B}$  there is a structure  $\mathcal{B}' \cong \mathcal{B}$  such that  $\psi \in D(\mathcal{C}_{\mathcal{A}}) \Leftrightarrow \psi \in D(\mathcal{B}')$  for all atomic sentences with code  $< n$ . Then  $\varphi_e^{D(\mathcal{B}')}(\psi) = 1$  for all  $\psi \in D(\mathcal{F})$  so that  $\mathcal{F} \subseteq \mathcal{B}'$  and hence  $f(\mathcal{B}) \models \theta$ .

( $\Leftarrow$ ) Let  $f : K_0 \rightarrow K_1$  be such function. For any  $\mathcal{C} \in I_0$  and any finite consistent set of atomic sentences  $\Delta$  in the language of class  $K_1$  we denote via  $\mathcal{E}_\Delta^{\mathcal{C}}$  a structure isomorphic to  $f(\mathcal{C})$  such that  $\Delta \subseteq D(\mathcal{E}_\Delta^{\mathcal{C}})$ .

Since  $f(\mathcal{C})$  is always a computable structure, for each  $\mathcal{C} \in I_0$  the correspondence  $\Delta \mapsto \mathcal{E}_\Delta^{\mathcal{C}}$  can be chosen partial computable in the sense that knowing a canonical index of a finite set  $\Delta$  we can effectively determine membership of any atomic sentence  $\psi$  in  $D(\mathcal{E}_\Delta^{\mathcal{C}})$  if  $\mathcal{E}_\Delta^{\mathcal{C}}$  exists, and the last condition is c.e.

Let  $\mathcal{A} \in K_0$  be given. Let  $s_0 < s_1 < s_2 < \dots$  be all integers  $s \in \omega$  such that for some  $\mathcal{A}' \in K_0$  we have  $D(\mathcal{A}) \upharpoonright s = D(\mathcal{A}')$ . Then for any  $n \in \omega$  we denote via  $\mathcal{C}_n$  the structure from the finite collection  $I_0$  such that for some  $\mathcal{A}' \cong \mathcal{C}_n$  we have  $D(\mathcal{A}) \upharpoonright s_n = D(\mathcal{A}')$ . Note that  $\mathcal{C}_n \subseteq \mathcal{C}_{n+1}$  for each  $n \in \omega$ .

Now we inductively construct a sequence  $\{\Delta_n\}_{n \in \omega}$  of finite consistent sets of atomic sentences in the language of class  $K_1$ :

$$\Delta_0 = \emptyset,$$

$$\Delta_{n+1} = D(\mathcal{E}_{\Delta_m}^{\mathcal{C}_{n+1}}) \upharpoonright n, \text{ where } m \leq n \text{ is the least integer such that } \mathcal{C}_k = \mathcal{C}_{n+1} \text{ for all } k, m < k \leq n.$$

Note that  $\mathcal{E}_{\Delta_m}^{\mathcal{C}_{n+1}}$  always exists since  $\mathcal{C}_m \subseteq \mathcal{C}_{n+1}$  and hence  $Th_{\exists}(f(\mathcal{C}_m)) \subseteq Th_{\exists}(f(\mathcal{C}_{n+1}))$ . Moreover,  $\bigcup_n \Delta_n = D(\mathcal{B})$  for some  $\mathcal{B} \cong f(\mathcal{A})$ . It remains to note that  $D(\mathcal{B}) = W_e^{D(\mathcal{A})}$  for some  $e$  and apply Proposition 2.  $\square$

### 3. EMBEDDING OF CLASSES WITH FINITELY MANY ISOMORPHIC TYPES

The following two theorem give a full descriptions of  $c$ - and  $tc$ -embeddings of classes with finitely many isomorphic types. These will give an example of classes of finite structures for which the  $c$ - and  $tc$ -embeddings differ.

**Theorem 8.** *FLO  $\leq_c$  K iff there is a computable Friedberg numbering  $\{\mathcal{X}_n\}_{n \in \omega}$  of a subclass of K such that  $\mathcal{X}_n \subseteq \mathcal{X}_{n+1}$  for each n.*

**Theorem 9.** *FLO  $\leq_{tc}$  K iff there is a computable Friedberg numbering  $\{\mathcal{X}_n\}_{n \in \omega}$  of a subclass of K such that  $Th_{\exists}(\mathcal{X}_n) \subseteq Th_{\exists}(\mathcal{X}_{n+1})$  for each n.*

**Theorem 10.** *There is a class K of undirected finite graphs with such that*

a) *there is a computable Friedberg numbering  $\{\mathcal{X}_n\}_{n \in \omega}$  of the class K such that each graph  $\mathcal{X}_n$ ,  $n \in \omega$ , is embeddable into the graph  $\mathcal{X}_{n+1}$ , and*

b) *there is no computable Friedberg numbering  $\{\mathcal{Y}_n\}_{n \in \omega}$  of a subclass of K such that  $\mathcal{Y}_n \subseteq \mathcal{Y}_{n+1}$  for each n.*

**Corollary 11.** *There is a class K of undirected finite graphs such that FLO  $\leq_{tc}$  K and FLO  $\not\leq_c$  K.*

*Proof of Theorem 8.* ( $\Rightarrow$ ) Suppose that FLO  $\leq_c$  K via an e-operator  $\Phi$ . Let  $\mathcal{L}_n$  be the standard linear ordering of natural numbers  $< n$ . Then for each  $n \in \omega$   $\Phi(D(\mathcal{L}_n)) = D(\mathcal{X}_n)$  for some  $\mathcal{X}_n \in K$ . It is easy to check that  $\{\mathcal{X}_n\}_{n \in \omega}$  is the computable Friedberg numbering such that  $\mathcal{X}_n \subseteq \mathcal{X}_{n+1}$  for each  $n$ .

( $\Leftarrow$ ) Let there exists such computable Friedberg numbering  $\{\mathcal{X}_n\}_{n \in \omega}$ . We define an e-operator  $\Phi$  via the c.e. set of all axioms  $\langle \varphi, D(\mathcal{A}) \rangle$ , where  $\mathcal{A}$  is a linear ordering with  $n$  elements and  $\varphi \in D(\mathcal{X}_n)$ ,  $n \in \omega$ . Then FLO  $\leq_c$  K via the e-operator  $\Phi$ .  $\square$

*Proof of Theorem 9.* ( $\Rightarrow$ ) Let FLO  $\leq_{tc}$  K via a Turing operator  $\varphi_e$  and let  $\mathcal{L}_n$  be the standard linear ordering of natural numbers  $< n$ . For any  $n \in \omega$  we define  $\mathcal{X}_n$  as the structure from K such that  $D(\mathcal{X}_n) = \varphi_e^{D(\mathcal{L}_n)}$ . It is easy to see that  $\{\mathcal{X}_n\}_{n \in \omega}$  is the computable Friedberg numbering.

We prove that  $Th_{\exists}(\mathcal{X}_n) \subseteq Th_{\exists}(\mathcal{X}_{n+1})$  for each  $n$ . Let  $\mathcal{X}_n \models \theta$  for some existensial sentence  $\theta$ . Let  $\mathcal{F}$  be a finite substructure of  $\mathcal{X}_n$  such that  $\mathcal{F} \models \theta$  and let  $k$  be such integer that  $\varphi_e^{D(\mathcal{L}_n) \upharpoonright k}(\psi) \downarrow = 1$  for all  $\psi \in D(\mathcal{F})$ . We choose a linear ordering  $\mathcal{L}'_{n+1}$  with  $n+1$  elements such that  $\psi \in D(\mathcal{L}_n) \Leftrightarrow \psi \in D(\mathcal{L}'_{n+1})$  for all atomic sentences with code  $< k$ . Then  $\varphi_e^{D(\mathcal{L}'_{n+1})}(\psi) = 1$  for all  $\psi \in D(\mathcal{F})$  and hence  $\mathcal{X}_{n+1} \models \theta$ .

( $\Leftarrow$ ) Let there exists such computable Friedberg numbering  $\{\mathcal{X}_n\}_{n \in \omega}$ . For any  $n \in \omega$  and any finite consistent set of atomic sentences  $\Delta$  in the language of class K we denote via  $\mathcal{E}_\Delta^n$  a structure isomorphic

to  $\mathcal{X}_n$  such that  $\Delta \subseteq D(\mathcal{E}_\Delta^n)$ . As in the proof of Theorem 5 we can choose a partially computable correspondence  $(n, \Delta) \mapsto \mathcal{E}_\Delta^n$ .

Let a finite linear ordering  $\mathcal{L}$  be given. Let  $s_0 < s_1 < s_2 < \dots$  be all integers  $s \in \omega$  such that  $D(\mathcal{L}) \upharpoonright s$  is a diagram of some linear ordering. We denote via  $c(n)$  the number of elements in the linear ordering with the diagram  $D(\mathcal{L}) \upharpoonright s_n$ . Note that  $c(n) \leq c(n+1)$  for each  $n \in \omega$ .

We inductively construct a sequence  $\{\Delta_n\}_{n \in \omega}$  of finite consistent sets of atomic sentences in the language of class  $K$ :

$$\Delta_0 = \emptyset,$$

$$\Delta_{n+1} = D(\mathcal{E}_{\Delta_m}^{c(n+1)}) \upharpoonright n, \text{ where } m \leq s \text{ is the least integer such that } c(k) = c(n+1) \text{ for all } k, m < k \leq n.$$

Note that  $\mathcal{E}_{\Delta_m}^{c(n+1)}$  exists since  $Th_{\exists}(\mathcal{X}_{c(m)}) \subseteq Th_{\exists}(\mathcal{X}_{c(n+1)})$ . Moreover,  $\bigcup_s \Delta_s = D(\mathcal{B})$  for some  $\mathcal{B} \cong \mathcal{X}_{card(\mathcal{L})}$ . It remains to note that  $D(\mathcal{B}) = W_e^{D(\mathcal{L})}$  for some  $e$  and apply Proposition 2.  $\square$

*Proof of Corollary 11.* Let the language of undirected graphs contain one binary predicate  $R$  ( $R(a, b)$  means that vertices  $a$  and  $b$  are connected by an arc). For each  $m \geq 4$  define the following finite undirected graphs:

$\mathcal{A}_m$  is the graph with vertices  $a_1, \dots, a_{m+1}$

and with arcs  $\{a_1, a_2\}, \{a_2, a_3\}, \{a_3, a_4\}, \dots, \{a_{m-1}, a_m\}, \{a_m, a_1\}, \{a_m, a_{m+1}\}$ ;

$\mathcal{B}_m$  is the graph with vertices  $a_1, \dots, a_{m+2}$

and with arcs  $\{a_1, a_2\}, \{a_2, a_3\}, \{a_3, a_4\}, \dots, \{a_{m-1}, a_m\}, \{a_m, a_1\}, \{a_m, a_{m+1}\}, \{a_{m+1}, a_{m+2}\}$ ;

$\mathcal{C}_m$  is the graph with vertices  $a_1, \dots, a_{m+2}$  and with arcs

$\{a_1, a_2\}, \{a_2, a_3\}, \{a_3, a_4\}, \dots, \{a_{m-1}, a_m\}, \{a_m, a_1\}, \{a_m, a_{m+1}\}, \{a_{m+1}, a_{m+2}\}, \{a_m, a_{m+2}\}$ ;

$\mathcal{D}_m$  is the graph with vertices  $a_1, \dots, a_{m+4}$  and with arcs

$\{a_1, a_2\}, \{a_2, a_3\}, \{a_3, a_4\}, \dots, \{a_{m-1}, a_m\}, \{a_m, a_1\}, \{a_m, a_{m+1}\}, \{a_{m+1}, a_{m+2}\}, \{a_m, a_{m+2}\}, \{a_m, a_{m+3}\}, \{a_{m+3}, a_{m+4}\}$ .

Then for each  $i, j \in \omega$  we set  $\mathcal{F}_{i,j} = \mathcal{C}_{\langle i,j \rangle + 4}$  and  $\mathcal{G}_{i,j} = \mathcal{D}_{\langle i,j \rangle + 4}$  if there are integers  $x_1, \dots, x_{\langle i,j \rangle + 6}$  such that

- 1)  $\langle i, R(x_k, x_{k+1}) \rangle \in W_i$  for each  $k, 1 \leq k \leq \langle i, j \rangle + 3$ , and
- 2)  $\langle i, R(x_1, x_{\langle i,j \rangle + 4}) \rangle \in W_i$ ;
- 3)  $\langle i, R(x_{\langle i,j \rangle + 4}, x_{\langle i,j \rangle + 5}) \rangle \in W_i$ ;
- 4)  $\langle i + 1, R(x_{\langle i,j \rangle + 5}, x_{\langle i,j \rangle + 6}) \rangle \in W_i$  and  $\langle i + 1, \neg R(x_{\langle i,j \rangle + 4}, x_{\langle i,j \rangle + 6}) \rangle \in W_i$ .

Otherwise we set  $\mathcal{F}_{i,j} = \mathcal{A}_{\langle i,j \rangle + 4}$  and  $\mathcal{G}_{i,j} = \mathcal{B}_{\langle i,j \rangle + 4}$ .

Let finite undirected graph  $\mathcal{H}_n, n \in \omega$ , be the disjoint and disconnected union of all graphs  $\mathcal{F}_{i,n}, i \leq n$ , and all graphs  $\mathcal{G}_{i,j}, i \leq j < n$ . It is easy to see that for all  $n$  the graph  $\mathcal{H}_n$  is finite and  $\mathcal{H}_n$  is embeddable into  $\mathcal{H}_{n+1}$  since each  $\mathcal{A}_m$  is embeddable into  $\mathcal{B}_m$  and each  $\mathcal{C}_m$  is embeddable into  $\mathcal{D}_m$ . Moreover, there is a computable Friedberg numbering of finite undirected graphs  $\{\mathcal{X}_n\}_{n \in \omega}$  such that  $\mathcal{X}_n \cong \mathcal{H}_n$  for each  $n$  since the conditions 1)–4) are  $\Sigma_1$  and since each  $\mathcal{A}_m$  is embeddable into  $\mathcal{C}_m$  and each  $\mathcal{B}_m$  is embeddable into  $\mathcal{D}_m$ .

Let  $K$  be the class containing all graphs isomorphic to  $\mathcal{H}_n$  for some  $n \in \omega$ . Then  $K$  satisfies the condition a) of the theorem. Suppose that there is a computable Friedberg numbering  $\{\mathcal{Y}_n\}_{n \in \omega}$  of a subclass of  $K$  such that  $\mathcal{Y}_n \subseteq \mathcal{Y}_{n+1}$  for each  $n$ . Then there is a c.e. set  $W_i$  of pairs  $\langle n, \varphi \rangle$ , where  $n \in \omega$ ,  $\varphi$  is either atomic sentence, or its negation, such that  $D(\mathcal{Y}_n) = \{\varphi : \langle n, \varphi \rangle \in W_i\}$ .

Since  $\mathcal{Y}_n \subsetneq \mathcal{Y}_{n+1}$  for each  $n$  we have  $\mathcal{Y}_i \cong \mathcal{H}_j$  for some  $j \geq i$  and  $\mathcal{Y}_{i+1} \cong \mathcal{H}_{j'}$  for some  $j' > j$ . Then  $\mathcal{Y}_i$  contains a subgraph  $\mathcal{Y}'_i$  isomorphic to  $\mathcal{F}_{i,j}$  and  $\mathcal{Y}_{i+1}$  contains a subgraph  $\mathcal{Y}'_{i+1}$  isomorphic to  $\mathcal{G}_{i,j}$  such that  $\mathcal{Y}'_i \subseteq \mathcal{Y}'_{i+1}$ .

If  $\mathcal{F}_{i,j} = \mathcal{A}_{\langle i,j \rangle + 4}$  and  $\mathcal{G}_{i,j} = \mathcal{B}_{\langle i,j \rangle + 4}$  then obviously there are integers  $x_1, \dots, x_{\langle i,j \rangle + 5} \in \text{Supp}(\mathcal{Y}'_i)$  and  $x_{\langle i,j \rangle + 6} \in \text{Supp}(\mathcal{Y}'_{i+1})$  satisfying the conditions 1)–4), and hence  $\mathcal{F}_{i,j} = \mathcal{C}_{\langle i,j \rangle + 4}$ ,  $\mathcal{G}_{i,j} = \mathcal{D}_{\langle i,j \rangle + 4}$ , contradiction.

Thus,  $\mathcal{Y}'_i \cong \mathcal{F}_{i,j} = \mathcal{C}_{\langle i,j \rangle+4}$  and  $\mathcal{Y}'_{i+1} \cong \mathcal{G}_{i,j} = \mathcal{D}_{\langle i,j \rangle+4}$ . Hence there are integers  $x_1, \dots, x_{\langle i,j \rangle+6}$  satisfying the conditions 1)-4). Then  $x_1, \dots, x_{\langle i,j \rangle+5} \in \text{Supp}(\mathcal{Y}'_i)$  and  $x_{\langle i,j \rangle+6} \in \text{Supp}(\mathcal{Y}'_{i+1})$ . Since  $\mathcal{Y}'_i \cong \mathcal{C}_{\langle i,j \rangle+4}$ , there is an integer  $y \in \text{Supp}(\mathcal{Y}'_i)$  such that the atomic sentences  $R(x_{\langle i,j \rangle+4}, y)$ ,  $R(x_{\langle i,j \rangle+5}, y)$  belong to  $D(\mathcal{Y}'_i) \subseteq D(\mathcal{Y}'_{i+1})$ . By the condition 4) we have  $R(x_{\langle i,j \rangle+5}, x_{\langle i,j \rangle+6}) \in D(\mathcal{Y}'_{i+1})$  and  $\neg R(x_{\langle i,j \rangle+4}, x_{\langle i,j \rangle+6}) \in D(\mathcal{Y}'_{i+1})$  so that  $y \neq x_{\langle i,j \rangle+6}$ , and we get a contradiction with  $\mathcal{Y}'_{i+1} \cong \mathcal{D}_{\langle i,j \rangle+4}$ .

#### ACKNOWLEDGMENTS

This work was partially supported by the Ministry of Education and Science of the Russian Federation, project no. 1.451.2016/1.4.

#### REFERENCES

1. W. Calvert, D. Cummins, S. Miller, and J. F. Knight, “Comparing classes of finite structures,” *Algebra Logic* **43**, 365–373 (2004).
2. M. V. Boom, S. Miller, and J. F. Knight, “Turing computable embeddings,” *J. Symbolic Logic* **72**, 901–918 (2007).
3. I. Sh. Kalimullin, “Algorithmic reducibilities of algebraic structures,” *J. Logic Comput.* **22**, 831–845 (2012).