



# Weak $\Xi_{QQ} \rightarrow \Sigma_Q \ell^+ \ell^-$ decays induced by FCNC in QCD

T. M. Aliev<sup>1,a</sup>, M. Savci<sup>1,b</sup>, S. Bilmis<sup>1,2,c</sup>

<sup>1</sup> Department of Physics, Middle East Technical University, Ankara 06800, Turkey

<sup>2</sup> TUBITAK ULAKBIM, Ankara 06510, Turkey

Received: 20 June 2022 / Accepted: 25 September 2022 / Published online: 2 October 2022  
© The Author(s) 2022

**Abstract** With the discovery of the doubly heavy  $\Xi_{cc}$  baryon, comprehensive studies of the properties of the doubly heavy baryons are started. In the present work, we examine the  $\Xi_{bb} \rightarrow \Sigma_b \ell^+ \ell^-$  and  $\Xi_{cc} \rightarrow \Sigma_c \ell^+ \ell^-$  decays induced by flavor-changing neutral currents (FCNC) in the framework of the light-cone sum rules. After obtaining the sum rules for the form factors induced by the tensor current, the branching ratios of the relevant transitions are estimated. We found that the branching ratio for the  $c \rightarrow u$  transition is around five orders smaller than the  $b \rightarrow d$  transition. Our findings are also compared with other approaches.

## 1 Introduction

The quark model has been quite successful in the classification of the hadrons. However, up to now, only the hadron state  $\Xi_{cc}^{++}$  has been discovered among all the baryons containing double heavy quarks anticipated by the quark model [1–3]. The detailed analysis to determine the properties of these hadrons is crucial to precisely testing the Standard Model (SM) as well as looking for new physics effects. Weak decays induced by flavor-changing neutral current (FCNC) of doubly heavy baryons are an ideal framework to check SM predictions at the loop level. The new physics effects can manifest themselves in these interactions either by modifying the so-called Wilson coefficients existing in the SM without introducing new operators or by introducing new effective operators.

The observation of the doubly heavy hadrons triggered many theoretical studies on this subject (see [4] and the references therein). In this context, a comprehensive analysis of the weak decays of doubly heavy baryons occupies a spe-

cial place. The main ingredient of weak decays is the transition matrix elements between the initial and final states due to the weak currents of quarks. These matrix elements are parametrized in terms of the form factors. Calculation of the form factors is the main ingredient of studying the weak decays which belong to the non-perturbative domain of the QCD. For this reason, some non-perturbative methods are needed for their calculation. Among various non-perturbative methods, sum rules method that is based on the fundamental QCD Lagrangian occupies an exceptional place. Form factors of some of the doubly heavy baryons due to the charged current are already studied with the traditional and light cone version of the sum rules in [5], and [6,7], respectively. It should be noted here that form factors of the doubly heavy baryons decaying to single heavy baryons are studied in the works [8–11] in the framework of the light-front quark model. Moreover, FCNC processes of the doubly heavy baryons are studied within this approach in [12] and [10]. It should also be noted that the FCNC-induced decay of  $\Xi_{QQ} \rightarrow \Lambda_Q \ell^+ \ell^-$  decay within the light-cone sum rules are studied in [13].

Before we delve into our analysis, we would like to say a few words about the SU(3) classification of the doubly heavy baryons. A doubly heavy baryon contains two heavy and one light quark. Doubly heavy baryons with  $J^P = \frac{1}{2}^+$  in the  $cc$  sector are  $\Xi_{cc}^{++}$ ,  $\Xi_{cc}^+$  and  $\Omega_{cc}^+$ , and those in the  $bb$  sector are  $\Xi_{bb}^0$ ,  $\Xi_{bb}^-$  and  $\Omega_{bb}^-$ . Additionally, there are two sets of baryons in the  $bc$  sector which are antisymmetric or symmetric under the interchange of  $b$  and  $c$  quarks. While the single heavy baryons  $\Lambda_Q$ ,  $\Xi_Q$  belong to the triplet(anti) representation,  $\Sigma_Q$ ,  $\Sigma_Q'$ , and  $\Omega_Q$  baryons lie in the sextet representation of SU(3).

In the present work, we study the  $\Xi_{cc}^+ \rightarrow \Sigma_c^+ \ell^+ \ell^-$ , and  $\Xi_{bb}^0 \rightarrow \Sigma_b^0 \ell^+ \ell^-$  decay in the framework of the light cone QCD sum rules method (LCSR). This method is an extension of the traditional QCD sum rules method [14], and to the light cone [15, 16]. Within the framework of this method,

<sup>a</sup> e-mail: taliev@metu.edu.tr

<sup>b</sup> e-mail: savci@metu.edu.tr

<sup>c</sup> e-mail: sbilmis@metu.edu.tr (corresponding author)

many aspects of the hadron physics are studied (see the review [17]). In the framework of the LCSR method, instead of the local operator product expansion (OPE), the light cone expansion of the non-local operators is used. Moreover, in this method, light-cone distribution amplitudes appear instead of the local condensates, and OPE is performed over twists rather than the dimensions of the local operators.

The paper is organized as follows. In Sect. 2, we derive the sum rules for the relevant form factors for the  $\Sigma_{QQ} \rightarrow \Sigma_Q \ell^+ \ell^-$  decay in framework of the LCSR method. Numerical analysis of these form factors is presented in Sect. 3. In this section, we also estimate the branching ratios of the corresponding decay using these form factors. Conclusions and discussions of the obtained results are presented in the last Sect. 4.

### 2 Sum rules of the transition form factors for the $\Xi_{QQ} \rightarrow \Sigma_Q \ell^+ \ell^-$ decays

The flavor-changing neutral  $b \rightarrow q (d \text{ or } s) \ell^+ \ell^-$  transitions up to mass dimension six is described by the standard weak effective field theory [18]. The effective Hamiltonian for this transition can be written as

$$H_{eff} = -\frac{2G_F \alpha}{\sqrt{2}} \frac{1}{4\pi} V_{tb} V_{tq}^* \sum C_i^{eff}(\mu) \mathcal{O}_i \tag{1}$$

where  $G_F$  is the Fermi constant,  $V_{ij}$  are the elements of Cabibbo–Kobayashi–Maskawa (CKM) matrix elements,  $\alpha$  is the electromagnetic coupling,  $C_i^{eff}(\mu)$  are the short-distance Wilson coefficients and  $\mathcal{O}_i$  are the local effective field operators.

For the decays under consideration, only operators  $\mathcal{O}_7, \mathcal{O}_9, \mathcal{O}_{10}$

$$\begin{aligned} \mathcal{O}_7 &= m_b \bar{q} \sigma_{\mu\nu} (1 + \gamma_5) b F^{\mu\nu} \\ \mathcal{O}_9 &= \bar{q} \gamma_\mu (1 - \gamma_5) b \bar{l} \gamma^\mu l \\ \mathcal{O}_{10} &= \bar{q} \gamma_\mu (1 - \gamma_5) b \bar{l} \gamma^\mu \gamma_5 l \end{aligned} \tag{2}$$

are significant at the scale  $\mu = m_Q$ . It should be noted that the four-quark operators induced by the W-boson exchange (or penguin annihilation) can also contribute to the considered transition. However, these contributions have not been estimated systematically. So-called ‘‘charm-loop effects’’ studied for B-meson decays [19]. These effects might also be important for baryon counterparts. These contributions should be accurately calculated for precise determination of the form factors. However, the effects of these contributions are beyond the scope of this work.

At the quark level, the  $\Xi_{QQ} \rightarrow \Sigma_Q \ell^+ \ell^-$  decays take place through the  $c \rightarrow u$  or  $b \rightarrow d$  transitions. The hadronic matrix elements for the  $\Xi_{cc} \rightarrow \Sigma_c \ell^+ \ell^-$  and  $\Xi_{bb} \rightarrow \Sigma_b \ell^+ \ell^-$  decays are determined by sandwiching the transition currents between the initial and final hadron states.

For example, the matrix element for the  $b \rightarrow q \ell^+ \ell^-$  transition amplitude between the initial and final hadron states can be written

$$\begin{aligned} \mathcal{M} &= \frac{G_F \alpha}{2\sqrt{2}\pi} V_{tb} V_{tq}^* \left[ (C_9^{eff} \langle \Sigma_b | \bar{q} \gamma_\mu (1 - \gamma_5) b | \Xi_{bb} \rangle \right. \\ &\quad \left. - \frac{2m_b}{q^2} C_7^{eff} \langle \Sigma_b | \bar{q} i \sigma_{\mu\nu} q^\nu (1 + \gamma_5) b | \Xi_{bb} \rangle \right) \bar{l} \gamma^\mu l \\ &\quad + C_{10}^{eff} \langle \Sigma_b | \bar{q} \gamma_\mu (1 - \gamma_5) b | \Xi_{bb} \rangle \bar{l} \gamma^\mu \gamma_5 l \Big]. \end{aligned} \tag{3}$$

The effective Wilson coefficient  $C_9^{eff}$  for the  $b \rightarrow q \ell^+ \ell^-$  transition is,

$$\begin{aligned} C_9^{eff} &= \left\{ C_9 + h(m_c, q^2) C_0 \right. \\ &\quad + \lambda_u \left[ h(m_c, q^2) - h(m_u, q^2) \right] (3C_1 + C_2) \\ &\quad - \frac{1}{2} h(m_b, q^2) (4C_3 + 4C_4 + 3C_5 + C_6) \\ &\quad - \frac{1}{2} h(0, q^2) (C_3 + 3C_4) \\ &\quad \left. + \frac{2}{9} (3C_3 + C_4 + 3C_5 + C_6) \right\}, \end{aligned} \tag{4}$$

where,

$$\begin{aligned} \lambda_u &= \frac{V_{ud}^* V_{ub}}{V_{tb}^* V_{tq}}, \text{ and, } C_0 = 3C_1 + C_2 \\ &\quad + 3C_3 + C_4 + 3C_5 + C_6, \tag{5} \\ h(m_q, q^2) &= -\frac{8}{9} \ln \frac{m_q}{m_c} + \frac{8}{27} + \frac{4}{9} x - \frac{2}{9} (2+x) \sqrt{|1-x|} \\ &\quad \times \left[ \Theta(1-x_q) \left( \ln \frac{1+\sqrt{1-x_q}}{1-\sqrt{1-x_q}} - i\pi \right) \right. \\ &\quad \left. + \Theta(x_q - 1) 2 \arctan \frac{1}{\sqrt{x_q - 1}} \right], \tag{6} \\ h(0, q^2) &= \frac{8}{27} - \frac{4}{9} \ln \frac{q^2}{m_c^2} + \frac{4}{9} i\pi. \tag{7} \end{aligned}$$

In the last equation,  $x_q = 4m_q^2/q^2$  and  $\Theta(x)$  is the Heaviside step function. Both  $h(m_c, q^2), h(m_b, q^2)$  can be obtained from  $h(m_q, q^2)$  by making the replacements  $m_q \rightarrow m_c$  and  $m_q \rightarrow m_b$ , respectively. The numerical values of the Wilson coefficients  $C_7^{eff}$  and  $C_{10}^{eff}$  as well as the other  $C_i$  for the  $b \rightarrow d$  transition can be found in [20].

The matrix element for the  $\Xi_{cc} \rightarrow \Sigma_c \ell^+ \ell^-$  can be obtained from Eq. (3) with the help of the following replacements:

$$V_{tb} V_{tq}^* \rightarrow V_{cd} V_{ud}^* + V_{cs} V_{bs}^*; m_b \rightarrow m_c; \Xi_{bb} \rightarrow \Xi_{cc}, \tag{8}$$

and replace  $C_i^{eff}$  for the b-quark case with the corresponding c-quark counterparts given as below.

The effective Wilson coefficient  $C_9^{eff}$  for  $c \rightarrow u$  transition is given as [21],

$$C_9^{eff} = C_9 + h_c(m_c, q^2) \left( 7C_3 + \frac{4}{3}C_4 + 76C_5 + \frac{64}{3}C_6 \right) - h_c(m_c, q^2)(3C_3 + 30C_5) + \frac{4}{3}h(0, q^2) \left( 3C_3 + C_4 + \frac{69}{2}C_5 + 16C_6 \right) + \frac{8}{3}(C_3 + 10C_5) - \left[ V_{cd}^* V_{ud} h(0, q^2) + V_{cs}^* V_{us} h(m_q, q^2) \right] \left( \frac{2}{3}C_1 + \frac{1}{2}C_2 \right), \tag{9}$$

The values of the Wilson coefficients are presented in [22], and the effective Wilson coefficient  $C_7^{eff}$  is given in [21], which we shall use in further numerical analysis. Note also that, due to the GIM cancellation,  $C_{10}^{eff}$  is zero.

It should be noted here that,  $C_9^{eff}$ , which appears in the  $c \rightarrow u$  and  $b \rightarrow d$  transitions, receives contributions also from vector mesons (long-distance effects). Long-distance contributions are only significant when  $q^2$  is close to the mass of the corresponding vector mesons. However, far from these points, these effects are small; hence, we take only short-distance effects into account. After these preliminary remarks, we now proceed to calculate the form factors in the framework of the QCD sum rules.

The matrix elements entering into Eq. (3) are parametrized in terms of the form factors in the following way,

$$\langle \Sigma_Q(p) | \bar{q} \gamma_\mu (1 - \gamma_5) Q | \Xi_{QQ}(p+q) \rangle = \bar{u}_{\Sigma_Q}(p) \left[ \gamma_\mu f_1(q^2) + \frac{i\sigma_{\mu\nu} q^\nu}{m_{\Xi_{QQ}}} f_2(q^2) + \frac{q_\mu}{m_{\Xi_{QQ}}} f_3(q^2) - \gamma_\mu \gamma_5 g_1(q^2) - \frac{i\sigma_{\mu\nu} \gamma_5 q^\nu}{m_{\Xi_{QQ}}} g_2(q^2) - \frac{\gamma_5 q_\mu}{m_{\Xi_{QQ}}} g_3(q^2) \right] u_{\Xi_{QQ}}(p+q), \tag{10}$$

$$\langle \Sigma_Q(p) | \bar{q} i\sigma_{\mu\nu} q^\nu (1 + \gamma_5) Q | \Xi_{QQ}(p+q) \rangle = \bar{u}_{\Sigma_Q}(p) \left[ (\gamma_\mu q^2 - q_\mu \not{q}) \frac{f_1^T(q^2)}{m_{\Xi_{QQ}}} + i\sigma_{\mu\nu} q^\nu f_2^T(q^2) + (\gamma_\mu q^2 - q_\mu \not{q}) \gamma_5 \frac{g_1^T(q^2)}{m_{\Xi_{QQ}}} + i\sigma_{\mu\nu} \gamma_5 q^\nu g_2^T(q^2) \right] u_{\Xi_{QQ}}(p+q). \tag{11}$$

where  $u_{\Sigma_Q}$  and  $u_{\Xi_{QQ}}$  are the spinors of the single and doubly heavy baryons. The form factors  $f_i$  and  $g_i$  are estimated in the framework of the light cone sum rules method in [7], and for this reason, we pay attention to the calculation of the form factors  $f_i^T$  and  $g_i^T$  using the LCSR method only.

In order to calculate the form factors  $f_i^T$  and  $g_i^T$  in the framework of the LCSR method, we start with the following

form correlation function.

$$\Pi_\mu(p, q) = i \int d^4x e^{iq \cdot x} \langle \Sigma_Q(p) | T \{ \bar{q} i\sigma_{\mu\nu} q^\nu (1 + \gamma_5) Q \bar{j}_{\Xi_{QQ}} \} | 0 \rangle, \tag{12}$$

where  $Q = c$  or  $b$ . The interpolating current of the  $\Xi_{QQ}$  baryon is,

$$j_{\Xi_{QQ}}(0) = \varepsilon^{abc} (Q^{aT} C \gamma^\mu Q^b) \gamma_\mu \gamma_5 q^c, \tag{13}$$

where  $a, b, c$  are the color indices. In the LCSR method, the expression of the correlation function is obtained in two different ways. One of the representations can be written in terms of the hadrons, and the other is from the QCD side, i.e. in terms of the quarks and gluons. On the hadronic side, the correlation function is obtained by inserting hadronic states with the quantum numbers of  $\Xi_{QQ}$  baryon. Then, isolating the ground state contributions of the  $\Xi_{QQ}$  baryon, we get,

$$\Pi_\mu(p, q) = \frac{\langle \Sigma_Q(p) | \bar{q} i\sigma_{\mu\nu} q^\nu (1 + \gamma_5) Q | \Xi_{QQ}(p+q) \rangle \langle \Xi_{QQ}(p+q) | \bar{j}_{\Xi_{QQ}}(0) \rangle}{m_{\Xi_{QQ}}^2 - (p+q)^2}. \tag{14}$$

The second matrix element can be written as,

$$\langle \Xi_{QQ}(p+q) | \bar{j}_{\Xi_{QQ}}(0) \rangle = f_{\Xi_{QQ}} \bar{u}_{\Xi_{QQ}}(p+q),$$

and the first matrix element which is defined in terms of the form factors, is given in Eq.(11).

Using the completeness condition of the Dirac bispinors, we get the following result for the correlation function in terms of hadrons

$$\begin{aligned} \Pi_\mu(p, q) &= \frac{f_{\Xi_{QQ}}}{m_{\Xi_{QQ}}^2 - (p+q)^2} \bar{u}(p) \left\{ \frac{f_1^T}{m_{\Xi_{QQ}}} (\gamma_\mu q^2 - q_\mu \not{q}) + i f_2^T \sigma_{\mu\nu} q^\nu + \frac{g_1^T}{m_{\Xi_{QQ}}} (\gamma_\mu q^2 - q_\mu \not{q}) \gamma_5 \right. \\ &\quad \left. + i g_2^T \sigma_{\mu\nu} q^\nu \gamma_5 \right\} (\not{p} + \not{q} + m_{\Xi_{QQ}}) \\ &= \frac{f}{m_1^2 - (p+q)^2} \bar{u}(v) \left\{ \left[ \left( \frac{m_2}{m_1} - 1 \right) f_1^T + f_2^T \right] \not{q} q_\mu + 2m_2 f_2^T \not{q} v_\mu + \left[ \left( \frac{m_2}{m_1} + 1 \right) g_1^T + g_2^T \right] \not{q} \gamma_5 q_\mu - 2m_2 g_2^T \not{q} \gamma_5 v_\mu \right\} + \text{other structures}. \tag{15} \end{aligned}$$

In the last step of the derivation, we used the heavy quark limit, i.e.,  $p_\mu \rightarrow m_{\Sigma_Q} v_\mu$ , and for brevity we replaced  $m_{\Xi_{QQ}}$  by  $m_1$ ,  $m_{\Sigma_Q}$  by  $m_2$ , and  $f_{\Xi_{QQ}}$  by  $f$ . Having obtained the

expression of the correlation function from the hadronic side, let us turn our attention to the calculation of the correlation function from the QCD side.

After using the Wick theorem for the correlation function from the QCD side, we get

$$\begin{aligned} \Pi_\mu(p, q) = & \int d^4x e^{i(q \cdot x)} \\ & \times \left\{ [i\sigma_{\mu\nu} q^\nu (1 + \gamma_5)]_{\alpha\beta} (\gamma_5 \gamma_\tau)_{\rho\gamma} (C\gamma_\tau)_{\sigma\phi} \epsilon^{abc} \right. \\ & \times (\Sigma_Q(p) | (-\bar{q}_\alpha^e(x) \bar{q}_\rho^c(0) \bar{Q}_\phi^a(0) S_{\beta\sigma}^{eb}(x) \\ & \left. + \bar{q}_\alpha^e(x) \bar{q}_\rho^c(0) \bar{Q}_\phi^b(0) S_{\beta\sigma}^{ea}(x) ) | 0 \right\}. \end{aligned} \tag{16}$$

The matrix element  $\epsilon^{abc} \langle \Sigma_c(p) | \bar{q}_\alpha^a(t_1) q_\beta^b(t_2) \bar{Q}_\rho^c(0) | 0 \rangle$  appearing in Eq. (16) is determined in terms of the heavy baryon DAs. The light cone distribution amplitudes are studied in [23]. In determining the parameters appearing in DAs, the standard sum rules method in heavy quark mass limit is considered. The distribution amplitudes of the  $\Sigma_Q$  baryon in the sextet representation of SU(3) are determined in the following way,

$$\begin{aligned} & \frac{\bar{v}^\mu}{v_+} \epsilon^{abc} \langle 0 | q_1^{aT} C \not{n} q_2^b(t_2) h_\gamma^c(0) | \Sigma_Q(v) \rangle \\ & = \frac{1}{\sqrt{3}} \psi_2(t_1, t_2) f^{(1)} \varepsilon_\parallel^\mu u_\gamma, \\ & i \frac{\bar{v}^\mu}{v_+} \epsilon^{abc} \langle 0 | q_1^{aT} C \sigma_{\alpha\beta} q_2^b(t_2) h_\gamma^c(0) | \Sigma_Q(v) \rangle \\ & = \frac{1}{\sqrt{3}} \psi_3^a(t_1, t_2) f^{(2)} \varepsilon_\parallel^\mu u_\gamma, \\ & \bar{v}^\mu \epsilon^{abc} \langle 0 | q_1^{aT} C q_2^b(t_2) h_\gamma^c(0) | \Sigma_Q(v) \rangle \\ & = \frac{1}{\sqrt{3}} \psi_3^\sigma(t_1, t_2) f^{(2)} \varepsilon_\parallel^\mu u_\gamma, \\ & -v_+ \bar{v}^\mu \epsilon^{abc} \langle 0 | q_1^{aT} C \not{n} q_2^b(t_2) h_\gamma^c(0) | \Sigma_Q(v) \rangle \\ & = \frac{1}{\sqrt{3}} \psi_4(t_1, t_2) f^{(1)} \varepsilon_\parallel^\mu u_\gamma, \end{aligned}$$

where  $\psi_i$  are the distribution amplitudes with definite twist,  $t_i$  are the distance between the  $i$ th light quark and the origin along the direction of  $n$ ,  $n^\mu$  and  $\bar{n}^\mu$  are the two light vectors,  $\bar{v}^\mu = \frac{1}{2} \left( \frac{n^\mu}{v_+} - v_+ \bar{n}^\mu \right)$ ,  $v^\mu = \frac{1}{2} \left( \frac{n^\mu}{v_+} + v_+ \bar{n}^\mu \right)$ , and the space coordinates are taken as  $t_i n^\mu$ . In further discussion we will work in the rest frame of the  $\Sigma_Q$  heavy baryons, i.e.,  $v_+ = 1$ . Here,  $u_\gamma(v)$  is the heavy baryon spinor and  $h_\gamma$  is the static heavy quark field in HQET.

Here we would like to make the following remark. The light cone DAs of heavy baryons are obtained in the HQET in terms of four-velocity and heavy quark field,  $h$ . However, in QCD, heavy baryon state,  $|\Sigma\rangle$ , is described by the momentum  $p$  and heavy quark field by  $Q$  (see Eq. 16). Therefore, these quantities should be transformed to the

HQET counterparts. The heavy quark field,  $Q$ , should be replaced by the corresponding heavy quark effective field  $h(0)$ , i.e.  $Q(0) \rightarrow h(0)$ . In addition, the heavy baryon state can be written in terms of the HQET baryon state by using  $|\Sigma_Q(p)\rangle = \sqrt{m_2} |\Sigma(v)\rangle + \mathcal{O}(1/m_2)$ . In HQET, since the higher order of the inverse heavy quark mass terms can be neglected, we obtain  $\Sigma_Q(p)\rangle = \sqrt{m_2} |\Sigma(v)\rangle$ . Applying this transformation to both sides of the correlation function, we see that the replacement  $|\Sigma(p)\rangle \rightarrow |\Sigma(v)\rangle$  can be made safely. However, this transformation is only valid for tree-level calculations. When  $\mathcal{O}(\alpha_s)$  corrections are taken into account, the matching relations among the QCD currents and HQET currents should be used (see [24]). In this work, we neglected the NLO corrections.

As a result, the matrix element  $\epsilon^{abc} \langle \Sigma_Q(v) | q_{1\alpha}^a(t_1) \bar{q}_{2\beta}^b(t_2) h_\gamma^c(0) | 0 \rangle$  in terms of  $\Sigma_Q$  distribution amplitudes can be written as

$$\begin{aligned} & \epsilon^{abc} \langle \Sigma_Q(v) | \bar{q}_{1\alpha}^a(t_1) \bar{q}_{2\beta}^b(t_2) \bar{h}_\gamma^c(0) | 0 \rangle \\ & = \sum_{i=1}^4 A_i (\bar{u}_\Sigma \not{v} \gamma_5)_\gamma (C^{-1} \Gamma_i)_{\alpha\beta}, \end{aligned} \tag{17}$$

where

$$\begin{aligned} A_1 &= \frac{f^{(1)}}{8} \psi_2(t_1, t_2), & A_3 &= \frac{f^{(2)}}{4} \psi_3^{(s)}(t_1, t_2), \\ A_2 &= -\frac{f^{(2)}}{8} \psi_3^{(\sigma)}(t_1, t_2), & A_4 &= \frac{f^{(1)}}{8} \psi_4(t_1, t_2), \end{aligned} \tag{18}$$

and  $\Gamma_1 = \not{n}$ ,  $\Gamma_2 = i\sigma_{\alpha\beta} n^\alpha \bar{n}^\beta$ ,  $\Gamma_3 = I$ ,  $\Gamma_4 = \not{n}$ . The distribution amplitudes  $\psi$  are defined as,

$$\psi(t_1, t_2) = \int_0^\infty dw w \int_0^1 du e^{-iw(t_1 u + t_2 \bar{u})} \psi(t_1, t_2),$$

where  $\bar{u} = 1 - u$ ,  $t_i = v x_i$ ,  $w_2 = \bar{u} w$  and  $w$  is the total light diquark momentum. Although the DAs are presented only for the bottomed baryons in [23, 25], one can use these DAs for the baryons containing charm quarks as well in the heavy quark mass limit. In the present work, both  $\Sigma_c$  and  $\Sigma_b$  are described by the same DAs given in [23]. Their explicit forms are,

$$\begin{aligned} \psi_2(u, w) &= w^2 \bar{u} u \sum_{n=0}^2 \frac{a_n C_n^{3/2} (2u-1)}{\varepsilon_n^4 |C_n^{3/2}|^2} e^{-w/\varepsilon_n}, \\ \psi_4(u, w) &= w^2 \bar{u} u \sum_{n=0}^2 \frac{a_n C_n^{1/2} (2u-1)}{\varepsilon_n^2 |C_n^{1/2}|^2} e^{-w/\varepsilon_n}, \\ \psi_3^{(\sigma,s)}(u, w) &= \frac{w}{2} \bar{u} u \sum_{n=0}^2 \frac{a_n C_n^{1/2} (2u-1)}{\varepsilon_n^3 |C_n^{1/2}|^2} e^{-w/\varepsilon_n}. \end{aligned} \tag{19}$$

The values of the parameters  $a_0, a_1, a_2$ , and  $\varepsilon_0, \varepsilon_1, \varepsilon_2$  are given in [23], and  $C_n^\lambda(2u-1)$  is the Gegenbauer polynomial.

Substituting the light-cone distribution amplitudes Eq. (17) into Eq. (16) and using the heavy quark propagator in momentum representation and performing integration over  $x$ , the correlation function at the QCD level can be written as,

$$\begin{aligned} \Pi_{\mu}^{QCD}[(p+q)^2, q^2] = & \int du \int dw \sum_{n=1}^3 \left\{ \frac{\rho_n^{(1)}(u, w)}{(\Delta - m_Q^2)^n} q_{\mu} \not{q} \right. \\ & + \frac{\rho_n^{(2)}(u, w)}{(\Delta - m_Q^2)^n} \not{q} v_{\mu} \\ & + \frac{\rho_n^{(3)}(u, w)}{(\Delta - m_Q^2)^n} q_{\mu} \not{q} \gamma_5 \\ & + \frac{\rho_n^{(4)}(u, w)}{(\Delta - m_Q^2)^n} \not{q} v_{\mu} \gamma_5 \\ & \left. + \text{other structures} \right\}, \end{aligned} \tag{20}$$

where the invariant functions  $\rho_n^{(i)}$ , ( $i = 1, 2, 3, 4$ ) are presented in Appendix 1, and

$$\Delta = \frac{\bar{u}w}{m_{\Sigma_Q}}(p+q)^2 + q^2 \left(1 - \frac{\bar{u}w}{m_{\Sigma_Q}}\right) - \bar{u}w(m_{\Sigma} - \bar{u}w). \tag{21}$$

Matching the coefficients of the structures  $\not{q}q_{\mu}$ ,  $\not{q}v_{\mu}$ ,  $\not{q}\gamma_5q_{\mu}$ , and  $\not{q}\gamma_5v_{\mu}$  in both representations of the correlation function and applying the Borel transformation with respect to the variable  $-(p+q)^2$  in order to enhance the contributions of the ground states, and suppress the higher states and continuum contributions, the desired sum rules for the  $f_i^T$  and  $g_i^T$  form factors are obtained from the following equations,

$$\begin{aligned} f \left\{ \left(\frac{m_2}{m_1} - 1\right) f_1^T + f_2^T \right\} e^{-m_1^2/M^2} &= \Pi_1^B, \\ 2m_2 f e^{-m_1^2/M^2} f_2^T &= \Pi_2^B, \\ f \left\{ \left(\frac{m_2}{m_1} + 1\right) g_1^T + g_2^T \right\} e^{-m_1^2/M^2} &= \Pi_3^B, \\ -2m_2 f e^{-m_1^2/M^2} g_2^T &= \Pi_4^B, \end{aligned} \tag{22}$$

where  $\Pi_i^B$  are the Borel transformed coefficients of the structures mentioned above, and  $M^2$  is the Borel mass parameter.

The Borel transformation and continuum subtraction is performed with the help of the following master formula.

$$\begin{aligned} & \int dw \frac{\rho(u, w)}{(\Delta - m_Q^2)^n} \\ &= \sum_{n=1}^{\infty} \left\{ (-1)^n \int_0^{w_0} dw \frac{e^{-s/M^2}}{(n-1)!(M^2)^{n-1}} I_n \right. \\ & \quad \left. - \left[ \frac{(-1)^{n-1}}{(n-1)!} e^{-s/M^2} \sum_{j=1}^{n-1} \frac{1}{(M^2)^{n-j-1}} \frac{1}{s'} \left(\frac{d}{dw} \frac{1}{s'}\right)^{j-1} I_n \right]_{w=w_0} \right\}, \end{aligned}$$

where

$$s = \frac{m_Q^2 - \bar{u}w(\bar{u}w - m_2) - \left(1 - \frac{\bar{u}w}{m_2}\right) q^2}{\frac{\bar{u}w}{m_2}},$$

$$I_n = \frac{\rho(u, w)}{\left(\frac{\bar{u}w}{m_2}\right)^n} \tag{23}$$

Note that  $w = w_0$  is the solutions of the equations  $s = s_{th}$  (and also  $s = 4m_Q^2$ ),  $s' = \frac{ds}{dw}$ , and the differential operator is defined as,

$$\left(\frac{d}{dw} \frac{1}{s'}\right)^{j-1} I_n \rightarrow \frac{d}{dw} \left[ \frac{d}{dw} \frac{1}{s'} \cdots I_n \right], \tag{24}$$

where  $\cdots$  means the operation should be repeated  $j - 1$  times.

### 3 Numerical analysis

The primary aim of this section is to determine the  $q^2$  dependence of the form factors  $f_1^T, f_2^T, g_1^T$ , and  $g_2^T$ , whose LCSR are derived in the previous section. Then we estimate the branching ratios of the  $\Xi_{QQ} \rightarrow \Sigma_Q \ell^+ \ell^-$  decays.

The LCSR of the form factors contain numerous input parameters. In further numerical analysis, we choose the masses of the heavy quarks in the  $\overline{MS}$  scheme, i.e.,  $m_c(\overline{m}_c) = (1.28 \pm 0.03)$  GeV, and  $m_b(\overline{m}_b) = (4.18 \pm 0.03)$  GeV [26]. The masses, lifetime and decay constants  $f$  of the doubly heavy baryons are given in Table 1 (see also [27–31]).

The mass and decay constants of the  $\Sigma_Q$  baryon are chosen as  $m_{\Sigma_c} = 2.454$  GeV,  $m_{\Sigma_b} = 5.814$  GeV, and  $f^{(1)} = f^{(2)} = 0.38$  [36]

In addition to these input values, two extra auxiliary parameters, continuum threshold  $s_{th}$ , and the Borel mass parameter  $M^2$ , appear in the LCSR method. These parameters are determined with the following criteria. The working region of  $M^2$  is determined by requiring that the power corrections and continuum contributions both should be suppressed compared to the leading twist-2 contribution. The continuum threshold  $s_{th}$  is determined so that the mass sum

**Table 1** The mass, decay constants and lifetimes of the doubly heavy  $\Xi_{QQ}$  baryons

Baryons	Mass (GeV)	Life time ( $f$ s)	$f$ (GeV) <sup>3</sup> [32]
$\Xi_{cc}^{++}$	3.621 [1]	256 [3]	$0.109 \pm 0.021$
$\Xi_{cc}^+$	3.621 [1]	45 [33]	$0.109 \pm 0.021$
$\Xi_{bb}^0$	10.143 [34]	370 [35]	$0.281 \pm 0.071$
$\Xi_{bb}^-$	10.143 [34]	370 [35]	$0.281 \pm 0.071$



**Table 2** The values of the fit parameters for the form factors of the  $\Xi_{QQ} \rightarrow \Sigma_Q$  transition

	$f(0)$	$m_{fit}$	$\delta$
Form factors (c-sector)			
$f_1^T(q^2)$	$3.05 \pm 0.7$	1.69	0.43
$f_2^T(q^2)$	$0.49 \pm 0.1$	1.15	4.21
$g_1^T(q^2)$	$1.44 \pm 0.26$	1.43	0.42
$g_2^T(q^2)$	$0.53 \pm 0.13$	–	–
Form factors (b-sector)			
$f_1^T(q^2)$	$-1.72 \pm 0.35$	3.60	0.31
$f_2^T(q^2)$	$0.32 \pm 0.06$	3.49	0.75
$g_1^T(q^2)$	$0.21 \pm 0.04$	–	–
$g_2^T(q^2)$	$0.32 \pm 0.06$	2.93	0.42

rule reproduces the experimentally measured value of mass to within  $\pm 5\%$  accuracy.

Based on these conditions imposed by the LCSR method, we obtain the following working regions of the parameters  $s_{th}$  and  $M^2$  for the transitions under consideration, i.e.,  $s_{th} = (16 \pm 1) \text{ GeV}^2$ ,  $M^2 = (10 \pm 2) \text{ GeV}^2$  for the  $\Xi_{cc} \rightarrow \Sigma_c$  transition, and  $s_{th} = (112 \pm 2) \text{ GeV}^2$ ,  $M^2 = (20 \pm 2) \text{ GeV}^2$  for the  $\Xi_{bb} \rightarrow \Sigma_b$  transition, respectively. It should be emphasized here that these working regions are more or less in the same range as those determined for the transitions induced by the charged current [7].

It should be reminded that LCSR predictions are reliable in the low-energy region. Our calculations show that the sum rule for the form factors is meaningful in the domains  $q^2 \leq 0.5 \text{ GeV}^2$  for the  $\Xi_{cc} \rightarrow \Sigma_c$  transition and  $q^2 \leq 10 \text{ GeV}^2$  for the  $\Xi_{bb} \rightarrow \Sigma_b$  transition, respectively.

Having determined the working regions for the QCD results of the form factors, we can extend the LCSR predictions to the entire physical region. For this goal, we extrapolate these form factors to the physical region in such a way that in the region where LCSR is reliable, the result of the fit formula and the result LCSR method coincide with each other. Our analysis shows that the best-fit formula which satisfies the required restrictions is given as,

$$F_i^T(q^2) = \frac{f_i^T(0)}{1 - \frac{q^2}{m_{fit}^2} + \delta \left( \frac{q^2}{m_{fit}^2} \right)^2}, \tag{25}$$

**Table 3** The branching ratios of the  $\Xi_{QQ} \rightarrow \Sigma_Q \ell^+ \ell^-$  decays

Branching ratios	Ours	[10]	[12]
$\Xi_{cc}^{++} \rightarrow \Sigma_c^{++} \ell^+ \ell^-$	$1.62(1 \pm 0.45) \times 10^{-13}$	–	–
$\Xi_{bb}^0 \rightarrow \Sigma_b^0 e^+ e^-$	$2.51(1 \pm 0.35) \times 10^{-8}$	$9.00 \times 10^{-9}$	$5.91 \times 10^{-9}$
$\Xi_{bb}^0 \rightarrow \Sigma_b^0 \mu^+ \mu^-$	$1.73(1 \pm 0.30) \times 10^{-8}$	$7.94 \times 10^{-9}$	$4.89 \times 10^{-9}$
$\Xi_{bb}^0 \rightarrow \Sigma_b^0 \tau^+ \tau^-$	$1.50(1 \pm 0.31) \times 10^{-9}$	$1.18 \times 10^{-9}$	$1.91 \times 10^{-10}$

where the values of the fit parameters  $f_i^T(0)$ ,  $m_{fit}$ , and  $\delta$  are presented in Table 2.

The errors presented in the values of  $f_i^T(q^2 = 0)$  point are due to the uncertainties in the mass of the heavy quark, Borel mass parameter, continuum threshold  $s_{th}$ , as well as from the input parameters appearing in the DAs of the  $\Sigma_b$  baryon.

Having the results for the form factors, we now proceed to calculate the corresponding branching ratios of the  $\Xi_{cc}^{++} \rightarrow \Sigma_c^{++} \ell^+ \ell^-$  and  $\Xi_{bb}^0 \rightarrow \Sigma_b^0 \ell^+ \ell^-$  decays. Using the definition of the matrix element of the  $\Xi_{bb} \rightarrow \Sigma_b \ell^+ \ell^-$  the decay width is obtained as

$$\frac{d\Gamma(s)}{ds} = \frac{G_F^2 \alpha^2 m_1}{4096\pi^5} |V_{tb} V_{td}^*|^2 v \sqrt{\lambda(1, r, s)} \left[ T_1(s) + \frac{1}{3} T_2(s) \right]$$

where  $v = \sqrt{1 - \frac{4m_\ell^2}{q^2}}$  is the lepton velocity,  $\lambda(1, r, s) = 1 + r^2 + s^2 - 2r - 2s - 2rs$ ,  $s = \frac{q^2}{m_1^2}$ , and  $r = \frac{m_2^2}{m_1^2}$ . The lengthy expressions  $T_1(s)$  and  $T_2(s)$  can be found in [13].

Performing integration over the parameter  $s$  in the domain  $\frac{4m_\ell^2}{m_1^2} \leq s \leq (1 - \sqrt{r})$ , and using the lifetimes of  $\Xi_{cc}^{++}$ ,  $\Xi_c^+$ , and  $\Xi_{bb}^0$ , we calculate the branching ratios of the  $\Xi_{QQ} \rightarrow \Sigma_Q$  decays, whose numerical results are all presented in Table 3. For comparison, we also present the corresponding branching ratios predicted by the Light Front approach [10, 11]. We observe from this comparison that our prediction for the  $\Xi_{QQ} \rightarrow \Sigma_Q \ell^+ \ell^-$  ( $\ell = e, \mu$ ) transition is larger than the predictions of the Light Front approach. Considering the results summarized in Table 3, one can conclude that the branching ratios of the  $\Xi_{bb} \rightarrow \Sigma_b \ell^+ \ell^-$  decays could be measured in future experiments at LHCb, while the measurement of the branching ratios of the  $\Xi_{cc} \rightarrow \Sigma_c \ell^+ \ell^-$  decays presents quite a complex problem.

### 4 Conclusion

The decays induced by the flavor-changing neutral currents  $b \rightarrow d$  and  $c \rightarrow u$  of the doubly heavy baryons are studied in the framework of the Light Cone Sum Rules method. We derive the LCSR of the form factors induced by the tensor

current. Using the results of the form factors obtained, we estimated the corresponding branching ratios. We found out that the branching ratios for  $\Xi_{bb} \rightarrow \Sigma_b \ell^+ \ell^-$  ( $\ell = e, \mu$ ) is at the order of  $\sim 10^{-8}$ . Moreover, for  $\Xi_{cc} \rightarrow \Sigma_c \ell^+ \ell^-$  decay, the branching ratios are much smaller and at the order of  $\sim 10^{-13}$ . The relatively large value of the branching ratio  $\Xi_{bb} \rightarrow \Sigma_b \ell^+ \ell^-$  indicates the possibility of being observed in future experiments at LHCb.

Future improvements for the DA's of the  $\Sigma_Q$  baryon and the inclusion of the gluon radiative corrections to the correlation function could pave the way to more accurate sum rules and numerical predictions.

**Data Availability Statement** This manuscript has no associated data or the data will not be deposited. [Authors' comment: This manuscript has no associated data or the data will not be deposited.]

**Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

Funded by SCOAP<sup>3</sup>. SCOAP<sup>3</sup> supports the goals of the International Year of Basic Sciences for Sustainable Development.

**Appendix: The expression of the invariant functions**

$$\rho_n^{(i)}(u, w)$$

$$\begin{aligned} \rho_1^{(1)} &= f^{(2)} w \psi_3^{(s)}(u, w), \\ \rho_2^{(1)} &= \bar{u} \left\{ \left[ f^{(1)} m_Q \widehat{\psi}_4(u, w) \right. \right. \\ &\quad \left. \left. + 2f^{(2)} [\bar{u}w - (q \cdot v)] \widehat{\psi}_3^{(\sigma)}(u, w) \right. \right. \\ &\quad \left. \left. - f^{(1)} [m_Q - 4(q \cdot v)] \widehat{\psi}_2(u, w) \right. \right. \\ &\quad \left. \left. - 4f^{(2)} \bar{u} \left[ w \widehat{\psi}_3^{(s)}(u, w) + 2\widehat{\psi}_3^{(\sigma)}(u, w) \right] \right\}, \\ \rho_3^{(1)} &= 8f^{(2)} \bar{u}^2 [q^2 - \bar{u}^2 w^2 - 2m_Q (q \cdot v)] \widehat{\psi}_3^{(\sigma)}(u, w), \\ \rho_1^{(2)} &= 2f^{(2)} \bar{u} [w^2 \psi_3^{(s)}(u, w) + \widehat{\psi}_3^{(\sigma)}(u, w)], \\ \rho_2^{(2)} &= -4\bar{u} \left\{ f^{(1)} q^2 \widehat{\psi}_2(u, w) + f^{(2)} \bar{u} [w(q \cdot v) \widehat{\psi}_3^{(\sigma)}(u, w) \right. \\ &\quad \left. + 2m_Q \widehat{\psi}_3^{(\sigma)}(u, w) + \bar{u}w (w \widehat{\psi}_3^{(s)}(u, w) \right. \\ &\quad \left. + 4\widehat{\psi}_3^{(\sigma)}(u, w)) \right\}, \\ \rho_3^{(2)} &= 16f^{(2)} \bar{u}^2 [q^2 (m_Q + \bar{u}w) + \bar{u}^2 w^2 (q \cdot v)] \widehat{\psi}_3^{(\sigma)}(u, w), \end{aligned}$$

$$\begin{aligned} \rho_1^{(3)} &= -f^{(2)} w \psi_3^{(s)}(u, w), \\ \rho_2^{(3)} &= \bar{u} \left\{ f^{(1)} m_Q \widehat{\psi}_4(u, w) - 2f^{(2)} [\bar{u}w - (q \cdot v)] \widehat{\psi}_3^{(\sigma)}(u, w) \right. \\ &\quad \left. - f^{(1)} [m_Q - 4(q \cdot v)] \widehat{\psi}_2(u, w) + 4f^{(2)} \bar{u} [w \widehat{\psi}_3^{(s)}(u, w) \right. \\ &\quad \left. + 2\widehat{\psi}_3^{(\sigma)}(u, w) \right\}, \\ \rho_3^{(3)} &= -8f^{(2)} \bar{u}^2 [q^2 - \bar{u}^2 w^2 - 2m_Q (q \cdot v)] \widehat{\psi}_3^{(\sigma)}(u, w), \\ \rho_1^{(4)} &= -2f^{(2)} \bar{u} [w^2 \psi_3^{(s)}(u, w) + \widehat{\psi}_3^{(\sigma)}(u, w)], \\ \rho_2^{(4)} &= -4\bar{u} \left\{ \left[ f^{(1)} q^2 \widehat{\psi}_2(u, w) \right] - f^{(2)} \bar{u} [w(q \cdot v) \widehat{\psi}_3^{(\sigma)}(u, w) \right. \\ &\quad \left. + 2m_Q \widehat{\psi}_3^{(\sigma)}(u, w) + \bar{u}w (w \widehat{\psi}_3^{(s)}(u, w) \right. \\ &\quad \left. + 4\widehat{\psi}_3^{(\sigma)}(u, w)) \right\}, \\ \rho_3^{(4)} &= -16f^{(2)} \bar{u}^2 [q^2 (m_Q + \bar{u}w) + \bar{u}^2 w^2 (q \cdot v)] \widehat{\psi}_3^{(\sigma)}(u, w). \end{aligned} \tag{A.1}$$

The functions  $\widehat{\psi}(u, w)$  and  $\widehat{\psi}(u, w)$  are defined as,

$$\begin{aligned} \widehat{\psi}(u, w) &= \int_0^w d\tau' \psi(u, \tau') \tau', \\ \widehat{\psi}(u, w) &= \int_0^w d\tau' \widehat{\psi}(u, \tau'). \end{aligned}$$

**References**

1. LHCb Collaboration, R. Aaij et al., *Observation of the doubly charmed baryon  $\Xi_{cc}^{++}$* . Phys. Rev. Lett. **119**(11), 112001 (2017). [arXiv:1707.01621](https://arxiv.org/abs/1707.01621)
2. LHCb Collaboration, R. Aaij et al., *First observation of the doubly charmed baryon decay  $\Xi_{cc}^{++} \rightarrow \Xi_c^+ \pi^+$* . Phys. Rev. Lett. **121**(16), 162002 (2018) [arXiv:1807.01919](https://arxiv.org/abs/1807.01919)
3. LHCb Collaboration, R. Aaij et al., *Measurement of the lifetime of the doubly charmed baryon  $\Xi_{cc}^{++}$* . Phys. Rev. Lett. **121**(5), 052002 (2018). [arXiv:1806.02744](https://arxiv.org/abs/1806.02744)
4. T. Aliev, S. Bilmis, *Properties of doubly heavy baryons in QCD*. Turk. J. Phys. **46**(1), 1–26 (2022). [arXiv:2203.02965](https://arxiv.org/abs/2203.02965)
5. Y.-J. Shi, W. Wang, Z.-X. Zhao, *QCD Sum Rules Analysis of Weak Decays of Doubly-Heavy Baryons*. Eur. Phys. J. C **80**(6), 568 (2020). [arXiv:1902.01092](https://arxiv.org/abs/1902.01092)
6. Y.-J. Shi, Y. Xing, Z.-X. Zhao, *Light-cone sum rules analysis of  $\Xi_{QQ'q} \rightarrow \Lambda_{Q'}$  weak decays*. Eur. Phys. J. C **79**(6), 501 (2019). [arXiv:1903.03921](https://arxiv.org/abs/1903.03921)
7. X.-H. Hu, Y.-J. Shi, *Light-cone sum rules analysis of  $\Xi_{QQ'} \rightarrow \Sigma_{Q'}$  weak decays*. Eur. Phys. J. C **80**(1), 56 (2020). [arXiv:1910.07909](https://arxiv.org/abs/1910.07909)
8. W. Wang, F.-S. Yu, Z.-X. Zhao, *Weak decays of doubly heavy baryons: the 1/2  $\rightarrow$  1/2 case*. Eur. Phys. J. C **77**(11), 781 (2017). [arXiv:1707.02834](https://arxiv.org/abs/1707.02834)
9. Z.-X. Zhao, *Weak decays of doubly heavy baryons: the 1/2  $\rightarrow$  3/2 case*. Eur. Phys. J. C **78**(9), 756 (2018). [arXiv:1805.10878](https://arxiv.org/abs/1805.10878)
10. Z.-P. Xing, Z.-X. Zhao, *Weak decays of doubly heavy baryons: the FCNC processes*. Phys. Rev. D **98**(5), 056002 (2018). [arXiv:1807.03101](https://arxiv.org/abs/1807.03101)

11. H.-W. Ke, F. Lu, X.-H. Liu, X.-Q. Li, Study on  $\Xi_{cc} \rightarrow \Xi_c$  and  $\Xi_{cc} \rightarrow \Xi'_c$  weak decays in the light-front quark model. *Eur. Phys. J. C* **80**(2), 140 (2020). [arXiv:1912.01435](https://arxiv.org/abs/1912.01435)
12. X.-H. Hu, R.-H. Li, Z.-P. Xing, A comprehensive analysis of weak transition form factors for doubly heavy baryons in the light front approach. *Eur. Phys. J. C* **80**(4), 320 (2020). [arXiv:2001.06375](https://arxiv.org/abs/2001.06375)
13. T. M. Aliev, S. Bilmis, M. Savci, “Analysis of FCNC  $\Xi_{QQ} \rightarrow \Lambda_Q l^+ l^-$  decay in light-cone sum rules,” *Phys. Rev. D* **106**(3) 034017 (2022)
14. M.A. Shifman, A.I. Vainshtein, V.I. Zakharov, QCD and resonance physics: theoretical foundations. *Nucl. Phys. B* **147**, 385–447 (1979). [https://doi.org/10.1016/0550-3213\(79\)90022-1](https://doi.org/10.1016/0550-3213(79)90022-1)
15. V.L. Chernyak, A.R. Zhitnitsky, Asymptotic behavior of exclusive processes in QCD. *Phys. Rept.* **112**, 173 (1984). [https://doi.org/10.1016/0370-1573\(84\)90126-1](https://doi.org/10.1016/0370-1573(84)90126-1)
16. I.I. Balitsky, V.M. Braun, A.V. Kolesnichenko, Radiative decay  $\sigma^+ \rightarrow p \gamma$  in quantum chromodynamics. *Nucl. Phys. B* **312**, 509–550 (1989). [https://doi.org/10.1016/0550-3213\(89\)90570-1](https://doi.org/10.1016/0550-3213(89)90570-1)
17. P. Colangelo, A. Khodjamirian, QCD sum rules, a modern perspective. [arXiv:hep-ph/0010175](https://arxiv.org/abs/hep-ph/0010175)
18. A.J. Buras, M. Munz, Effective Hamiltonian for  $B \rightarrow X(s) e^+ e^-$  beyond leading logarithms in the NDR and HV schemes. *Phys. Rev. D* **52**, 186–195 (1995). [arXiv:hep-ph/9501281](https://arxiv.org/abs/hep-ph/9501281)
19. A. Khodjamirian, T. Mannel, A.A. Pivovarov, Y.M. Wang, Charm-loop effect in  $B \rightarrow K^{(*)} \ell^+ \ell^-$  and  $B \rightarrow K^* \gamma$ . *JHEP* **09**, 089 (2010). [arXiv:1006.4945](https://arxiv.org/abs/1006.4945)
20. W. Altmannshofer, P. Ball, A. Bharucha, A.J. Buras, D.M. Straub, M. Wick, Symmetries and Asymmetries of  $B \rightarrow K^* \mu^+ \mu^-$  Decays in the Standard Model and Beyond. *JHEP* **01**, 019 (2009). [arXiv:0811.1214](https://arxiv.org/abs/0811.1214)
21. S. de Boer, G. Hiller, Flavor and new physics opportunities with rare charm decays into leptons. *Phys. Rev. D* **93**(7), 074001 (2016). [arXiv:1510.00311](https://arxiv.org/abs/1510.00311)
22. S. de Boer, B. Müller, D. Seidel, Higher-order Wilson coefficients for  $c \rightarrow u$  transitions in the standard model. *JHEP* **08**, 091 (2016). [arXiv:1606.05521](https://arxiv.org/abs/1606.05521)
23. A. Ali, C. Hambrock, A.Y. Parkhomenko, W. Wang, Light-Cone Distribution Amplitudes of the Ground State Bottom Baryons in HQET. *Eur. Phys. J. C* **73**(2), 2302 (2013). [arXiv:1212.3280](https://arxiv.org/abs/1212.3280)
24. B. Grinstein, D. Pirjol, Exclusive rare  $B \rightarrow K^* \ell^+ \ell^-$  decays at low recoil: Controlling the long-distance effects. *Phys. Rev. D* **70**, 114005 (2004). [arXiv:hep-ph/0404250](https://arxiv.org/abs/hep-ph/0404250)
25. G. Bell, T. Feldmann, Y.-M. Wang, M.W.Y. Yip, Light-cone distribution amplitudes for heavy-quark hadrons. *JHEP* **11**, 191 (2013). [arXiv:1308.6114](https://arxiv.org/abs/1308.6114)
26. Particle Data Group Collaboration, M. Tanabashi et al., Review of particle physics. *Phys. Rev. D* **98**, 030001 (2018). <https://doi.org/10.1103/PhysRevD.98.030001>
27. M. Karliner, J.L. Rosner, Baryons with two heavy quarks: Masses, production, decays, and detection. *Phys. Rev. D* **90**(9), 094007 (2014). [arXiv:1408.5877](https://arxiv.org/abs/1408.5877)
28. T.M. Aliev, K. Azizi, M. Savci, Doubly Heavy Spin-1/2 Baryon Spectrum in QCD. *Nucl. Phys. A* **895**, 59–70 (2012). [arXiv:1205.2873](https://arxiv.org/abs/1205.2873)
29. Z. Shah, K. Thakkar, A.K. Rai, Excited State Mass spectra of doubly heavy baryons  $\Omega_{cc}$ ,  $\Omega_{bb}$  and  $\Omega_{bc}$ . *Eur. Phys. J. C* **76**(10), 530 (2016). [arXiv:1609.03030](https://arxiv.org/abs/1609.03030)
30. Z. Shah, A.K. Rai, Excited state mass spectra of doubly heavy  $\Xi$  baryons. *Eur. Phys. J. C* **77**(2), 129 (2017). [arXiv:1702.02726](https://arxiv.org/abs/1702.02726)
31. V.V. Kiselev, A.K. Likhoded, Baryons with two heavy quarks. *Phys. Usp.* **45**, 455–506 (2002). [arXiv:hep-ph/0103169](https://arxiv.org/abs/hep-ph/0103169)
32. X.-H. Hu, Y.-L. Shen, W. Wang, Z.-X. Zhao, Weak decays of doubly heavy baryons: “decay constants”. *Chin. Phys. C* **42**(12), 123102 (2018). [arXiv:1711.10289](https://arxiv.org/abs/1711.10289)
33. H.-Y. Cheng, Y.-L. Shi, Lifetimes of Doubly Charmed Baryons. *Phys. Rev. D* **98**(11), 113005 (2018). [arXiv:1809.08102](https://arxiv.org/abs/1809.08102)
34. Z.S. Brown, W. Detmold, S. Meinel, K. Orginos, Charmed bottom baryon spectroscopy from lattice QCD. *Phys. Rev. D* **90**(9), 094507 (2014). [arXiv:1409.0497](https://arxiv.org/abs/1409.0497)
35. M. Karliner, J.L. Rosner, Discovery of doubly-charmed  $\Xi_{cc}$  baryon implies a stable ( $bb\bar{u}\bar{d}$ ) tetraquark. *Phys. Rev. Lett.* **119**(20), 202001 (2017). [arXiv:1707.07666](https://arxiv.org/abs/1707.07666)
36. S. Groote, J.G. Korner, O.I. Yakovlev, QCD sum rules for heavy baryons at next-to-leading order in  $\alpha_s$ . *Phys. Rev. D* **55**, 3016–3026 (1997). [arXiv:hep-ph/9609469](https://arxiv.org/abs/hep-ph/9609469)