

RESEARCH ARTICLE

Open Access

ER β 1: characterization, prognosis, and evaluation of treatment strategies in ER α -positive and -negative breast cancer

Jordan M Reese^{1,2}, Vera J Suman³, Malayannan Subramaniam¹, Xianglin Wu¹, Vivian Negron⁴, Anne Gingery¹, Kevin S Pitel¹, Sejal S Shah⁴, Heather E Cunliffe⁵, Ann E McCullough⁶, Barbara A Pockaj⁷, Fergus J Couch⁴, Janet E Olson⁸, Carol Reynolds⁴, Wilma L Lingle⁴, Thomas C Spelsberg¹, Matthew P Goetz^{2,9}, James N Ingle⁹ and John R Hawse^{1*}

Abstract

Background: The role and clinical value of ER β 1 expression is controversial and recent data demonstrates that many ER β antibodies are insensitive and/or non-specific. Therefore, we sought to comprehensively characterize ER β 1 expression across all sub-types of breast cancer using a validated antibody and determine the roles of this receptor in mediating response to multiple forms of endocrine therapy both in the presence and absence of ER α expression.

Methods: Nuclear and cytoplasmic expression patterns of ER β 1 were analyzed in three patient cohorts, including a retrospective analysis of a prospective adjuvant tamoxifen study and a triple negative breast cancer cohort. To investigate the utility of therapeutically targeting ER β 1, we generated multiple ER β 1 expressing cell model systems and determined their proliferative responses following anti-estrogenic or ER β -specific agonist exposure.

Results: Nuclear ER β 1 was shown to be expressed across all major sub-types of breast cancer, including 25% of triple negative breast cancers and 33% of ER-positive tumors, and was associated with significantly improved outcomes in ER α -positive tamoxifen-treated patients. In agreement with these observations, ER β 1 expression sensitized ER α -positive breast cancer cells to the anti-cancer effects of selective estrogen receptor modulators (SERMs). However, in the absence of ER α expression, ER β -specific agonists potently inhibited cell proliferation rates while anti-estrogenic therapies were ineffective.

Conclusions: Using a validated antibody, we have confirmed that nuclear ER β 1 expression is commonly present in breast cancer and is prognostic in tamoxifen-treated patients. Using multiple breast cancer cell lines, ER β appears to be a novel therapeutic target. However, the efficacy of SERMs and ER β -specific agonists differ as a function of ER α expression.

Keywords: Estrogen receptor beta, Breast cancer, Estrogen receptor alpha, Triple negative breast cancer, Therapy

Background

The global incidence of breast cancer has grown from 1980 to 2010 at an annual rate of 3.1%. In 2010, there were 1.65 million women diagnosed with breast cancer and 425,000 deaths caused by this disease [1]. Despite the substantial advances in understanding breast cancer biology, the clinical management of women with this disease continues to rely almost solely on the tumoral

expression of estrogen receptor alpha (ER α), progesterone receptor (PR) and epidermal growth factor receptor 2 (HER2). ER α is expressed in approximately 70% of all breast tumors and is the basis for the use of selective estrogen receptor modulators (SERMs) and aromatase inhibitors (AIs), which substantially reduce the risk for disease recurrence and prolong patient survival. Despite the discovery of a second form of the ER, ER β 1, more than 15 years ago [2,3], the endocrine sensitivity and ER status of breast tumors continues to be clinically defined exclusively by ER α expression [4-6].

* Correspondence: hawse.john@mayo.edu

¹Department of Biochemistry and Molecular Biology, Mayo Clinic, 16-01B Guggenheim Building, 200 First St. SW, Rochester, MN 55905, USA
Full list of author information is available at the end of the article

Like ER α , ER β 1 is a member of the nuclear receptor superfamily of proteins that functions as a ligand-mediated transcription factor [3]. The DNA binding domains of ER α and ER β 1 share 96% homology at the amino acid level, however, the remainder of the protein domains are highly divergent with the hinge region, AF1 domain, and ligand binding domain sharing only 30%, 30% and 53% conservation respectively [3,7]. A number of microarray studies from our laboratory and others have demonstrated that these two proteins function differently in response to both estrogen and anti-estrogens [8-14]. Consistent with these data, the genome wide chromatin binding profiles, or cistromes, of ER α and ER β 1 share only 40% overlap following short term estrogen treatment [14].

While ER β is highly expressed in normal breast tissue [15-21], a number of immunohistochemistry-based studies have demonstrated conflicting data with regard to ER β expression in breast tumors. For example, the frequency of ER β expression in breast tumors has been reported to range from 17-100% [15,18,21-35] and from 13-83% in ER α negative breast cancer [17,24,29,30,33,36]. With regard to the biological functions of ER β , a number of studies have shown that the presence of this receptor correlates with improved rates of recurrence, disease-free survival and overall survival [22,24-27,37-41] while others indicate little to no correlation [28,30,38] or even worse prognosis [33,42-44]. Lastly, several studies have reported that the presence of ER β in breast tumors increases the effectiveness of tamoxifen therapy [36,45-48] or aromatase inhibitor therapy [47,49]. For these reasons, the expression profiles and biological functions of ER β in human breast tumors remains unclear and has limited its utility as a prognostic and/or predictive biomarker for this disease. A potential reason for the conflicting data relates to the known existence of at least 4 different ER β splice variants (ER β 2-5) in human breast tumors whose biological functions largely remain unknown. Additionally, a recent report by our laboratory and others suggests that some of the inconsistencies regarding the prevalence of ER β in breast tumors may be related to the use of non-specific and/or insensitive ER β antibodies [20,50].

For these reasons, we sought to further characterize the expression patterns of ER β 1 across multiple breast cancer sub-types using a validated antibody. This particular antibody (PPG5/10) has been shown by us and others to detect only the full-length form of this receptor and is highly sensitive and specific in immunohistochemical studies [20,50,51]. Here, we have examined nuclear and cytoplasmic ER β 1 expression levels in over 400 breast tumors and have correlated these levels with other prognostic biomarkers and/or known patient outcomes. Our results demonstrate that ER β 1 is expressed across all tumor sub-types, including triple negative

breast cancers (TNBC), and is significantly associated with improved patient outcomes in women taking tamoxifen for adjuvant therapy of resected, ER α -positive, early stage breast cancer. Based on these observations, we explored the utility of therapeutically targeting ER β 1 using ER β -specific agonists and multiple anti-estrogenic compounds in both ER α -positive and ER α -negative breast cancers using a number of cell model systems. Our results demonstrate that targeting this receptor results in potent anti-proliferative effects in multiple breast cancer sub-types. However, the effectiveness of these two classes of drugs varies dramatically as a function of ER α status.

Methods

Study cohorts

For this study, 3 distinct patient cohorts were utilized to examine the prevalence of ER β 1 expression across multiple breast tumor sub-types and to determine its association with other prognostic biomarkers and response to endocrine therapy. The first cohort (C1) is a retrospectively assembled cohort of 184 women who underwent primary breast cancer surgery at Mayo Clinic Rochester between 2001 and 2008. The second cohort (C2) is a retrospectively assembled cohort of 68 patients who underwent primary breast cancer surgery between 1998 and 2011 at Mayo Clinic Scottsdale, selected for the presence of TNBC on central pathology testing. The third cohort (C3) is a secondary analysis of a prospective adjuvant tamoxifen study in postmenopausal women with early stage ER α -positive breast cancer (North Central Cancer Treatment Group (NCCTG) Trial 89-30-52)) who were randomized to adjuvant treatment with tamoxifen (20 mg per day orally for 5 years) plus fluoxymesterone (10 mg orally twice per day for 1 year) and who had a tumor specimen available from their primary surgery (177 of 258 eligible patients) [52]. All patients enrolled in this study provided informed consent and the use of patient tumor samples for immunohistochemical analysis was approved by the Institutional Review Board at Mayo Clinic (protocol #: 13-000585). Patient characteristics within these three cohorts are shown in Table 1 and the molecular and histologic subtypes represented within each cohort is shown in Table 2.

Tissue microarrays and IHC testing of patient samples

Tissue microarrays (TMAs) were constructed for cohorts C1 and C2 using three 0.6 mm tissue cores collected from areas of invasive breast cancer on each tissue block. Five micron sections were cut for immunostaining and analysis as previously described [20]. Full tumor sections from cohort C3 were processed in an identical manner. For HER2 staining, the HercepTest kit (Dako, Carpinteria, CA) was utilized following the

Table 1 Patient characteristics and clinicopathological variables for each of three cohorts

Patient characteristics	Cohort 1 n = 184	Cohort 2 n = 68	Cohort 3 n = 177
median age (range)	58 (28–87)	60 (27–82)	68 (48–89)
Histology			
Ductal	138 (75.0%)	52 (76.5%)	143 (80.8%)
Lobular	28 (15.2%)	0	16 (9.0%)
Other	18 (9.8%)	16 (23.5%)	18 (10.2%)
Receptor status			
ERpos/PRpos or unknown	143 (77.3%)	0	177 (100%)
ERpos/PRneg	27 (14.6%)	0	0
ERneg/PRneg	14 (8.1%)	68 (100%)	0
Her2 status			
positive	27 (14.7%)	0	15 (8.5%)
negative	145 (78.8%)	68 (100%)	160 (90.4%)
unknown	12 (6.5%)	0	2 (1.1%)
Max tumor dimension			
0.1-2.0 cm	115 (62.5%)	42 (61.8%)	
2.1-5.0 cm	51 (27.6%)	21 (30.9%)	
5.1+ cm	18 (9.7%)	5 (7.4%)	*
Number of positive nodes			
0	112 (60.9%)	49 (73.5%)	110 (62.1%)
1-3	46 (25.0%)	14 (20.6%)	47 (26.6%)
4-9	16 (8.7%)	3 (4.4%)	13 (7.3%)
10+	10 (5.4%)	1 (1.5%)	7 (4.0%)
unknown	0	1 (1.5%)	0
Nuclear Grade 3	45 (24.5%)	55 (80.9%)	41 (23.2%)
max Ki67 across all cores			
not done	3 (1.6%)	6 (8.7%)	177 (100%)
0 – 10%	61 (33.2%)	16 (23.2%)	
10.1 – 25%	59 (32.1%)	9 (13.0%)	
25.1 – 50%	40 (21.7%)	6 (8.7%)	
50.1-100%	21 (11.4%)	32 (46.4%)	
ERβ1 nuclear expression			
negative/low (0–2)	121 (65.7%)	51 (75.0%)	32 (18.1%)
moderate (3–5)	59 (32.1%)	17 (25.0%)	96 (54.2%)
high (6–7)	4 (2.2%)	0	49 (27.7%)
ERβ1 cytoplasmic expression			
negative/low (0–2)	164 (89.1%)	45 (66.2%)	1 (0.6%)
moderate (3–5)	20 (10.9%)	21 (30.9%)	52 (29.4%)
high (6–7)	0	2 (2.9%)	124 (70.1%)

*tumor size collected as < 3 m vs. ≥ 3 cm: 140 (79.1%) vs. 37 (20.5%).

manufacturers protocol. All other IHC stains were performed on a Leica Bond III stainer using the following antibodies: 1) A monoclonal ERβ1 PPG5/10 antibody;

1:75 dilution (Thermo Scientific, Waltham, MA), 2) a monoclonal ERα 1D5 antibody; 1:300 dilution (Dako, Carpinteria, CA), 3) a monoclonal PgR 636 antibody; 1:800 dilution (Dako) and 4) a monoclonal Ki67 MIB-1 antibody; 1:300 dilution (Dako). ERα and PgR positivity was determined using standard procedures. Ki67 was scored as previously described [53]. The monoclonal ERβ1 antibody used in this study has been shown to be highly specific and sensitive for detection of only the full-length form of this receptor in IHC studies [20,50,51]. Specifically, we have utilized multiple cell model systems which either transiently express ERβ1, or stably express this receptor under the control of a doxycycline inducible promoter, to fully characterize the detection methods and optimal dilution of the PPG5/10 antibody for IHC purposes [20]. Additionally, we have shown that this antibody does not cross-react with ERα or any of the ERβ splice variant forms [20]. Finally, this antibody was compared to multiple other commercially available ERβ specific antibodies and was shown to be one of the best for use in IHC studies using human breast tissue [50]. All slides were reviewed by a dedicated breast cancer pathologist and ERβ1 protein levels were evaluated in both nuclei and cytoplasm. Pathological categorization of ERβ1 levels was determined as a sum of the extent and intensity scores. The extent of staining was scored as follows: 0: less than 1% positive cells, 1: 1%-25%, 2: 26%-50%, 3: 51%-75% and 4: 76%-100%. Intensity of staining was scored as none (0), weak (1), moderate (2) or strong (3). The resulting scores were grouped into 3 categories, namely, ERβ1-negative/low (0–2), ERβ1-moderate (3–5) and ERβ1-high (6–7) and the percentage of tumors falling into these three groups for both nuclear and cytoplasmic staining are indicated throughout this manuscript. A representative tumor determined to be ERβ1-negative, moderate and high is shown in Figure 1 for both nuclear and cytoplasmic localization.

Cell culture, chemicals and reagents

Parental and ERβ1-expressing MCF7 cells [12] and doxycycline-inducible Hs578T-ERβ1 cells [8] were cultured as previously described. Doxycycline-inducible ERβ1-expressing MDA-MB-231 cell lines were established using the T-REx™ System (Invitrogen) as previously described [9] and were maintained in DMEM/F12 medium supplemented with 10% FBS, 1% AA, 5 mg/L blasticidin S and 500 mg/L zeocin. Charcoal-stripped fetal bovine serum (CS-FBS) was purchased from Gemini Bio-Products (West Sacramento, CA). 17β-estradiol (E2), (Z)-tamoxifen, (Z)-4-hydroxy-tamoxifen and doxycycline (Dox) were purchased from Sigma-Aldrich (St. Louis, MO). (Z)-endoxifen was synthesized by the National Cancer Institute (Bethesda, MD). The ERβ-specific agonists; DPN, WAY200070, FERb 033 and Liquiritigenin, as well

Table 2 ERβ1 expression levels by morphology and subtype

	ERβ1 expression	Cohort 1 n = 184**		Cohort 2 n = 68		n = 177	
		Nucleus	Cytoplasm	Nucleus	Cytoplasm	Nucleus	Cytoplasm
Molecular Subtype							
Luminal A (ERα +/ HER2 -/ Ki67 ≤ 10)	Neg/low	31 (18.3)	47 (25.5)				
	Moderate	18 (10.7)	3 (1.6)				
	High	1 (0.6)	0				
Luminal B (ERα +/ HER2 -/ Ki67 > 10)	Neg/low	59 (34.9)	76 (41.3)				
	Moderate	25 (14.8)	11 (6.0)				
	High	3 (1.8)	0				
Her2+	Neg/low	20 (11.8)	24 (13.0)				
	Moderate	7 (4.1)	3 (1.6)				
	High	0	0				
Triple Negative (ERα -/PR- / HER2 -)	Neg/low	1 (0.6)	5 (2.7)	51 (75.0)	45 (66.2)		
	Moderate	4 (2.4)	0	17 (25.0)	21 (30.9)		
	High	0	0	0	2 (2.9)		
Histologic Subtype							
Ductal	Neg/low	93 (50.5)	120 (65.2)	39 (57.4)	34 (50.0)	28 (15.8)	0
	Moderate	42 (22.8)	18 (9.8)	13 (19.1)	16 (23.5)	79 (44.6)	45 (25.4)
	High	3 (1.6)	0	0	2 (2.9)	36 (20.3)	98 (55.4)
Lobular	Neg/low	18 (9.8)	26 (14.1)	0	0	1 (0.6)	0
	Moderate	9 (4.9)	2 (1.1)	0	0	10 (5.7)	3 (1.7)
	High	1 (0.5)	0	0	0	5 (2.8)	13 (7.3)
Other	Neg/low	10 (5.4)	18 (9.8)	12 (17.6)	11 (16.2)	3 (1.7)	1 (0.6)
	Moderate	8 (4.4)	0	4 (5.9)	5 (7.4)	7 (4.0)	4 (2.3)
	High	0	0	0	0	8 (4.5)	13 (7.3)

*ki67 not performed **unable to determine molecular subtype in 15 Cohort 1 pts.




Figure 1 Immunohistochemical staining for ERβ1 in human breast tumors. Representative images depicting tumors with negative/low, moderate or high expression of nuclear and cytoplasmic ERβ1 as detected using the PPG5/10 antibody.

as the pure ER antagonist ICI 182,780, were purchased from Tocris Bioscience (Bristol, United Kingdom).

Real-time reverse transcription polymerase chain reaction

To confirm stable integration and doxycycline inducibility of ER β 1 in the MDA-MB-231 clonal cell lines, cells were plated in 6-well tissue culture plates in the presence and absence of doxycycline (0.1 μ g/ml). Following 24 hours of culture, total RNA was isolated using Trizol reagent (Invitrogen), cDNA was synthesized and real-time PCR using ER β specific primers was performed as previously described [54] and two clones (#4 and 12) exhibiting substantial expression of ER β 1 were chosen for further analysis. To confirm functionality of ER β 1, cells were plated as described above using phenol red-free 10% CS-FBS containing media and treated with ethanol or estradiol (1nM) for 24 hours. RT-PCR was performed using primers specific for the progesterone receptor (PR), PS2 and KLF10 as previously described [12].

Western blotting

MDA-MB-231-ER β 1 cell lines #4 and #12 were plated in 6-well plates in the presence and absence of doxycycline for 24 hours. Cell lysates were harvested using NETN buffer (150 mM NaCl, 1 mM EDTA, 20 mM Tris [pH 8.0], 0.5% Nonidet P-40), protein concentrations were determined using Bradford Reagent and western blots were performed using Flag (M2, Sigma-Aldrich) and α -Tubulin (DM 1A, Sigma-Aldrich) specific antibodies as previously described [12].

Proliferation assays

In order to assess anchorage dependent cell proliferation, a crystal violet assay was utilized. This method is well accepted to be reflective of cell number and does not rely on measurements related to mitochondrial activity or intracellular ATP levels that could be compromised by treatments targeting ER β which is known to be expressed in mitochondria [55-59]. Briefly, cells were plated in replicates of 8 at a density of 1000 cells per well in 96-well tissue culture plates using 10% CS-FBS containing phenol red-free medium. Twenty-four hours after plating, cells were treated with indicated ligands. Cell culture media was replaced every 3 days and crystal violet staining was performed following 12 days of treatment. Crystal violet staining was quantitated using a plate reader set at a wavelength of 550 nm and replicates were averaged among treatment groups.

Statistical analyses

Descriptive statistics were used to summarize nuclear and cytoplasmic ER β 1 expression levels in each patient cohort. The primary outcome of interest was the recurrence-free interval defined as the time from randomization to

documentation of a local, regional, or distant breast recurrence. A stratified log-rank test with strata defined by whether tumor size was ≥ 3 cm and lymph nodes were positive for disease was used to determine whether the recurrence-free interval differed with respect to nuclear or cytoplasmic ER β 1 expression. For all real-time PCR and proliferation assays, a two-sided Student's *t*-test was utilized. P-values < 0.05 were considered to be statistically significant.

Results

Association of ER β 1 with other prognostic biomarkers and tumor grade in an unselected patient cohort

In a cohort of 184 women with primary breast cancer (C1), nuclear ER β 1 expression was determined to be low/negative in 121 (65.7%), moderate in 59 (32.1%) and high in 4 (2.2%) women (Table 1). This is in contrast to cytoplasmic ER β 1 expression that was low/negative in 164 (89.1%) and moderate in 20 (10.9%) women with no tumor exhibiting high cytoplasmic expression (Table 1). The concordance between nuclear and cytoplasmic ER β 1 expression was 66.3% (122/184). ER β 1 was detected across all molecular and histologic subtypes of breast cancer within this patient cohort (Table 2). Moderate to high levels of nuclear ER β 1 expression were detected in 56 of the 170 (32.9%) ER α -positive cases and 7 of the 14 (50.0%) ER α -negative cases (Table 3). In contrast, cytoplasmic ER β 1 expression was similar between the ER α -positive and ER α -negative cancers with approximately 10% of these tumors having moderate to high expression (Table 3). The distributions of nuclear and cytoplasmic ER β 1 expression were similar between HER2 positive and negative tumors; Ki67 low and high tumors; high and low grade tumors; and cases with node positive or negative disease (Table 3).

Expression of ER β 1 in triple negative breast cancers

Due to the low number of ER α -negative tumors in our unselected patient cohort (C1), we leveraged another cohort of 68 cases (C2) with confirmed primary TNBC. Nuclear ER β 1 expression was determined to be low/negative in 51 (75.0%) and moderate in 17 (25.0%) tumors (Table 1). This is similar to cytoplasmic ER β 1 expression that was low/negative in 45 (66.2%), moderate in 21 (30.9%) and high in 2 (2.9%) tumors (Table 1). The concordance between nuclear and cytoplasmic ER β 1 expression was 70.6% (48/68). Ki67 results were available in 63 cases. Among the 16 cases whose Ki67 level was not elevated ($\leq 10\%$), 1 case had moderate levels of both nuclear and cytoplasmic ER β 1 a second case had moderate nuclear expression but negative/low cytoplasmic expression (Table 4). The remaining 14 cases with low Ki67 levels had negative/low nuclear and cytoplasmic ER β 1 expression (Table 4). In contrast, 25 (54.3%) of the

Table 3 ERβ1 expression levels in a population of breast cancer patients diagnosed at Mayo Clinic Rochester (cohort 1) and its association with other biomarkers, tumor grade and nodal status

Disease characteristics		ERβ1 status	Nucleus # of Pts. (%)	Cytoplasm # of Pts. (%)
ERα	Positive (n = 170)	Negative/Low	114 (67.1)	151 (88.8)
		Moderate	52 (30.6)	19 (11.2)
		High	4 (2.4)	0 (0)
	Negative (n = 14)	Negative/Low	7 (50.0)	13 (92.9)
		Moderate	7 (50.0)	1 (7.1)
		High	0 (0)	0 (0)
HER2	Positive (n = 27)	Negative/Low	20 (74.1)	24 (88.9)
		Moderate	7 (25.9)	3 (11.1)
		High	0 (0)	0 (0)
	Negative (n = 145)	Negative/Low	94 (64.8)	131 (90.4)
		Moderate	47 (32.4)	14 (9.7)
		High	4 (2.8)	0
Ki67	≤ 10% (61)	Negative/Low	39 (63.9)	57 (93.4)
		Moderate	21 (34.4)	4 (6.6)
		High	1 (1.6)	0 (0)
	> 10% (120)	Negative/Low	79 (65.8)	104 (86.7)
		Moderate	38 (31.7)	16 (13.3)
		High	3 (2.5)	0 (0)
Tumor grade	Grade 1–2 (136)	Negative/Low	89 (65.4)	122 (89.7)
		Moderate	44 (32.4)	14 (10.3)
		High	3 (2.2)	0
	Grade 3 (45)	Negative/Low	31 (68.9)	41 (91.1)
		Moderate	13 (28.9)	4 (8.9)
		High	1 (2.2)	0 (0)
Nodal disease	Present (72)	Negative/Low	44 (61.1)	65 (90.3)
		Moderate	24 (33.3)	7 (9.7)
		High	4 (5.6)	0 (0)
	Not present (112)	Negative/Low	77 (68.8)	99 (88.4)
		Moderate	35 (31.3)	13 (11.6)
		High	0 (0)	0 (0)

46 cases with elevated Ki67 levels had moderate to high ERβ1 expression in the nucleus and/or cytoplasm (Table 4).

ERβ and outcomes with adjuvant endocrine therapy

A cohort of 177 postmenopausal women with early stage ERα-positive breast cancer enrolled onto NCCTG 89-30-52 who were randomized to the adjuvant treatment with tamoxifen plus fluoxymesterone arm (C3) was used to assess whether ERβ1 expression is associated with the likelihood of a breast cancer event (local, regional or distant recurrence). With a median length of follow-up of 19.5 years, 56 women are currently alive without disease recurrence, 11 are alive having had disease recurrence

and/or a second primary cancer, 49 have died following disease recurrence and/or a second primary cancer and 61 have died without disease recurrence or a second primary disease. Nuclear ERβ1 expression was determined to be low/negative in 32 (18.1%), moderate in 96 (54.2%) and high in 49 (27.7%) women (Table 1). In contrast, cytoplasmic ERβ1 expression was determined to be low/negative in 1 (0.6%), moderate in 52 (29.3%) and high in 124 (70.1%) women (Table 1). As was the case with the other two cohorts, ERβ1 expression was detected across all histologic subtypes of breast cancer (Table 2). The recurrence-free interval (time to local, regional, distant progression) was found to differ with respect to degree

Table 4 ERβ1 expression levels in triple negative breast tumors and its association with Ki67 expression levels

Ki67 expression	ERβ1 Status	ERβ1 Status cytoplasm		
		Negative/Low	Moderate	High
Nucleus				
Ki67 > 10% (46)	Negative/Low	21 (45.6%)	11 (23.9%)	0
	Moderate	4 (8.7%)	8 (17.4%)	2 (4.4%)
	High	0	0	0
Ki67 ≤ 10% (16)	Negative/Low	14 (87.5%)	0	0
	Moderate	1 (6.3%)	1 (6.3%)	0
	High	0	0	0

of nuclear ERβ1 expression (stratified log-rank test, adjusted for tumor size and node metastasis $p = 0.023$) with 10 year recurrence-free rates of 74%, 84%, and 88% for patients whose cancers had negative/low, moderate and high levels of ERβ1, respectively (Figure 2). However, the recurrence-free interval was not found to differ with respect to degree of cytoplasmic ERβ1 expression (stratified log-rank test $p = 0.623$) with 10 year recurrence-free rates of 82% and 84% for patients whose cancers had moderate and high cytoplasmic expression of ERβ1, respectively (Additional file 1: Figure S1).

Therapeutic targeting of ERβ1 in ERα positive breast cancer cells

Based on the observation that ERβ1 expression is associated with lower rates of recurrence in ERα positive breast cancer, we sought to further characterize the effects of multiple targeted therapies using a breast cancer

cell line designed to mimic this tumor sub-type. Therefore, we utilized parental and ERβ1-expressing MCF7 cells previously developed in our laboratory [12]. As a first step, we analyzed the role of ERβ1 in mediating the pro-proliferative effects of 17-beta estradiol (estrogen) and the anti-proliferative effects of anti-estrogenic compounds. As expected, estrogen treatment was shown to induce proliferation in both cell lines; however, the magnitude of induction was decreased in ERβ1 expressing cells (Figure 3A). Tamoxifen had no effect on estrogen-induced proliferation rates regardless of ERβ1 expression (Figure 3A). Interestingly, a low dose (10 nM) of 4HT increased the proliferation rate of parental MCF7 cells above that of estrogen treatment alone, an effect that was not observed in cells expressing ERβ1 (Figure 3A). Higher doses (100 nM) of endoxifen and 4HT, as well as a low dose (10 nM) of ICI, resulted in almost complete blockade of estrogen-induced proliferation in ERβ1-expressing cells but not in parental cells expressing only ERα (Figure 3A).

We next sought to determine if ERβ-specific agonists modulated the proliferation rates of these cells in both the presence and absence of estrogen treatment. In the absence of estrogen (Figure 3B), low (10 nM) and moderate (100 nM) doses of DPN induced proliferation in both parental and ERβ1-expressing MCF7 cells. The magnitude of induction following DPN treatment was nearly identical to that of estrogen treatment in parental MCF7 cells but less than that of estrogen in ERβ1-expressing cells (Figure 3B). Low doses of WAY200070 and FERb 033 had little to no effect on the proliferation rates of parental or ERβ1-expressing cells while higher doses induced proliferation (Figure 3B). A similar pattern was observed following treatment with liquiritigenin with





Figure 2 Increased nuclear ERβ1 expression is associated with prolonged recurrence-free interval in women treated with adjuvant tamoxifen and fluoxymesterone therapy. Kaplan-Meier estimates of breast cancer recurrence-free interval in patients with negative/low, moderate or high nuclear expression of ERβ1 are shown.



(See figure on previous page.)

Figure 3 Effects of anti-estrogenic (A) and ER β agonist (B) treatment on the proliferation rates of MCF7 and MCF7-ER β 1 expressing cells. Crystal violet assays were used to determine proliferation rates following indicated treatments for 12 days. P-values < 0.05 were considered to be statistically significant. *Denotes significant difference between indicated treatment and vehicle control treated cells and # between indicated treatment and estrogen treated cells.

the exception that low doses of this compound were inhibitory regardless of ER β 1 expression (Figure 3B). When each ER β -specific agonist was administered in the presence of estrogen, the observed dose-dependent effects were abrogated in both cell lines and the proliferation rates of parental and ER β 1-expressing cells were either equivalent or slightly greater than that of estrogen treatment alone (Additional file 2: Figure S2).

Development and characterization of MDA-MB-231-ER β 1 cell lines

Since approximately 25% of TNBC were shown to express nuclear ER β 1 (Table 1; cohort 2), we next sought to determine whether expression of ER β 1 in MDA-MB-231 cells, a well-characterized model of TNBC, would alter the cell's response to ER β targeting treatments. Two clonal cell lines (#4 and #12) exhibiting robust doxycycline induced expression of ER β 1 mRNA and protein were chosen for further analysis (Figure 4A). To confirm that expression of ER β 1 was exclusively limited to the presence of doxycycline and that the expressed receptor was functional, cells were treated with vehicle, estrogen (1 nM) or estrogen plus ICI (100 nM) for 24 hours and the expression profiles of known ER β 1 target genes were examined. As shown in Figure 4B, these genes were significantly induced following estrogen treatment in the presence of doxycycline, an effect that was completely blocked by the addition of ICI. However, these genes were not induced by estrogen in the absence of doxycycline confirming that these cells do not endogenously express any form of the estrogen receptor (Figure 4B).

Effects of anti-estrogens and ER β -specific agonists on the proliferation rates of ER β 1-positive triple negative breast cancer cells


We next performed a series of proliferation assays to determine which therapeutic strategies may be most effective for the treatment of ER β 1 positive TNBC. Interestingly, estrogen treatment (1 nM) was shown to substantially inhibit the proliferation rates of MDA-MB-231-ER β 1 cells (Figure 5), an effect that was not observed in the absence of doxycycline (data not shown). The addition of multiple anti-estrogens significantly reversed the inhibitory effect of estrogen in MDA-MB-231-ER β 1 cells (Figure 5A). In order to ensure that these effects were not unique to the MDA-MB-231 cell line, identical assays were performed using Hs578T-ER β 1 expressing cells [8]. Estrogen treatment

significantly repressed proliferation of Hs578T-ER β 1 cells, effects that were reversed following the addition of endoxifen, 4HT or ICI (Figure 5A). Similar responses were observed in the MDA-MB-231-ER β 1 clonal cell line #12 (Additional file 3: Figure S3A).

Since estrogen treatment resulted in substantial reductions in the proliferation rates of ER β 1-expressing TNBC cells, we next analyzed the effects of multiple ER β -specific agonists in these two cell lines. All of the ER β -specific agonists tested significantly inhibited the proliferation rates of MDA-MB-231-ER β 1 and Hs578T-ER β 1 cells with DPN and WAY200070 eliciting the greatest responses (Figure 5B). Nearly identical responses were observed in the MDA-MB-231-ER β 1 clonal cell line #12 (Additional file 3: Figure S3B). Combinatorial treatment with 1 nM concentrations of estrogen plus ER β -specific agonists did not result in greater anti-proliferative effects (data not shown).

Discussion

In this study, we have compared the nuclear and cytoplasmic expression profiles of ER β 1 across multiple sub-types of breast cancer and in a population of well annotated patients treated with adjuvant endocrine therapy. Our results have revealed that ER β 1 expression, while present in nearly all normal breast epithelium, is lost in many breast cancers. However, the expression of ER β 1 is associated with substantially improved anti-tumor effects in ER α -positive tamoxifen treated breast cancer, as well as potent anti-proliferative effects *in vitro*, confirming its role as a tumor suppressor. Interestingly, the biological effects of therapeutically targeting ER β appear to be critically correlated with the presence of ER α . In ER α -positive cell lines, expression of ER β 1 enhanced the anti-proliferative effects of anti-estrogenic therapies including endoxifen, 4HT and ICI. However, targeting ER β with specific agonists in MCF7 cells was not an effective treatment strategy and led to growth stimulation in most instances, likely due to the known cross-reactivity of these compounds with ER α at higher concentrations (100 nM). In contrast, activation of ER β 1 with estrogen or ER β -specific agonists was shown to substantially repress TNBC cell proliferation rates while the use of anti-estrogens was ineffective and in some cases resulted in stimulation of cell proliferation. Taken together, our studies have comprehensively analyzed the protein expression profiles of ER β 1 across multiple breast cancer sub-types and demonstrated critical roles for this




receptor in mediating the effectiveness of multiple therapeutic treatment strategies for breast cancer patients that are related in part to the presence and absence of ERα expression.

Using a well-validated and highly specific antibody and a large cohort of unselected breast cancer patients, we

have shown that ERβ1 expression is lost in most cancers as approximately 65% of all breast tumors were determined to be ERβ1-negative. When ERβ1 is expressed, it can exhibit both nuclear and cytoplasmic localization in tumor cells. These data are in agreement with the largest study conducted to date that reported a frequency of

A



B




Figure 5 Effects of anti-estrogenic (A) and ERβ agonist (B) treatment on the proliferation rates of MDA-MB-231-ERβ1 and Hs578t-ERβ1 cells. Crystal violet assays were used to determine proliferation rates following indicated treatments for 12 days. P-values < 0.05 were considered to be statistically significant. *Denotes significant difference between indicated treatment and vehicle control treated cells and # between indicated treatment and estrogen treated cells.

39% for nuclear ER β 1 expression in ER α -positive invasive breast cancers using the same antibody as was used in this study (PPG5/10) [46]. Similar to our findings, this study also did not find an association between ER β 1 expression and other clinicopathological factors [46]. However, it should be noted that others have reported somewhat higher frequencies of ER β 1 positivity using this antibody [26,29,30,45,60], and one study concluded that ER β 1 was significantly associated with expression of ER α and PR and inversely associated with HER2 overexpression [60]. We also detected ER β 1 expression across all molecular and histologic subtypes of breast cancer. This is in contrast to a recent publication by Huang et al., in which they demonstrated that ER β 1 is only expressed in lobular, and not ductal, carcinomas [21]. These discrepancies may be explained by the use of different antibody dilutions and detection techniques as well as scoring criteria. In alignment with the study by Novelli and colleagues [45], we utilized a conservative approach and categorized tumors exhibiting low expression of ER β 1 as negative.

Although many studies have examined the expression profiles of ER β 1 in ER α -positive tumors, fewer have reported the frequencies of ER β 1 in ER α -negative tumors. To our knowledge, the data reported here are the first to analyze ER β 1 expression in a patient cohort of confirmed TNBC. Of the 68 TNBCs analyzed, 24% expressed nuclear ER β 1 with approximately 34% exhibiting cytoplasmic localization of this full-length receptor. These results are consistent with previous reports, which have suggested that between 24% and 44% of ER α negative tumors, but not necessarily TNBC, are ER β positive [23,30,33,61,62]. However, two other publications have reported higher frequencies of ER β in ER α -negative tumors [32,63] although it should be noted that in both of these studies, PR and HER2 status were not analyzed and ER α status was determined by ligand binding assays, not immunohistochemistry. None of these studies commented on cytoplasmic expression of ER β and some studies utilized antibodies that are not specific for full length ER β and instead can cross-react with its splice variant forms. In contrast to our data in ER α -positive tumors, ER β 1 expression in TNBC was associated with higher expression of Ki67 as has been reported by others [30,32,33,64]. Taken together, these data suggest that the functions of ER β 1 in the absence of ER α expression may be substantially different. Additionally, it is possible that Ki67 levels may vary in ER β 1-positive TNBC based on menopausal status, a possibility that has yet to be examined.

Using a well-annotated cohort of 177 ER α -positive breast cancer patients who were treated with adjuvant tamoxifen (20 mg/day for 5 years) plus fluoxymesterone (10 mg orally twice per day for 1 year), we found that

increased expression of nuclear ER β 1 was associated with prolonged recurrence-free interval. To our knowledge, our results are the first prospective-retrospective study of the prognostic value of ER β 1 in patients treated with tamoxifen in the adjuvant setting. These data are consistent with previous studies that have utilized various antibodies specific for full length ER β 1 and have demonstrated that high expression correlates with increased response to tamoxifen therapy [40,46,65], improved disease free survival [22,25-27], longer overall survival [27,36] and no relapse within 5 years [38]. Although all of the patients included in this cohort were ER α positive, a recent publication has also suggested that ER β 1 may have additional predictive value for tamoxifen responsiveness in ER α negative tumors which express high levels of SRAP [48]. In contrast to prior literature, we also analyzed cytoplasmic staining for ER β 1 and demonstrated no association with the risk of recurrence. While the basis for cytoplasmic localization of ER β 1 is not well understood and remains somewhat controversial, a number of reports have demonstrated that ER β 1 is expressed in mitochondria [55-59] and it is possible that this in part explains the detection of this hormone receptor within this sub-cellular compartment. Interestingly, a recent study has suggested that tamoxifen resistance may develop by agonizing mitochondrial ER β 1 resulting in up-regulation of MnSOD activity and ultimately enhancing cell survival and growth [59]. This could partially explain why cytoplasmic expression of ER β 1 did not correlate with improved tamoxifen responsiveness, as was the case for nuclear ER β 1, in the cohort of patients analyzed here. Additionally, these observations may explain why some studies (which did not distinguish between nuclear and cytoplasmic expression) did not detect an association between ER β expression and outcomes in tamoxifen treated patients [28] while one study actually reported a non-significant trend towards increased recurrence rates in women with high ER β expression [30]. Overall, these results suggest that determination of nuclear ER β 1 status will improve our ability to predict an individual's likelihood of response to adjuvant tamoxifen therapy, effects which may be magnified in the absence of cytoplasmic ER β 1.

Given that ER β 1 is expressed in both ER α -positive and ER α -negative breast tumors, we sought to compare a number of different therapeutic strategies to determine which might be most effective for the treatment of patients with ER β 1-positive breast cancer. Using MCF7-ER β 1 and Hs578T-ER β 1 cell lines previously developed in our laboratory [8,12], as well as a newly developed MDA-MB-231-ER β 1 cell line whose characterization is described in the present manuscript, we performed cell proliferation assays using anti-estrogenic compounds as well as ER β -specific agonists. In the ER α -positive MCF7

cell line, ER β 1 expression was shown to diminish the pro-proliferative effects of estrogen, a phenomenon reported previously in ER α -positive cells [66-68]. These results are consistent with the observation that ER α -positive/ER β 1-positive tumors typically have reduced expression of Ki67 relative to ER α -positive/ER β 1-negative tumors. Additionally, ER β 1 expression in these cells enhanced the anti-estrogenic effect of endoxifen, 4HT, and ICI. These data are also consistent with previous reports by our laboratory [12] and others [68-70] demonstrating that ER β 1 expression improves the anti-proliferative effects of 4HT, raloxifene and ICI *in vitro*. These data also correlate with our present studies demonstrating that moderate to high expression of ER β 1 in human breast cancers is associated with improved patient outcomes following tamoxifen therapy.

Treatment of MCF7-ER β 1 cells with 4 different ER β -specific agonists resulted in variable effects on cell proliferation. Moderate (100 nM) concentrations of all of the ER β -specific agonists led to induction of cell proliferation. These effects were also observed in the parental MCF7 cell line which does not express ER β 1 and are therefore highly likely to occur through the known activation of ER α at these doses [71,72]. Low (10 nM) concentrations of these compounds were shown to have minimal effects on cell proliferation rates with the exception liquiritigenin which actually inhibited MCF7-ER β 1 cell proliferation following 12 days of treatment. However, a nearly identical effect was observed in the parental cell line suggesting that ER β 1 is not responsible for mediating this inhibitory effect. Our results are consistent with previous reports demonstrating that low concentrations of WAY200070 and DPN have little to no effect on the proliferation rates of T47D parental or ER β 1 expressing cells [66,73,74] while higher concentrations of DPN were shown to stimulate proliferation rates above that of vehicle control treated cells [66]. Taken together, these studies suggest that the use of modern day ER β -specific agonists for ER β 1-positive tumors is highly unlikely to be beneficial in the treatment of breast tumors which also express ER α . Instead, the utilization of anti-estrogenic therapies is likely to remain a superior choice for this sub-type of breast cancer; at least until more specific and potent ER β -specific agonists are developed.

In contrast to the results observed in ER α -expressing breast cancer cells, estrogen treatment of two different TNBC lines which were engineered to express ER β 1 led to substantial reductions in cell proliferation rates. These results are consistent with previous reports from our laboratory and others demonstrating that expression of ER β 1 in TNBC cells can lead to suppression of both basal and/or estrogen-mediated proliferation rates [8,75,76]. As expected, the addition of anti-estrogens such as endoxifen,

4HT, and ICI blocked these estrogen-mediated effects. However, our present data are the first to demonstrate that the use of ER β -specific agonists in ER β 1-positive TNBC cells can elicit at least equivalent anti-proliferative effects compared to estrogen treatment alone. As might be expected, combinatorial treatments of estrogen plus ER β -specific agonists did not result in additive or synergistic effects confirming that these compounds function specifically through ER β 1 in our model systems. Overall, our results indicate that further study of ER β -targeted therapies is warranted for the treatment of patients with ER β 1-positive TNBC, a subgroup of patients with extremely poor outcomes and for which no form of a targeted cancer therapy is currently available.

Conclusions

In summary, we have examined the expression patterns of ER β 1 across all sub-types of breast cancer using a highly specific and sensitive monoclonal antibody and have reaffirmed the importance of ER β 1 as a tumor suppressor. Specifically, nuclear expression of ER β 1 is associated with significantly improved outcomes in women treated with adjuvant tamoxifen therapy and these observations were confirmed in cell proliferation assays which demonstrated that ER β 1 expression in ER α -positive MCF7 cells significantly improved their responsiveness to anti-estrogenic therapies. However, activation of ER β 1 with either estrogen or ER β -specific agonists was shown to result in substantial inhibition of cell proliferation in TNBC cells. These results lay the foundation for future studies aimed at analyzing the anti-tumor activity of ER β -agonists for the treatment of ER β 1-positive TNBC. The outcomes of such studies could have a dramatic impact on our ability to offer alternative therapies and more effectively treat individuals with this form of the disease.

Additional files

Additional file 1: Figure S1. Cytoplasmic expression of ER β 1 is not associated with differences in recurrence-free interval in women treated with adjuvant tamoxifen and fluoxymesterone therapy. Kaplan-Meier estimates of breast cancer recurrence-free interval in patients with moderate or high cytoplasmic expression of ER β are shown.

Additional file 2: Figure S2. Effects of ER β agonist + estrogen treatment on the proliferation rates of MCF7 and MCF7-ER β 1 expressing cells. Crystal violet assays were used to determine proliferation rates following indicated treatments for 12 days. P-values < 0.05 were considered to be statistically significant. *Denotes significant difference between indicated treatment and vehicle control treated cells and #between indicated treatment and estrogen treated cells.

Additional file 3: Figure S3. Effects of anti-estrogenic (A) and ER β agonist (B) treatment on the proliferation rates of MDA-MB-231-ER β 1 cells (clone #12). Crystal violet assays were used to determine proliferation rates following indicated treatments for 12 days. P-values < 0.05 were considered to be statistically significant. *Denotes significant difference between indicated treatment and vehicle control treated cells and #between indicated treatment and estrogen treated cells.

Competing interests

The authors declare that they have no competing interests.

Authors' contribution

Concept and design: JMR, MS, XW, WLL, TCS, MPG, JNI and JRH. Collection and assembly of data: JMR, VJS, MS, XW, VN, KSP, SSS, HEC, AEM, BAP, FJC, JEO, CR, WLL, MPG, JNI, JRH. Data analysis and interpretation: JMR, VJS, MS, AG, WLL, TCS, MPG, JNI, JRH. Manuscript writing: JMR, VJS, MS, MPG, JNI and JRH. All authors read and approved the final manuscript.

Authors' information

James N. Ingle and John R. Hawse are co-senior authors.

Acknowledgements

The authors would like to thank Dr. Thomas J. Flotte for his assistance with capturing the immunohistochemical staining images. The work presented here was supported by the Mayo Clinic Breast Cancer SPORC: P50CA116201 (JNI, MPG and JRH), two Breast Cancer Research Foundation grants (JNI and FJC), Susan G. Komen for the Cure KG100142 (JRH), a Paul Calabresi K12 award NCI CA90628 (JRH), a generous gift from Bruce and Martha Atwater (TCS, JRH, MPG, JNI), the David and Margaret T. Grohne Family Foundation (FJC), the Ting Tsung and Wei Fong Chao Foundation (FJC), the Mayo Clinic Graduate School (JMR) and the Mayo Foundation (TCS).

Author details

¹Department of Biochemistry and Molecular Biology, Mayo Clinic, 16-01B Guggenheim Building, 200 First St. SW, Rochester, MN 55905, USA. ²Department of Molecular Pharmacology and Experimental Therapeutics, Mayo Clinic, Rochester, MN, USA. ³Department of Biomedical Statistics and Informatics, Mayo Clinic, Rochester, MN, USA. ⁴Department of Laboratory Medicine and Pathology, Mayo Clinic, Rochester, MN, USA. ⁵The Department of Pathology, Dunedin School of Medicine, University of Otago, Dunedin, New Zealand. ⁶Department of Laboratory Medicine and Pathology, Mayo Clinic, Scottsdale, AZ, USA. ⁷Department of Surgery, Mayo Clinic, Scottsdale, AZ, USA. ⁸Department of Health Sciences Research, Mayo Clinic, Rochester, MN, USA. ⁹Department of Oncology, Mayo Clinic, Rochester, MN, USA.

Received: 21 July 2014 Accepted: 25 September 2014
Published: 7 October 2014

References

- Forouzanfar MH, Foreman KJ, Delossantos AM, Lozano R, Lopez AD, Murray CJ, Naghavi M: **Breast and cervical cancer in 187 countries between 1980 and 2010: a systematic analysis.** *Lancet* 2011, **378**(9801):1461–1484.
- Kuiper GG, Enmark E, Peltö-Huikko M, Nilsson S, Gustafsson JA: **Cloning of a novel receptor expressed in rat prostate and ovary.** *Proc Natl Acad Sci U S A* 1996, **93**(12):5925–5930.
- Mosselman S, Polman J, Dijkema R: **ER beta: identification and characterization of a novel human estrogen receptor.** *FEBS Lett* 1996, **392**(1):49–53.
- Osborne CK: **Steroid hormone receptors in breast cancer management.** *Breast Cancer Res Treat* 1998, **51**(3):227–238.
- Osborne CK, Schiff R: **Estrogen-receptor biology: continuing progress and therapeutic implications.** *J Clin Oncol* 2005, **23**(8):1616–1622.
- Davidson NE: **Biology of breast cancer and its clinical implications.** *Curr Opin Oncol* 1992, **4**(6):1003–1009.
- Pearce ST, Jordan VC: **The biological role of estrogen receptors alpha and beta in cancer.** *Crit Rev Oncol Hematol* 2004, **50**(1):3–22.
- Secreto FJ, Monroe DG, Dutta S, Ingle JN, Spelsberg TC: **Estrogen receptor alpha/beta isoforms, but not beta2, modulate unique patterns of gene expression and cell proliferation in Hs578T cells.** *J Cell Biochem* 2007, **101**(5):1125–1147.
- Monroe DG, Getz BJ, Johnsen SA, Riggs BL, Khosla S, Spelsberg TC: **Estrogen receptor isoform-specific regulation of endogenous gene expression in human osteoblastic cell lines expressing either ERalpha or ERbeta.** *J Cell Biochem* 2003, **90**(2):315–326.
- Kian Tee M, Rogatsky I, Tzagarakis-Foster C, Cvorovic A, An J, Christy RJ, Yamamoto KR, Leitman DC: **Estradiol and selective estrogen receptor modulators differentially regulate target genes with estrogen receptors alpha and beta.** *Mol Biol Cell* 2004, **15**(3):1262–1272.
- Stossi F, Barnett DH, Frasor J, Komm B, Lyttle CR, Katzenellenbogen BS: **Transcriptional profiling of estrogen-regulated gene expression via estrogen receptor (ER) alpha or ERbeta in human osteosarcoma cells: distinct and common target genes for these receptors.** *Endocrinology* 2004, **145**(7):3473–3486.
- Wu X, Subramaniam M, Grygo SB, Sun Z, Negron V, Lingle WL, Goetz MP, Ingle JN, Spelsberg TC, Hawse JR: **Estrogen receptor-beta sensitizes breast cancer cells to the anti-estrogenic actions of endoxifen.** *Breast Cancer Res* 2011, **13**(2):R27.
- Chang EC, Frasor J, Komm B, Katzenellenbogen BS: **Impact of estrogen receptor beta on gene networks regulated by estrogen receptor alpha in breast cancer cells.** *Endocrinology* 2006, **147**(10):4831–4842.
- Madak-Erdogan Z, Charn TH, Jiang Y, Liu ET, Katzenellenbogen JA, Katzenellenbogen BS: **Integrative genomics of gene and metabolic regulation by estrogen receptors alpha and beta, and their coregulators.** *Mol Syst Biol* 2013, **9**:676.
- Jarvinen TA, Peltö-Huikko M, Holli K, Isola J: **Estrogen receptor beta is coexpressed with ERalpha and PR and associated with nodal status, grade, and proliferation rate in breast cancer.** *Am J Pathol* 2000, **156**(1):29–35.
- Roger P, Sahla ME, Makela S, Gustafsson JA, Baldet P, Rochefort H: **Decreased expression of estrogen receptor beta protein in proliferative preinvasive mammary tumors.** *Cancer Res* 2001, **61**(6):2537–2541.
- Shaaban AM, O'Neill PA, Davies MP, Sibson R, West CR, Smith PH, Foster CS: **Declining estrogen receptor-beta expression defines malignant progression of human breast neoplasia.** *Am J Surg Pathol* 2003, **27**(12):1502–1512.
- Skloris GP, Carder PJ, Lansdown MR, Speirs V: **Immunohistochemical detection of ERbeta in breast cancer: towards more detailed receptor profiling?** *Br J Cancer* 2001, **84**(8):1095–1098.
- Skloris GP, Munot K, Bell SM, Carder PJ, Lane S, Horgan K, Lansdown MR, Parkes AT, Hanby AM, Markham AF, Speirs V: **Reduced expression of oestrogen receptor beta in invasive breast cancer and its re-expression using DNA methyl transferase inhibitors in a cell line model.** *J Pathol* 2003, **201**(2):213–220.
- Wu X, Subramaniam M, Negron V, Cicek M, Reynolds C, Lingle WL, Goetz MP, Ingle JN, Spelsberg TC, Hawse JR: **Development, characterization, and applications of a novel estrogen receptor beta monoclonal antibody.** *J Cell Biochem* 2012, **113**(2):711–723.
- Huang B, Omoto Y, Iwase H, Yamashita H, Toyama T, Coombes RC, Filipovic A, Warner M, Gustafsson JA: **Differential expression of estrogen receptor alpha, beta1, and beta2 in lobular and ductal breast cancer.** *Proc Natl Acad Sci U S A* 2014, **111**(5):1933–1938.
- Omoto Y, Inoue S, Ogawa S, Toyama T, Yamashita H, Muramatsu M, Kobayashi S, Iwase H: **Clinical value of the wild-type estrogen receptor beta expression in breast cancer.** *Cancer Lett* 2001, **163**(2):207–212.
- Fuqua SA, Schiff R, Parra I, Moore JT, Mohsin SK, Osborne CK, Clark GM, Allred DC: **Estrogen receptor beta protein in human breast cancer: correlation with clinical tumor parameters.** *Cancer Res* 2003, **63**(10):2434–2439.
- Mann S, Laucirica R, Carlson N, Younes PS, Ali N, Younes A, Li Y, Younes M: **Estrogen receptor beta expression in invasive breast cancer.** *Hum Pathol* 2001, **32**(1):113–118.
- Myers E, Fleming FJ, Crotty TB, Kelly G, McDermott EW, O'Higgins NJ, Hill AD, Young LS: **Inverse relationship between ER-beta and SRC-1 predicts outcome in endocrine-resistant breast cancer.** *Br J Cancer* 2004, **91**(9):1687–1693.
- Nakopoulou L, Lazaris AC, Panayotopoulou EG, Giannopoulou I, Givalos N, Markaki S, Keramopoulos A: **The favourable prognostic value of oestrogen receptor beta immunohistochemical expression in breast cancer.** *J Clin Pathol* 2004, **57**(5):523–528.
- Sugiyama H, Toyama T, Hara Y, Zhang Z, Kobayashi S, Fujii Y, Iwase H, Yamashita H: **Expression of estrogen receptor beta wild-type and its variant ERbeta2 is correlated with better prognosis in breast cancer.** *Jpn J Clin Oncol* 2007, **37**(11):820–828.
- Miller WR, Anderson TJ, Dixon JM, Saunders PT: **Oestrogen receptor beta and neoadjuvant therapy with tamoxifen: prediction of response and effects of treatment.** *Br J Cancer* 2006, **94**(9):1333–1338.
- Saunders PT, Millar MR, Williams K, Macpherson S, Bayne C, O'Sullivan C, Anderson TJ, Groome NP, Miller WR: **Expression of oestrogen receptor beta (ERbeta1) protein in human breast cancer biopsies.** *Br J Cancer* 2002, **86**(2):250–256.
- O'Neill PA, Davies MP, Shaaban AM, Innes H, Torevell A, Sibson DR, Foster CS: **Wild-type oestrogen receptor beta (ERbeta1) mRNA and protein**

- expression in Tamoxifen-treated post-menopausal breast cancers. *Br J Cancer* 2004, **91**(9):1694–1702.
31. Miyoshi Y, Taguchi T, Gustafsson JA, Noguchi S: Clinicopathological characteristics of estrogen receptor-beta-positive human breast cancers. *Jpn J Cancer Res* 2001, **92**(10):1057–1061.
 32. Skliris GP, Leygue E, Curtis-Snell L, Watson PH, Murphy LC: Expression of oestrogen receptor-beta in oestrogen receptor-alpha negative human breast tumours. *Br J Cancer* 2006, **95**(5):616–626.
 33. Jensen EV, Cheng G, Palmieri C, Saji S, Makela S, Van Noorden S, Wahlstrom T, Warner M, Coombes RC, Gustafsson JA: Estrogen receptors and proliferation markers in primary and recurrent breast cancer. *Proc Natl Acad Sci U S A* 2001, **98**(26):15197–15202.
 34. Poola I, Fuqua SA, De Witty RL, Abraham J, Marshallack JJ, Liu A: Estrogen receptor alpha-negative breast cancer tissues express significant levels of estrogen-independent transcription factors, ERbeta1 and ERbeta5: potential molecular targets for chemoprevention. *Clin Cancer Res* 2005, **11**(20):7579–7585.
 35. Umekita Y, Souda M, Ohi Y, Sagara Y, Rai Y, Takahama T, Yoshida H: Expression of wild-type estrogen receptor beta protein in human breast cancer: specific correlation with HER2/neu overexpression. *Pathol Int* 2006, **56**(8):423–427.
 36. Honma N, Horii R, Iwase T, Saji S, Younes M, Takubo K, Matsuura M, Ito Y, Akiyama F, Sakamoto G: Clinical importance of estrogen receptor-beta evaluation in breast cancer patients treated with adjuvant tamoxifen therapy. *J Clin Oncol* 2008, **26**(22):3727–3734.
 37. Esslimani-Sahla M, Simony-Lafontaine J, Kramar A, Lavaill R, Mollevi C, Warner M, Gustafsson JA, Rochefort H: Estrogen receptor beta (ER beta) level but not its ER beta cx variant helps to predict tamoxifen resistance in breast cancer. *Clin Cancer Res* 2004, **10**(17):5769–5776.
 38. Fleming FJ, Hill AD, McDermott EW, O'Higgins NJ, Young LS: Differential recruitment of coregulator proteins steroid receptor coactivator-1 and silencing mediator for retinoid and thyroid receptors to the estrogen receptor-estrogen response element by beta-estradiol and 4-hydroxytamoxifen in human breast cancer. *J Clin Endocrinol Metab* 2004, **89**(1):375–383.
 39. Hopp TA, Weiss HL, Parra IS, Cui Y, Osborne CK, Fuqua SA: Low levels of estrogen receptor beta protein predict resistance to tamoxifen therapy in breast cancer. *Clin Cancer Res* 2004, **10**(22):7490–7499.
 40. Iwase H, Zhang Z, Omoto Y, Sugiura H, Yamashita H, Toyama T, Iwata H, Kobayashi S: Clinical significance of the expression of estrogen receptors alpha and beta for endocrine therapy of breast cancer. *Cancer Chemother Pharmacol* 2003, **52**(Suppl 1):S34–38.
 41. Murphy LC, Leygue E, Niu Y, Snell L, Ho SM, Watson PH: Relationship of coregulator and oestrogen receptor isoform expression to de novo tamoxifen resistance in human breast cancer. *Br J Cancer* 2002, **87**(12):1411–1416.
 42. Speirs V, Parkes AT, Kerin MJ, Walton DS, Carleton PJ, Fox JN, Atkin SL: Coexpression of estrogen receptor alpha and beta: poor prognostic factors in human breast cancer? *Cancer Res* 1999, **59**(3):525–528.
 43. Speirs V, Malone C, Walton DS, Kerin MJ, Atkin SL: Increased expression of estrogen receptor beta mRNA in tamoxifen-resistant breast cancer patients. *Cancer Res* 1999, **59**(21):5421–5424.
 44. Guo L, Meng J, Yilamu D, Jakulin A, Fu M, Wang B, Abulajiang G: Significance of ERbeta expression in different molecular subtypes of breast cancer. *Diagn Pathol* 2014, **9**:20.
 45. Shaaban AM, Green AR, Karthik S, Alizadeh Y, Hughes TA, Harkins L, Ellis IO, Robertson JF, Paish EC, Saunders PT, Groome NP, Speirs V: Nuclear and cytoplasmic expression of ERbeta1, ERbeta2, and ERbeta5 identifies distinct prognostic outcome for breast cancer patients. *Clin Cancer Res* 2008, **14**(16):5228–5235.
 46. Novelli F, Milella M, Melucci E, Di Benedetto A, Sperduti I, Perrone-Donnorso R, Perracchio L, Ventura I, Nistico C, Fabi A, Buglioni S, Natali PG, Mottolose M: A divergent role for estrogen receptor-beta in node-positive and node-negative breast cancer classified according to molecular subtypes: an observational prospective study. *Breast Cancer Res* 2008, **10**(5):R74.
 47. Madeira M, Mattar A, Logullo AF, Soares FA, Gebrim LH: Estrogen receptor alpha/beta ratio and estrogen receptor beta as predictors of endocrine therapy responsiveness—a randomized neoadjuvant trial comparison between anastrozole and tamoxifen for the treatment of postmenopausal breast cancer. *BMC Cancer* 2013, **13**:425.
 48. Yan Y, Li X, Blanchard A, Bramwell VH, Pritchard KI, Tu D, Shepherd L, Myal Y, Penner C, Watson PH, Leygue E, Murphy LC: Expression of both estrogen receptor-beta 1 (ER-beta1) and its co-regulator steroid receptor RNA activator protein (SRAP) are predictive for benefit from tamoxifen therapy in patients with estrogen receptor-alpha (ER-alpha)-negative early breast cancer (EBC). *Ann Oncol* 2013, **24**(8):1986–1993.
 49. Motomura K, Ishitobi M, Komoike Y, Koyama H, Nagase H, Inaji H, Noguchi S: Expression of estrogen receptor beta and phosphorylation of estrogen receptor alpha serine 167 correlate with progression-free survival in patients with metastatic breast cancer treated with aromatase inhibitors. *Oncology* 2010, **79**(1–2):55–61.
 50. Weitsman GE, Skliris G, Ung K, Peng B, Younes M, Watson PH, Murphy LC: Assessment of multiple different estrogen receptor-beta antibodies for their ability to immunoprecipitate under chromatin immunoprecipitation conditions. *Breast Cancer Res Treat* 2006, **100**(1):23–31.
 51. Wimberly H, Han G, Pinnaduwaage D, Murphy LC, Yang XR, Andrulis IL, Sherman M, Figueroa J, Rimm DL: ERbeta splice variant expression in four large cohorts of human breast cancer patient tumors. *Breast Cancer Res Treat* 2014, **146**(3):657–667.
 52. Ingle JN, Suman VJ, Mailliard JA, Kugler JW, Krook JE, Michalak JC, Pisansky TM, Wold LE, Donohue JH, Goetz MP, Perez EA: Randomized trial of tamoxifen alone or combined with fluoxymesterone as adjuvant therapy in postmenopausal women with resected estrogen receptor positive breast cancer. North Central Cancer Treatment Group Trial 89-30-52. *Breast Cancer Res Treat* 2006, **98**(2):217–222.
 53. Ali HR, Dawson SJ, Blows FM, Provenzano E, Leung S, Nielsen T, Pharoah PD, Caldas C: A Ki67/BCL2 index based on immunohistochemistry is highly prognostic in ER-positive breast cancer. *J Pathol* 2012, **226**(1):97–107.
 54. Hawse JR, Subramaniam M, Cicek M, Wu X, Gingery A, Grygo SB, Sun Z, Pitel KS, Lingle WL, Goetz MP, Ingle JN, Spelsberg TC: Endoxifen's molecular mechanisms of action are concentration dependent and different than that of other anti-estrogens. *PLoS One* 2013, **8**(1):e54613.
 55. Pedram A, Razandi M, Wallace DC, Levin ER: Functional estrogen receptors in the mitochondria of breast cancer cells. *Mol Biol Cell* 2006, **17**(5):2125–2137.
 56. Milanese L, Vasconsuelo A, de Boland AR, Boland R: Expression and subcellular distribution of native estrogen receptor beta in murine C2C12 cells and skeletal muscle tissue. *Steroids* 2009, **74**(6):489–497.
 57. Chen JQ, Delannoy M, Cooke C, Yager JD: Mitochondrial localization of ERalpha and ERbeta in human MCF7 cells. *Am J Physiol Endocrinol Metab* 2004, **286**(6):E1011–1022.
 58. Yang SH, Liu R, Perez EJ, Wen Y, Stevens SM Jr, Valencia T, Brun-Zinkernagel AM, Prokai L, Will Y, Dykens J, Koulen P, Simpkins JW: Mitochondrial localization of estrogen receptor beta. *Proc Natl Acad Sci U S A* 2004, **101**(12):4130–4135.
 59. Razandi M, Pedram A, Jordan VC, Fuqua S, Levin ER: Tamoxifen regulates cell fate through mitochondrial estrogen receptor beta in breast cancer. *Oncogene* 2013, **32**(27):3274–3285.
 60. Marotti JD, Collins LC, Hu R, Tamimi RM: Estrogen receptor-beta expression in invasive breast cancer in relation to molecular phenotype: results from the Nurses' Health Study. *Mod Pathol* 2010, **23**(2):197–204.
 61. Litwiniuk MM, Roznowski K, Filas V, Godlewski DD, Stawicka M, Kaleta R, Breborowicz J: Expression of estrogen receptor beta in the breast carcinoma of BRCA1 mutation carriers. *BMC Cancer* 2008, **8**:100.
 62. Tonetti DA, Gao W, Escarzaga D, Walters K, Szafran A, Coon JS: PKCalpha and ERbeta Are Associated with Triple-Negative Breast Cancers in African American and Caucasian Patients. *Int J Breast Cancer* 2012, **2012**:740353.
 63. Poola I, Abraham J, Baldwin K, Saunders A, Bhatnagar R: Estrogen receptors beta4 and beta5 are full length functionally distinct ERbeta isoforms: cloning from human ovary and functional characterization. *Endocrine* 2005, **27**(3):227–238.
 64. Gruvberger-Saal SK, Bendahl PO, Saal LH, Laakso M, Hegardt C, Eden P, Peterson C, Malmstrom P, Isola J, Borg A, Ferno M: Estrogen receptor beta expression is associated with tamoxifen response in ERalpha-negative breast carcinoma. *Clin Cancer Res* 2007, **13**(7):1987–1994.
 65. Yan M, Rayoo M, Takano EA, Fox SB: Nuclear and cytoplasmic expressions of ERbeta1 and ERbeta2 are predictive of response to therapy and alters prognosis in familial breast cancers. *Breast Cancer Res Treat* 2011, **126**(2):395–405.
 66. Sotoca AM, van den Berg H, Vervoort J, van der Saag P, Strom A, Gustafsson JA, Rietjens I, Murk AJ: Influence of cellular ERalpha/ERbeta ratio on the ERalpha-agonist induced proliferation of human T47D breast cancer cells. *Toxicol Sci* 2008, **105**(2):303–311.

67. Paruthiyil S, Parmar H, Kerekatte V, Cunha GR, Firestone GL, Leitman DC: **Estrogen receptor beta inhibits human breast cancer cell proliferation and tumor formation by causing a G2 cell cycle arrest.** *Cancer Res* 2004, **64**(1):423–428.
68. Strom A, Hartman J, Foster JS, Kietz S, Wimalasena J, Gustafsson JA: **Estrogen receptor beta inhibits 17beta-estradiol-stimulated proliferation of the breast cancer cell line T47D.** *Proc Natl Acad Sci U S A* 2004, **101**(6):1566–1571.
69. Hodges-Gallagher L, Valentine CD, El Bader S, Kushner PJ: **Estrogen receptor beta increases the efficacy of antiestrogens by effects on apoptosis and cell cycling in breast cancer cells.** *Breast Cancer Res Treat* 2008, **109**(2):241–250.
70. Tu Z, Li H, Ma Y, Tang B, Tian J, Akers W, Achilefu S, Gu Y: **The enhanced antiproliferative response to combined treatment of trichostatin A with raloxifene in MCF-7 breast cancer cells and its relevance to estrogen receptor beta expression.** *Mol Cell Biochem* 2012, **366**(1–2):111–122.
71. Harris HA, Albert LM, Leathurby Y, Malamas MS, Mewshaw RE, Miller CP, Kharode YP, Marzolf J, Komm BS, Winneker RC, Frail DE, Henderson RA, Zhu Y, Keith JC Jr: **Evaluation of an estrogen receptor-beta agonist in animal models of human disease.** *Endocrinology* 2003, **144**(10):4241–4249.
72. Malamas MS, Manas ES, McDevitt RE, Gunawan I, Xu ZB, Collini MD, Miller CP, Dinh T, Henderson RA, Keith JC Jr, Harris HA: **Design and synthesis of aryl diphenolic azoles as potent and selective estrogen receptor-beta ligands.** *J Med Chem* 2004, **47**(21):5021–5040.
73. Lattrich C, Schuler S, Haring J, Skrzypczak M, Ortmann O, Treeck O: **Effects of a combined treatment with tamoxifen and estrogen receptor beta agonists on human breast cancer cell lines.** *Arch Gynecol Obstet* 2014, **289**(1):163–171.
74. Cotrim CZ, Fabris V, Doria ML, Lindberg K, Gustafsson JA, Amado F, Lanari C, Helguero LA: **Estrogen receptor beta growth-inhibitory effects are repressed through activation of MAPK and PI3K signalling in mammary epithelial and breast cancer cells.** *Oncogene* 2013, **32**(19):2390–2402.
75. Lazennec G, Bresson D, Lucas A, Chauveau C, Vignon F: **ER beta inhibits proliferation and invasion of breast cancer cells.** *Endocrinology* 2001, **142**(9):4120–4130.
76. Shanle EK, Zhao Z, Hawse J, Wisinski K, Keles S, Yuan M, Xu W: **Research Resource: Global identification of estrogen receptor beta target genes in triple negative breast cancer cells.** *Mol Endocrinol* 2013, **27**(10):1762–1775.

doi:10.1186/1471-2407-14-749

Cite this article as: Reese et al.: ER β 1: characterization, prognosis, and evaluation of treatment strategies in ER α -positive and -negative breast cancer. *BMC Cancer* 2014 **14**:749.

**Submit your next manuscript to BioMed Central
and take full advantage of:**

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

