

RESEARCH

Open Access

Korovkin type approximation theorem for functions of two variables through statistical A -summability

Mohammad Mursaleen^{1*} and Abdullah Alotaibi²

* Correspondence:

mursaleenm@gmail.com

¹Department of Mathematics,
Aligarh Muslim University, Aligarh
202002, India

Full list of author information is
available at the end of the article

Abstract

In this article, we prove a Korovkin type approximation theorem for a function of two variables by using the notion of statistical A -summability. We also study the rate of statistical A -summability of positive linear operators. Finally we construct an example by Bleimann et al. operators to show that our result is stronger than those of previously proved by other authors.

AMS Subject Classification 2000: 41A10; 41A25; 41A36; 40A30; 40G15.

Keywords: density, statistical convergence, A -statistical convergence, statistical A -summability, positive linear operator, Korovkin type approximation theorem

1 Introduction and preliminaries

The concept of statistical convergence for sequences of real numbers was introduced by Fast [1] and further studied many others.

Let $K \subseteq \mathbb{N}$ and $K_n = \{k \leq n : k \in K\}$. Then the *natural density* of K is defined by $\delta(K) = \lim_n n^{-1} |K_n|$ if the limit exists, where $|K_n|$ denotes the cardinality of K_n .

A sequence $x = (x_k)$ of real numbers is said to be *statistically convergent* to L provided that for every $\varepsilon > 0$ the set $K_\varepsilon := \{k \in \mathbb{N} : |x_k - L| \geq \varepsilon\}$ has natural density zero, i.e. for each $\varepsilon > 0$,

$$\lim_n \frac{1}{n} |\{k \leq n : |x_k - L| \geq \varepsilon\}| = 0.$$

In this case we write $st\text{-}\lim x = L$. Note that if $x = (x_k)$ is convergent then it is statistically convergent but not conversely. The idea of statistical convergence of double sequences has been introduced and studied in [2,3].

Let $A = (a_{nk})$, $n, k \in \mathbb{N}$, be an infinite matrix and $x = (x_k)$ be a sequence. Then the (transformed) sequence, $Ax := (y_n)$, is denoted by

$$y_n := \sum_{k=1}^{\infty} a_{nk} x_k,$$

where it is assumed that the series on the right converges for each $n \in \mathbb{N}$. We say that a sequence x is A -summable to the limit ℓ if $y_n \rightarrow \ell$ as $n \rightarrow \infty$.

A matrix transformation is said to be regular if it maps every convergent sequence into a convergent sequence with the same limit. The well-known conditions for two dimensional matrix to be regular are known as Silverman-Toeplitz conditions.

In [4], Edely and Mursaleen have given the notion of statistical A -summability for single sequences and statistical A -summability for double sequences has recently been studied in [5].

Let $A = (a_{nk})$ be a nonnegative regular summability matrix and $x = (x_k)$ be a sequence of real or complex sequences. We say that x is statistically A -summable to L if for every $\varepsilon > 0$,

$$\delta(\{n \in \mathbb{N} : |y_n - L| \geq \varepsilon\}) = 0.$$

So, if x is statistically A -summable to L then for every $\varepsilon > 0$,

$$\lim_m \frac{1}{m} |\{n \leq m : |y_n - L| \geq \varepsilon\}| = 0.$$

Note that if a sequence is bounded and A -statistically convergent to L , then it is A -summable to L ; hence it is statistically A -summable to L but not conversely (see [4]).

Example 1.1. Let $A = (C, 1)$, the Cesàro matrix and the sequence $u = (u_k)$ be defined by

$$u_k = \begin{cases} 1 & \text{if } k \text{ is odd,} \\ 0 & \text{if } k \text{ is even.} \end{cases} \tag{1.1}$$

Then u is A -summable to $\frac{1}{2}$ (and hence statistically A -summable to $1/2$) but not statistically (and not A -statistically as well) convergent.

Let $I := [0, \infty)$ and $C(I)$ denote the space of all continuous real valued functions on I . Let $C_B(I) := \{f \in C(I) : f \text{ is bounded on } I\}$. $C(I)$ and $C_B(I)$ are equipped with norm

$$\|f\|_{C(I)} := \sup_{x \in I} |f(x)|.$$

Let $H_\omega(I)$ denote the space of all real valued functions f on I such that

$$|f(s) - f(x)| \leq \omega\left(f; \left| \frac{s}{1+s} - \frac{x}{1+x} \right| \right),$$

where ω is the modulus of continuity, i.e.

$$\omega(f; \delta) = \sup_{s, x \in I} \{|f(s) - f(x)| : |s - x| \leq \delta\}.$$

It is to be noted that any function $f \in H_\omega(I)$ is continuous and bounded on I .

The following Korovkin type theorem (see [6]) was proved by Çakar and Gadjiev [7].

Theorem A. Let (L_n) be a sequence of positive linear operators from $H_\omega(I)$ into $C_B(I)$. Then for all $f \in H_\omega(I)$

$$\lim_{n \rightarrow \infty} \|L_n(f; x) - f(x)\|_{C_B(I)} = 0$$

if and only if

$$\lim_{n \rightarrow \infty} \|L_n(f_i; x) - g_i\|_{C_B(I)} = 0 \quad (i = 0, 1, 2),$$

where

$$g_0(x) = 1, g_1(x) = \frac{x}{1+x}, g_2(x) = \left(\frac{x}{1+x}\right)^2.$$

Erkuş and Duman [8] have given the st_A -version of the above theorem for functions of two variables. Quite recently, Korovkin type of approximation theorems have been proved in [9,10] by using almost convergence; in [11-15] by using variants of statistical convergence and in [16-19] for functions of two variables by using statistical convergence, A -statistical convergence and statistical A -summability of double sequences. In this article, we use the notion of statistical A -summability to prove a Korovkin type approximation theorem for functions of two variables with the help of test functions $1, \frac{x}{1+x}, \frac{y}{1+y}, \left(\frac{x}{1+x}\right)^2 + \left(\frac{y}{1+y}\right)^2$.

2 Main result

Let $I = [0, \infty)$ and $K = I \times I$. We denote by $C_B(K)$ the space of all bounded and continuous real valued functions on K equipped with norm

$$\|f\|_{C_B(K)} := \sup_{(x,y) \in K} |f(x,y)|, \quad f \in C_B(K).$$

Let $H_{\omega^*}(K)$ denote the space of all real valued functions f on K such that

$$|f(s,t) - f(x,y)| \leq \omega^* \left(f; \sqrt{\left(\frac{s}{1+s} - \frac{x}{1+x}\right)^2 + \left(\frac{t}{1+t} - \frac{y}{1+y}\right)^2} \right),$$

where ω^* is the modulus of continuity, i.e.

$$\omega^*(f; \delta) = \sup_{(s,t),(x,y) \in K} \left\{ |f(s,t) - f(x,y)| : \sqrt{(s-x)^2 + (t-y)^2} \leq \delta \right\}.$$

It is to be noted that any function $f \in H_{\omega^*}(K)$ is bounded and continuous on K , and a necessary and sufficient condition for $f \in H_{\omega^*}(K)$ is that

$$\lim_{\delta \rightarrow 0} \omega^*(f; \delta) = 0.$$

We prove the following result:

Theorem 2.1. Let $A = (a_{nk})$ be nonnegative regular summability matrix. Let (T_k) be a sequence of positive linear operators from $H_{\omega^*}(K)$ into $C_B(K)$. Then for all $f \in H_{\omega^*}(K)$

$$st - \lim_{n \rightarrow \infty} \left\| \sum_{k=1}^{\infty} a_{nk} T_k(f; x, y) - f(x, y) \right\|_{C_B(K)} = 0 \tag{2.0}$$

if and only if

$$st - \lim_{n \rightarrow \infty} \left\| \sum_{k=1}^{\infty} a_{nk} T_k(1; x, y) - 1 \right\|_{C_B(K)} = 0 \tag{2.1}$$

$$st - \lim_{n \rightarrow \infty} \left\| \sum_{k=1}^{\infty} a_{nk} T_k\left(\frac{s}{1+s}; x, y\right) - \frac{x}{1+x} \right\|_{C_B(K)} = 0 \tag{2.2}$$

$$st - \lim_{n \rightarrow \infty} \left\| \sum_{k=1}^{\infty} a_{nk} T_k\left(\frac{t}{1+t}; x, y\right) - \frac{y}{1+y} \right\|_{C_B(K)} = 0 \tag{2.3}$$

$$st - \lim_{n \rightarrow \infty} \left\| \sum_{k=1}^{\infty} a_{nk} T_k \left(\left(\frac{s}{1+s} \right)^2 + \left(\frac{t}{1+t} \right)^2; x, \gamma \right) - \left(\left(\frac{x}{1+x} \right)^2 + \left(\frac{\gamma}{1+\gamma} \right)^2 \right) \right\|_{C_B(K)} = 0. \quad (2.4)$$

Proof. Since each of the functions $f_0(x, \gamma) = 1, f_1(x, \gamma) = \frac{x}{1+x}, f_2(x, \gamma) = \frac{\gamma}{1+\gamma}, f_3(x, \gamma) = \left(\frac{x}{1+x} \right)^2 + \left(\frac{\gamma}{1+\gamma} \right)^2$ belongs to $H_{\omega^*}(K)$, conditions (2.1)-(2.4) follow immediately from (2.0).

Let $f \in H_{\omega^*}(K)$ and $(x, \gamma) \in K$ be fixed. Then for $\varepsilon > 0$ there exist $\delta_1, \delta_2 > 0$ such that $|f(s, t) - f(x, \gamma)| < \varepsilon$ holds for all $(s, t) \in K$ satisfying $|\frac{s}{1+s} - \frac{x}{1+x}| < \delta_1$ and $|\frac{t}{1+t} - \frac{\gamma}{1+\gamma}| < \delta_2$.

Let

$$K(\delta) := \left\{ (s, t) \in K : \sqrt{\left(\frac{s}{1+s} - \frac{x}{1+x} \right)^2 + \left(\frac{t}{1+t} - \frac{\gamma}{1+\gamma} \right)^2} < \delta = \min\{\delta_1, \delta_2\} \right\}.$$

Hence

$$\begin{aligned} |f(s, t) - f(x, \gamma)| &= |f(s, t) - f(x, \gamma)| \chi_{K(\delta)}(s, t) + |f(s, t) - f(x, \gamma)| \chi_{K \setminus K(\delta)}(s, t) \\ &\leq \varepsilon + 2N \chi_{K \setminus K(\delta)}(s, t) \end{aligned} \quad (2.5)$$

Where χ_D denotes the characteristic function of the set D and $N = \|f\|_{C_B(K)}$. Further we get

$$\chi_{K \setminus K(\delta)}(s, t) \leq \frac{1}{\delta_1^2} \left(\frac{s}{1+s} - \frac{x}{1+x} \right)^2 + \frac{1}{\delta_2^2} \left(\frac{t}{1+t} - \frac{\gamma}{1+\gamma} \right)^2. \quad (2.6)$$

Combining (2.5) and (2.6), we get

$$|f(s, t) - f(x, \gamma)| \leq \varepsilon + \frac{2N}{\delta^2} \left\{ \left(\frac{s}{1+s} - \frac{x}{1+x} \right)^2 + \left(\frac{t}{1+t} - \frac{\gamma}{1+\gamma} \right)^2 \right\}, \quad (2.7)$$

After using the properties of f , a simple calculation gives that

$$\begin{aligned} |T_k(f; x, \gamma) - f(x, \gamma)| &\leq \varepsilon + M \{ |T_k(f_0; x, \gamma) - f_0(x, \gamma)| + |T_k(f_1; x, \gamma) - f_1(x, \gamma)| \\ &\quad + |T_k(f_2; x, \gamma) - f_2(x, \gamma)| + |T_k(f_3; x, \gamma) - f_3(x, \gamma)| \}, \end{aligned} \quad (2.8)$$

where

$$M := \varepsilon + N + \frac{4N}{\delta^2}.$$

Now replacing $T_k(f; x, \gamma)$ by $\sum_{k=1}^{\infty} a_{nk} T_k(f; x, \gamma)$ and taking $\sup_{(x, \gamma) \in K}$, we get

$$\begin{aligned} \left\| \sum_{k=1}^{\infty} a_{nk} T_k(f; x, \gamma) - f(x, \gamma) \right\|_{C_B(K)} &\leq \varepsilon + M \left(\left\| \sum_{k=1}^{\infty} a_{nk} T_k(f_0; x, \gamma) - f_0(x, \gamma) \right\|_{C_B(K)} \right. \\ &\quad + \left\| \sum_{k=1}^{\infty} a_{nk} T_k(f_1; x, \gamma) - f_1(x, \gamma) \right\|_{C_B(K)} + \left\| \sum_{k=1}^{\infty} a_{nk} T_k(f_2; x, \gamma) - f_2(x, \gamma) \right\|_{C_B(K)} \\ &\quad \left. + \left\| \sum_{k=1}^{\infty} a_{nk} T_k(f_3; x, \gamma) - f_3(x, \gamma) \right\|_{C_B(K)} \right). \end{aligned} \quad (2.9)$$

For a given $r > 0$ choose $\varepsilon > 0$ such that $\varepsilon < r$. Define the following sets

$$\begin{aligned}
 D &:= \left\{ n : \left\| \sum_{k=1}^{\infty} a_{nk} T_k(f; x, \gamma) - f(x, \gamma) \right\|_{C_B(K)} \geq r \right\}, \\
 D_1 &:= \left\{ n : \left\| \sum_{k=1}^{\infty} a_{nk} T_k(f_0; x, \gamma) - f_0(x, \gamma) \right\|_{C_B(K)} \geq \frac{r - \varepsilon}{4K} \right\}, \\
 D_2 &:= \left\{ n : \left\| \sum_{k=1}^{\infty} a_{nk} T_k(f_1; x, \gamma) - f_1(x, \gamma) \right\|_{C_B(K)} \geq \frac{r - \varepsilon}{4K} \right\}, \\
 D_3 &:= \left\{ n : \left\| \sum_{k=1}^{\infty} a_{nk} T_k(f_2; x, \gamma) - f_2(x, \gamma) \right\|_{C_B(K)} \geq \frac{r - \varepsilon}{4K} \right\}, \\
 D_4 &:= \left\{ n : \left\| \sum_{k=1}^{\infty} a_{nk} T_k(f_3; x, \gamma) - f_3(x, \gamma) \right\|_{C_B(K)} \geq \frac{r - \varepsilon}{4K} \right\}.
 \end{aligned}$$

Then from (2.9), we see that $D \subset D_1 \cup D_2 \cup D_3 \cup D_4$ and therefore $\delta(D) \leq \delta(D_1) + \delta(D_2) + \delta(D_3) + \delta(D_4)$. Hence conditions (2.1)-(2.4) imply the condition (2.0).

This completes the proof of the theorem.

If we replace the matrix A in Theorem 2.1 by identity matrix, then we immediately get the following result which is due to Erkuş and Duman [8]:

Corollary 2.2. Let $A = (a_{nk})$ be nonnegative regular summability matrix. Let (T_k) be a sequence of positive linear operators from $H_{\omega^*}(K)$ into $C_B(K)$. Then for all $f \in H_{\omega^*}(K)$

$$st - \lim_{k \rightarrow \infty} \|T_k(f; x, \gamma) - f(x, \gamma)\|_{C_B(K)} = 0 \tag{2.10}$$

if and only if

$$st - \lim_{k \rightarrow \infty} \|T_k(1; x, \gamma) - 1\|_{C_B(K)} = 0, \tag{2.11}$$

$$st - \lim_{k \rightarrow \infty} \left\| T_k\left(\frac{s}{1+s}; x, \gamma\right) - \frac{x}{1+x} \right\|_{C_B(K)} = 0, \tag{2.12}$$

$$st - \lim_{k \rightarrow \infty} \left\| T_k\left(\frac{t}{1+t}; x, \gamma\right) - \frac{\gamma}{1+\gamma} \right\|_{C_B(K)} = 0, \tag{2.13}$$

$$st - \lim_{k \rightarrow \infty} \left\| T_k\left(\left(\frac{s}{1+s}\right)^2 + \left(\frac{t}{1+t}\right)^2; x, \gamma\right) - \left(\left(\frac{x}{1+x}\right)^2 + \left(\frac{\gamma}{1+\gamma}\right)^2\right) \right\|_{C_B(K)} = 0. \tag{2.14}$$

3 Statistical rate of convergence

In this section, using the concept of statistical A -summability we study the rate of convergence of positive linear operators with the help of the modulus of continuity. Let us recall, for $f \in H_{\omega^*}(K)$

$$|f(s, t) - f(x, \gamma)| \leq \omega^* \left(f; \sqrt{\left(\frac{s}{1+s} - \frac{x}{1+x}\right)^2 + \left(\frac{t}{1+t} - \frac{\gamma}{1+\gamma}\right)^2} \right),$$

where

$$\omega^*(f; \delta) = \sup_{(s,t),(x,y) \in K} \left\{ |f(s, t) - f(x, y)| : \sqrt{(s-x)^2 + (t-y)^2} \leq \delta \right\}.$$

We have the following result:

Theorem 3.1. Let $A = (a_{nk})$ be nonnegative regular summability matrix. Let (T_k) be a sequence of positive linear operators from $H_{\omega^*}(K)$ into $C_B(K)$. Assume that

$$(i) \quad st\text{-}\lim_{n \rightarrow \infty} \left\| \sum_{k=1}^{\infty} a_{nk} T_k(f_0) - f_0 \right\|_{C_B(K)} = 0,$$

(ii) $st\text{-}\lim_{n \rightarrow \infty} \omega^*(f; \delta_n) = 0$, where

$$\delta_n = \sqrt{\left\| \sum_{k=1}^{\infty} a_{nk} T_k(\psi) \right\|_{C_B(K)}} \quad \text{with } \psi = \psi(s, t) = \left(\frac{s}{1+s} - \frac{x}{1+x} \right)^2 + \left(\frac{t}{1+t} - \frac{y}{1+y} \right)^2.$$

Then for all $f \in H_{\omega^*}(K)$

$$st\text{-}\lim_{n \rightarrow \infty} \left\| \sum_{k=1}^{\infty} a_{nk} T_k(f) - f \right\|_{C_B(K)} = 0.$$

Proof. Let $f \in H_{\omega^*}(K)$ be fixed and $(x, y) \in K$ be fixed. Using linearity and positivity of the operators T_k for all $n \in \mathbb{N}$, we have

$$\begin{aligned} & \left| \sum_{k=1}^{\infty} a_{nk} T_k(f; x, y) - f(x, y) \right| \leq \sum_{k=1}^{\infty} a_{nk} T_k(|f(s, t) - f(x, y)|; x, y) \\ & \quad + |f(x, y)| \left| \sum_{k=1}^{\infty} a_{nk} T_k(f_0; x, y) - f_0(x, y) \right| \\ & \leq \sum_{k=1}^{\infty} a_{nk} T_k \left(\omega^* \left(f; \delta \frac{\sqrt{\left(\frac{s}{1+s} - \frac{x}{1+x} \right)^2 + \left(\frac{t}{1+t} - \frac{y}{1+y} \right)^2}}{\delta} \right); x, y \right) \\ & \quad + \|f\|_{C_B(K)} \left| \sum_{k=1}^{\infty} a_{nk} T_k(f_0; x, y) - f_0(x, y) \right| \\ & \leq \sum_{k=1}^{\infty} a_{nk} T_k \left(\left(1 + \left[\frac{\sqrt{\left(\frac{s}{1+s} - \frac{x}{1+x} \right)^2 + \left(\frac{t}{1+t} - \frac{y}{1+y} \right)^2}}{\delta} \right]^2 \right) \omega^*(f; \delta); x, y \right) \\ & \quad + \|f\|_{C_B(K)} \left| \sum_{k=1}^{\infty} a_{nk} T_k(f_0; x, y) - f_0(x, y) \right| \\ & \leq \sum_{k=1}^{\infty} a_{nk} \omega^*(f; \delta) T_k \left(\left(1 + \frac{\left(\frac{s}{1+s} - \frac{x}{1+x} \right)^2 + \left(\frac{t}{1+t} - \frac{y}{1+y} \right)^2}{\delta^2} \right); x, y \right) \\ & \quad + \|f\|_{C_B(K)} \left| \sum_{k=1}^{\infty} a_{nk} T_k(f_0; x, y) - f_0(x, y) \right| \\ & \leq \omega^*(f; \delta) \left| \sum_{k=1}^{\infty} a_{nk} T_k(f_0; x, y) - f_0(x, y) \right| + \|f\|_{C_B(K)} \left| \sum_{k=1}^{\infty} a_{nk} T_k(f_0; x, y) - f_0(x, y) \right| \\ & \quad + \omega^*(f; \delta) + \frac{\omega^*(f; \delta)}{\delta^2} \sum_{k=1}^{\infty} a_{nk} T_k \left(\left(\left(\frac{s}{1+s} - \frac{x}{1+x} \right)^2 + \left(\frac{t}{1+t} - \frac{y}{1+y} \right)^2 \right); x, y \right). \end{aligned}$$

Hence

$$\begin{aligned} & \left\| \sum_{k=1}^{\infty} a_{nk} T_k(f) - f \right\|_{C_B(K)}. \\ \leq & \|f\|_{C_B(K)} \left\| \sum_{k=1}^{\infty} a_{nk} T_k(f_0) - f_0 \right\|_{C_B(K)} + \omega^*(f; \delta) \left\| \sum_{k=1}^{\infty} a_{nk} T_k(f_0) - f_0 \right\|_{C_B(K)} \\ & + \frac{\omega^*(f; \delta)}{\delta^2} \left\| \sum_{k=1}^{\infty} a_{nk} T_k(\psi) \right\|_{C_B(K)} + \omega^*(f; \delta). \end{aligned}$$

Now if we choose $\delta := \delta_n := \sqrt{\left\| \sum_{k=1}^{\infty} a_{nk} T_k(\psi) \right\|_{C_B(K)}}$ then

$$\begin{aligned} & \left\| \sum_{k=1}^{\infty} a_{nk} T_k(f) - f \right\|_{C_B(K)}. \\ \leq & \|f\|_{C_B(K)} \left\| \sum_{k=1}^{\infty} a_{nk} T_k(f_0) - f_0 \right\|_{C_B(K)} \\ & + \omega^*(f; \delta_n) \left\| \sum_{k=1}^{\infty} a_{nk} T_k(f_0) - f_0 \right\|_{C_B(K)} + 2\omega^*(f; \delta_n). \end{aligned}$$

Therefore

$$\begin{aligned} & \left\| \sum_{k=1}^{\infty} a_{nk} T_k(f) - f \right\|_{C_B(K)}. \\ \leq & M \left\{ \left\| \sum_{k=1}^{\infty} a_{nk} T_k(f_0) - f_0 \right\|_{C_B(K)} \right. \\ & \left. + \omega^*(f; \delta_n) \left\| \sum_{k=1}^{\infty} a_{nk} T_k(f_0) - f_0 \right\|_{C_B(K)} + \omega^*(f; \delta_n) \right\}, \end{aligned} \tag{3.1}$$

where $M = \max\{2, \|f\|_{C_B(K)}\}$. Now, for a given $r > 0$, choose $\varepsilon > 0$ such that $\varepsilon > r$. Let us write

$$\begin{aligned} E & := \left\{ n : \left\| \sum_{k=1}^{\infty} a_{nk} T_k(f; x, \gamma) - f(x, \gamma) \right\|_{C_B(K)} \geq r \right\}, \\ E_1 & := \left\{ n : \left\| \sum_{k=1}^{\infty} a_{nk} T_k(f_0; x, \gamma) - f_0(x, \gamma) \right\|_{C_B(K)} \geq \frac{r}{3K} \right\}, \\ E_2 & := \left\{ n : \omega^*(f; \delta_n) \geq \frac{r}{3K} \right\}, \\ E_3 & := \left\{ n : \omega^*(f; \delta_n) \left\| \sum_{k=1}^{\infty} a_{nk} T_k(f_0; x, \gamma) - f_0(x, \gamma) \right\|_{C_B(K)} \geq \frac{r}{3K} \right\}. \end{aligned}$$

Then $E \subset E_1 \cup E_2 \cup E_3$ and therefore $\delta(E) \leq \delta(E_1) + \delta(E_2) + \delta(E_3)$. Using conditions (i) and (ii) we conclude

$$st - \lim_{n \rightarrow \infty} \left\| \sum_{k=1}^{\infty} a_{nk} T_k(f) - f \right\|_{C_B(K)} = 0.$$

This completes the proof of the theorem.

4 Example and the concluding remark

We show that the following double sequence of positive linear operators satisfies the conditions of Theorem 2.1 but does not satisfy the conditions of Corollary 2.2 and Theorem 2.1 of [8].

Example 4.1. Consider the following Bleimann et al. [20] (of two variables) operators:

$$B_n(f; x, y) := \frac{1}{(1+x)^n(1+y)^n} \sum_{j=0}^n \sum_{k=0}^n f\left(\frac{j}{n-j+1}, \frac{k}{n-k+1}\right) \binom{n}{j} \binom{n}{k} x^j y^k, \quad (4.1)$$

where $f \in H_\omega(K)$, $K = [0, \infty) \times [0, \infty)$ and $n \in \mathbb{N}$.

Since

$$(1+x)^n = \sum_{j=0}^n \binom{n}{j} x^j,$$

it is easy to see that

$$B_n(f_0; x, y) \rightarrow 1 = f_0(x, y).$$

Also by simple calculation, we obtain

$$B_n(f_1; x, y) = \frac{n}{n+1} \left(\frac{x}{1+x}\right) \rightarrow \frac{x}{1+x} = f_1(x, y),$$

and

$$B_n(f_2; x, y) = \frac{n}{n+1} \left(\frac{y}{1+y}\right) \rightarrow \frac{y}{1+y} = f_2(x, y).$$

Finally, we get

$$\begin{aligned} & B_n(f_3; x, y) \\ &= \frac{n(n-1)}{(n+1)^2} \left(\frac{x}{1+x}\right)^2 + \frac{n}{(n+1)^2} \left(\frac{x}{1+x}\right) \\ & \quad + \frac{n(n-1)}{(n+1)^2} \left(\frac{y}{1+y}\right)^2 + \frac{n}{(n+1)^2} \left(\frac{y}{1+y}\right) \\ & \rightarrow \left(\frac{x}{1+x}\right)^2 + \left(\frac{y}{1+y}\right)^2 = f_3(x, y). \end{aligned}$$

Now, take $A = C(1, 1)$ and define $u = (u_n)$ by (1.1). Let the operator $L_n : H_\omega(K) \rightarrow C_B(K)$ be defined by

$$L_n(f; x, y) = (1 + u_n)B_n(f; x, y).$$

It is easy to see that the sequence (L_n) satisfies the conditions (2.1)-(2.4). Hence by Theorem 2.1, we have

$$\begin{aligned} & st - \lim_{m \rightarrow \infty} \left\| \sum_{n=1}^{\infty} a_{mn} L_n(f; x, \gamma) - f(x, \gamma) \right\|_{C_b(K)} \\ &= st - \lim_{m \rightarrow \infty} \left\| \frac{1}{m} \sum_{n=1}^m L_n(f; x, \gamma) - f(x, \gamma) \right\|_{C_b(K)} = 0. \end{aligned}$$

On the other hand, the sequence (L_n) does not satisfy the conditions of Theorem A and Corollary 2.2 and Theorem 2.1 of [8], since (L_n) is neither convergent nor statistically (nor A -statistically) convergent. That is, Theorem A, Corollary 2.2 and Theorem 2.1 of [8] do not work for our operators L_n . Hence our Theorem 2.1 is stronger than Corollary 2.2 and Theorem 2.1 of [8].

Author details

¹Department of Mathematics, Aligarh Muslim University, Aligarh 202002, India ²Department of Mathematics, King Abdulaziz University, Jeddah 21589, Kingdom of Saudi Arabia

Authors' contributions

The main result of the paper was proposed and proved by AA, Section 3 was given by MM, and Section 4 was designed by both the authors jointly. All authors read and approved the final manuscript.

Competing interests

The authors declare that they have no competing interests.

Received: 13 March 2012 Accepted: 23 May 2012 Published: 23 May 2012

References

1. Fast, H: Sur la convergence statistique. *Colloq Math.* **2**, 241–244 (1951)
2. Moricz, F: Statistical convergence of multiple sequences. *Arch Math.* **81**, 82–89 (2003). doi:10.1007/s00013-003-0506-9
3. Mursaleen, M, Edely, OHH: Statistical convergence of double sequences. *J Math Anal Appl.* **288**, 223–231 (2003). doi:10.1016/j.jmaa.2003.08.004
4. Edely, OHH, Mursaleen, M: On statistical A -summability. *Math Comput Model.* **49**, 672–680 (2009). doi:10.1016/j.mcm.2008.05.053
5. Belen, C, Mursaleen, M, Yildirim, M: Statistical A -summability of double sequences and a Korovkin type approximation theorem. *Bull Korean Math Soc.* **49**(4), 851–861 (2012)
6. Korovkin, PP: *Linear Operators And Approximation Theory*. Hindustan Publ. Co., Delhi (1960)
7. Çakar, Ö, Gadjevi, AD: On uniform approximation by Bleimann, Butzer and Hahn on all positive semiaxis. *Tras Acad Sci Azerb Ser Phys Tech Math Sci.* **19**, 21–26 (1999)
8. Erkuş, E, Duman, O: A -Statistical extension of the Korovkin type approximation theorem. *Proc Indian Acad Sci (Math Sci).* **115**(4), 499–508 (2005). doi:10.1007/BF02829812
9. Anastassiou, GA, Mursaleen, M, Mohiuddine, SA: Some approximation theorems for functions of two variables through almost convergence of double sequences. *J Comput Anal Appl.* **13**, 37–40 (2011)
10. Mohiuddine, SA: An application of almost convergence in approximation theorems. *Appl Math Lett.* **24**, 1856–1860 (2011). doi:10.1016/j.aml.2011.05.006
11. Edely, OHH, Mohiuddine, SA, Noman, AK: Korovkin type approximation theorems obtained through generalized statistical convergence. *Appl Math Lett.* **23**, 1382–1387 (2010). doi:10.1016/j.aml.2010.07.004
12. Mursaleen, M, Alotaibi, A: Statistical summability and approximation by de la Vallée-Poussin mean. *Appl Math Lett.* **24**, 320–324 (2011). doi:10.1016/j.aml.2010.10.014
13. Mursaleen, M, Alotaibi, A: Statistical lacunary summability and a Korovkin type approximation theorem. *Ann Univ Ferrara.* **57**(2), 373–381 (2011). doi:10.1007/s11565-011-0122-8
14. Mursaleen, M, Karakaya, V, Ertürk, M, Gürsoy, F: Weighted statistical convergence and its application to Korovkin type approximation theorem. *Appl Math Comput.* **218**, 9132–9137 (2012). doi:10.1016/j.amc.2012.02.068
15. Srivastava, HM, Mursaleen, M, Khan, A: Generalized equi-statistical convergence of positive linear operators and associated approximation theorems. *Math Comput Mod.* **55**, 2040–2051 (2012). doi:10.1016/j.mcm.2011.12.011
16. Demirci, K, Karakuş, S: Statistical A -summability of positive linear operators. *Math Comput Model.* **53**, 189–195 (2011). doi:10.1016/j.mcm.2010.08.003
17. Demirci, K, Karakuş, S: Korovkin-type approximation theorem for double sequences of positive linear operators via statistical A -summability. *Results Math* (2011). doi:10.1007/s00025-011-0140-y
18. Dirik, F, Demirci, K: Korovkin type approximation theorem for functions of two variables in statistical sense. *Turk J Math.* **34**, 73–83 (2010)
19. Dirik, F, Demirci, K: A Korovkin type approximation theorem for double sequences of positive linear operators of two variables in A -statistical sense. *Bull Korean Math Soc.* **47**, 825–837 (2010). doi:10.4134/BKMS.2010.47.4.825

20. Bleimann, G, Butzer, PL, Hahn, L: A Bernstein type operator approximating continuous functions on semiaxis. *Indag Math.* **42**, 255–262 (1980)

doi:10.1186/1687-1847-2012-65

Cite this article as: Mursaleen and Alotaibi: Korovkin type approximation theorem for functions of two variables through statistical **A**-summability. *Advances in Difference Equations* 2012 **2012**:65.

Submit your manuscript to a SpringerOpen[®] journal and benefit from:

- ▶ Convenient online submission
- ▶ Rigorous peer review
- ▶ Immediate publication on acceptance
- ▶ Open access: articles freely available online
- ▶ High visibility within the field
- ▶ Retaining the copyright to your article

Submit your next manuscript at ▶ springeropen.com
