

RESEARCH ARTICLE

Open Access

The binding characteristics of isoniazid with copper–zinc superoxide dismutase and its effect on enzymatic activity

Nana Du¹, Liangquan Sheng^{1,2*}, Zhaodi Liu¹, Xiaojuan Hu^{1,2}, Huajie Xu¹ and Shuisheng Chen¹

Abstract

Background: Isoniazid (INH) is front-line anti-tuberculosis (TB) drugs, which are usually prescribed to TB patients for a total period of 6 months. Antituberculosis drug-induced hepatotoxicity (ATDH) is a serious adverse reaction of TB treatment. It is reported that INH-induced hepatotoxicity is associated with oxidative stress. Superoxide dismutase (SOD, EC 1.15.1.1) is the key enzyme for the protection of oxidative stress, which catalyzes the removal of superoxide radical anion, thereby raising the need to better understand the interaction between INH and SOD.

Results: The experimental results showed that the fluorescence intensity of Cu/Zn-SOD regularly decreased owing to form a 1:1 INH-SOD complex. According to the corresponding association constants (K_{SV}) between INH and SOD obtained from Stern–Volmer plot, it is shown that values of K_A are 1.01×10^4 , 5.31×10^3 , 3.33×10^3 , $2.20 \times 10^3 \text{ L} \cdot \text{mol}^{-1}$ at four different temperatures, respectively. The binding constants, binding sites and the corresponding thermodynamic parameters (ΔH , ΔG and ΔS) were calculated. A value of 3.93 nm for the average distance between INH and chromophore of Cu/Zn-SOD was derived from Förster theory of non-radiation energy transfer. The conformational investigation showed that the presence of INH resulted in the microenvironment and conformational changes of Cu/Zn-SOD. In addition, Effects of INH on superoxide dismutase activity was examined.

Conclusions: The results show that the hydrogen bonding and van der Waals forces play major roles in stabilizing the 1:1 INH-SOD complex. After addition of INH during the range of the experiment, the conformation and microenvironment of Cu/Zn-SOD are changed, but the activity of Cu/Zn-SOD is not changed.

Keywords: Superoxide dismutase, Isoniazid, Interaction, Antioxidation

Background

Tuberculosis (TB) is one of the leading causes of death due to a single disease, accounting for up to 2 million lives each year [1]. Isoniazid (INH) is the foremost first-line antibiotic used to treat TB, which continues to be the cornerstone of all antituberculosis regimens and remains the only agent recommended for tuberculosis chemoprophylaxis for children [2].

Despite numerous and intensive studies, we have a limited knowledge of the action mechanism of INH [3–6]. The consensus opinion is that in the presence of a slow flux of H_2O_2 or superoxide, KatG converts INH

into a radical species which is subsequently coupled to NADH to form an INH-NAD(P) adduct. The INH-NAD adduct is a potent inhibitor of *Mtb* InhA, an enoyl reductase required for the elongation steps in mycolic acid biosynthesis [7,8]. It has been recently suggested that the superoxide reactivity affects antitubercular activity of INH to some extent, and thereby raises the needs to better understand the interaction between INH and superoxide [9].

Furthermore, with increasing occurrence of TB all over the world, a growing number of patients may be at risk for severe adverse drug reactions (ADRs) such as antituberculosis drug-induced hepatotoxicity (ATDH) when treated with antituberculosis chemotherapy [10,11]. ATDH can be fatal if it is not recognised at an early stage, after which therapy should be interrupted timely. Moreover, ATDH has a negative impact on

* Correspondence: shenglq@fync.edu.cn

¹College of Chemistry and Chemical Engineering, Fuyang Normal College, Fuyang 236041, People's Republic of China

²College of Chemistry and Chemical Engineering, Anhui University, Hefei 230039, People's Republic of China

therapy adherence, decreases success rates of treatment and eventually increases treatment failure, relapse or the emergence of drug resistance. The occurrence of hepatotoxicity related to INH has been well-defined [12] and shown to increase as a result of drug-drug pharmacokinetic or pharmacodynamic interactions [13]. It is generally acknowledged that INH-induced hepatotoxicity is associated with oxidative stress [14].

Superoxide dismutase (SOD, EC 1.15.1.1) is an antioxidant enzyme in animals, plants, fungi and bacteria, which catalyzes the removal of superoxide radical anion ($O_2^{\bullet-}$) to hydrogen peroxide (H_2O_2) that can be subsequently converted to water by the enzyme catalase [15,16]. Therefore, it is the key enzyme for the protection of oxidative stress and could be a suitable candidate drug to protect liver from ATDH [17-19]. To better understand the protective mechanism of SOD, the interaction between SOD and INH should be investigated.


Protein–drug interaction is a hot topic in fields of medicine, chemistry and biology. Also, due to the effects on the drug of pharmacokinetics when bounded to protein, there is an increasing interest in protein from clinical and pharmaceutical perspectives [20-22]. Furthermore, the binding of drugs to protein in vitro is considered as a model in protein chemistry to study the binding behavior of protein. Consequently, the investigation of the binding between drugs and SOD is of fundamental importance in pharmacology and pharmacodynamics. In addition, the analysis of the interaction between SOD and small drug molecules is interesting to some extent, because the binding of specific small molecules to SOD has only been characterized in detail for a few examples [23,24].

In this paper, the interaction between INH and Cu/Zn-SOD has been investigated with fluorescence and ultraviolet spectroscopy. The binding mechanism, binding constants, binding sites and binding distance were obtained. The nature of the binding force was analyzed based on the thermodynamic parameters. Also, the effects of INH on the conformation of Cu/Zn-SOD were examined by synchronous fluorescence and three-dimensional fluorescence spectroscopy. What is more, the effects of INH on the activity of Cu/Zn-SOD were also explored. We hope that this work can provide useful information for pharmacology of INH.

Results and discussion

Effects of INH on Cu/Zn-SOD fluorescence

The fluorescence of Cu/Zn-SOD comes from the tryptophan, tyrosine and phenylalanine residues [25]. With 280 nm of excited wavelength, the maximum emission peak of Cu/Zn-SOD could be achieved at 306 nm [26]. Figure 1 shows the fluorescence spectra of Cu/Zn-SOD upon the addition of INH. Obviously, with the increasing concentration of INH, the fluorescence intensity



decreased regularly but without significant position-shift. The strong quenching of the fluorescence clearly indicated that the interaction between INH and Cu/Zn-SOD changed the microenvironment of chromophore or the tertiary structure of Cu/Zn-SOD.

The quenching mechanism of Cu/Zn-SOD fluorescence by INH

Fluorescence quenching refers to any processes which decrease the fluorescence intensity of a sample such as excited state reactions, energy transfers, ground-state complexes formation and collisional process [27]. Mechanisms of quenching are usually divided into dynamic quenching and static quenching [27]. Different quenching mechanisms can be distinguished by their different dependence on temperature and excited-state life time. As higher temperatures could result in larger diffusion coefficients, the dynamic quenching constants are expected to be larger with an increasing temperature. In contrast, the increase of temperature is likely to result in a decreased stability of complexes; thus, the values of static quenching constants are expected to be smaller. In the present work, the fluorescence-quenching spectra of Cu/Zn-SOD with presence of different concentrations of INH at four different temperatures (the temperatures used were 292, 295, 299 and 303 K) were measured. The fluorescence quenching was usually analyzed by the well-known Stern–Volmer equation [28,29]:

$$F_0/F = 1 + K_{SV}[Q] = 1 + K_q\tau_0[Q] \quad (1)$$

where F_0 and F are fluorescence intensities of Cu/Zn-SOD in the absence and presence of the quencher (INH), respectively, K_q is the quenching rate constant, K_{SV} is the Stern–Volmer constant, τ_0 is the average life time of the

molecule without quencher and the fluorescence lifetime of the biopolymer is 10^{-8} s, $[Q]$ is the quencher concentration.

Figure 2 shows the Stern-Volmer plots of the Cu/Zn-SOD quenching fluorescence by INH. The values of K_{SV} obtained from the slope of plots are 5.06×10^3 , 4.45×10^3 , 3.89×10^3 and 3.41×10^3 L·mol⁻¹, respectively. It is evident that the K_{SV} values are inversely correlated with temperature (inset in Figure 2), and K_q is much greater than the value of the maximum scatter collision quenching constant (2.0×10^{10} L·mol⁻¹·s⁻¹), which indicates the static mechanism of quenching and reveals that the fluorescence quenching is caused by specific interaction and complex formation between Cu/Zn-SOD and INH molecules [27].

Calculation of binding parameters

For static quenching, the following equation was employed to calculate the binding constant and binding sites [30,31]:

$$\log \frac{F_0 - F}{F} = \log K_A + n \log [Q] \quad (2)$$

where F_0 , F and $[Q]$ are the same as in Eq.(1), K_A and n are the binding constant and the number of binding sites respectively, K_A and n can be obtained from the plots of $\log[(F_0 - F)/F]$ vs. $\log[Q]$ for the INH-SOD system at four temperatures. The calculated results are shown in Table 1. The results show that K_A decreases with the increase of temperature, which probably indicates the formation of a compound partially decomposes at higher temperatures. The value of n is equal to approximately 1, indicating the involvement of a single binding site in the INH-SOD interaction.




Figure 2 Stern-Volmer plots for the quenching of SOD by INH at four different temperatures, (pH = 7.4). Inset: K_{SV} - T .

In order to confirm the probable quenching mechanism of Cu/Zn-SOD by INH, the absorption spectra of Cu/Zn-SOD in the presence and absence of INH at 303 K were recorded and presented in Figure 3. As seen in Figure 3, with the addition of INH the absorbance peak around 213 nm, which was mainly caused by the transition π - π^* of peptide bonds in SOD decreased regularly with peak red-shift (from 213 to 220). Thus, the ground state complex between SOD and INH was possibly formed. The absorbance peak caused by the transition π - π^* gets red-shifted with the increase of solvent polarity, so we can draw a conclusion that the polarity of the microenvironment of SOD is altered, leading to a conformational change that the peptide strands of SOD are extended and its hydrophobicity is decreased [28,30,31].

According to literature [32,33], when one small organic molecule binds to biological macromolecule, the relationship between absorbance intensity of biological macromolecules and the small organic molecule concentration can be written as follows:

$$(A_0 - A)^{-1} = A_0^{-1} + K_A^{-1} A_0^{-1} [Q]^{-1} \quad (3)$$

where A_0 and A are the absorbance intensities of Cu/Zn-SOD in the absence and presence of the INH, respectively, K_A and $[Q]$ are the binding constant and INH concentration. As shown in the inset in Figure 3, plot of $1/(A_0 - A)$ against $1/[Q]$ gives a straight line, the correlation coefficient of which is 0.9986 and the binding constant (K_A) is equal to 2.32×10^3 L·mol⁻¹. The results further prove that only one molecule of INH binds Cu/Zn-SOD.

Thermodynamic analysis and intermolecular forces

The intermolecular forces of interaction between small molecules and biomolecules mainly include hydrogen bonding, van der Waals forces, electro static forces and hydrophobic forces. Within a small temperature range, the enthalpy of interaction can be regarded as a constant leading to the following equations [34,35]:

$$\ln(K_{A2}/K_{A1}) = \Delta H(1/T_1 - 1/T_2)/R \quad (4)$$

$$\Delta G = \Delta H - T\Delta S \quad (5)$$

$$\Delta S = -(\Delta G - \Delta H)/T \quad (6)$$

where R is the gas constant, T is the absolute temperature and K_A is the apparent binding constant at corresponding temperature T . If $\Delta S > 0$ and $\Delta H > 0$, the main force of interaction is the hydrophobic force, if $\Delta S > 0$, $\Delta H < 0$, the main force is electro static attraction, if $\Delta S < 0$, $\Delta H < 0$, the main force includes both van der Waals forces and hydrogen bonding. Table 1 shows the values of ΔH , ΔS and ΔG at four temperatures. The negative values of ΔG reveal that the interaction process is spontaneous. As $\Delta S < 0$ and $\Delta H < 0$, which indicates that

Table 1 Binding parameters, number of binding sites and thermodynamic parameters of INH-SOD at four temperatures

T(K)	K_{SV} (L · mol ⁻¹)	K_A (L · mol ⁻¹)	n	ΔG (kJ · mol ⁻¹)	ΔH (kJ · mol ⁻¹)	ΔS (J · mol ⁻¹ · K ⁻¹)
292	5.06×10^3	1.01×10^4	1.069	-22.38 (s = 1.69)	-105.70 (s = 9.54)	-284.66 (s = 18.50)
295	4.45×10^3	5.31×10^3	1.019	-21.04 (s = 1.73)		-286.31 (s = 15.67)
299	3.89×10^3	3.33×10^3	0.995	-20.16 (s = 1.54)		-285.42 (s = 19.92)
303	3.41×10^3	2.20×10^3	0.999	-19.36 (s = 1.26)		-284.29 (s = 19.31)

the interaction between INH and Cu/Zn-SOD is mainly due to hydrogen bonding and van der Waals forces [34,35].

Energy transfer between Cu/Zn-SOD and INH

The energy transfer in biochemistry is important, because the efficiency of transfer can be used to evaluate the distance between the ligand and the chromophore in protein. The overlap of the UV-Vis absorption spectrum of INH with the fluorescence emission spectrum of Cu/Zn-SOD, as is shown in Figure 4. According to Förster nonradiative energy-transfer theory, the efficiency of energy transfer is E , the critical distance for 50% energy transfer is R_0 , and the actual distance of separation is r . These values were calculated by Eqs. (7, 8, 9) [35-37]:

$$E = 1 - \frac{F}{F_0} = \frac{R_0^6}{R_0^6 + r^6} \quad (7)$$

$$R_0^6 = 8.8 \times 10^{-25} K^2 \Phi N^{-4} J \quad (8)$$

where r is the distance from the ligand to the chromophore of the protein, and R_0 is the Förster critical



distance at which 50% of the excitation energy is transferred to the acceptor.

$$J = \frac{\sum F(\lambda)\epsilon(\lambda)\lambda^4\Delta\lambda}{\sum F(\lambda)\Delta\lambda} \quad (9)$$

where $F(\lambda)$ is the fluorescence intensity of the donor at the wavelength range λ , and $\epsilon(\lambda)$ is the molar absorption coefficient of the acceptor at wavelength λ . In this case, $K_2 = 2/3$, $N = 1.336$ and $\Phi = 0.13$ [36]. According to the equation (7), (8), (9), we could calculate that $J = 2.3 \times 10^{-13} \text{ cm}^3 \cdot \text{L} \cdot \text{mol}^{-1}$, $E = 0.022$, $R_0 = 2.58 \text{ nm}$, and $r = 3.93 \text{ nm}$. The binding distance between INH and Cu/Zn-SOD is shorter than 7 nm, which indicates that the energy transfer from Cu/Zn-SOD to INH occurs with high possibility [35-37]. It is also suggested that the binding of INH to Cu/Zn-SOD is conducted through energy transfer, which is consistent with a static quenching mechanism.


Conformation investigation of Cu/Zn-SOD

The influence of INH on conformational changes of Cu/Zn-SOD was assessed by synchronous fluorescence. Synchronous fluorescence measurement provides information about the molecular microenvironment in the vicinity of the fluorophore functional groups [38]. When the wavelength shift ($\Delta\lambda$) between the emission and




excitation wavelengths was stabilized at 20 or 80 nm [39], the synchronous fluorescence gives the characteristic information of tyrosine or tryptophan residues, respectively. Figure 5A and Figure 5B show the effects of INH on the synchronous fluorescence of Cu/Zn-SOD. As seen in Figure 5, the fluorescence intensities of tryptophan residue and tyrosine residue decreased with the increase of the concentration of INH. The maximum emission wavelength of tyrosine residues does not undergo a significant shift in the investigated concentration range. In contrast, an obvious red shift was observed for tryptophan residues, which indicated that the polarity around tryptophan residues increased. In other words, tryptophan residues were placed in a less hydrophobic environment and were more exposed to the solvent.

In recent years, the three-dimensional fluorescence technique has been widely used, because it allows fluorescence characteristics to be acquired by changing the excitation and emission wavelengths simultaneously. We studied the conformational and microenvironmental changes of Cu/Zn-SOD by comparing the spectral changes in the absence and presence of INH, as shown



in Figure 6. In these figures, peak a is the Rayleigh scattering peak ($\lambda_{ex} = \lambda_{em}$), the peak intensity of which increased with the addition of INH. A reasonable explanation may be that an INH-SOD complex formed after the addition of INH, increased the diameter of the macromolecule, and thus enhanced the scattering effects [40]. Peak b is the second-ordered scattering peak ($\lambda_{em} = 2\lambda_{ex}$). Peak 1 ($\lambda_{ex} = 280.0$ nm, $\lambda_{em} = 306.0$ nm) mainly reveals the spectral behavior of the Trp and Tyr residues since excitation of SOD at this wavelength mainly reveals their intrinsic fluorescence, and the fluorescence of the phenylalanine residue can be negligible [25]. Besides peak 1, there is another strong fluorescence peak (peak 2, $\lambda_{ex} = 232.0$ nm, $\lambda_{em} = 306.0$ nm), which mainly exhibits the fluorescence spectral behavior of polypeptide backbone structures [40]. Both peaks were involved in



fluorescence quenching of Cu/Zn-SOD by INH to different degrees. By analyzing the intensity changes of peak 1 and peak 2, in the absence and presence of INH, the fluorescence intensity ratio of peak 1 to peak 2 is 1.16:1 and 1.18:1, respectively. All of these phenomena and analyses revealed that the binding of INH to SOD induced some microenvironmental and conformational changes in Cu/Zn-SOD.

Effects of INH on Cu/Zn-SOD activity

The activity of Cu/Zn-SOD after the addition of INH was studied to investigate the effects of INH on superoxide dismutase activity. Figure 7a shows the relationship between antioxidant effect of INH-SOD and INH concentration. Figure 7b shows the relationship between antioxidant effect of dissociative INH and its concentration. Due to antioxidant effect of INH in the range of the experiment, the antioxidation of INH-SOD was strengthened with increasing INH concentration. The enzyme activity was determined by subtracting the antioxidant effect of INH from antioxidant effect of INH-SOD (Figure 7c). The results show that the enzyme activity is almost the same throughout the experiment. The observation leads to the conclusion that the INH does not affect the activity of Cu/Zn-SOD.

Experimental

Materials

Cu/Zn-SOD was purchased from Doulai Biotechnology Company (Nanjing, China), INH was purchased from Taizhou Medical Corporation (Zhejiang, China). SOD was directly dissolved in redistilled water to prepare the stock solution (0.12 mM), and the stock solution was kept in the dark at 0–4°C. INH solution was obtained by dissolving it in redistilled water (1.2 mM). 0.05 M Tris-HCl buffer solution of pH 7.4 and 8.2 were prepared. Other

reagents are local products of analytical grade. The water used was redistilled and ion-free.

Fluorescence measurements were carried out in 1.0 cm quartz cells on a FluoroMax-4 spectrophotometer (Horiba, Japan) equipped with a thermostat bath. UV-vis spectra were recorded on an 8000A spectrophotometer (Beijing Purkinje General Corporation, China). All pH measurements were made using a pH-3 digital pH-meter (Chengdu Sanke Instrument Corporation, China) with a combined glass electrode.

Assay

A 3.0 mL Tris-HCl (pH 7.4, 0.05 M) buffer solution containing 4 μ M SOD was titrated by successive additions of 1.2 mM INH solution and the concentration of INH varied from 0 to 40 μ M. Titrations were done manually by using the microinjectors. After reaction for 10 min, fluorescence spectra were measured in the range of 290–450 nm at the excitation wavelength of 280 nm. The fluorescence spectra were performed at four temperatures (292, 295, 299 and 303 K). The excitation and emission slits were 5 nm.

The synchronous fluorescence spectra were recorded at 299 K from 260 nm to 350 nm (for tyrosine residues) at $\Delta\lambda = 20$ and from 280 to 400 at $\Delta\lambda = 80$ nm (for tryptophan residues). The excitation and emission slits were 10 nm.

Three-dimensional fluorescence spectra of SOD were recorded at 299 K in the presence and absence of INH with an excitation wavelength in the range 230–300 nm and an emission wavelength in the range 260–500 nm. The excitation and emission slits were 5 nm.

A 3.0 mL Tris-HCl (pH 7.4, 0.05 M) buffer solution containing 4 μ M SOD was titrated by successive additions of 1.2 mM INH solution and the concentration of INH varied from 20 to 200 μ M. Titrations were done manually by using the microinjectors. After reaction for 10 min at 303 K, absorption spectra of SOD-INH were recorded in the range 190–400 nm.

The activity of Cu/Zn-SOD was assayed by using the pyrogallol autoxidation method. The assay mixture contained 5.0 ml Tris-HCl (pH 8.2, 0.05 M) buffer, 0.1 mM pyrogallol solution, 2.5 nM SOD solution and different concentrations of INH. The concentration of INH varied from 0 to 2.4 mM. Absorption at 325 nm at 303 K against time was recorded using a spectrophotometer. The SOD activity was determined as pyrogallol autoxidation rate.

Conclusions

In this paper, the interaction between INH and SOD was studied using fluorescence and ultraviolet spectroscopy at different temperatures under imitated physiological conditions. The results show that the quenching




Figure 7 Effect of INH on the activity of superoxide dismutase (SOD) (pH = 7.4, T = 303 K). **a** antioxidant effect of INH-SOD, **b** antioxidant effect of INH, **c** enzyme relative activity of SOD.

mechanism of fluorescence of SOD by INH is a static quenching process. The binding constants, binding sites and the corresponding thermodynamic parameters (ΔH , ΔG and ΔS) were determined, which indicates that hydrogen bonding and van der Waals forces play a major role in the binding process. According to Förster theory of nonradiation energy transfer, the binding distance between Cu/Zn-SOD and INH is 3.93 nm and the energy transfer occurs between SOD and INH with high probability. UV-vis, synchronous fluorescence and three-dimensional fluorescence studies indicates that the interaction leads to a change in the conformation and microenvironment of Cu/Zn-SOD. Moreover, determination of SOD activity in different concentration INH leads to the conclusion that the binding of INH with SOD does not affect the activity of Cu/Zn-SOD. This report has special significance in pharmacology and clinical medicine as well as methodology.

Competing interests

The authors declare that they have no competing interests.

Authors' contributions

ND made a significant contribution to acquisition of data, analysis and manuscript preparation. XH has made a substantial contribution to experimental design and data analysis. ZL and HX participated in partial experiments. LS made a significant contribution to experimental design, data analysis, and manuscript revision. SC participated in study design and manuscript revision. All authors read and approved the final manuscript.

Acknowledgement

This work was financially supported by the International Sea Area Resources Survey and Development of the 12th Five-year Plan of China (DY125-15-E-01), the Nature Science Foundation of China (20971024) and the Higher Education Institutions Key Nature Science Foundation of Anhui (kj2009A127).

Received: 17 April 2013 Accepted: 30 May 2013

Published: 6 June 2013

References

1. Iyawoo K: Tuberculosis in Malaysia: problems and prospect of treatment and control. *Tuberculosis* 2004, **84**:4–7.
2. Prasad B, Singh S: LC-MS/TOF and UHPLC-MS/MS study of *in vivo* fate of rifamycin isonicotinyl hydrazone formed on oral co-administration of rifampicin and isoniazid. *J Pharmaceut Biomed* 2010, **52**:377–383.
3. Verma RK, Mukker JK, Singh RSP, Kumar K, Verma PRP, Misra A: Partial biodistribution and pharmacokinetics of isoniazid and rifabutin following pulmonary delivery of inhalable microparticles to rhesus macaques. *Mol Pharm* 2012, **9**:1011–1016.
4. Ràfols C, Bosch E, Ruiz R, Box KJ, Reis M, Ventura C, Santos S, Araújo ME, Martins F: Acidity and hydrophobicity of several new potential antitubercular drugs: isoniazid and benzimidazole derivatives. *J Chem Eng Data* 2012, **57**:330–338.
5. Vilchère C, Wang F, Arai M, Hazbón MH, Colangeli R, Kremer L, Weisbrod TR, Alland D, Sacchetti JC, Jacobs WR: Transfer of a point mutation in MT *inhA* resolves the target for isoniazid. *Nat Med* 2006, **12**:1027–1029.
6. Argyrou A, Vetting MW, Blanchard JS: New insight into the mechanism of action of and resistance to isoniazid: interaction of mycobacterium tuberculosis enoyl-ACP reductase with INH-NADP. *J Am Chem Soc* 2007, **129**:9582–9583.
7. Zhao XB, Yu SW, Magliozzo RS: Characterization of the binding of isoniazid and analogues to Mycobacterium tuberculosis catalase-peroxidase. *Biochemistry* 2007, **46**:3161–3170.
8. Wahab HA, Choong YS, Ibrahim P, Sadikun A, Scior T: Elucidating isoniazid resistance using molecular modeling. *J Chem Inf Model* 2009, **49**:97–107.
9. Ghiladi RA, Medzihradsky KF, Rusnak FM, Ortiz de Montellano PR: Correlation between isoniazid resistance and superoxide reactivity in *Mycobacterium tuberculosis* KatG. *J Am Chem Soc* 2005, **127**:13428–13442.
10. Nicolas V, Michel FR, Alberto C, Florence D, Pierre D, Bernadette M, Laurent N, Jules D, Denis H: CYP2E1 genotype and isoniazid-induced hepatotoxicity in patients treated for latent tuberculosis. *Eur J Clin Pharmacol* 2006, **62**:423–429.
11. Alma T, Martin JB, Wilbert HMP, Hennie MJR, Rob EA, André JAM, van der Ven PN, Dekhuijzen R: Isoniazid and its toxic metabolite hydrazine induce *in vitro* pyrazinamide toxicity. *Int J Antimicrob Ag* 2008, **31**:577–580.
12. Yee D, Valiquette C, Pelletier M, Parisien I, Rocher I, Menzies D: Incidence of serious side effects from first-line antituberculosis drugs among patients treated for active tuberculosis. *Am J Respir Crit Care Med* 2003, **167**:1472–1477.
13. Schaberg T, Rebhan K, Lode H: Risk factors for side effects of isoniazid, rifampin and pyrazinamide in patients hospitalized for pulmonary tuberculosis. *Eur Respir J* 1996, **9**:2026–2030.
14. Yakup E, Tulay E, Hafize U, Habibe G, Tuncay A, Ethem E: Effect of vitamin C on oxidative liver injury due to isoniazid in rats. *Pediatr Int* 2010, **52**:69–74.
15. McCord JM, Fridovich I: Superoxide dismutase. An enzymic function for erythrocyte hemocypretein (hemocypretein). *J Biol Chem* 1969, **244**:6049–6055.
16. Maurizi CP: Alzheimer's disease: roles for mitochondrial damage, the hydroxyl radical, and cerebrospinal fluid deficiency of melatonin. *Med Hypotheses* 2001, **57**:156–160.
17. Wang XF, Lv Y, Okamura T, Kawaguchi H, Wu G, Sun WY, Ueyama N: Structure variation of mercury(II) halide complexes with different imidazole-containing ligands. *Cryst Growth Des* 2007, **7**:1125–1133.
18. Hayakawa N, Asayama S, Noda Y, Shimizu T, Kawakami H: Pharmaceutical effect of manganese porphyrins on manganese superoxide dismutase deficient mice. *Mol Pharm* 2012, **9**:2956–2959.
19. Axthelm F, Casse O, Koppenol WH, Nauser T, Meier W, Palivan CG: Antioxidant nanoreactor based on superoxide dismutase encapsulated in superoxide-permeable vesicles. *J Phys Chem B* 2008, **112**:8211–8217.
20. Maji SK, Amsden JJ, Rothschild KJ, Condrón MM, Teplow DB: Conformational dynamics of amyloid β -protein assembly probed using intrinsic fluorescence. *Biochemistry-US* 2005, **44**:13365–13376.
21. Xu XL, Zhang LY, Shen DK, Wu H, Peng LL, Li JH: Effect of metal ion substitutions in anticoagulation factor I from the venom of *Agkistrodon acutus* on the binding of activated coagulation factor X and on structural stability. *J Biol Inorg Chem* 2009, **14**:559–571.
22. Xu XL, Zhang LY, Luo ZF, Shen DK, Wu H, Peng LL, Song JJ, Zhang Y: Metal ions binding to NAD-glycohydrolase from the venom of *Agkistrodon acutus*: regulation of multicatalytic activity. *Metallomics* 2010, **2**:480–489.
23. Prasanth GK, Divya LM, Sadasivan C: Effects of mono and di(*n*-butyl) phthalate on superoxide dismutase. *Toxicology* 2009, **262**:38–42.
24. Banci L, Bertini I, Blažević O, Calderone V, Cantini F, Mao J, Trapananti A, Viero M, Amori I, Cozzolino M, Carri MT: Interaction of cisplatin with human superoxide dismutase. *J Am Chem Soc* 2012, **134**:7009–7014.
25. Sulkowska A, Bojko B, Równicka J, Sulkowski WW: Paracetamol and cytarabine binding competition in high affinity binding sites of transporting protein. *J Mol Struct* 2006, **792**:249–256.
26. Sheng LQ, Zheng XY, Tong HW, Liu SM, Liu QL: Study on interaction between copper ion and Hog liver CuZnSOD by spectroscopy. *Acta Chim Sinica* 2005, **63**:1759–1764.
27. Xu XL, Zhang LY, Shen DK, Wu H, Liu QL: Oxygen-dependent oxidation of Fe(II) to Fe(III) and interaction of Fe(III) with bovine serum albumin, leading to a hysteretic effect on the fluorescence of bovine serum albumin. *J Fluoresc* 2008, **18**:193–201.
28. Shen DK, Xu XL, Wu H, Peng LL, Zhang Y, Song JJ, Su QD: Metal ion binding to anticoagulation factor II from the venom of *Agkistrodon acutus*: stabilization of the structure and regulation of the binding affinity to activated coagulation factor X. *J Biol Inorg Chem* 2011, **16**:523–537.
29. Li YF, Yang GZ, Mei ZN: Spectroscopic and dynamic light scattering studies of the interaction between pterodonic acid and bovine serum albumin. *Acta Pharm Sin B* 2012, **2**:53–59.
30. Zhang LY, Xu XL, Luo ZF, Zhang Y, Shen DK, Peng LL, Song JJ: Cu(II)- and disulfide bonds-induced stabilization during the guanidine hydrochloride- and thermal-induced denaturation of NAD-glycohydrolase from the venom of *Agkistrodon acutus*. *Metallomics* 2012, **4**:166–173.

31. Markarian SA, Aznauryan MG: Study on the interaction between isoniazid and bovine serum albumin by fluorescence spectroscopy: the effect of dimethylsulfoxide. *Mol Biol Rep* 2012, **39**:7559–7567.
32. Benesi HA, Hildebrand JH: A spectrophotometric investigation of the interaction of iodine with aromatic hydrocarbons. *J Am Chem Soc* 1949, **71**:2703–2707.
33. Liu Y, Xie MX, Kang J: Influence of total saponins of panax notoginseng on the conformation of SOD. *Acta Chim Sinica* 2003, **61**:1305–1310.
34. Wu H, Xu XL, Shen DK, Peng LL, Song JJ, Yan Z: Binding of Ca^{2+} and Zn^{2+} to factor IX/X-binding protein from venom of *Agkistrodon Halys Pallas*: stabilization of the structure during GdnHCl- and thermal-induced denaturation. *J Biol Inorg Chem* 2011, **16**:69–79.
35. Zhang YZ, Zhang XP, Hou HN, Dai J, Liu Y: Study on the interaction between $\text{Cu}(\text{phen})_3^{2+}$ and bovine serum albumin by spectroscopic methods. *Biol Trace Elem Res* 2008, **121**:276–287.
36. Cheng ZJ: Interaction of ergosterol with bovine serum albumin and human serum albumin by spectroscopic analysis. *Mol Biol Rep* 2012, **39**:9493–9508.
37. Zhang YZ, Li HR, Dai J, Chen WJ, Zhang J, Liu Y: Spectroscopic studies on the binding of cobalt(II) 1,10-phenanthroline complex to bovine serum albumin. *Biol Trace Elem Res* 2010, **135**:136–152.
38. Hu YJ, Liu Y, Pi ZB, Qu SS: Interaction of cromolyn sodium with human serum albumin: a fluorescence quenching study. *Bioorg Med Chem* 2005, **13**:6609–6614.
39. Sheng LQ, Liu SM, Jia LL, Xu XL, Xie RS, Liu QL: The interaction of CuZn-superoxide dismutase and salicylic acid. *Chinese J Anal Chem* 2003, **31**:283–286.
40. Zhang YZ, Zhou B, Liu YX, Zhou CX, Ding XL, Liu Y: Fluorescence study on the interaction of bovine serum albumin with P-aminoazobenzene. *J Fluoresc* 2008, **18**:109–118.

doi:10.1186/1752-153X-7-97

Cite this article as: Du et al.: The binding characteristics of isoniazid with copper–zinc superoxide dismutase and its effect on enzymatic activity. *Chemistry Central Journal* 2013 **7**:97.

Publish with **ChemistryCentral** and every scientist can read your work free of charge

“Open access provides opportunities to our colleagues in other parts of the globe, by allowing anyone to view the content free of charge.”

W. Jeffery Hurst, The Hershey Company.

- available free of charge to the entire scientific community
- peer reviewed and published immediately upon acceptance
- cited in PubMed and archived on PubMed Central
- yours — you keep the copyright

Submit your manuscript here:
<http://www.chemistrycentral.com/manuscript/>



ChemistryCentral