

RESEARCH

Open Access

Cysteine- rich secretory protein 3 (CRISP3), ERG and PTEN define a molecular subtype of prostate cancer with implication to patients' prognosis

Samir Al Bashir^{1,6}, Mohammed Alshalalfa⁴, Samar A Hegazy¹, Michael Dolph¹, Bryan Donnelly^{4,5} and Tarek A Bismar^{1,2,3,4*}

Abstract

Cysteine- rich secretory protein 3 (CRISP3) prognostic significance in prostate cancer (PCA) has generated mixed result. Herein, we investigated and independently validated CRISP3 expression in relation to *ERG* and *PTEN* genomic aberrations and clinical outcome. CRISP3 protein expression was examined by immunohistochemistry using a cohort of patients with localized PCA (n = 215) and castration resistant PCA (CRPC) (n = 46). The Memorial Sloan Kettering (MSKCC) and Swedish cohorts were used for prognostic validation. Results showed, CRISP3 protein intensity to be significantly associated with neoplastic epithelium, being highest in CRPC vs. benign prostate tissue ($p < 0.0001$), but was not related to Gleason score (GS). CRISP3 mRNA was significantly associated with higher GS ($p = 0.022$ in MSKCC, $p = 1.1 \times 10^{-4}$ in Swedish). Significant association between CRISP3 expression and clinical outcome was documented at the mRNA but not the protein expression levels. CRISP3 mRNA expression was related to biochemical recurrence in the MSKCC ($p = 0.038$) and lethal disease in the Swedish cohort ($p = 0.0086$) and retained its prognostic value in the subgroup of patients with GS 6 & 7. Furthermore, CRISP3 protein and mRNA expression was significantly associated with positive *ERG* status and with *PTEN* deletions. Functional biology analysis documented phenylalanine metabolism as the most significant pathway governing high CRISP3 and *ERG* expression in this subtype of PCA. In conclusion, the combined status of CRISP3, *ERG* and *PTEN* define a molecular subtype of PCA with poorest and lethal outcome. Assessing their combined value may be of added value in stratifying patients into different prognostic groups and identify those with poorest clinical outcome.

Keywords: *CRISP3*, *ERG*, *PTEN*, Biological pathways, Prostate cancer, Poorest outcome, Molecular subtypes

Introduction

Prostate cancer (PCA) remains a major cancer in the western countries and ranks as the second leading cause of cancer related deaths among men [1]. Although PCA shows different biological behaviour in its disease progression and clinical course, there is strong correlation between pathological prognostic parameters and clinical outcome. In the past decade, several genomic studies have implemented methods to mark specific genomic alterations associated with disease progression and

categorize PCA into molecular subtypes with proposed prognostic and therapeutic subgroups. However, currently, those road maps are still being built and refined to better allow physicians to accurately offer better targeted treatment options for PCA patients.

Cysteine- rich secretory protein 3 (CRISP3) is a member of large family of cysteine- rich secretory proteins that are expressed in vertebrates, insects, plants, fungi, and yeast [2-6]. In humans, CRISP3 mRNA is expressed predominantly in salivary gland, prostate, and pancreas, and at much lower levels in epididymis, thymus, ovary, testis, colon, and lacrimal glands [7,8].

ERG gene rearrangement is the most abundant genomic aberrations in prostate cancer that accounts for around 50% of clinically localized disease. Furthermore, several studies have implicated *ERG* gene rearrangements with

* Correspondence: tarek.bismar@cls.ab.ca

¹Department of Pathology and Laboratory Medicine, University of Calgary and Calgary Laboratory Services, Calgary, AB, Canada

²Departments of Oncology, Biochemistry and Molecular Biology, Calgary, AB, Canada

Full list of author information is available at the end of the article

increased rate of *PTEN* deletion. The presence of both *ERG* and *PTEN* aberration has been suggested to signify a distinct molecular subtype of PCA. However, its prognostic implication remains unclear.

Recent studies have associated high CRISP3 with *ERG*-gene rearrangements and *PTEN* deletions and upregulation of transcript levels of CRISP3 has been previously documented in PCA compared to benign prostate tissue using in situ hybridization [2]. This upregulation was also confirmed by few other studies [7,9]. However, data related to the prognostic significance of CRISP 3 has been mixed with conflicting results [10-14] The aim of the current study was to investigate CRISP3 protein expression in large clinical cohort of localized disease and to assess its expression in relation to *ERG* gene rearrangement and *PTEN* deletions and across various stages of disease progression including castration resistant prostate cancer (CRPC). We also aimed to validate the prognostic value of CRISP3 in conjunction to the molecular subtype of PCA with *ERG* and *PTEN* genomic aberrations and to utilize bioinformatics to delineate possible genetic connections and pathways related to this molecular subtype of prostate cancer.

Material and methods

Study population and Tissue microarray construction

We utilized two cohorts in this study. The first cohort consisted of 215 patients treated by retro-pubic radical prostatectomy for localized prostate cancer between 1992–2004 with a mean follow-up of 4.8 years (range 0–15.8). Clinical progression was defined as a post-operative serum PSA elevation of >0.2 ng/ml assessed on two occasions following decrease to non detectable levels. Three tissue microarray (TMA) blocks were created from this cohort using a manual tissue arrayer (Beecher Instruments, Silver Spring, MD). Each block was assembled without prior knowledge of any clinical or pathological staging information. From each case, one to nine cores (average 3.3), 0.6 mm in diameter, were sampled from the paraffin embedded tissue blocks, containing benign, high grade intraepithelial neoplasia (HGPIN) and prostate cancer (PCA). We sampled both Gleason pattern originally present in the radical prostatectomy to reflect the average GS from each case. After construction, 4 µm sections were cut and stained with hematoxylin and eosin on the initial slides to confirm the histological diagnosis and grading of Gleason score. The patients' demographics of this cohort have been previously described [15,16]. The second cohort consisted of 46 patients with castration resistant prostate cancer assembled onto one TMA block. Tissues from those samples were obtained by transurethral resections of prostate performed to relieve clinical obstructive symptoms thus representing locally advanced disease. The study was approved by the University of Calgary ethics review board who waived informed

patients consent. All Clinical and pathological information of patients were collected under this approval.

Independent validation of mRNA expression data

Two independent prostate cancer expression data were used to validate the findings from the protein expression study. The first data is the Memorial Sloan Kettering Cancer Center (MSKCC) data [17]. The MSKCC sample data were downloaded from Gene Expression Omnibus (GEO) database series GSE21032. All tissues were collected during Radical prostatectomy surgery, snap frozen in liquid nitrogen, and stored at -80°C. Care was taken to identify regions of tumor free from contaminating stroma. The data has the gene expression of 29 normal samples, 131 primary tumor samples after Radical prostatectomy, 19 metastasis samples and 6 cell lines (DU145, LNCaP, LNCaP104R, LNCaP104S, PC3 and VCaP). Samples had a follow up time till biochemical relapse for up to 120 months and are mainly low risk samples. The second data is the Swedish cohort that was downloaded from GEO series GSE16560 that has expression of 281 localized cancer samples; mainly Gleason score 7, and follow up time of overall survival for up to 250 months. In the MSKCC data *ERG*, *PTEN* status was determined from expression data using PAM clustering implemented in cluster R package. In the Swedish data, *ERG* was determined by FISH and *PTEN* was determined from expression data.

CRISP3 protein expression by immunohistochemistry (IHC)

4 µm thick TMA sections from formalin-fixed paraffin tissue were cut and stained with Leica auto-stainer Bond-Max. The slides were pretreated for deparaffinization followed by heat induced antigen retrieval by Epitope retrieval solution 1 (Leica Microsystems, Buffalo Grove, IL, USA) for 20 minutes, and then incubated for 15 minutes at room temperature with CRISP3 rabbit polyclonal antibody (ProteinTech Group, Inc., Chicago, IL, USA) at 1:100 dilution. Bond polymer Refine Detection kit (Leica Microsystems, Buffalo Grove, IL, USA) was used for DABi detection and counter stain. A multi-tissue control TMA slide containing samples of various tissue organs was previously stained and utilized as positive and negative control to confirm antibody specificity.

Pathological analysis

All TMA cores were assigned a diagnosis (i.e. benign, HGPIN or PCA) and confirmed independently by three pathologists (SAB, SAH and TAB). Gleason scoring was assessed according to the 2005 ISUP criteria [18]. For each patient, the two predominant patterns were sampled and included on the TMAs blocks for analysis. CRISP3 protein expression was examined by two study pathologists (SAB, SAH) and the intensity of staining

was categorized using 4-tiered system according to the following parameters: The staining intensity (0, 1+, 2+, and 3+) for negative, weak, moderate and high intensity plus the fraction of positive tumor cells were documented for each tissue spot. A final score was achieved from these two parameters as follows: Negative(0): absence of CRISP3 staining in 100% of cells; weak (1): intensity of 1+ in >70% of tumor cells or staining intensity of 2+ in ≤30% of tumor cells; moderate(2): intensity of 1+ in >70% of tumor cells, or staining intensity of 2+ in >30% but ≤70% of tumor cells or staining intensity of 3+ ≤30% of tumor cells; strong(3): intensity of 2+ in >70% of tumor cells or staining intensity of 3+ in >30% of tumor cells. As CRISP3 was noticed to be weakly expressed in adjacent stromal cells, when we assessed CRISP3 expression in our study, we used the expression of adjacent stromal cells as a control for measuring the relative expression of epithelial cells.

ERG expression was previously evaluated on this cohort based on positive vs. negative expression as validated and correlated to *ERG* gene rearrangements using the break apart FISH probes (data not shown). *PTEN* deletions were previously evaluated and recorded as not deleted, hemizygous or homozygous deletions by counting 100 nuclei in each core as previously described using a four color FISH probe [19].

Statistical analysis

Statistical analysis was done to assess various clinical and pathological variables, means and ranges for continuous variables. Chi-square tests were conducted to test for associations between CRISP3 protein expression and Gleason score, surgical margin and pathological stage, *ERG* expression and *PTEN* genomic deletions. Wilcoxon statistical test was used to assess the significance of the difference of CRISP3 mRNA expression across Gleason score, *ERG* and *PTEN* status in the MSKCC and Swedish cohorts. CRISP3 protein expression in clinically localized prostate cancer cases association to disease free survival time was assessed using Cox Proportional Hazards regression analysis and the Kaplan-Meier curves along with the log-rank test. In all statistical tests a p value <0.05 was considered significant.

Results

Expression of CRISP3 protein in benign, HGPIN, localized and castration resistant prostate cancer

A total of 1194 core samples were available for evaluation in the localized cohort; [benign (n = 175), HGPIN (n = 77), PCA (n = 942)]. CRISP3 protein expression (reported as mean ± SD) showed significant increase in PCA (1.71 ± 0.63), compared to benign prostate tissue (1.04 ± 0.42) (p < 0.0001). In HGPIN, mean CRISP3 protein expression was comparable to PCA (1.65 ± 0.56) (p = 0.84) and still significantly higher than benign prostate tissue (P < 0.0001). In

the advanced/castration resistant prostate cancer samples (n = 92), CRISP3 protein mean expression levels were the highest with mean intensity of (2.09 ± 0.75) compared to all other types of neoplastic and benign epithelium (p < 0.0001) (Figure 1).

CRISP3 protein expression in association to Gleason score, surgical margin, pathological stage and PSA biochemical relapse


To investigate the association of CRISP3 protein expression in relation to Gleason score, we grouped samples based on Gleason score sum of individual TMA cores. To enable better correlation of CRISP3 protein expression with GS, CRISP3 protein expression was assessed as binary value, grouping moderate/high expression vs. negative/weak. In this cohort, we did not observe association between CRISP3 protein expression and GS. Moderate/high CRISP3 protein expression was noted in 25/68 (36%) of GS < 7 compared with 52/145 (35.8%) of GS ≥ 7.

Similarly, we did not observe significant relation to any other clinical or pathological parameters. However, a non-significant trend toward inverse relation between moderate/high CRISP3 protein expression and higher pathological stage was noted. Herein, 19/68 (27%) moderate/high CRISP3 protein expression were noted in pT3 vs. 54/135 (40%) in pT2 (p = 0.09). CRISP3 protein was not related to disease free survival as measured by post serum PSA elevation of ≥ 0.2 ng/ml (Figure 2). Table 1 illustrates patients' demographics in relation to CRISP3 protein expression.

CRISP3 protein expression in relation to ERG expression and PTEN genomic deletions

To assess the association between CRISP3 protein and *ERG* expression, 812 cores of localized prostate cancer were assessed for the dual expression of both markers. There was significant association between higher CRISP3 protein expression and *ERG* expression. Moderate/high CRISP3 protein expression was present in 217/316 (69%) in *ERG* positive vs. 287/496 (58%) in *ERG* negative tumor cores (p = 0.002) (Figure 3A).

Since it was previously documented that *ERG* genomic rearrangements are enriched for *PTEN* deletions [20], we sought to investigate the association of CRISP3 with *PTEN* genomic deletions based on the type of *PTEN* deletion present within the individual core sample. Overall, there was significant association between CRISP3 protein expression and *PTEN* deletions (71% vs. 58%) (p = 0.01). To further investigate the relation of the type of *PTEN* genomic deletions and CRISP3 protein expression, we grouped the samples into *PTEN* intact, hemizygous *PTEN* deletions and homozygous *PTEN* deletions. In this cohort, higher CRISP3 protein expression was



significantly and positively associated with the degree of *PTEN* genomic aberrations (not deleted vs. hemizygous vs. homozygous deletions) (Table 2). Moderate/high CRISP3 protein expression was present in 279/477 (58%) *PTEN* intact tumors vs. 92/136 (68%) *PTEN* hemizygous deletions vs. 40/52 (77%) in *PTEN* homozygous deleted PCA ($P = 0.01$) (Figure 3B).

CRISP3 expression in PCA Cell lines

To validate CRISP3 mRNA expression relative to *ERG* status, we investigated the mRNA expression of CRISP3 and *ERG* in multiple cell lines from the MSKCC data (GSE21032). Both CRISP3 and *ERG* showed two-fold increase in VCaP cell line which harbors *TMPRSS2-ERG* rearrangement (Figure 3C). Using VCaP cell line to validate whether CRISP3 mRNA expression is a direct target of the *ERG* transcription factor, Ribeiro et al. detected three putative ETS-binding-sites containing-regions of the CRISP3 promoter in the *ERG*-bound chromatin, supporting direct link between *ERG* and CRISP3 [13].

CRISP3 expression in independent gene expression data of the MSKCC and Swedish prostate cancer cohorts

We next characterized the expression of CRISP3 with prostate cancer subtypes and clinical variables at the mRNA level. We used two public mRNA gene expression data MSKCC (GSE21032) and the Swedish cohort (GSE16560) and used Wilcox test to assess the statistical significance of the association between CRISP3 and other clinical variables. Analyzing the mRNA expression of CRISP3 in normal and PCA showed that CRISP3 mRNA expression is significantly upregulated in PCA ($p = 0.0053$) and metastasis ($p = 4.4e-5$) (Figure 3A). CRISP3 mRNA expression was also significantly associated with *ERG* status as determined by mRNA *ERG* expression ($p = 2.9e-9$) and *PTEN* deletions as determined by mRNA *PTEN* expression ($p = 0.053$) (Figure 3B,C) in the MSKCC data, and significantly associated with *ERG* gene rearrangements in Swedish cohort determined by FISH ($p = 2.5e-7$) (Figure 4A).

Investigating relation to Gleason score and grouping cases into <7 vs. ≥ 7 , high CRISP3 mRNA expression was associated with high GS in both MSKCC ($p = 0.022$)




Table 1 Patients demographics of the localized cohort stratified based on CRISP3 expression

Parameter	Negative/weak (n = 137)	Moderate/high (n = 78)	p-value
Age (years) (mean; range)	64.53 (43–81)	63.58 (47–78)	0.280
Pre-PSA level (ng/ml)			0.212
= < 10	43 (68%)	28 (80%)	
> 10	20 (32%)	7 (20%)	
Gleason summary			0.750
< 7	43 (32%)	25 (33%)	
3 + 4	43 (32%)	24 (31%)	
4 + 3	31 (22%)	20 (26%)	
> 7	19 (14%)	8 (10%)	
pT-stage			0.091
pT2	81 (62%)	54 (74%)	
pT3	49 (38%)	19 (26%)	
Surgical margin			0.285
Negative	69 (53%)	45 (61%)	
Positive	61 (47%)	29 (39%)	




Figure 3 Expression of CRISP3 mRNA in MSKCC data. **A.** Association of CRISP3 to prostate cancer progression, **B.** CRISP3 association to *ERG*, **C.** CRISP3 association to *PTEN*. **D.** Association of CRISP3 to Gleason score. **E.** Kaplan Meier of CRISP3 mRNA expression to time till biochemical relapse in MSKCC Cohort.

(Figure 3D) and the Swedish ($p = 1e^{-4}$) (Figure 4B). High CRISP3 mRNA expression, determined by PAM clustering, also was associated with time to biochemical recurrence (BCR) after Radical prostatectomy ($p = 0.038$, HR: 1.15) (Figure 3E), but not BCR event (data not shown). However, when investigating association to lethal disease, high CRISP3 mRNA expression was associated with time to death ($p = 0.0066$) (Figure 4C) in all samples of the Swedish cohort and the subsamples in Gleason score 7 ($p = 0.029$) (Figure 4D). Additional analysis supported the implications of high CRISP3 mRNA

expression in prostate cancer lethality ($p = 0.0086$) (Figure 4E).


The biological pathways of genes deregulated in PCA subset with *ERG* positive and high CRISP3 mRNA expression

To analyze the cross-talk between *ERG* and CRISP3, we identified the differentially expressed genes from MSKCC data between *ERG* positive vs. *ERG* negative, high CRISP3 mRNA expression vs. low CRISP3 mRNA expression and *ERG* positive/ high CRISP3 mRNA expression vs. others

Table 2 Association between CRISP3 immunostaining and *PTEN* intact, hemizygous *PTEN* deletions and homozygous *PTEN* deletions

	Negative/weak CRISP 3 expression n (%)	Moderate/high CRISP 3 expression n (%)	Total (n)	p-value
<i>PTEN</i> not deleted (n)	304(50%)	310(50%)	614	
<i>PTEN</i> hemizygous deletion (n)	53(35%)	97(65%)	150	<0.0001
<i>PTEN</i> homozygous deletion (n)	13(24%)	42(76%)	55	
Total (n)	370(45%)	449(55%)	819	

Cases assessed as individual TMA cores.



(Figure 5A). To select differentially expressed genes, we used student *t*-test, adjusting for multiple corrections using bonferroni method. Herein, we identified a group of 12 genes that are implicated in both CRISP3 and *ERG* subgroups. These 12 genes are highly correlated and interacting with AKT1, TP53, CTNBN1 and FOS network of genes extracted from the FI reactome functional protein networks implemented in Cytoscape (Figure 5B) suggesting a critical role of CRISP3 in cell proliferation [21].


Expression of *ERG* and *CRISP3* mRNA expression is significantly correlated. Low CRISP3 mRNA expression was associated with *ERG* negative, while high CRISP3 mRNA expression was associated with *ERG* positive. However, 50% of *ERG* positive samples has low CRISP3 mRNA expression, suggesting that CRISP3 expression is conditionally dependent on the expression of *ERG* or CRISP3 overexpression is a consequence event of *ERG* rearrangement.

We further investigated the functional biology of the 84 genes that are implicated in the *ERG* positive –high CRISP3 PCA subtype. These 84 genes were differentially expressed in only samples with high CRISP3 mRNA

expression and *ERG* positive samples, and define a distinct molecular profile of samples. SPINK1 is found to be differentially expressed in the *ERG*-positive-high CRISP3 PCA subtype. Using Gene set enrichment analysis [22], the 84 genes were enriched in ribosomal proteins ($p = 4.1e-4$) and translational elongation process ($p = 0.018$) (Table 3). Using network-based enrichment analysis tool (<http://www.enrichnet.org/>) to characterize the biological pathways governing these genes, phenylalanine metabolism was the most significant pathway that governs the 84 genes. This depicts the molecular mechanism underlying the combination of *ERG* and CRISP3 regulation.

Potential clinical utility of combining *ERG*, CRISP3 and PTEN

To confirm the significant association of CRISP3, *ERG* and PTEN, we assessed the prognostic value of combining *ERG*, PTEN and CRISP3 expression. The mRNA expression levels of the three genes were extracted from the Swedish cohort and hierarchical clustering was used to group all samples based on the three genes' expression. Clustering



confirmed that samples enriched with high ERG, high CRISP3 and low PTEN have marginally poor outcome ($p = 0.053$). We then focused on patients with low risk tumors; with Gleason score 6 or (3 + 4) to investigate whether this signature is able to stratify patients within this prognostic ambiguous group (Figure 5C), patients with the signature of high ERG, high CRISP3 and low PTEN were

Table 3 Top deregulated genes in high CRISP3 expression, ERG positive samples and the pathways they are involved in

Pathway	Top genes in the 84 set
mTORC mediated signaling	EIF4B
Unwinding of DNA	GIN52
Steroid hormone biosynthesis	HSD17B3
Peptide chain elongation	RPL4, RPL5, RPS3, RPL17
Protein translation and ribosomes	EIF4B, RPL4, RPL5, RPS3, RPL17
Deadenylation of mRNA	EIF4B
Signaling by BMP	BMPRI1B

significantly associated with poorest clinical outcome ($p = 0.0057$) (Figure 5D).

Discussion

CRISP3 belongs to a large family of cysteine –rich secretory proteins which are poorly characterized extracellular proteins [14]. This family is highly expressed in salivary glands and in the male reproductive tract, most of which are under strong androgen-dependency [13]. In humans, CRISP3 was first described in neutrophils, but later it was shown to be highly expressed in several exocrine glands, including salivary glands, pancreas and prostate [2-6] and within different cells of the immune system including pre-B cells, neutrophils, and eosinophils [23,24].

CRISP3 has been previously reported to be overexpressed in prostate cancer [7,9,25]. Using in situ hybridization, Kosari et al. reported that CRISP3 mRNA expression is epithelial-specific and is up-regulated in prostate cancer compared with benign

prostate tissue [2]. Ernst et al. found that majority of *CRISP3* production came from the prostatic epithelium, rather than the surrounding stromal tissue [9]. Others noted that *CRISP3* expression increases more than 50 fold in prostate cancer [7]. In our immunohistochemistry study we document that *CRISP3* protein levels are highly expressed in all neoplastic prostate epithelium compared with benign prostate tissue. We also report for the first time that *CRISP3* protein expression is highest in the CRPC category. This suggests a key role of *CRISP3* in AR-independent mediated transcriptional program.

The association between *CRISP3* and clinico-pathological parameters has been investigated in several previous publications, but with mixed results. Table 4. Summarizes the results of *CRISP3* protein expression in correlation with clinico-pathological parameters from five previous studies in addition to our own [10-14]. Bjartell et al. [26] found increased *CRISP3* protein expression in higher grade PCA (Gleason 4 and 5). However in a subsequent study by the same author, no significant difference in GS between patients with positive and negative *CRISP3* was noted [12]. The association between GS and *CRISP3* protein expression was only evident in one more study by Grupp et al., but was not demonstrated in three other. In our study, we were not able to document any association between *CRISP3* and GS at the protein level; however, at the mRNA levels we document significant positive association with higher GS. One possible explanation to this discrepancy could be related to the inclusion of adjacent stromal cells in the mRNA expression assessment (Taylor et al.) where stromal cells had indeed expressed lower levels of *CRISP3*, though the authors dissected the samples carefully to ensure minimal contamination of tumor samples with non-neoplastic tissue. Moreover, the relation of *CRISP3* to other pathological parameters, was only documented in one study (pathological stage and surgical margin) with all other 4 studies including ours failing to show such association. These data suggest either that

morphological assessment of *CRISP3* protein expression is inaccurate due to potential heterogeneity or protein abundance or that *CRISP3* is not actually related to GS and other pathological parameters and hence provide added value beyond those of current pathological parameters. The latter is more likely given that *CRISP3* mRNA expression was significantly associated to time of biochemical relapse and lethal disease in the MSKCC and Swedish cohorts.

CRISP3 has been reported to be one of the top genes that are linked to *TMPRSS2-ERG* positive prostate cancer with about 53-fold increase when compared with fusion negative prostate cancer [13]. Our study confirms this strong association.

It is well known that up to 40% of human prostate cancers show loss of at least one copy of the *PTEN* gene via submicroscopic deletion [27-29] and that the percent of tumors harboring *PTEN* deletions increases with disease progression and in advanced castration resistant disease [19,20]. Moreover, the complexity of *PTEN* deletions increases with higher Gleason score and is tightly associated with *ERG* rearrangement [8,20,28-31]. In this study, we carried out a detailed investigation between *CRISP3* protein expression and the type of *PTEN* deletions which was only partially addressed in the study by Grupp et al. [14], where they documented significant association between *PTEN* deletions and *CRISP3* protein expression. However, the relation between the type of *PTEN* deletion and *CRISP3* protein expression was not addressed. Herein, we confirm significant increase in *CRISP3* protein expression not only in *PTEN* deleted tumors, but also document that this expression is also connected and increases with the degree and severity of *PTEN* deletion (hemizygous vs. homozygous) (65% vs. 76%) respectively.

We also investigated the potential linkage between *CRISP3* mRNA expression and *ERG* gene rearrangements and identified a 12-gene signature that is implicated in both *CRISP3* and *ERG* subgroup of PCA. We also investigated the functional biology of an 84 genes

Table 4 Results of *CRISP3* expression in correlation with clinico-pathological parameters from comparatives studies

Study	Correlation of <i>CRISP3</i> expression with clinico-pathological parameters			Outcome
	Gleason score	Pathological stage	Surgical margins	
Ribeiro et al. [13]	No	No †	Not Tested	N/A
Grupp et al. [14]	Yes	Yes	Yes	Univariate
Dahlman et al. [10]	No?	No?	No?	Trends*
Hoogland et al. [11]	No	No	No	N/A
Bjartell et al. [12]	No	No	No	Yes [‡]
Al-Bashir et al.	No	No	No	No [‡]

*There was a trend that patients with high *CRISP3* expression had higher risk of biochemical recurrence.

[‡]Adding *CRISP3* expression to current parameters does not provide any added significance to the prediction.

[†]Using non-parametric tests on the qRT-PCR data from the validation series.

[‡]Based on protein expression. However, mRNA *CRISP3* levels were prognostic based on the validation within Taylor and Swedish cohorts.

that are implicated in the *ERG* positive –high *CRISP3* mRNA expression PCA subtype, documenting phenylalanine metabolism as the most significant and likely pathway that governs these genes. This correlation suggests that *CRISP3* is a major target of *ERG* that is strongly overexpressed in PCA.

Although our study did not provide any prognostic implication of *CRISP3* expression in prostate cancer at the protein level, we were able to confirm significant prognostic value for *CRISP3*, in the MSKCC and Swedish cohorts. This may be due to heterogeneity or the methods used to assess for protein expression and highlight the need for more accurate measurement of protein expression, likely through the use of image analysis systems. The tight link between *CRISP3* expression, *ERG* gene rearrangements and/ or *PTEN* deletions in the three cohorts of prostate cancer studied, suggest the presence of distinct molecular alterations in this subgroup of tumors that may have additional clinical implication if assessed collectively.

In this study, we validated the prognostic value of the combined status of the three genes by clustering the samples in the Swedish data into multiple clusters. The clustering showed that the status of the three genes can stratify patients into distinct subgroups with distinct outcome. The cluster that is enriched with high *ERG*, high *CRISP3* and low *PTEN* demonstrated the poorest survival outcome and this was evident in the group of patients with GS6 & (3 + 4), which is the most group likely to benefit of signature biomarker. This finding is novel and has not been shown in any previous study, including the study by Grupp et al. in which association of *CRISP3* was noted with *ERG* and *PTEN*, but it failed to document any added prognostic value for the three markers combined [14].

In Summary, in this study we confirmed significant association between *CRISP3* expression and prostate cancer progression as well as a strong link between *CRISP3* expression and each of *ERG* expression and *PTEN* genomic deletions. In addition, we documented and detailed potential genes that are tightly connecting this molecular subtype of PCA. More importantly, we confirm and for the first time, that tumors defined by *CRISP3*, *ERG* and *PTEN* signify a molecular subtype of PCA with the poorest clinical outcome. This suggests that the combination of the three genes could be used as a molecular test to stratify patients' risk. Further studies to elucidate the molecular network connecting *CRISP3*, *ERG* and *PTEN* and to validate their combined prognostic role may be of added value in stratifying patients with prostate cancer, specifically those with Gleason score 6&7.

Competing interests

M. A is employer of GenomeDx Bioscience Inc. The other authors declare that they have no competing interests.

Authors' contributions

SA carried out immunostaining, pathological analysis and wrote initial manuscript. MA performed the bioinformatics analysis. SH carried out pathological analysis. MD performed statistical and survival analysis. BD designed the study. TB designed the study, carried out pathological analysis and wrote the initial manuscript. All authors read and approved the final manuscript.

Acknowledgement

This work was supported by the Young Investigator Award of the Prostate Cancer Foundation, USA, Prostate Cancer Canada, and is proudly funded by the Movember foundation, Grant #132013-01.

Author details

¹Department of Pathology and Laboratory Medicine, University of Calgary and Calgary Laboratory Services, Calgary, AB, Canada. ²Departments of Oncology, Biochemistry and Molecular Biology, Calgary, AB, Canada. ³Southern Alberta Cancer Institute and Tom Baker Cancer Center, Calgary, AB, Canada. ⁴The Prostate Cancer Center, Calgary, AB, Canada. ⁵Department of Urology, University of Calgary, Calgary, AB, Canada. ⁶Department of Pathology and Laboratory Medicine, Jordan University of Science and Technology, Irbid, Jordan.

Received: 16 January 2014 Accepted: 3 March 2014

Published: 7 March 2014

References

1. Hsing AW, Tsao L, Devesa SS: **International trends and patterns of prostate cancer incidence and mortality.** *Int J Cancer* 2000, **85**(1):60–67.
2. Carswell BM, Woda BA, Wang X, Li C, Dresser K, Jiang Z: **Detection of prostate cancer by alpha-methylacyl CoA racemase (P504S) in needle biopsy specimens previously reported as negative for malignancy.** *Histopathology* 2006, **48**(6):668–673.
3. Lu G, Villalba M, Coscia MR, Hoffman DR, King TP: **Sequence analysis and antigenic cross-reactivity of a venom allergen, antigen 5, from hornets, wasps, and yellow jackets.** *J Immunol* 1993, **150**(7):2823–2830.
4. Fritig B, Heitz T, Legrand M: **Antimicrobial proteins in induced plant defense.** *Curr Opin Immunol* 1998, **10**(1):16–22.
5. Schuren FH, Asgeirsdottir SA, Kothe EM, Scheer JM, Wessels JG: **The Sc7/Sc14 gene family of Schizophyllum commune codes for extracellular proteins specifically expressed during fruit-body formation.** *J Gen Microbiol* 1993, **139**(9):2083–2090.
6. Miosga T, Schaaff-Gerstenschlager I, Chalwatzis N, Baur A, Boles E, Fournier C, Schmitt S, Velten C, Wilhelm N, Zimmermann FK: **Sequence analysis of a 33.1 kb fragment from the left arm of Saccharomyces cerevisiae chromosome X, including putative proteins with leucine zippers, a fungal Zn(II) 2-Cys6 binuclear cluster domain and a putative alpha 2-SCB-alpha 2 binding site.** *Yeast* 1995, **11**(7):681–689.
7. Asmann YW, Kosari F, Wang K, Cheville JC, Vasmatzis G: **Identification of differentially expressed genes in normal and malignant prostate by electronic profiling of expressed sequence tags.** *Cancer Res* 2002, **62**(11):3308–3314.
8. Han B, Mehra R, Lonigro RJ, Wang L, Suleman K, Menon A, Palanisamy N, Tomlins SA, Chinnaiyan AM, Shah RB: **Fluorescence in situ hybridization study shows association of PTEN deletion with ERG rearrangement during prostate cancer progression.** *Mod Pathol* 2009, **22**(8):1083–1093.
9. Ernst T, Hergenbahn M, Kenzelmann M, Cohen CD, Bonrouhi M, Weninger A, Klaren R, Grone EF, Wiesel M, Gudemann C, Küster J, Schott W, Staehler G, Kretzler M, Hollstein M, Gröne HJ: **Decrease and gain of gene expression are equally discriminatory markers for prostate carcinoma: a gene expression analysis on total and microdissected prostate tissue.** *Am J Pathol* 2002, **160**(6):2169–2180.
10. Dahlman A, Rexhepaj E, Brennan DJ, Gallagher WM, Gaber A, Lindgren A, Jirstrom K, Bjartell A: **Evaluation of the prognostic significance of MSMB and CRISP3 in prostate cancer using automated image analysis.** *Mod Pathol* 2011, **24**(5):708–719.
11. Hoogland AM, Dahlman A, Vissers KJ, Wolters T, Schroder FH, Roobol MJ, Bjartell AS, van Leenders GJ: **Cysteine-rich secretory protein 3 and beta-microseminoprotein on prostate cancer needle biopsies do not have predictive value for subsequent prostatectomy outcome.** *BJU Int* 2011, **108**(8):1356–1362.

12. Bjartell AS, Al-Ahmadie H, Serio AM, Eastham JA, Eggener SE, Fine SW, Udby L, Gerald WL, Vickers AJ, Lilja H, Reuter VE, Scardino PT: **Association of cysteine-rich secretory protein 3 and beta-microseminoprotein with outcome after radical prostatectomy.** *Clin Cancer Res* 2007, **13**(14):4130–4138.
13. Ribeiro FR, Paulo P, Costa VL, Barros-Silva JD, Ramalho-Carvalho J, Jeronimo C, Henrique R, Lind GE, Skotheim RI, Lothe RA, Teixeira MR: **Cysteine-rich secretory protein-3 (CRISP3) is strongly Up-regulated in prostate carcinomas with the TMPRSS2-ERG fusion gene.** *PLoS One* 2011, **6**(7):e22317.
14. Grupp K, Kohl S, Sirma H, Simon R, Steurer S, Becker A, Adam M, Izbicki J, Sauter G, Minner S, Schlomm T, Tsourlakis MC: **Cysteine-rich secretory protein 3 overexpression is linked to a subset of PTEN-deleted ERG fusion-positive prostate cancers with early biochemical recurrence.** *Mod Pathol* 2012, **26**(5):733–742.
15. Darnel AD, Behmoaram E, Vollmer RT, Corcos J, Bijian K, Sircar K, Su J, Jiao J, Alaoui-Jamali MA, Bismar TA: **Fascin regulates prostate cancer cell invasion and is associated with metastasis and biochemical failure in prostate cancer.** *Clin Cancer Res* 2009, **15**(4):1376–1383.
16. Brimo F, Vollmer RT, Friszt M, Corcos J, Bismar TA: **Syndecan-1 expression in prostate cancer and its value as biomarker for disease progression.** *BJU Int* 2010, **106**(3):418–423.
17. Taylor BS, Schultz N, Hieronymus H, Gopalan A, Xiao Y, Carver BS, Arora VK, Kaushik P, Cerami E, Reva B, Antipin Y, Mitsiades N, Landers T, Dolgalev I, Major JE, Wilson M, Socci ND, Lash AE, Heguy A, Eastham JA, Scher HI, Reuter VE, Scardino PT, Sander C, Sawyers CL, Gerald WL: **Integrative genomic profiling of human prostate cancer.** *Cancer cell* 2010, **18**(1):11–22.
18. Epstein JI, Allsbrook WC Jr, Amin MB, Egevad LL: **The 2005 International Society of Urological Pathology (ISUP) consensus conference on gleason grading of prostatic carcinoma.** *Am J Surg Pathol* 2005, **29**(9):1228–1242.
19. Bismar TA, Yoshimoto M, Duan Q, Liu S, Sircar K, Squire JA: **Interactions and relationships of PTEN, ERG, SPINK1 and AR in castration-resistant prostate cancer.** *Histopathology* 2012, **60**(4):645–652.
20. Bismar TA, Yoshimoto M, Vollmer RT, Duan Q, Friszt M, Corcos J, Squire JA: **PTEN genomic deletion is an early event associated with ERG gene rearrangements in prostate cancer.** *BJU Int* 2011, **107**(3):477–485.
21. Saito R, Smoot ME, Ono K, Ruschinski J, Wang PL, Lotia S, Pico AR, Bader GD, Ideker T: **A travel guide to Cytoscape plugins.** *Nat Methods* 2012, **9**(11):1069–1076.
22. da Huang W, Sherman BT, Tan Q, Collins JR, Alvord WG, Roayaei J, Stephens R, Baseler MW, Lane HC, Lempicki RA: **The DAVID Gene Functional Classification Tool: a novel biological module-centric algorithm to functionally analyze large gene lists.** *Genome Biol* 2007, **8**(9):R183.
23. Udby L, Calafat J, Sorensen OE, Borregaard N, Kjeldsen L: **Identification of human cysteine-rich secretory protein 3 (CRISP-3) as a matrix protein in a subset of peroxidase-negative granules of neutrophils and in the granules of eosinophils.** *J Leukoc Biol* 2002, **72**(3):462–469.
24. Udby L, Cowland JB, Johnsen AH, Sorensen OE, Borregaard N, Kjeldsen L: **An ELISA for SGP28/CRISP-3, a cysteine-rich secretory protein in human neutrophils, plasma, and exocrine secretions.** *J Immunol Methods* 2002, **263**(1–2):43–55.
25. Kratzschmar J, Haendler B, Eberspaecher U, Roosterman D, Donner P, Schleuning WD: **The human cysteine-rich secretory protein (CRISP) family. Primary structure and tissue distribution of CRISP-1, CRISP-2 and CRISP-3.** *Eur J Biochem* 1996, **236**(3):827–836.
26. Bjartell A, Johansson R, Bjork T, Gadaleanu V, Lundwall A, Lilja H, Kjeldsen L, Udby L: **Immunohistochemical detection of cysteine-rich secretory protein 3 in tissue and in serum from men with cancer or benign enlargement of the prostate gland.** *Prostate* 2006, **66**(6):591–603.
27. Yoshimoto M, Cunha IW, Coudry RA, Fonseca FP, Torres CH, Soares FA, Squire JA: **FISH analysis of 107 prostate cancers shows that PTEN genomic deletion is associated with poor clinical outcome.** *Br J Cancer* 2007, **97**(5):678–685.
28. Yoshimoto M, Joshua AM, Cunha IW, Coudry RA, Fonseca FP, Ludkovski O, Zielenska M, Soares FA, Squire JA: **Absence of TMPRSS2:ERG fusions and PTEN losses in prostate cancer is associated with a favorable outcome.** *Mod Pathol* 2008, **21**(12):1451–1460.
29. Reid AH, Attard G, Ambroisine L, Fisher G, Kovacs G, Brewer D, Clark J, Flohr P, Edwards S, Berney DM, Foster CS, Fletcher A, Gerald WL, Møller H, Reuter VE, Scardino PT, Cuzick J, de Bono JS, Cooper CS: **Molecular characterisation of ERG, ETV1 and PTEN gene loci identifies patients at low and high risk of death from prostate cancer.** *Br J Cancer* 2010, **102**(4):678–684.
30. Attard G, Swennenhuis JF, Olmos D, Reid AH, Vickers E, A'Hern R, Levink R, Coumans F, Moreira J, Riisnaes R, Oommen NB, Hawche G, Jameson C, Thompson E, Sipkema R, Carden CP, Parker C, Dearnaley D, Kaye SB, Cooper CS, Molina A, Cox ME, Terstappen LW, de Bono JS: **Characterization of ERG, AR and PTEN gene status in circulating tumor cells from patients with castration-resistant prostate cancer.** *Cancer Res* 2009, **69**(7):2912–2918.
31. Gopalan A, Leversha MA, Satagopan JM, Zhou Q, Al-Ahmadie HA, Fine SW, Eastham JA, Scardino PT, Scher HI, Tickoo SK, Reuter VE, Gerald WL: **TMPRSS2-ERG gene fusion is not associated with outcome in patients treated by prostatectomy.** *Cancer Res* 2009, **69**(4):1400–1406.

doi:10.1186/1756-8722-7-21

Cite this article as: Al Bashir *et al.*: Cysteine-rich secretory protein 3 (CRISP3), ERG and PTEN define a molecular subtype of prostate cancer with implication to patients' prognosis. *Journal of Hematology & Oncology* 2014 **7**:21.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

