

RESEARCH

Open Access

# Activation tagging in *Salvia miltiorrhiza* can cause increased leaf size and accumulation of tanshinone I and IIA in its roots

Hsin-Shueh Ho<sup>1</sup>, Rishi Kishore Vishwakarma<sup>1</sup>, Emily Chin-Fun Chen<sup>1</sup> and Hsin-Sheng Tsay<sup>1,2\*</sup>

## Abstract

**Background:** *Salvia miltiorrhiza* Bunge (Danshen), an important herb in traditional Chinese medicine, is commonly used for treatment of cardiovascular diseases. One of the major bioactive constituents of Danshen, diterpenoid tanshinone, has been proved with pharmacological properties and have the potential to be a new drug candidate against various diseases. In our previous study, we have established an activation tagging mutagenesis (ATM) population of callus lines of *S. miltiorrhiza* Bunge by *Agrobacterium*- mediated transformation.

**Results:** In the present study, we have identified ATM transgenic *Salvia* plant (SH41) with different leaf morphology and more tanshinones in its roots. The transgenic background of SH41 was identified by PCR (using *hptII* primers) and Southern blots. PCR analysis showed a single band of *hptII* gene and Southern blot analysis showed single insertion in SH41. External appearance of ATM transgenic SH41 was observed with broader leaves comparing to non-transformed plants. More healthy trichomes as well as bigger and wobbly guard cells and stomata were observed in SH41 by scanning electron microscopy (SEM). Quantitative analysis of active compounds in SH41 roots revealed a significant increase in tanshinone I (3.7 fold) and tanshinone IIA (2 fold) contents as compared to the wild plant.

**Conclusions:** We have generated an activation tagged transgenic *Salvia* plant (SH41) with different leaf morphology and high diterpenes content in its roots. The increased amount of tanshinones in SH41 will definitely offer a route for maximizing the benefits of this plant in traditional Chinese herbal medicines. The present report may also facilitate the application of ATM for genetic manipulation of other medicinal crops and subsequent improved metabolite contents.

**Keywords:** Activation-tagging; HPLC; Tanshinones; *Salvia miltiorrhiza*; Scanning electron microscopy

## Background

*Salvia miltiorrhiza* Bunge (Danshen) is a recognized traditional Chinese medicine and is broadly used for the treatment of various cardiovascular and cerebrovascular diseases. This wide range of medicinal uses is mainly due to its property of stimulating blood circulation and removing blood stasis (Zhou et al. 2012). Danshen contained water soluble phenolics including danshensu, protocatechuic aldehyde, protocatechuic acid, caffeic acid, rosmarinic acid and salvianolic acid A & B as well as lipid soluble diterpene quinones: cryptotanshinone, tanshinone I and tanshinone IIA (Liu et al. 2007). Phenolics and quinones have alike

individual pharmacological properties such as antioxidant, anti-apoptosis and vasodilation (Zhou et al. 2005; Lam et al. 2007; Lam et al. 2008a; Wang et al. 2011a).

Many experimental and clinical investigations have reported that tanshinones and their derivatives can prevent or slow the development of a various diseases including hypoxia/reoxygenation injury, cardiovascular diseases, cancer, neonatal hypoxic ischemic encephalopathy, hepatic fibrosis as well as neuro degenerative diseases (Wu et al. 1993; Yagi et al. 1994; Yoon et al. 1999; Takahashi et al. 2002; Han et al. 2008; Shu et al. 2010).

Due to its extensive medicinal values, *S. miltiorrhiza* has been studied by various research groups over the world, especially in biotechnological and *in vitro* propagation field. Through several strategies like genetic manipulation of biosynthetic pathway via genetic transformation (Chen et al. 1997; Lee et al. 2008; Kai et al. 2011), hairy

\* Correspondence: hstsay@cyut.edu.tw

<sup>1</sup>Department of Applied Chemistry, Chaoyang University of Technology, 168, Jifong E Road, Taichung, Wufong 41349, Taiwan

<sup>2</sup>Department of Agronomy, National Chung Hsing University, Taichung, Taiwan

root cultures (Hu et al. 1993), use of plant growth regulators and elicitors (Wu et al. 2003; Zhang et al., 2004; Ge and Wu 2005a, 2005b; Gupta et al. 2011), success has been reported in enhancing the production of secondary metabolites in *S. miltiorrhiza*.

Although, several reports represented the micropropagation and genetic transformation methods for *S. miltiorrhiza* by other means (Zhang et al. 1995; Chen et al. 1997; Zhang et al. 1997; Yan and Wang 2007; Lee et al. 2008), but there is a limited report available on activation tagging mutagenesis. Therefore, the overall objective of the current research was to develop and investigate the effect of ATM on morphological as well as histological changes and tanshinones content in transgenic *Salvia* plant.

## Methods

### Plant material

*S. miltiorrhiza* used for this study was collected from Zhengzhou City, Henan Province Jiyuan County mountains, China. Activation tagged mutagenic (ATM) *Salvia* plants are maintained at greenhouse in Chaoyang University of Technology.

### Gene constructs and bacterial strain

*A. tumefaciens* strain EHA105 harboring a binary vector pTAG8 (Hsing et al. 2007; Chen et al. 2009; Tsay et al. 2012) was used for the genetic transformation of *S. miltiorrhiza*. The T-DNA region of the binary vector contained a selectable marker coding for the hygromycin phosphotransferase (*hptII*) gene under control of CaMV 35S promoter (provided by Dr. Su-May Yu, Academia Sinica, Taipei, Taiwan).

### Genetic transformation and confirmation of ATM insertion in *Salvia* plant

*Salvia* transgenic raised by Lee et al. (2008) was used for this study. ATM transgenic *S. miltiorrhiza* plant (SH41) with an altered morphological appearance was investigated. The transformation was confirmed by PCR detection of hygromycin phosphotransferase gene (*hptII*) gene using genomic DNA and gene specific primers *hptII* F 5'-GTCGTGGCGATCCTGCAAGC-3' and *hptII* R 5'-CC TGCGGGTAAATAGCTGCGC-3'. Total genomic DNA was isolated from young leaves of SH41 and control plants using ZR plant/seed DNA MiniPrep kit (Zymo Research). The PCR reaction contained 50 ng genomic DNA, 0.2 mM dNTPs mix, 500 nM each forward and reverse primers, and 1U of Taq DNA polymerase. The PCR was carried out in a thermal cycler (BIO-RAD, USA) under the following conditions: 1 cycle of 94°C (5 min), 35 cycles of 94°C (30 s), 55°C (30 s) and 72°C (1 min), and final extension at 72°C (7 min). The plasmid pTAG8 was used as a positive control. The amplification products were analyzed on 1%

agarose gel by electrophoresis and visualized under UV light after ethidium bromide staining.

The insertion of the gene was further confirmed by Southern blot analysis. Genomic DNA of SH41 and control plants was isolated using the modified cetyl trimethyl ammonium bromide (CTAB) method (Doyle and Doyle 1987). Isolated genomic DNA (15 µg) was digested overnight with restriction enzymes *Pst*I, *Hind*III and *Eco*RI in appropriate buffer at 37°C. The digested DNA was electrophoresed on 0.8% agarose gel in 1X Tris-acetate/EDTA buffer. The gel was then treated with dupurination, denaturation and neutralization buffers and DNA was blotted onto a positively charged nylon membrane (Bio-Rad) as described by Sambrook et al. (1989). A PCR-amplified *hptII* gene fragment was labelled with digoxigenin-9-dUTP using Dig High Prime Labeling and Detection Starter Kit I (Roche, USA) and used as a probe. After prehybridization for 2 h at 52°C, labeled probe was added and kept for hybridization at the same temperature for 16 h in hybridization chamber. The membrane was subjected to a series of stringency washes (high stringency at 56°C) to eliminate nonspecific probe binding. The signals were immuno-detected with anti-digoxigenin-AP and visualized with the colorimetric substrates BCIP/NBT according to manufacturer's instruction.

### Scanning electron microscopy (SEM) analysis

In order to investigate the differences observed in the morphology of the wild type (WT) and transgenic (SH41) *Salvia* plants, scanning electron microscopy (SEM) was performed. *Salvia* leaf and root samples were cut into an appropriate size, adhered to the metal base, frozen in liquid nitrogen and scanned in low-vacuum SEM (JEOL JSM-6330F) with chamber pressure of 30 Pa and an accelerated voltage 15 kV.

### Quantitative (HPLC) analysis

High performance liquid chromatography (HPLC) was performed to detect the water soluble metabolites danshensu, protocatechuic acid, protocatechuic aldehyde, caffeic acid, rosmarinic acid, salvianolic acid A and salvinic acid B, as well as lipid soluble tanshinone I, tanshinone IIA and cryptotanshinone in roots. All the standards were obtained from Sigma and Formosa Kingstone Bioproducts International Corp.

First, extraction was optimized using different solvent system with commercially available *Salvia* plant roots (CK; purchased from Joint Pharmacy, Taichung City, Taiwan). For further analysis, 50 mg dried roots of 4 months old transgenic (SH41) and wild type *Salvia* plants was finely ground by mortar pestle using liquid nitrogen and extracted with 10 ml of 60% ethanol containing 0.05% formic acid by sonication for 10 min along with and commercially available (CK) *Salvia* root

powder. The extracts were filtered through filter paper and vacuum dried. Further, the dried samples were dissolved in methanol/water (1:1), filtered through 0.22  $\mu$ m membrane and subjected to HPLC analysis. The HPLC system consisted of a pump (Hitachi L-2130), automatic injector (Hitachi L-2200) and Diode Array Detector (Hitachi L-2450). The column used was C18 (Waters; 5  $\mu$ m, 4.6  $\times$  250 mm) and UV detection was carried out at 280 nm. The mobile phase acetonitrile/water containing 0.05% H<sub>3</sub>PO<sub>4</sub> was pumped at a flow rate was 1.0 mL/min and programmed as follows: 10% acetonitrile for 0 min; 40% acetonitrile for 15 min; 60% acetonitrile for 15 min and 75% acetonitrile for 30 min. Dilutions (1.512, 3.125, 6.25, 12.5, 25, 50, 100 mg/mL) of standard compounds were prepared and 10  $\mu$ L aliquots were subjected to HPLC. This was repeated thrice and calibration plots were constructed from the peak areas of standard compounds.

#### Statistical analysis

Each result shown in tables was the mean of at least three replicated experiments and the standard deviation ( $\pm$ SD) value was calculated.

#### Results

##### Morphological investigation of activation tagged *Salvia* plant


As shown in Figure 1, we can clearly see the morphological difference between SH41 and wild type plant.

SH41 plant exhibited larger sized leaves as well as thick and smaller petiole than wild type (Figure 1).


The microscopic (SEM) analysis showed that activation tag mutagenesis affected the morphological development of leaf tissues during growth. Cross section of leaf blades showed thickening of the palisade and the spongy tissues were slightly larger in SH41 as compared to wild type (Figures 2A, 2B). No significant differences were observed in the morphology of roots (Figures 2C, 2D). Adaxial and abaxial surface of SH41 leaf showed more trichomes than WT with bigger and wobbly guard cells and stomata (Figures 2E, 2F, 2G, 2H). Two types of hairy assemblies were found in the microscopic observation of leaves, one similar to flagpole-shaped head (Figures 3A, 3B) and another bamboo shoot shaped (Figures 3C, 3D), the latter is mostly deposited in the blade surface in the SH41.

##### PCR identification and DNA blot analysis of ATM transgenic *Salvia*

PCR analysis with *hptII* gene specific primers was carried out to confirm the presence of the activation tagging T-DNA insertion in the genome of the putative transformant. The 650 bp expected *hptII* fragment was found in the positive control (pTAG8 plasmid) as well as in transformed *Salvia* SH41. No amplification was observed in wild type plant (control) (Figure 4A).



**Figure 1** *S. miltiorrhiza* plant morphology. (A) Wild type (WT)- above ground. (B) SH41- above ground. (C) WT- roots. (D) SH41- roots. Scale bar = 1 cm.




**Figure 2 SEM images showing morphological differences.** (A & B) Leaf blade cross-sections of WT and SH41 respectively. (C & D) Root cross sections of WT and SH41 respectively. (E & F) Adaxial leaf surface of WT and SH41 respectively. (G & H) Abaxial leaf surface of WT and SH41 respectively. (a & g) Trichomes, (b) Palisade tissue, (c) Spongy tissue, (d) Cork layer, (e) Cortex, (f) Formation of the layer, and (h) Guard cells.

Southern hybridization analysis was performed in order to estimate the number of inserted loci. Figure 4B showed the Southern hybridization pattern obtained with *hptII* probe. The result showed single insertion in SH41 and no band was observed in the non-transformed control plant (Figure 4B).


#### Compound extraction and HPLC analysis

For optimal extraction of active constituents from the *Salvia* plant, different solvent system was tested. The

addition of formic acid improved the extraction efficiency of salvianolic acid B and Danshensu. As the result, 60% ethanol with 0.05% formic acid was the best solvent system (Table 1). The tanshinone I and tanshinone IIA detected in SH41 was 3.7 and 2 fold higher than wild type plant, respectively. These compounds were also greater than CK (2 fold and 3 fold higher, respectively). Salvianolic acid B was also found 1.4 and 2.4 fold higher in SH41 as compared to wild plant and CK, respectively. Other compounds like Danshensu, rosmarinic acid and



**Figure 3 SEM image showing two forms of hairy structure (trichomes).** (A & B) Flagpole-shaped trichome of WT and SH41 respectively, and (C & D) Bamboo shoot shaped trichome of WT and SH41 respectively.



**Figure 4** Molecular analysis of ATM transformed *Salvia miltiorrhiza*. **(A)** PCR analysis of transformed *Salvia* plant. Genomic DNA was amplified using *hptII* gene specific primers. Approx. 650 bp amplification was observed in transformed plant. Lanes: (M) Molecular weight marker, (1) Positive control (pTAG8 plasmid), (2) Wild type *Salvia*; and (3) Active mutagenic *Salvia* plant (SH41). **(B)** Confirmation of activation-tagged *Salvia miltiorrhiza* Bunge by Southern blot analysis. Genomic DNA was extracted from leaves and *hptII* gene used as a probe. Hybridization was performed at 52°C. V- pTAG-8; SP- SH41 and WP- WT digested by *Pst* I; SH- SH41 and WH- WT digested by *Hind* III; SE- SH41 and WE- WT digested by *Eco*R I. Arrows indicate hybridization signals.

salvianolic acid A were slightly higher in SH41, however, salvianolic acid A was less in SH41 as compared to CK. On the other hand, protocatechuic, protocatechuic aldehyde and caffeic acid were not detected in SH41 (Table 2).

## Discussion

A new biotechnological method of mutagenesis, called activation tagging mutagenesis (ATM), has increased the understanding of development in various plants by intensifying the range of gene expressions and accumulation of secondary metabolites. This technique exploits the transferred DNA (T-DNA) from *Agrobacterium tumefaciens* to introduce a viral CaMV35S enhancer region randomly into the genome. The enhancer may dominantly “activate” and/or widen the pattern of expression of a gene that is near the enhancer of the introduced T-DNA (Hsing et al. 2007). It has been reported that the CaMV35S enhancer activate the expression of

genes located upstream or downstream of T-DNA in *Arabidopsis* and rice (Ichikawa et al. 2003; Jeong et al. 2006). The insertion of an enhancer sequence in the vicinity of an endogenous gene can alter the transcriptional pattern of the gene, resulting in a mutant phenotype. Activation tagging has been undertaken extensively in a number of dicot and monocot plants (Michael and Anthony 2011).

*Salvia miltiorrhiza* (Danshen) is an important Chinese medicinal herb and diterpene tanshinones are its major active compounds. In previous studies, *in vitro* methods such as hairy root cultures (Hu et al. 1993; Ge and Wu 2005a; Yan et al. 2005), cell suspension cultures (Chen et al. 1997), addition of elicitors and optimization of media (Wu et al. 2003; Zhang et al. 2004; Ge and Wu 2005b; Gupta et al. 2011), have been reported. In our previous study, ATM transgenic callus lines of *Salvia* with greater tanshinones content were developed (Lee

**Table 1** Different solvent extractions of *Salvia miltiorrhiza* root (commercially available plant)

Compounds	100% CH	70% CH	70% CH/F	100% Et	60% Et	60% Et/F
Danshensu	0.63 ± 0.06	1.05 ± 0.12	1.22 ± 0.02	0.36 ± 0.04	0.98 ± 0.04	1.30 ± 0.04
Protocatechuic	0.10 ± 0.01	0.13 ± 0.01	0.12 ± 0.01	0.09 ± 0.03	0.13 ± 0.13	0.14 ± 0.14
Protocatechuic aldehyde	0.06 ± 0.05	0.06 ± 0.01	0.06 ± 0.03	0.06 ± 0.02	0.06 ± 0.06	0.06 ± 0.06
Caffeic acid	0.06 ± 0.05	0.06 ± 0.02	0.06 ± 0.03	0.05 ± 0.02	0.06 ± 0.06	0.06 ± 0.06
Rosmarinic acid	0.65 ± 0.07	1.08 ± 0.08	1.20 ± 0.02	0.36 ± 0.03	0.96 ± 0.02	1.21 ± 0.04
Salvianolic acid B	3.35 ± 0.09	5.43 ± 0.04	5.97 ± 0.08	1.68 ± 0.08	4.87 ± 0.07	6.08 ± 0.06
Salvianolic acid A	0.25 ± 0.03	0.36 ± 0.04	0.39 ± 0.04	0.16 ± 0.01	0.33 ± 0.03	0.41 ± 0.02
Cryptotanshinone	0.03 ± 0.01	0.06 ± 0.02	0.07 ± 0.02	0.07 ± 0.03	0.06 ± 0.01	0.07 ± 0.02
Tanshinone I	0.08 ± 0.03	0.10 ± 0.04	0.11 ± 0.03	0.18 ± 0.04	0.07 ± 0.02	0.14 ± 0.05
Tanshinone IIA	0.04 ± 0.01	0.07 ± 0.02	0.08 ± 0.02	0.15 ± 0.04	0.07 ± 0.01	0.14 ± 0.03

CH, Methanol; CH/F, Methanol with 0.05% formic acid; Et, Ethanol; Et/F, Ethanol with 0.05% formic acid. Data expressed as mean ± SD for 3 repeats (mg/g).

**Table 2 HPLC-UV detection of *Salvia* roots extracted with 60% ethanol containing 0.05% formic acid for seven water-soluble and three fat-soluble ingredients, (n = 3) (mean ± SD; mg/g extract)**

Constituents	WTR	SHR	CK
Danshensu	0.19 ± 0.07	0.28 ± 0.08	1.30 ± 0.17
Protocatechuic	ND	ND	0.14 ± 0.04
Protocatechuic aldehyde	0.08 ± 0.01	ND	0.06 ± 0.01
Caffeic acid	ND	ND	0.06 ± 0.04
Rosmarinic acid	2.11 ± 0.21	2.87 ± 0.74	1.21 ± 0.47
Salvianolic acid B	10.44 ± 0.75	14.45 ± 0.37	6.08 ± 0.39
Salvianolic acid A	0.12 ± 0.07	0.16 ± 0.09	0.41 ± 0.04
Cryptotanshinone	0.59 ± 0.13	0.34 ± 0.08	0.07 ± 0.05
Tanshinone I	0.08 ± 0.01	0.30 ± 0.05	0.14 ± 0.06
Tanshinone IIA	0.20 ± 0.09	0.42 ± 0.09	0.14 ± 0.04

WTR Wild type root, SHR SH41 root, CK Commercially available *Salvia miltiorrhiza* root, ND Non detectable.

et al. 2008). In the present study, one of the 4 months old ATM transgenic (SH41) with an altered phenotype was identified and subjected to quantitative analysis of medicinally important ingredients. Insertion of T-DNA was confirmed by PCR with hygromycin phosphotransferase (*hptII*) gene specific primers and Southern blot analysis. PCR analysis showed amplification of *hptII* gene which confirmed the presence of an activation tagging T-DNA in SH41. DNA blot analysis showed single copy insertion of transgene in SH41. Single or low-copy integration of transgenes through *Agrobacterium* have been observed in several plant species (Find et al. 2005; Maghuly et al. 2006; Tsay et al. 2012). Microscopic (SEM) analysis showed adaxial and abaxial surface of transgenic *Salvia* leaves has more trichomes as well as bigger and wobbly guard cells and stomata. In contrast, wild type plant showed the less number of trichomes and smaller guard cells and stomata. Genetic modification in plants leads to change in morphology. Ectopic expression of gene in *Arabidopsis* showed different morphology and abnormal flowers containing wrinkled organs (Liu et al. 2008). Weigel et al. (2000) reported that the 35S enhancer element might increase the expression of nearby genes without altering the original expression pattern. Since, incorporation of CaMV35S enhancers of pTAG8 is random in the genome, so changes in leaf morphology of SH41 is may be due to the activation of genes responsible for leaf development, which remains to be confirmed.

Based on previous studies, phenolic acids and diterpene contents vary with different solvent systems and extraction methods. In the present study, the combinations of methanol and ethanol with water were used to recover greater yield of both water as well as lipid soluble constituents of *Salvia* roots by sonication. It was observed that, 60% ethanol containing 0.05% formic acid

was the best extraction system and the addition of formic acid increased the extraction efficiency (Table 1). Although, the effect of temperature on extraction were reported in other studies, such as highest content of salvianolic acid B ( $48.37 \pm 1.21$  mg/g) was yielded in microwave assisted extraction with water (MAE-W) at 75°C, or the highest contents of tanshinones including dihydrotanshinone, cryptotanshinone, tanshinone I, and tanshinone IIA were obtained in MAE-W at 100°C (Zhou et al. 2012), but we needed a method to extract salvianolic acid B and tanshinones simultaneously. Therefore, 60% ethanol containing 0.05% formic acid was the best choice for this study.

The active compound of *S. miltiorrhiza*, tanshinone, was proved with lots of biological activities. For instance, tanshinone I could enhance learning and memory (Kim et al. 2009), while tanshinone IIA has anti-oxidative (Wang et al. 2003; Yang et al. 2008), anti-inflammatory (Jang et al. 2006), anti-proliferative (Liu et al. 2006) and anti-tumor properties (Dong et al. 2007). Therefore, *S. miltiorrhiza* has been widely used to produce a number of traditional Chinese medicine preparations. However, because of low content, the supplement of tanshinone for the increasing needs of clinical applications has become a research attention. Therefore, development of transgenic *Salvia* with altered tanshinone content could overcome the shortage of traditional Chinese medicine preparations. Manipulating multiple genes at multiple control sites in desired metabolite production may cause metabolic flux alteration (Nims et al. 2006). The activation tagging has the ability to activate whole branches of biochemical pathways leading to the accumulation of several subclasses of natural product usually present in low concentration (Tani et al. 2004). In this study ATM transgenic of *Salvia* showed significant increase in diterpene tanshinone contents in roots as compared to wild plant. Tanshinone I, tanshinone IIA and salvianolic acid B were 3.7, 2 and 1.4 fold higher in SH41 as compared to wild plant. In other study it was reported that the over-expression of genes encoding 3-hydroxy-3-methylglutaryl CoA reductase (*SmHMGR*), 1-deoxy-D-xylulose-5-phosphatesynthase (*SmDXS*) and geranylgeranyl diphosphate synthase (*SmGGPPS*) in *S. miltiorrhiza* can significantly enhance the production of tanshinones in roots to levels higher than that of the control. *S. miltiorrhiza* plants over-expressing *SmHMGR*, *SmGGPPS* and *SmDXS* produced 2.1 to 5.7 fold higher tanshinones than control plants (Kai et al. 2011).

## Conclusions

In conclusion, the current study provides good basement and helpful information for commercial large-scale production of tanshinones in the roots of *Salvia* by developing ATM transgenic lines with altered phenotype. Also,

the extraction procedure and quantitative HPLC procedures were developed for determining liposoluble as well as water-soluble phenolic acid compounds.

#### Abbreviations

ATM: Activation tagging mutagenesis; HPLC: High performance liquid chromatography; *hptII*: Hygromycin phosphotransferase II gene; SEM: Scanning electron microscopy.

#### Competing interests

The authors declare that they have no competing interests.

#### Authors' contributions

Prof. HS Tsay designed experiment and reviewed the manuscript. Mr. HS Ho performed experiments. Dr. RK Vishwakarma prepared the manuscript. Prof. Emily CF Chen helped in analyzing data and proof reading of the manuscript. All authors read and approved the final manuscript.

#### Acknowledgements

The authors wish to thank Dr. Chiou-Ing Yuan (From Taiwan Agricultural Chemical and Toxic Substances Research Institute, Council of Agriculture) for providing SEM facility. This work was supported by National Science Council of Taiwan under the project grant NSC-97-2317-B-324-004.

Received: 9 July 2013 Accepted: 5 September 2013

Published: 24 September 2013

#### References

- Chen H, Yuan JP, Chen F, Zhang YL, Song JY (1997) Tanshinone production in Ti-transformed *Salvia miltiorrhiza* cell suspension cultures. *J Biotechnol* 58:147–156
- Chen ECF, Su YH, Kanagarajan S, Agrawal DC, Tsay HS (2009) Development of an activation tagging system for the basidiomycetous medicinal fungus *Antrodia cinnamomea*. *Mycol Res* 113:290–297
- Dong XR, Dong JH, Peng G, Hou XH, Wu G (2007) Growth-inhibiting and apoptosis-inducing effects of Tanshinone IIA on human gastric carcinoma cells. *J Huazhong Univ Sci Technol* 27:706–709
- Doyle JJ, Doyle JL (1987) A rapid DNA isolation procedure for small quantities of fresh leaf tissue. *Electrophoresis* 8:11–15
- Find JI, Charity JA, Grace LJ, Kritensen MMMH M, Krogstrup P, Walter C (2005) Stable genetic transformation of embryogenic cultures of *Abies nordmanniana* (Nordmann fir) and regeneration of transgenic plants. *In Vitro Cell Dev Biol Plant* 41:725–730
- Ge X, Wu J (2005a) Induction and potentiation of diterpenoid tanshinone accumulation in *Salvia miltiorrhiza* hairy roots by beta-aminobutyric acid. *Appl Microbiol Biotechnol* 68:183–188
- Ge X, Wu J (2005b) Tanshinone production and isoprenoid pathways in *Salvia miltiorrhiza* hairy roots induced by Ag+ and yeast elicitor. *Plant Sci* 168:487–491
- Gupta SK, Liu RB, Liaw SY, Chan HS, Tsay HS (2011) Enhanced tanshinone production in hairy roots of *Salvia miltiorrhiza* Bunge under the influence of plant growth regulators in liquid culture. *Bot Stud* 52:435–443
- Han JY, Fan JY, Horie Y, Miura S, Cui DH, Ishii H, Hibi T, Tsuneki H, Kimura I (2008) Ameliorating effects of compounds derived from *Salvia miltiorrhiza* root extract on microcirculatory disturbance and target organ injury by ischemia and reperfusion. *Pharmacol Ther* 117:280–295
- Hsing YI, Chen CG, Fan MJ, Lu PC, Chen KT, Lo SF, Sun PK, Ho SL, Lee KW, Wang YC, Huang WL, Ko SS, Chen S, Chen JL, Chung CI, Lin YC, Hour AL, Wang YW, Chang YC, Tsai MW, Lin YS, Chen YC, Yen HM, Li CP, Wey CK, Tseng CS, Lai MH, Huang SC, Chen LJ, Yu SM (2007) A rice gene activation/knockout mutant resource for high throughput functional genomics. *Plant Mol Biol* 63:351–364
- Hu ZB, Alfermann AW (1993) Diterpenoid production in hairy roots cultures of *Salvia miltiorrhiza*. *Phytochemistry* 32:99–103
- Ichikawa T, Nakazawa M, Kawashima M, Muto S, Gohda K, Suzuki K, Ishikawa A, Kobayashi H, Yoshizumi T, Tsumoto Y, Tsubara Y, Iizumi H, Goto Y, Matsui M (2003) Sequence database of 1172 T-DNA insertion sites in *Arabidopsis* activation-tagging lines that showed phenotypes in T1 generation. *Plant J* 36:421–429
- Jang SI, Kim HJ, Kim YJ (2006) Tanshinone IIA inhibits LPS-induced NF- $\kappa$ B activation in RAW264.7 cells: possible involvement of the NIK-IKK, ERK1/2, p38 and JNK pathways. *Eur J Pharmacol* 542:1–7
- Jeong DH, An S, Park S, Kang HG, Park GG, Kim SR, Sim J, Kim YO, Kim MK, Kim SR, Kim J, Shin M, Jung M, An G (2006) Generation of a flanking sequence-tag database for activation-tagging lines in japonica rice. *Plant J* 45:123–132
- Kai G, Xu H, Zhou C, Liao P, Xiao J, Luo X, You L, Zhang L (2011) Metabolic engineering tanshinone biosynthetic pathway in *Salvia miltiorrhiza* hairy root cultures. *Metab Eng* 13:319–327
- Kim DH, Kim S, Jeon SJ, Son KH, Lee S, Yoon BH, Cheong JH, Ko KH, Ryu JH (2009) Tanshinone I enhances learning and memory, and ameliorates memory impairment in mice via the extracellular signal-regulated kinase signalling pathway. *Br J Pharmacol* 158:1131–1142
- Lam FF, Yeung JH, Chan KM, Or PM (2007) Relaxant effects of danshen aqueous extract and its constituent danshensu on rat coronary artery are mediated by inhibition of calcium channels. *Vascul Pharmacol* 46:271–277
- Lam FF, Yeung JH, Chan KM, Or PM (2008a) Dihydro-tanshinone, a lipophilic component of *Salvia miltiorrhiza* (danshen), relaxes rat coronary artery by inhibition of calcium channels. *J Ethnopharmacol* 119:318–321
- Lee CY, Agrawal DC, Wang CS, Yu SM, Chen JW, Tsay HS (2008) T-DNA activation tagging as a tool to isolate *Salvia miltiorrhiza* Bunge transgenic lines for higher yields of tanshinones. *Planta Med* 74:1–7
- Liu JJ, Lin DJ, Liu PQ, Huang M, Li XD, Huang RW (2006) Induction of apoptosis and inhibition of cell adhesive and invasive effects by tanshinone IIA in acute promyelocytic leukemia cells *in vitro*. *J Biomed Sci* 13:813–823
- Liu AH, Guo H, Ye M, Lin YH, Sun JH, Xu M, Guo DA (2007) Detection, characterization and identification of phenolic acids in Danshen using high-performance liquid chromatography with diode array detection and electrospray ionization mass spectrometry. *J Chromatogr* 1161:170–182
- Liu J, Ha D, Xie Z, Wang C, Wang H, Zhang W, Zhang J, Chen S (2008) Ectopic expression of soybean GmKNT1 in *Arabidopsis* results in altered leaf morphology and flower identity. *J Genet Genomics* 35:441–449
- Maghuly F, Leopold S, da Câmara MA, Borroto Fernandez E, Khan MA, Gambino G, Gribaudo I, Scharl A, Laimer M (2006) Molecular characterization of grapevine plants transformed with GFLV resistance genes: II. *Plant Cell Rep* 25:546–553
- Michael AA, Anthony JP (2011) Activation tagging and insertional mutagenesis in barley. *Methods Mol Biol* 678:107–128
- Nims E, Dubois C, Roberts SC, Walker EL (2006) Expression profiling of genes involved in paclitaxel biosynthesis for targeted metabolic engineering. *Metab Eng* 8:385–394
- Sambrook J, Fritsch EF, Maniatis T (1989) Molecular cloning: A laboratory manual (2nd ed). Cold Spring Harbor Press, New York, p 2344. ISBN 0-87969-136-0
- Shu ZY, Zhang K, Ye XY, et al. (2010) Combination of naloxone, tanshinone and touch treatment on hypoxic ischemic encephalopathy of neonate. *Res Int Trad Chin West Med* 3:133–134
- Takahashi K, Ouyang X, Komatsu K, Nakamura N, Hattori M, Baba A, Azuma J (2002) Sodium tanshinone IIA sulfonate derived from Danshen (*Salvia miltiorrhiza*) attenuates hypertrophy induced by angiotensin II in cultured neonatal rat cardiac cells. *Biochem Pharmacol* 64:745–749
- Tani H, Chen X, Nurnberg P, Grant JJ, Maria MS, Chini A, Gilroy E, Birch PR, Loake GJ (2004) Activation tagging in plants: a tool for gene discovery. *Funct Integr Genomics* 4:258–266
- Tsay HS, Ho HM, Gupta SK, Wang CS, Chen PT, Chen ECF (2012) Development of pollen mediated activation tagging system for *Phalaenopsis* and *Doritaenopsis*. *Electron J Biotechnol*. doi:10.2225/vol15-issue4-fulltext-1
- Wang AM, Sha SH, Lesniak W, Schacht J (2003) Tanshinone (*Salvia miltiorrhiza* Extract) Preparations attenuate aminoglycoside-induced free radical formation *in vitro* and ototoxicity *in vivo*. *Antimicrob Agents Chemother* 47:1836–1841
- Wang X, Wang Y, Jiang M, Zhu Y, Hu L, Fan G, Li X, Gao X (2011a) Differential cardioprotective effects of salvianolic acid and tanshinone on acute myocardial infarction are mediated by unique signaling pathways. *J Ethnopharmacol* 135:662–671
- Weigel D, Ahn JH, Blazquez MA, Borevitz JO, Christensen SK, Fankhauser C, Ferrandiz C, Kardailsky I, Malancharuvil EJ, Neff MM, Nguyen JT, Sato S, Wang ZY, Xia Y, Dixon RA, Harrison MJ, Lamb CJ, Yanofsky MF, Chory J (2000) Activation tagging in *Arabidopsis*. *Plant Physiol* 122:1003–1013
- Wu TW, Zeng LH, Fung KP, Wu J, Pang H, Grey AA, Weisel RD, Wang JY (1993) Effect of sodium tanshinone IIA sulfonate in the rabbit myocardium and on human cardiomyocytes and vascular endothelial cells. *Biochem Pharmacol* 46:2327–2332
- Wu CT, Mulabagal V, Nalawade SM, Chen CL, Yang TF, Tsay HS (2003) Isolation and quantitative analysis of cryptotanshinone, an active quinoid diterpene formed in callus of *Salvia miltiorrhiza* Bunge. *Biol Pharm Bull* 26:845–848

- Yagi A, Okamura N, Tanonaka K, Takeo S (1994) Effects of tanshinone VI derivatives on post-hypoxic contractile dysfunction of perfused rat hearts. *Planta Med* 60:405–409
- Yan YP, Wang ZZ (2007) Genetic transformation of the medicinal plant *Salvia miltiorrhiza* by *Agrobacterium tumefaciens*-mediated method. *Plant Cell Tiss Org* 88:175–184
- Yan Q, Hu ZD, Tan RX, Wu JY (2005) Efficient production and recovery of diterpenoid tanshinonein *Salvia miltiorrhiza* hairy root cultures with in situ adsorption, elicitation and semi-continuous operation. *J Biotechnol* 119:416–424
- Yang R, Liu A, Ma X, Li L, Su D, Liu J (2008) Sodium tanshinone IIA sulfonate protects cardiomyocytes against oxidative stress-mediated apoptosis through inhibiting JNK activation. *J Cardiovasc Pharmacol* 51:396–401
- Yoon Y, Kim YO, Jeon WK, Park HJ, Sung HJ (1999) Tanshinone IIA isolated from *Salvia miltiorrhiza* BUNGE induced apoptosis in HL60 human promyelocytic leukemia cell line. *J Ethnopharmacol* 68:121–127
- Zhang Y, Song J, Zhao B, Liu H (1995) Crown gall culture and production of tanshinone in *Salvia miltiorrhiza*. *Chin J Biotechnol* 11:137–141
- Zhang Y, Song J, Qi J, Lu G (1997) The plant regeneration of *Salvia miltiorrhiza* Bge. transformed by *Agrobacterium*. *Zhongguo Zhong Yao Za Zhi* (in Chinese) 22:274–275
- Zhang C, Yan Q, Cheuk WK, Wu J (2004) Enhancement of tanshinone production in *Salvia miltiorrhiza* hairy root culture by Ag + elicitation and nutrient feeding. *Planta Med* 70:147–151
- Zhou L, Zuo Z, Chow MS (2005) Danshen: an overview of its chemistry, pharmacology, pharmacokinetics, and clinical use. *J Clin Pharmacol* 45:1345–1359
- Zhou X, Chan K, Yeung JH (2012a) Herb–drug interactions with Danshen (*Salvia miltiorrhiza*): a review on the role of cytochrome P450 enzymes. *Drug Metabol Drug Interact* 27:9–18
- Zhou X, Chan SW, Tseng HL, Deng Y, Hoi PM, Choi PS, Or PM, Yang JM, Lam FF, Lee SM, Leung GP, Kong SK, Ho HP, Kwan YW, Yeung JH (2012b) Danshensu is the major marker for the antioxidant and vasorelaxation effects of Danshen (*Salvia miltiorrhiza*) water-extracts produced by different heat water-extractions. *Phytomedicine* 19:1263–1269

doi:10.1186/1999-3110-54-37

**Cite this article as:** Ho et al.: Activation tagging in *Salvia miltiorrhiza* can cause increased leaf size and accumulation of tanshinone I and IIA in its roots. *Botanical Studies* 2013 **54**:37.

Submit your manuscript to a SpringerOpen<sup>®</sup> journal and benefit from:

- Convenient online submission
- Rigorous peer review
- Immediate publication on acceptance
- Open access: articles freely available online
- High visibility within the field
- Retaining the copyright to your article

Submit your next manuscript at ► [springeropen.com](http://springeropen.com)