

## Research article

## Open Access

**Methotrexate enhances the anti-inflammatory effect of CF101 via up-regulation of the A<sub>3</sub> adenosine receptor expression**

Avivit Ochaion<sup>1,2</sup>, Sara Bar-Yehuda<sup>1</sup>, Shira Cohn<sup>1,2</sup>, Luis Del Valle<sup>3</sup>, Georginia Perez-Liz<sup>3</sup>, Lea Madi<sup>1</sup>, Faina Barer<sup>1</sup>, Motti Farbstein<sup>1</sup>, Sari Fishman-Furman<sup>1</sup>, Tatiana Reitblat<sup>4</sup>, Alexander Reitblat<sup>4</sup>, Howard Amital<sup>5</sup>, Yair Levi<sup>5</sup>, Yair Molad<sup>6</sup>, Reuven Mader<sup>7</sup>, Moshe Tishler<sup>8</sup>, Pnina Langevitz<sup>9</sup>, Alexander Zabutti<sup>1</sup> and Pnina Fishman<sup>1</sup>

<sup>1</sup>Can-Fite Biopharma Ltd., 10 Baret Street, Kiryat-Matalon, Petah-Tikva, 49170, Israel

<sup>2</sup>The Mina and Everard Goodman Faculty of Life Sciences, Bar-Ilan University, Heren Hyesod Street, Ramat-Gan, 5200, Israel

<sup>3</sup>Department of Neuroscience, Neuropathology Core & Center for NeuroVirology, Temple University School of Medicine, North 12th Street, Philadelphia, PA 19122, USA

<sup>4</sup>Internal Department D, The Barzilai Medical Center, Hahistadrut Street, Ashkelon, 78278, Israel

<sup>5</sup>Internal Department D/E, Meir Medical Center, Tshernihovsky Street, Kfar Saba, 44281, Israel

<sup>6</sup>Rheumatology Department, Rabin Medica Center, Zabutinsky Street, Petah-Tikva, 49100, Israel

<sup>7</sup>Medical Clinic of Rheumatology, Ha'Emek Medical Center, Afula, 18101 Israel

<sup>8</sup>Rheumatology Department, Assaf Harofeh Medical Center, Zerifin, Beer Yakov, 70300, Israel

<sup>9</sup>Internal Department F, The Chaim Sheba Medical Center, Tel-Hashomer, 52621 Israel

Corresponding author: Pnina Fishman, [pnina@canfite.co.il](mailto:pnina@canfite.co.il)

Received: 27 Jul 2006 Revisions requested: 10 Aug 2006 Revisions received: 24 Oct 2006 Accepted: 13 Nov 2006 Published: 13 Nov 2006

*Arthritis Research & Therapy* 2006, **8**:R169 (doi:10.1186/ar2078)

This article is online at: <http://arthritis-research.com/content/8/6/R169>

© 2006 Ochaion *et al*; licensee BioMed Central Ltd.

This is an open access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/2.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

**Abstract**

Methotrexate (MTX) exerts an anti-inflammatory effect via its metabolite adenosine, which activates adenosine receptors. The A<sub>3</sub> adenosine receptor (A<sub>3</sub>AR) was found to be highly expressed in inflammatory tissues and peripheral blood mononuclear cells (PBMCs) of rats with adjuvant-induced arthritis (AIA). CF101 (IB-MECA), an A<sub>3</sub>AR agonist, was previously found to inhibit the clinical and pathological manifestations of AIA. The aim of the present study was to examine the effect of MTX on A<sub>3</sub>AR expression level and the efficacy of combined treatment with CF101 and MTX in AIA rats. AIA rats were treated with MTX, CF101, or both agents combined. A<sub>3</sub>AR mRNA, protein expression and exhibition were tested in paw and PBMC extracts from AIA rats utilizing immunohistochemistry staining, RT-PCR and Western blot analysis. A<sub>3</sub>AR level was tested in PBMC extracts from patients chronically treated with MTX and healthy individuals. The effect of CF101, MTX and combined treatment on A<sub>3</sub>AR expression level was also tested in PHA-

stimulated PBMCs from healthy individuals and from MTX-treated patients with rheumatoid arthritis (RA). Combined treatment with CF101 and MTX resulted in an additive anti-inflammatory effect in AIA rats. MTX induced A<sub>2A</sub>AR and A<sub>3</sub>AR over-expression in paw cells from treated animals. Moreover, increased A<sub>3</sub>AR expression level was detected in PBMCs from MTX-treated RA patients compared with cells from healthy individuals. MTX also increased the protein expression level of PHA-stimulated PBMCs from healthy individuals. The increase in A<sub>3</sub>AR level was counteracted *in vitro* by adenosine deaminase and mimicked *in vivo* by dipyridamole, demonstrating that receptor over-expression was mediated by adenosine. In conclusion, the data presented here indicate that MTX induces increased A<sub>3</sub>AR expression and exhibition, thereby potentiating the inhibitory effect of CF101 and supporting combined use of these drugs to treat RA.

**Introduction**

Low-dose methotrexate (MTX) is the most widely used

antirheumatic drug and it is the 'gold standard' against which other systemic medications are compared [1]. It has as its

A<sub>3</sub>AR = A<sub>3</sub> adenosine receptor; ADA = adenosine deaminase; AIA = adjuvant-induced arthritis; BSA = bovine serum albumin; IB-MECA = 1-deoxy-1-[6-[[[(3-iodophenyl)methyl]amino]-9H-9-yl]-N-methyl-β-d-ribofura-nuronamide; MTX = methotrexate; NF-κB = nuclear factor-κB; PBMC = peripheral blood mononuclear cell; PBS = phosphate-buffered saline; PHA = phytohemagglutinin; RA = rheumatoid arthritis; RT-PCR = reverse transcription polymerase chain reaction; TNF = tumour necrosis factor.

target the enzyme dihydrofolate reductase, which is required for reduction of dihydrofolate to tetrahydrofolate. It is presumed that cells exposed to MTX die as a result of reduced folate depletion [2]. Adenosine, an additional active metabolite of MTX, has been found to have potent anti-inflammatory effects, and earlier studies [3,4] strongly support the notion that the anti-inflammatory effect of MTX is attributed more to adenosine than to tetrahydrofolate. MTX increases the extracellular concentration of adenosine, where it is known to exert its anti-inflammatory effect via suppression of inflammatory cytokines such as tumour necrosis factor (TNF)- $\alpha$ , interleukin-6, or macrophage inhibitory protein-1 $\alpha$  [5-7]. It was further found that the anti-inflammatory effect of adenosine is mediated via A<sub>2A</sub> and the A<sub>3</sub> adenosine receptors [8,9].

The highly selective A<sub>3</sub> adenosine receptor (A<sub>3</sub>AR) agonist IB-MECA (1-deoxy-1-[6-[[[3-iodophenyl)methyl]amino]-9H-9-yl]-N-methyl- $\beta$ -D-ribofuranonamide) had an anti-inflammatory effect in collagen-induced arthritis in DBA1 mice and adjuvant-induced arthritis (AIA) in rats [10,11]. Interestingly, A<sub>3</sub>AR was found to be over-expressed in the synovial and paw inflammatory tissues as compared with corresponding tissues in normal, healthy animals. Moreover, receptor upregulation was also identified in the peripheral blood mononuclear cells (PBMCs) of AIA rats compared with control animals. Mechanistically, on treatment with IB-MECA, downregulation of A<sub>3</sub>AR expression level was noted in cells derived from the synovial tissue, most probably due to receptor internalization and degradation. Subsequently, decreased levels of expression of phosphatidylinositol-3 kinase and protein kinase B/Akt were observed. The latter of these proteins is known to control the nuclear factor- $\kappa$ B (NF- $\kappa$ B) signal transduction pathway. The decreased levels of PKB/Akt resulted in failure to phosphorylate IKK, which in turn resulted in inability to release NF- $\kappa$ B from its I $\kappa$ B complex. These events led to decreased expression of NF- $\kappa$ B and TNF- $\alpha$ , resulting in apoptosis of synovial cells. Remarkably, the PBMCs of AIA rats responded to IB-MECA treatment in the same manner as did the synovial cells, namely with receptor downregulation, suggesting that PBMCs reflect the receptor situation in inflammatory tissues and may have utility as a biomarker for monitoring response to IB-MECA [12]. Furthermore, Gessi and coworkers [13] recently noted upregulation of A<sub>3</sub>AR expression in phytohemagglutinin (PHA)-stimulated PBMCs from healthy individuals. It thus seems that A<sub>3</sub>AR expression correlates with cell activation or pathogenicity.

Recently, IB-MECA (commercially known as CF101) was tested in phase I clinical trials in healthy individuals. CF101, in single and multiple oral dose studies, was found to be safe and well tolerated, and the pharmacokinetics were linearly proportional to dose [14]. In an early phase II clinical trial of CF101 conducted in patients with rheumatoid arthritis (RA), the drug was well tolerated and conferred benefit as monotherapy [15].

Most biological disease-modifying antirheumatic drugs currently are given in combination with MTX [16]. Therefore, the aim of the present study was to evaluate the efficacy of combined MTX+CF101 treatment. We found that MTX treatment increased expression of A<sub>3</sub>AR, rendering inflammatory cells more susceptible to CF101. Combined treatment of AIA rats with MTX and CF101 enhanced the anti-inflammatory effect of each drug. In addition, we found A<sub>3</sub>AR to be over-expressed in PBMCs of MTX-treated patients and in activated cells from healthy individuals. The molecular mechanisms involved are explored.

## Materials and methods

### Reagents

The A<sub>3</sub>AR agonist IB-MECA was synthesized for Can-Fite BioPharma Ltd (Petah-Tikva, Israel) by Albany Molecular Research Inc (Albany, NY, USA) and is referred to as CF101 in the following text. A stock solution of 10 mmol/l was prepared in DMSO and further dilutions were performed. RPMI, foetal bovine serum and antibiotics for cell cultures were obtained from Beit Haemek (Haifa, Israel). The A<sub>3</sub>AR antagonist MRS1523 was purchased from Sigma (St. Louis, MO, USA) and diluted in the same manner as for CF101. Rabbit polyclonal antibodies against rat and human A<sub>3</sub>AR as well as rat A<sub>2A</sub>AR were purchased from Santa Cruz Biotechnology Inc. (Santa Cruz, CA, USA). MTX, PHA, adenosine and adenosine deaminase (ADA) were purchased from ABIC (Beit Shemesh, Israel). Dipyridamole was purchased from Sigma.

### Effect of CF101 and MTX on development of AIA

Female Lewis rats, aged 9 weeks, were obtained from Harlan Laboratories (Jerusalem, Israel). Rats were maintained on a standardized pellet diet and supplied with tap water. Experiments were performed in accordance with the guidelines established by the Institutional Animal Care and Use Committee at Can-Fite BioPharma, Petach Tikva, Israel. The rats were injected subcutaneously at the tail base with 100  $\mu$ l suspension composed of incomplete Freund's adjuvant with 10 mg/ml heat killed *Mycobacterium tuberculosis* H37Ra (Difco, Detroit, MI, USA). Each group included 10 animals.

For the prophylactic treatment study, MTX treatment (0.25 mg/kg intraperitoneally) was administered once weekly, starting 3 days after disease induction. CF101 (100  $\mu$ g/kg) was orally administered by gavage, twice daily, starting at disease onset. The study also included a group treated with a combination of MTX and CF101. The control group received vehicle only (DMSO at a dilution corresponding to that of CF101).

For the therapeutic treatment study, MTX (0.75 mg/kg intraperitoneally, once weekly) as well as CF101 (100  $\mu$ g/kg orally, twice daily) were initiated on disease onset. As in the prophylactic study, a group treated with a combination of MTX and CF101 was included.

Clinical disease activity score was assessed as follows. The animals were inspected every day for clinical arthritis. The scoring system ranged from 0 to 4 of each limb: 0 indicates no arthritis, 1 indicates redness or swelling of one toe/finger joint, 2 indicates redness and swelling of more than one toe/finger joints, 3 indicates involvement of ankle and tarsal-metatarsal joints, and 4 indicates redness or swelling of the entire paw. The four individual leg scores were added together to yield a total clinical score.

At the end of the study the legs were removed up to the knees, fixed in 10% formaldehyde, decalcified, dehydrated, paraffin embedded and cut into 4  $\mu$ m sections. They were then stained by haematoxylin and eosin, and morphopathological assessment was performed.

Pathological assessments were performed using semiquantitative grading scales from 0 to 4 for the following parameters: the extent of inflammatory cell infiltration into the joint tissues, synovial lining cell hyperplasia, pannus formation, joint cartilage layer destruction and bone damage and erosion: 0 indicates normal; 1 indicates minimal loss of cortical bone at a few sites; 2 indicates mild loss of cortical trabecular bone; 3 indicates moderate loss of bone at many sites; 4 indicates marked loss of bone at many sites; and 5 indicates marked loss of bone at many sites, with fragmenting and full thickness penetration of inflammatory process or pannus into the cortical bone. The means of all of the histological parameter scores were summated to yield an overall 'histology score'.

In addition, blood samples were collected and subjected to Ficoll hypaque gradient. The PBMCs were then washed with phosphate-buffered saline (PBS), and protein extracts were prepared as detailed below.

The hind paws were dissected above the ankle joint. The bony tissue was broken into pieces, snap frozen in liquid nitrogen and stored at -80°C until use. The paw tissues were added to RIPA extraction buffer (4 ml/g tissue) containing 150 mmol/l NaCl, 50 mmol/l Tris, 1% NP40, 0.5% deoxycholate and 0.1% SDS. Tissues were homogenized on ice with a polytron, centrifuged and the supernatants were subjected to Western Blot analysis.

#### Effect of dipyridamole on the expression of A<sub>3</sub>AR in PBMCs from naïve rats

Dipyridamole (5 mg/kg intraperitoneally) was administered once to naïve rats and blood samples were drawn 2, 6, 24 and 48 hours after dipyridamole administration and subjected to Ficoll hypaque gradient. The PBMCs were then washed with PBS and protein extracts were prepared as detailed below.

#### Immunohistochemistry staining of paraffin-embedded slides of paws tissues derived from AIA rats

The paraffin on the slides was melted from the sections, which were placed in xylene three times for 30 min each. The tissues were hydrated with serial dilutions of ethanol and then antigen was retrieved by heating with citrate buffer at 95°C for 30 min. The slides were allowed to cool down and then washed three times in PBS. Endogenous peroxidase quenching was performed by washing the sections with fresh 20% hydrogen peroxide in methanol for 20 min. The sections were then blocked by incubating in 5% normal goat serum in PBS-BSA 0.1% for 2 hours.

The primary antibody (Novus Biologicals Inc., Littleton, CO, USA) was diluted in 0.1% PBS-BSA and incubated overnight at room temperature. After three washes in 1 × PBS, the slides were incubated in 0.5% biotinylated secondary antibody in

**Table 1**

#### Characteristic of patients newly and chronically treated with MTX

Characteristic	RA patients	
	Newly MTX treated	Chronically MTX treated
Age (years)	48.5 ± 5.89	59.58 ± 2.42
Disease duration (years)	4.37 ± 2.07	6.6 ± 0.97
CRP (mg/l)	10.75 ± 2.01	9.04 ± 1.65
ESR (mm/hour)	31.75 ± 0.63	33.3 ± 4.39
Swollen joint count	13.25 ± 2.56	3.19 ± 0.88
Tender joint count	10.5 ± 2.63	5 ± 0.86
Patient global assessment (VAS)		32.26 ± 4.41
CRP (DAS28-4)		3.23 ± 0.31
ESR (DAS28-4)		4.11 ± 0.3

Values are expressed as mean ± standard error. CRP, C-reactive protein; DAS, Disease Activity Score; ESR, erythrocyte sedimentation rate; MTX, methotrexate; RA, rheumatoid arthritis; VAS, visual analogue scale.

PBS-BSA 0.1% for 1 hour at room temperature and then subjected to an avidin-biotin complex. After another wash the slides were incubated with DAB substrate, washed in tap water (to dispose of the DAB) and inactivated with bleach. A light haematoxylin counterstain was performed and the haematoxylin was then removed by quick dip in acid alcohol and the slides were washed in ammonia water, dehydrated and mounted with Permount.

#### Human blood sample collection and separation

Blood samples were collected from healthy individuals and from patients with RA who were either newly and chronically treated with MTX. Protocols for the study were approved by the hospitals' ethical committees as was the blood sample collection. Healthy individuals and RA patients provided signed, informed consent prior to blood withdrawal. Patient characteristics are summarized in Table 1.

To separate PBMCs, heparanized blood (20 ml) was subjected to Ficoll hypaque gradient. The PBMCs were then washed with PBS and subjected to the various assays.

#### Double immunofluorescence staining of PBMCs of RA patients

PBMCs ( $1 \times 10^6$  cells) were smeared on electrophorized glass slide and fixed with 75% ethanol for 2 min. The slides were then blocked with normal horse serum for 1 hour at room temperature in a humidified chamber, after which a mouse monoclonal CD4 (Santa Cruz Biotechnology; Clone MT310, 1:50 dilution) primary antibody was incubated overnight at room temperature. After rinsing thoroughly with PBS, a rhodamine-tagged secondary antibody (Vector Laboratories, Burlingame, CA, USA; 1:200 dilution) was incubated overnight in the dark. Then the slides were rinsed with PBS, blocked with normal goat serum and incubated with a rabbit polyclonal anti- $A_3AR$  antibody (Chemicon International, Temecula, CA, USA; AB9111, 1:100 dilution) overnight. Cells were rinsed with PBS and a second, fluorescein-tagged anti-rabbit antibody was incubated for 1 hour in the dark. Finally, slides were rinsed, coverslipped with an aqueous based mounting media (Vectashield; Vector Laboratories) and visualized in a Nikon ultraviolet inverted microscope and processed with a deconvolution software (Slidebook 4.0; Intelligent Imaging, Denver, CO, USA).

#### Activation of PBMCs

PBMCs ( $2 \times 10^6$  cells/ml) from healthy individuals or from MTX-treated RA patients were incubated in cell cultures with RPMI 1640 supplemented with 10% FBS. PBMCs from healthy individuals were activated with PHA (5  $\mu$ g/ml) for 24 hours. MTX (1  $\mu$ mol/l) was added for an additional 27 hours. ADA (1 unit/ml), CF101 (10 nmol/l), or MRS1353 (10 nmol/l) was added to the culture system for the last 3 hours. In another set of experiments PBMCs from healthy individuals were incubated for 27 hours with adenosine (25  $\mu$ mol/l), and ADA (1

unit/ml) was added to the culture system for the last 3 hours. PBMCs from MTX-treated RA patients were incubated for 1 hour with CF101 (10 nmol/l). At the end of the incubation time in all of the above experiments, the PBMCs were collected from the culture plates and protein extracts were prepared.

#### Western blot analysis of $A_3AR$ and additional signalling proteins in PBMCs

Western blot analyses were carried out according to the following protocol. Samples were rinsed with ice-cold PBS and transferred to ice-cold lysis buffer (TNN buffer: 50 mmol/l Tris buffer [pH 7.5], 150 mmol/l NaCl and NP40). Cell debris was removed by centrifugation for 10 min at 7500 *g*. Protein concentrations were determined using the BioRad protein assay dye reagent (BioRad Laboratories, Hercules, CA, USA). Equal amounts of the sample (50  $\mu$ g) were separated by SDS-PAGE, using 12% polyacrylamide gels. The resolved proteins were then electroblotted onto nitrocellulose membranes (Schleicher & Schuell, Keene, NH, USA). Membranes were blocked with 1% BSA and incubated with the desired primary antibody (dilution 1:1000) for 24 hours at 4°C. Blots were then washed and incubated with a secondary antibody for 1 hour at room temperature. Bands were recorded using BCIP/NBT color development kit (Promega, Madison, WI, USA).

Analysis of  $A_3AR$  protein expression level in patient PBMCs was performed as follows. Samples from each of four RA patients were run in the same gel with a pool of samples from four healthy individuals, designated the standard. The blots were quantified by densitometric analysis and the ratio of RA patient/standard was calculated. Blots of mitogen stimulated cells were quantified against  $\beta$ -actin. The data presented in the figures are representative of at least three different experiments.

#### RT-PCR analysis of formalin-fixed paraffin-embedded paw tissue slides

Tissue sections of paws derived from AIA rats treated with vehicle, MTX, CF101, or MTX+CF101 were mounted on slides and then deparaffinized in xylene and rehydrated by washing in serial dilutions of ethanol. Slides were used immediately or stored at -80°C until use. After rehydration, 20  $\mu$ l of solution A (1.25 $\times$  PCR buffer [200 mmol/l Tris-HCl, 500 mmol/l KCl], 6.25 mmol/l  $MgCl_2$ , 5 U RNasin [Promega], 2 mmol/l DTT, 1 U RQ1 RNase-free DNase [Promega]) was directly applied to the marked area. The marked area was completely scraped off the slide using a pipette tip, and neoplastic tissue or normal tissue was collected into different microcentrifuge tubes. The samples were treated with proteinase K at a final concentration of 0.1 mg/ml. The samples were incubated at 37°C for 1 hour to allow for DNA digestion. Cells lysate were heated to 95°C for 15 min in order to inactivate DNase and proteinase K. Following centrifugation at 14,000 rpm for 5 min, 17  $\mu$ l of the supernatant was transferred to a separate tube and 4  $\mu$ l of RT mixture (5 mmol/l dNTPs, 2.5  $\mu$ mol/l ran-

dom hexamer, 5 U RNasin, 100 U SuperScript One Step RT-PCR with Platinum Taq (Invitrogene, San Diego, CA, USA) and the primers for rat A<sub>3</sub>AR (245 up CTA GCA CTG GCA GAC and 245 down CAG CAG AGG CCC AGG) were added.


The RT reaction was performed at 45°C for 45 min, followed by heating to 99°C for 5 min; then, 50 cycles at 94°C for 30 s, 50°C for 75 s and 73°C for 45 s, and an extension of 73°C for 7 min were performed. Products were electrophoresed on 2% agarose gels, stained with ethidium bromide and visualized with ultraviolet illumination. The specificity of the RT-PCR

reaction was confirmed by size determination on agarose gels in comparison with a positive control, from RNA extracted using standard techniques, and by sequencing the RT-PCR product and comparing the sequences with the known sequences (ADORA3-L77729, L77730). The optical density of the bands (Et-Br) was quantified using an image analysis system. This analysis was performed on tissues from three different experiments.

**Statistical analysis**

To analyze differences in clinical score between the four study

**Figure 1**



Effect of CF101 and prophylactic or therapeutic MTX treatment on AIA rats. Rats were injected subcutaneously at the tail base with 100 µl of a suspension composed of incomplete Freund's adjuvant and 10 mg/ml heat killed *Mycobacterium tuberculosis*. **(a)** Clinical score. Combined treatment with MTX (prophylactic treatment) + CF101 yield significantly lower values than treatment with each of the agents alone; also, all treatments yielded lower scores than control group. **(b)** The clinical score with combined therapeutic MTX treatment + CF101 was significantly lower than that in the other groups. **(c, d)** Morphopathological score. In AIA animals complete destruction of the cartilage and the bone, as well as severe inflammation in the hind paws, was noted. Treatment with a combination of MTX and CF101 preserved the normal features of the paw. AIA, adjuvant-induced arthritis; MTX, methotrexate.

groups (CF101, MTX, MTX+CF101 [combined treatment] and control), we used analysis of variance (ANOVA) with repeated measures from days 8 to 25 for prophylactic treatment and from days 14 to 20 for therapeutic treatment. To test differences in trends during the study between the four study groups, we used ANOVA using Dunnett method to evaluate differences between each of the study groups and the control group: for prophylactic treatment this was from days 20 to 25, adjusted to baseline values (day 8); and for therapeutic treatment this was from days 14 to day 20, adjusted to baseline values at day 14. ANOVA using Dunnett method was utilized to evaluate differences between each of the study groups and the combined treatment group (CF101+MTX): for prophylactic treatment this was from days 20 to 25, adjusted to baseline values (day 8); and for therapeutic treatment this was from days 14 to 20, adjusted to baseline values at day 14. The data were analyzed using SAS software (SAS Institute, Cary, NC, USA).

The Student's *t*-test was used in the Western blot analysis, and  $P < 0.05$  was considered statistically significant.

## Results

### Effect of MTX, CF101, and MTX+CF101 treatment on development of AIA in rats

In the prophylactic treatment study, about 21 days after immunization most of the vehicle-treated animals progressively developed arthritis. Comparing the four groups using ANOVA, CF101 treatment (100 µg/kg, given orally twice daily, starting on onset of disease) and MTX treatment (given once weekly, starting on day 3 after disease induction) resulted in a statistically significant difference between the study groups and the control group (Figure 1a;  $P < 0.001$ ). In order to identify the source of those differences, ANOVA with the Dunnett method was used. In general, up to day 20 a differences in trends were observed between the groups, but the differences were not statistically significant. From day 20 on, a clear trend toward lower values in the combined treatment group (MTX+CF101) was observed, as compared with the control group and each of the other treatment groups (Figure 1a).

In the therapeutic treatment study, in which all of the treatments were initiated at the onset of disease, clinical score in the MTX and CF101 combined treatment group was found to be statistically significantly lower than that in the control group. Moreover, clinical score in the group receiving combined treatment with MTX and CF101 was always lower than that the other groups. From day 17 the trend becomes statistically significant (Figure 1b).

Histological evaluation of the paws in the vehicle-treated arthritic animals revealed complete destruction of cartilage and bone, which were replaced by granulation tissue. Severe inflammation infiltration was also noted. In the MTX treated group bone destruction was also observed, with a few rem-

nants of bone spiculates seated in the granulation tissue. The degree of inflammatory infiltration was less than that observed in the control group, resulting in a 47% decrease in histological score. CF101 treatment resulted in a 75% decrease in histological score, which was manifested as a mild inflammatory infiltration. The articular space, cartilage, bone and bone marrow appeared normal. Treatment with MTX in combination with CF101 was associated with normal cartilage and bone architecture, and with an histological score that was almost zero (Figure 1c,d).

### Effect of MTX, CF101 and MTX+ CF101 treatment on exhibition and expression of A<sub>3</sub>AR in paw and PBMCs from AIA rats

Immunohistochemistry staining of paw sections derived from AIA rats revealed the of A<sub>3</sub>AR exhibition in the control paw, which was markedly increased with MTX treatment. In the CF101-treated group, A<sub>3</sub>AR exhibition was much lower and the receptor was barely detected at all in MTX+CF101-treated paw sections (Figure 2).

Analysis of mRNA and protein A<sub>3</sub>AR expression in paw extracts from the MTX-treated group revealed an increase compared with the vehicle-treated group. In the CF101-treatment and the MTX+CF101-treated groups, receptor downregulation was noted, supporting previous studies demonstrating that receptor downregulation represents a functional response to the agonist (Figure 3a,b). In PBMCs from the animals treated with MTX, CF101 and MTX+CF101, similar results to those noted in the paw extracts were recorded (Figure 3c).


We further examined the effect of MTX on expression of A<sub>3</sub>AR protein compared with A<sub>2A</sub>AR protein in paw extracts. Figure 3d shows that the expression levels of both receptors is increased with MTX treatment.

To explore the molecular mechanism involved in the increased expression of A<sub>3</sub>AR with MTX treatment, we assumed this phenomenon to be attributed to elevated adenosine levels in the cell microenvironment. We thus treated naïve rats with dipyrindamole, a nucleoside transporter inhibitor that increases extracellular adenosine concentration. Indeed, dipyrindamole induced A<sub>3</sub>AR over-expression in PBMCs from treated rats (Figure 4).

### A<sub>3</sub>AR expression in PBMCs from RA patients


Our next step was to examine A<sub>3</sub>AR expression level in PBMCs from MTX-treated RA patients. Figure 5a shows four blots from newly MTX treated RA patients. Upregulation of A<sub>3</sub>AR expression was observed in all patients after 10 weeks of MTX treatment. There was a marked statistically significant increase ( $P < 0.01$ ) in A<sub>3</sub>AR expression in PBMCs from RA patients undergoing chronic MTX treatment ( $n = 30$ ) in com-

**Figure 2**




Immunohistochemistry analysis of  $A_3AR$  expression in paws from AIA rats. We conducted immunohistochemistry staining of paraffin-embedded sections of paw from MTX-treated, CF101-treated and MTX+CF101-treated AIA rats. MTX treatment induced receptor expression, and treatment with CF101 alone or in combination with MTX resulted in receptor downregulation.  $A_3AR$ ,  $A_3$  adenosine receptor; AIA, adjuvant-induced arthritis; MTX, methotrexate.

**Figure 3**



*Ex vivo* analysis of  $A_3AR$  expression level in paw and PBMCs from AIA rats. **(a, b)** MTX treatment induced expression of  $A_3AR$  mRNA and protein, and CF101 treatment resulted in receptor downregulation. **(c)** This was reflected in the PBMCs. **(d)** Expression of  $A_{2A}AR$  was also induced in paw extracts derived AIA rats treated with MTX.  $A_3AR$ ,  $A_3$  adenosine receptor; AIA, adjuvant-induced arthritis; MTX, methotrexate; PBMC, peripheral blood mononuclear cell.

Figure 4



Effect of dipyridamole on the expression of A<sub>3</sub>AR in PBMCs from naïve rats. Dipyridamole (5 mg/kg intraperitoneally) was administered once to naïve rats and blood samples were drawn 2, 6, 24 and 48 hours after dipyridamole administration. Dipyridamole induced A<sub>3</sub>AR upregulation in PBMCs derived from treated rats. A<sub>3</sub>AR, A<sub>3</sub> adenosine receptor; PBMC, peripheral blood mononuclear cell.

parison with healthy individuals (Figure 5b). No correlation between A<sub>3</sub>AR expression level, MTX dose, or DAS28 (Disease Activity Score) was found.

In addition, we performed double immunofluorescence staining on PBMCs from RA patients to identify A<sub>3</sub>AR on the cell surface of CD4<sup>+</sup> T lymphocytes. Figure 5c shows massive A<sub>3</sub>AR staining on the CD4<sup>+</sup> T cells.

In an additional set of experiments we looked at the *in vitro* effect of CF101 on A<sub>3</sub>AR protein level in PBMCs from MTX-treated RA patients. The cells were cultured in the presence and absence of CF101 for 1 hour. Figure 5d shows A<sub>3</sub>AR over-expression in the RA samples compared with control samples. CF101 treatment induced A<sub>3</sub>AR downregulation, reflecting the response of PBMCs to the drug; this suggests that receptor expression may have utility as a biological prediction marker.

#### Effect of MTX+CF101 on A<sub>3</sub>AR expression level in PHA-stimulated PBMCs from healthy individuals

To further investigate the effect of MTX on A<sub>3</sub>AR expression, we used an *in vitro* system of PHA-stimulated PBMCs from healthy individuals. We found that A<sub>3</sub>AR expression level was increased on MTX treatment. Also, introduction of ADA to the culture system reverted this effect, suggesting that receptor over-expression was induced by adenosine, a metabolite of MTX (Figure 6a). Finally, the natural ligand adenosine induced an increase in A<sub>3</sub>AR expression when added to the culture system, supporting the notion presented above that adenosine acts as mediator of the MTX effect (Figure 6b).

Introduction of CF101 to MTX-treated, PHA-stimulated cells induced receptor downregulation, which was counteracted by the A<sub>3</sub>AR antagonist MRS1535 (Figure 6c).


## Discussion

The present study shows that combined treatment with CF101 and MTX has an additive anti-inflammatory effect, which is indicated by a decrease in the clinical and pathological manifestations of AIA. Mechanistically, MTX induced an increase in A<sub>3</sub>AR expression level in inflamed tissues and in PBMCs, thereby rendering the cells more susceptible to CF101 treatment. Interestingly, MTX given either in prophylactic or therapeutic mode in combination with CF101 resulted in the same additive effect.

To explore the mechanism underlying the efficacy of the combined treatment, we first examined the mode of action of MTX. It acts as an antagonist of folic acid, subsequently inhibiting the synthesis of purines and pyrimidines. It was further suggested that the anti-inflammatory effect of MTX is due to adenosine, which is known to exert potent anti-inflammatory effect [3,4]. More specifically, MTX polyglutamates inhibit the activity of the enzyme aminoimidazolecarbox-amidoadenosineribonucleotide transformylase [17,18]. This enzyme has a direct inhibitory effect on two additional enzymes: ADA, which metabolizes adenosine to inosine; and AMP deaminase, which converts adenosine to AMP. This chain of events results in intracellular accumulation of adenosine, which is then released into the extracellular environment [19]. The working hypothesis of the present study was that the increase in adenosine level may act via an autocrine pathway and induce the expression of its own receptors, in this case A<sub>3</sub>AR. Indeed, MTX treatment increased both mRNA and protein levels of A<sub>3</sub>AR in the inflammatory tissue (paw) of AIA rats, indicating that increased gene expression and translation took place. An increase in protein A<sub>3</sub>AR expression was also noted in the PBMCs from the animals. Interestingly, increased A<sub>2A</sub>AR protein levels were also noted in paw samples from MTX-treated animals. Ravid and coworkers [20] have shown that activation of A<sub>2A</sub>AR increases A<sub>3</sub>AR promoter activity. It could therefore be



Figure 5



A<sub>3</sub>AR expression level in PBMCs from RA patients. **(a)** Newly MTX treated RA patients. Western blots of the four patients at baseline and 10 weeks after MTX treatment are presented. Representative Western blots are shown at the bottom. **(b)** RA patients undergoing chronic treatment with MTX. Results are expressed as means  $\pm$  standard error for 30 patients. **(c)** Double-stained immunofluorescence analysis of PBMCs from RA patients. A<sub>3</sub>AR expression on the cell surface of the CD4<sup>+</sup> T cells was found. **(d)** Incubation of PBMCs from MTX-treated RA patients for 1 hour in RPMI 1640 supplemented with 10% foetal bovine serum in the presence of with CF101 (10 nmol/l) resulted in decreased expression of A<sub>3</sub>AR. A<sub>3</sub>AR, A<sub>3</sub> adenosine receptor; AIA, adjuvant-induced arthritis; MTX, methotrexate; PBMC, peripheral blood mononuclear cell; RA, rheumatoid arthritis.

suggested that over-expression of A<sub>2A</sub>AR plays a role in the increased level of A<sub>3</sub>AR that occurs with MTX treatment. Moreover, A<sub>3</sub>AR protein level was raised in RA patients treated with MTX. This was attributed to the increase in A<sub>3</sub>AR expression detected on CD4<sup>+</sup> T cells from RA patients compared with CD4<sup>+</sup> T cells from healthy individuals.


In this report we present two lines of experimental evidence that support a role for adenosine in modulating A<sub>3</sub>AR expression in response to MTX treatment. *In vitro* studies with PBMCs showed that MTX treatment induced an increase in A<sub>3</sub>AR expression that was reversed on treatment with ADA. Adenosine mimicked the effect of MTX and induced an elevation in A<sub>3</sub>AR expression. *In vivo* dipyridamole, an inhibitor of nucleoside transporters, known to increase adenosine levels, induced A<sub>3</sub>AR over-expression in PBMCs from treated rats. These data led to the conclusion that MTX, via the metabolite adenosine, induces elevation in A<sub>3</sub>AR expression.

Receptor density was previously reported as an important factor that controls cell response to a given agonist [21-23]. It

may be suggested that A<sub>3</sub>AR upregulation, mediated by MTX, preconditions cells to the effect of CF101, resulting in a more potent anti-inflammatory effect.

An additional important finding of the study is that inflammatory cells and PBMCs from CF101-treated AIA rats, as well as the PBMCs cultured *in vitro* with CF101, responded to agonist treatment and to combined treatment with receptor downregulation. The A<sub>3</sub>AR antagonist MRS 1353 counteracted the downregulation induced by the combined treatment. This finding supports the notion presented above that the effect of CF101 alone or in combination with MTX is solely mediated via A<sub>3</sub>AR, without the involvement of additional receptors. Our earlier studies in tumor cells showed that, on binding of an agonist such as CF101 to tumor cells, receptor is internalized and degraded within the cytoplasm [24,25]. This is followed by initiation of downstream signal transduction pathways, leading to inhibition of tumor growth. A few hours later receptor is resynthesized and recycled to the cell surface. Recently, we reported a similar chain of events in the synovial cells of AIA rats treated with CF101. A<sub>3</sub>AR receptor levels

Figure 6



Effect of MTX on A<sub>3</sub>AR expression level on PBMCs from healthy individuals. **(a)** PBMCs ( $2 \times 10^6$  cells/ml) from healthy individuals were incubated in RPMI 1640 supplemented with 10% foetal bovine serum and activated with PHA (5  $\mu$ g/ml) for 24 hours. MTX (1  $\mu$ mol/l) was added for an additional 27 hours and ADA (1 unit/ml) for the last 3 hours. A<sub>3</sub>AR expression level was induced by MTX treatment. The introduction of ADA to the culture system reverted this effect and induced downregulation of the receptor level. **(b)** PBMCs ( $2 \times 10^6$  cells/ml) were incubated for 27 hours with adenosine (25  $\mu$ mol/l) and ADA (1 unit/ml) was added to the culture system for the last 3 hours. Adenosine induced upregulation of A<sub>3</sub>AR expression level whereas the addition of ADA decreased the receptor level. **(c)** PBMCs from healthy individuals were incubated with 10% foetal bovine serum and activated with PHA (5  $\mu$ g/ml) for 24 hours. MTX (1  $\mu$ mol/l) was added for an additional 27 hours and CF101 (10 nmol/l), in the presence or absence of MRS1523 (10 nmol/l), was introduced into the culture system for the last 3 hours. CF101 introduction to MTX-treated, PHA-activated PBMCs induced receptor downregulation, and the MRS1523 counteracted this effect. A<sub>3</sub>AR, A<sub>3</sub> adenosine receptor; ADA, adenosine deaminase; MTX, methotrexate; PBMC, peripheral blood mononuclear cell; PHA, phytohemagglutinin; RA, rheumatoid arthritis.

were downregulated on agonist treatment, followed by modulation of cell signaling proteins that belong to the NF- $\kappa$ B signal transduction pathway, resulting in decreased expression of TNF- $\alpha$  and inhibition of the inflammatory process [12]. It thus seems that if A<sub>3</sub>AR upregulation is characteristic of inflammation, then CF101 treatment induces receptor downregulation, which reflects receptor functionality, and modulation of key signaling proteins that control the inflammatory process. Interestingly, introduction of CF101 to *in vitro* culture of PBMCs derived from MTX-treated RA patients resulted in A<sub>3</sub>AR downregulation. The ability to assess *in vitro* the response to CF101 may be utilized in the future to predict the response of individual patients to the drug before treatment.

The mechanism presented in this study by which MTX elevates A<sub>3</sub>AR expression may also account for the anti-inflammatory effect of this drug when given as a standalone therapy. Under physiological conditions A<sub>3</sub>AR is not activated because adenosine has the lowest affinity value to this receptor ( $A_{2A} > A_1 > A_{2B} > A_3$ ). On MTX treatment the adenosine level goes up, thereby activating A<sub>3</sub>AR and resulting in an anti-inflammatory effect [3,4,26].

High expression levels of A<sub>3</sub>AR in inflamed or activated cells were previously reported in other inflammatory conditions. A selective, approximately 10-fold upregulation of A<sub>3</sub>AR mRNA and protein was consistently found in nonpigmented ciliary epithelium of eyes of patients with pseudo-exfoliation syndrome, with and without glaucoma, as compared with normal and glaucomatous control eyes [27]. Blair and coworkers [28] showed that A<sub>3</sub>AR transcript abundance is greater in lung tissue and eosinophils from individuals with airway inflammation than in normal lung. A<sub>3</sub>AR elevation was also reflected in eosinophils derived from the peripheral blood of the same patients. Treatment of eosinophils with IB-MECA inhibited platelet-activating factor induced eosinophil chemotaxis [28]. Moreover, Gessi and coworkers [29] reported that A<sub>3</sub>AR is induced in activated PBMCs from healthy individuals, and further demonstrated that the CD4<sup>+</sup> T cells are the subpopulation of cells that over-express the receptor. Taken together, it may be concluded that A<sub>3</sub>AR upregulation is a characteristic of activated cells, and is noted in cells of inflammatory origin and is reflected in PBMCs.

To summarize, enhanced anti-inflammatory effect takes place on treatment of AIA rats with the combination of MTX and CF101. This effect is mediated via adenosine, which accumulates in cells on MTX treatment, leading to increased levels of A<sub>3</sub>AR, thereby rendering the cells more sensitive to the effect of CF101. We recently reported a positive response of RA patients to CF101 treatment (as a standalone therapy) in a phase IIa study [15]. The results of the present study provide justification for a phase IIb study combining CF101 with MTX treatment in a population of RA patients.

## Conclusion

Using an *in vivo* model of AIA in rats, we demonstrated that combined treatment with CF101 and MTX resulted in an additive anti-inflammatory effect. Mechanistically, we found that MTX induced an increase in the expression level of A<sub>3</sub>AR, rendering the inflammatory cells more susceptible to CF101 treatment. Adding further support to these findings were data from the MTX-treated RA patients, who exhibited A<sub>3</sub>AR over-expression in PBMCs.

## Competing interests

The authors declare that they have no competing interests.

## Authors' contributions

OA, CH, ML and BF conducted the molecular biology work. ZA conducted the *in vivo* studies. FM conducted data management for the patients. FFS conducted patient sample handling and patient assessment. RT, RA, AH, LY, MY, MR, TM and LP collected blood samples and patient data. SBY precipitate in study design and data analysis, and helped to draft the manuscript. FP conceived the study, participated in the study design and data analysis, and drafted the manuscript.

## References

- Weinblatt ME, Coblyn JS, Fox DA, Fraser PA, Holdsworth DE, Glass DN, Trentham DE: **Efficacy of low-dose methotrexate in rheumatoid arthritis.** *N Engl J Med* 1985, **312**:818-822.
- Chan L, Cronstein BN: **Molecular action of methotrexate in inflammatory diseases.** *Arthritis Res* 2002, **4**:266-273.
- Baggott JE, Morgan SL, Sams WM, Linden J: **Urinary adenosine and aminoimidazolcarboxamide excretion in methotrexate-treated patients with psoriasis.** *Arch Dermatol* 1999, **135**:813-817.
- Laghi Pasini F, Capecci PL, Di Perri T: **Adenosine plasma levels after low-dose methotrexate administration.** *J Rheumatol* 1997, **24**:2492-2493.
- Cronstein BN: **Adenosine, an endogenous anti-inflammatory agent.** *J Appl Physiol* 1994, **76**:5-13.
- Eigler A, Greten TF, Sinha B, Haslberger C, Sullivan GW, Endres S: **Endogenous adenosine curtails lipopolysaccharide-stimulated tumour necrosis factor synthesis.** *Scand J Immunol* 1997, **45**:132-139.
- Mabley J, Soriano F, Pacher P, Hasko G, Marton A, Wallace R, Salzman A, Szabo C: **The adenosine A3 receptor agonist, N6-(3-iodobenzyl)-adenosine-5'-N-methyluronamide, is protective in two murine models of colitis.** *Eur J Pharmacol* 2003, **466**:323-329.
- Sullivan GW: **Adenosine A2A receptor agonists as anti-inflammatory agents.** *Curr Opin Investig Drugs* 2003, **4**:1313-1319.
- Montesinos MC, Desai A, Delano D, Chen JF, Fink JS, Jacobson MA, Cronstein BN: **Adenosine A2A or A3 receptors are required for inhibition of inflammation by methotrexate and its analog MX-68.** *Arthritis Rheum* 2003, **48**:240-247.
- Szabo C, Scott GS, Virag L, Egnaczyk G, Salzman AL, Shanley TP, Hasko G: **Suppression of macrophage inflammatory protein (MIP)-1alpha production and collagen-induced arthritis by adenosine receptor agonists.** *Br J Pharmacol* 1998, **125**:379-387.
- Baharav E, Bar-Yehuda S, Madi L, Silberman D, Rath-Wolfson L, Halpren M, Ochaion A, Weinberger A, Fishman P: **The anti-inflammatory effect of A3 adenosine receptor agonists in murine autoimmune arthritis models.** *J Rheumatol* 2005, **32**:469-476.
- Fishman P, Bar-Yehuda S, Madi L, Rath-Wolfson L, Ochaion A, Cohen S, Baharav E: **The PI3K-NF-κB signal transduction pathway is involved in mediating the anti-inflammatory effect of IB-MECA in adjuvant induced arthritis.** *Arthritis Res Ther* 2006, **8**:R33.
- Gessi S, Cattabriga E, Avitabile A, Gafa' R, Lanza G, Cavazzini L, Bianchi N, Gambari R, Feo C, Liboni A, et al.: **Elevated expression of A3 adenosine receptors in human colorectal cancer is reflected in peripheral blood cells.** *Clin Cancer Res* 2004, **10**:5895-5901.
- van Troostenburg AR, Clark EV, Carey WD, Warrington SJ, Kerns WD, Cohn I, Silverman MH, Bar-Yehuda S, Fong KL, Fishman P: **Tolerability, pharmacokinetics and concentration-dependent hemodynamic effects of oral CF101, an A3 adenosine receptor agonist, in healthy young men.** *Int J Clin Pharmacol Ther* 2004, **42**:534-542.
- Nahir MA, Reitblat T, Molad Y, Rosner I, Mader R, Caspi D, Tishler M, Langevitz P, Rubinow A, Green L, et al.: **Targeting the A3 adenosine receptor to combat arthritis: early phase II study in RA patients treated with CF101.** In *Proceedings of the 2005 ACR/ARHP Annual Scientific Meeting; 12-17 November 2005; San Diego, CA, USA* Hoboken, NJ: John Wiley & Sons; 2005:1235.
- Weaver AL, Lautzenheiser RL, Schiff MH, Gibofsky A, Perruquet JL, Luetkemeyer J, Paulus HE, Xia HA, Leff JA: **Real-world effectiveness of select biologic and DMARD monotherapy and combination therapy in the treatment of rheumatoid arthritis: results from the RADIUS observational registry.** *Curr Med Res Opin* 2006, **22**:185-198.
- Allegra CJ, Drake JC, Jolivet J, Chabner BA: **Inhibition of phosphoribosylaminoimidazolecarboxamide transformylase by methotrexate and dihydrofolic acid polyglutamates.** *Proc Natl Acad Sci USA* 1985, **82**:4881-4885.
- Allegra CJ, Hoang K, Yeh GC, Drake JC, Baram J: **Evidence for direct inhibition of de novo purine synthesis in human MCF-7 breast cells as a principal mode of metabolic inhibition by methotrexate.** *J Biol Chem* 1987, **262**:13520-13526.
- Baggott JE, Vaughn WH, Hudson BB: **Inhibition of 5-aminoimidazole-4-carboxamide ribotide transformylase, adenosine deaminase and 5'-adenylate deaminase by polyglutamates of methotrexate and oxidized folates and by 5-aminoimidazole-4-carboxamide riboside and ribotide.** *Biochem J* 1986, **236**:193-200.
- Yaar R, Cataldo LM, Tzatsos A, Francis CE, Zhao Z, Ravid K: **Regulation of the A3 adenosine receptor gene in vascular smooth muscle cells: role of a cAMP and GATA element.** *Mol Pharmacol* 2002, **62**:1167-1176.
- Black RG Jr, Guo Y, Ge ZD, Murphree SS, Prabhu SD, Jones WK, Bolli R, Auchampach JA: **Gene dosage-dependent effects of cardiac-specific overexpression of the A3 adenosine receptor.** *Circ Res* 2002, **91**:165-172.
- Dougherty C, Barucha J, Schofield PR, Jacobson KA, Liang BT: **Cardiac myocytes rendered ischemia resistant by expressing the human adenosine A1 or A3 receptor.** *FASEB J* 1998, **12**:1785-1792.
- Dhalla AK, Shryock JC, Shreenivas R, Belardinelli L: **Pharmacology and Therapeutic application of A1 adenosine receptor ligands.** *Curr Top Med Chem* 2003, **3**:369-385.
- Madi L, Bar-Yehuda S, Barer F, Ardon E, Ochaion A, Fishman P: **A3 adenosine receptor activation in melanoma cells: association between receptor fate and tumor growth inhibition.** *J Biol Chem* 2003, **278**:42121-42130.
- Fishman P, Bar-Yehuda S, Ardon E, Rath-Wolfson L, Barrer F, Ochaion A, Madi L: **Targeting the A3 adenosine receptor for cancer therapy: inhibition of Prostate carcinoma cell growth by A3AR agonist.** *Anticancer Res* 2003, **23**:2077-2083.
- Fredholm BB, Ijzerman AP, Jacobson KA, Klotz KN, Linden J: **International Union of Pharmacology. XXV. Nomenclature and classification of adenosine receptors.** *Pharmacol Rev* 2001, **53**:527-552.
- Schlötzer-Schrehardt U, Zenkel M, Decking U, Haubs D, Kruse FE, Junemann A, Coca-Prados M, Naumann GO: **Selective upregulation of the A3 adenosine receptor in eyes with pseudoexfoliation syndrome and glaucoma.** *Invest Ophthalmol Vis Sci* 2005, **46**:2023-2034.
- Walker BA, Jacobson MA, Knight DA, Salvatore CA, Weir T, Zhou D, Bai TR: **Adenosine A3 receptor expression and function in eosinophils.** *Am J Respir Cell Mol Biol* 1997, **16**:531-537.
- Gessi S, Varani K, Merighi S, Cattabriga E, Avitabile A, Gavioli R, Fortini C, Leung E, Mac Lennan S, Borea PA: **Expression of A3**

adenosine receptors in human lymphocytes: up-regulation in T cell activation. *Mol Pharmacol* 2004, **65**:711-719.