

Plant polycistronic precursors containing non-homologous microRNAs target transcripts encoding functionally related proteins

Francisco Merchan^{*†}, Adnane Boualem[‡], Martin Crespi^{*} and Florian Frugier^{*}

Addresses: ^{*}Institut des Sciences du Végétal (ISV), Centre National de la Recherche Scientifique (CNRS), Avenue de la terrasse, 91198 Gif sur Yvette cedex, France. [†]Current address: Dto Microbiología y Parasitología, Facultad de Farmacia, Universidad de Sevilla, C/Profesor García González 2, 41012 Sevilla, Spain. [‡]Unité de recherche en Génomique Végétale (URGV), Institut National de la Recherche Agronomique (INRA), Rue Gaston Crémieux, 91057 Evry cedex, France.

Correspondence: Adnane Boualem. Email: boualem@evry.inra.fr. Martin Crespi. Email: crespi@isv.cnrs-gif.fr

Published: 1 December 2009

Genome **Biology** 2009, **10**:R136 (doi:10.1186/gb-2009-10-12-r136)

The electronic version of this article is the complete one and can be found online at <http://genomebiology.com/2009/10/12/R136>

Received: 28 May 2009

Revised: 14 September 2009

Accepted: 1 December 2009

© 2009 Merchan *et al.*; licensee BioMed Central Ltd.

This is an open access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/2.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Abstract

Background: MicroRNAs (miRNAs) are endogenous single-stranded small RNAs that regulate the expression of specific mRNAs involved in diverse biological processes. In plants, miRNAs are generally encoded as a single species in independent transcriptional units, referred to as *MIRNA* genes, in contrast to animal miRNAs, which are frequently clustered.

Results: We performed a comparative genomic analysis in three model plants (rice, poplar and *Arabidopsis*) and characterized miRNA clusters containing two to eight miRNA species. These clusters usually encode miRNAs of the same family and certain share a common evolutionary origin across monocot and dicot lineages. In addition, we identified miRNA clusters harboring miRNAs with unrelated sequences that are usually not evolutionarily conserved. Strikingly, non-homologous miRNAs from the same cluster were predicted to target transcripts encoding related proteins. At least four *Arabidopsis* non-homologous clusters were expressed as single transcriptional units. Overexpression of one of these polycistronic precursors, producing Ath-miR859 and Ath-miR774, led to the DCLI-dependent accumulation of both miRNAs and down-regulation of their different mRNA targets encoding F-box proteins.

Conclusions: In addition to polycistronic precursors carrying related miRNAs, plants also contain precursors allowing coordinated expression of non-homologous miRNAs to co-regulate functionally related target transcripts. This mechanism paves the way for using polycistronic *MIRNA* precursors as a new molecular tool for plant biologists to simultaneously control the expression of different genes.

Background

MicroRNAs (miRNAs) are endogenous approximately 21-nucleotide single-stranded small RNAs derived from *MIRNA* precursors that are able to fold-back into a stable secondary structure (stem loop or hairpin). miRNAs act in many developmental processes as well as environmental and pathogenic responses [1-4] through the post-transcriptional regulation of target mRNAs. These targets carry a sequence-specific miRNA recognition site, leading to transcript cleavage and/or inhibition of mRNA translation [1,5,6]. Primary miRNA transcripts (pri-*MIRNA*) are transcribed by RNA polymerase II, and several ribonucleoprotein (RNP) complexes are involved in their maturation, a process that differs between animals and plants [1,6-11]. In animals, formation of an approximately 21-bp miRNA-miRNA* duplex successively involves two RNase III enzymatic complexes: the Drosha enzyme, which cleaves long pri-*MIRNA* in the nucleus to generate short (approximately 70- to 80-nucleotide) hairpins (so called pre-*MIRNA*) and the Dicer enzyme, which produces the miRNA after cytoplasmic export of pre-*MIRNAs* through Exportin 5 [11]. In plants, however, both cleavages are likely nuclear localized and involve a single Dicer-like enzyme 1 (DCL1) complex [6,9,10]. The miRNA-miRNA* duplex is exported to the cytoplasm by HASTY, the plant ortholog of Exportin 5 [12,13]. Subsequently, these duplexes are converted into single-stranded miRNAs upon incorporation into an ARGONAUTE (AGO) ribonucleoprotein complex, referred to as the RNA-induced silencing complex (RISC). The miRNAs guide sequence-specific cleavage and/or translational repression of target transcripts into the RISC complex [6,9-11].

Recent deep sequencing of plant small RNA libraries has led to the identification of more than 1,300 miRNAs in various plants (miRBase, release 13.0, March 2009) [14]. Based on comparison of all available plant genomes (even partial ones; 16 genera referenced in miRBase), evolutionarily conserved and non-conserved miRNAs have been proposed. The non-conserved miRNAs have probably emerged in recent evolutionary time scales, and show a wide diversity compared to the restricted number of conserved miRNAs [15]. Indeed, only 5 miRNA families are found in more than 40 plant species whereas 25 exist in more than one plant genus [16]. The three higher plant models showing the most comprehensive description of their miRNome are rice (*Oryza sativa*; 377 *MIRNAs*), poplar (*Populus trichocarpa*; 234 *MIRNAs*) and *Arabidopsis* (*Arabidopsis thaliana*; 187 *MIRNAs*), with 22 families 'conserved' between them (indicated in bold in Additional data file 1 based on miRBase 13.0). The numerous non-conserved miRNAs are thus likely to play species-specific roles [15].

Plant and animal *MIRNA* genes differ in their genomic location and organization. Most plant miRNAs are encoded in intergenic loci, whereas animal miRNAs are also frequently encoded within introns of protein coding genes [17-19]. Plant

miRNAs are mainly generated from independent transcriptional units, whereas in *Drosophila*, nematodes, zebrafish and mammals, around 40 to 50% of the predicted *MIRNA* genes are located within clusters that are often evolutionarily conserved [18-27]. A maximal distance of 3 kb between two consecutive miRNAs has been used as a stringent criterion to estimate cluster numbers [18]. Clusters in animal genomes usually encode two to three miRNAs but some encode up to eight. Even larger miRNA clusters were predicted in human and zebrafish, containing more than 40 *MIRNA* loci [18,25,26]. In these clusters, miRNAs are encoded either in independent hairpins or sometimes in both arms of the same hairpin [28]. In plants, even though no systematic analysis of miRNA clusters has been performed in the different available genomes, a few miRNA clusters have been reported [16,29-33].

Clustered miRNAs can be either simultaneously transcribed into a single polycistronic transcript or independently transcribed [1,28,34]. Short distances between consecutive *MIRNA* loci and coordinated expression of clustered miRNAs are hallmarks of polycistronic transcription [18,22,34]. Most of the few reported plant miRNA clusters contain several copies of the same conserved miRNA (miR156, miR166, miR169, miR395 or miR399), in contrast to animals where miRNAs with unrelated sequences are often included in the same clusters [18,19,25,35]. Interestingly, certain animal miRNA clusters showing co-regulated expression can simultaneously target transcripts encoding different functionally related proteins. It has been proposed that this may coordinate the fine tuning of the regulation of specific molecular processes [1,18,19,25]. Recently, functional analysis of two human miRNA clusters revealed that the different encoded miRNAs co-regulate related cyclin dependent kinase inhibitors and facilitate cell cycle progression [27]. In plants, beyond the identification of a few expressed sequence tags (ESTs) spanning miRNA clusters [16,29-33], few experimental data indicate that clustered miRNAs are transcribed simultaneously. In the model legume *Medicago truncatula*, a miR166 tandem was shown to be encoded in a single transcriptional unit [32]. However, as both miRNAs are nearly identical, it is difficult to definitively conclude that this pri-*MIRNA* generates more than one miRNA.

In this study, we demonstrate that approximately 20% of plant miRNAs are clustered, and generally contain conserved miRNAs of the same family. Synteny analysis suggested a common evolutionary origin for certain clusters. Strikingly, a few clusters encode tandem non-conserved miRNAs with unrelated sequences, whose predicted targets correspond to transcripts encoding related proteins. In *Arabidopsis*, four of these clusters were transcribed as polycistronic precursors and we show that at least one cluster is processed to form both mature miRNA species in a DCL1-dependent manner. Accumulation of the mature miRNAs affected the stability of their respective predicted target transcripts. Consequently, plant

polycistronic *MIRNA* precursors can encode functional non-homologous miRNAs. This genomic organization may serve to co-regulate different mRNA targets post-transcriptionally.

Results

In silico identification of miRNA clusters in *Arabidopsis*, rice and poplar genomes

A systematic search for consecutive *MIRNA* loci was carried out in three model plant genomes that have an exhaustive description of their miRNA species (miRBase 13.0 [14]). Initially, a 3-kb distance between consecutive *MIRNA* was used as a stringent criterion to define miRNA clusters, similar to previous studies in animals [18,26]. As a result, 16, 10 and 9 clusters were identified in rice, *Arabidopsis*, and poplar, respectively, which represented 13%, 11% and 8% of the total *MIRNA* loci (Table 1; Additional data file 2). Co-expression studies and ESTs available in animal genomes have indicated that some miRNA clusters can be very large; therefore, the 3-kb criterion, which is useful to avoid overestimation of miRNA clusters, is probably too stringent [18,25,35]. Using a less stringent 10-kb cluster size criterion, the number of plant miRNA clusters increased to 18 to 24 in these genomes, thus representing up to 22% of the total *MIRNA* loci (Table 1).

Independently of the size threshold used, most of the clusters (61%, 75% and 90% in *Arabidopsis*, rice and poplar, respectively) contained several copies of the same miRNA family, generally two to three and a maximum of eight (the latter is the rice *Osa-MIR395m-s, x* cluster spanning 497 bp; Table 1; Additional data file 2), and were therefore called homologous clusters. These clusters frequently contained conserved miRNAs (Additional data file 1), and represent 90%, 54% and 44% of the clustered miRNAs in poplar, rice, and *Arabidopsis*, respectively (Table 1). Homologous clusters were found for

miR166, miR169 and miR395 families (based on the <10-kb threshold). This suggested a putative common origin of these clusters, involving successive gene duplications and losses as described for animal miRNA clusters [18,28]. Analysis of these miRNA clusters using VISTA Plot [36,37] (Figure 1) revealed that some were syntenic between monocot and dicot plants (Figure 1a, b, f; two rice miR395 clusters and the *Ath-MIR169i-n* locus) or only within monocot plants (Figure 1c; a third rice miR395 cluster). Surprisingly, no miR395 syntenic locus could be retrieved in the poplar genome (Figure 1a-d). Other miRNA clusters were specific to one plant genome analyzed (Figure 1d, e; the fourth rice miR395 cluster and the *Ath-MIR166c, d* locus). These results suggest that certain ancestral miRNA clusters appeared before the divergence of monocot and dicot lineages and showed differential expansions in the various plant genomes analyzed. Furthermore, clustering of specific miRNAs (for example, miR395, miR169, miR166) is evolutionarily conserved.

Non-homologous miRNAs are expressed as polycistronic pri-MiRNAs

When *MIRNA* loci at a distance of <1 kb on the same DNA strand were considered, 8, 9 and 14 miRNA clusters were identified in *Arabidopsis*, poplar, and rice, respectively (Table 1). In contrast, only four clusters encoding two miRNAs on opposite strands were found (Additional data file 2). These results indicate that short range clustering (based on the 1-kb threshold) is strongly biased (χ^2 test, $P = 9.4 \text{ E-}13$) towards 'same DNA-strand' clustering, suggesting an eventual co-transcription. As small clusters may correspond to polycistronic *MIRNA* precursors, we searched pri-MiRNAs containing several tightly linked miRNAs (Additional data file 2). For example, homologous rice miR395 clusters show the highest number of miRNAs, each encoded in independent stem-loops that are probably generated by successive dupli-

Table 1

Summary of clustered miRNAs in *Arabidopsis thaliana*, rice (*Oryza sativa*) and poplar (*Populus trichocarpa*) genomes²

	<i>Populus trichocarpa</i>	<i>Oryza sativa</i>	<i>Arabidopsis thaliana</i>
Number of clusters with consecutive miRNAs at a distant of (same strand miRNAs):			
<1 kb	9 (9)	16 (14)	10 (8)
<3 kb	14 (14)	16 (14)	13 (11)
<10 kb	20 (20)	24 (22)	18 (16)
% of clustered miRNAs (number of clustered miRNAs/total miRNAs)	20% (47/234)	19% (68/353)	22% (42/187)
Maximal number of miRNAs encoded in a cluster (cluster size)	4 (4,593 bp)	8 (497 bp)	6 (5,522 bp)
% of clusters with (number/total of <10 kb miRNA clusters):			
A single miRNA family	90% (18/20)	75% (18/24)	61% (11/18)
Conserved miRNAs	90% (18/20)	54% (13/24)	44% (8/18)

Analyses were based on miRNA described in miRBase version 13.0 (March 2009).




Figure 1
 Microsynteny between homologous miRNA clusters conserved between *A. thaliana*, rice (*O. sativa*) and poplar (*P. trichocarpa*). VISTA plots [37,70] shows the conservation of different clustered miRNAs in the three selected genomes (Table 1; Additional data file 2): **(a-d)** the four rice miR395 clusters; **(e)** the *Ath-MIR166c, d* cluster; **(f)** the *Ath-MIR169i-n* cluster. To analyze evolutionary conservation between monocots, the sorghum genome is indicated. In each graph, gene models (blue for protein coding gene exons, and red for miRNA clusters) are indicated above, and percentage (50 to 100%) on the right side indicates the level of identity between target and reference genomes, visualized through pink and blue peaks for intronic/non-protein coding, and exonic regions, respectively. The name of syntenic clusters in non-reference genomes is indicated, and syntenic clusters are delimited by black bars. Ath, *Arabidopsis thaliana*; Ptc, *Populus trichocarpa*; Osa, *Oryza sativa*, Sbi, *Sorghum bicolor*.

cations of an ancestral hairpin (Figure 2a; Figure S1 in Additional data file 3). Folding analyses also revealed additional hairpins in the rice *Osa-MIR395m-s* and *Osa-MIR395h-l* clusters containing new miR395 loci not yet listed in miRBase (*Osa-MIR395x* [MiRBase:MI0013350] and *Osa-MIR395y* [MiRBase:MI0013351]; Additional data file 2).

In addition to clusters encoding homologous miRNAs, several clusters (two in rice, three in poplar and five in *Arabidop-*

sis) consisting of miRNAs with unrelated sequences were identified (Table 2). These non-homologous clusters mainly corresponded to non-conserved miRNAs, and their size ranged between 271 and 1,192 bp, with predicted hairpins separated by 81 to 670 bp (median of 114 bp; Additional data file 2), strongly suggesting that these clusters were polycistronic. Most of them were encoded in regions located between protein-coding genes, with the exception of two rice clusters that derive from intronic regions of different putative ribos-




Figure 2
 Representative RNA secondary structure of putative polycistronic clustered miRNAs. Representative examples of (a) a homologous miRNA cluster, *Osa-MIR395m-s, x*, and (b) a non-homologous cluster, *Ath-MIR859-774*. Mfold software [66,67] was used to generate the most probable RNA secondary structures. Mature miRNA sequences are indicated with a line. In the case of the rice cluster, a new miR395 locus (*Osa-MIR395x*) not yet present in miRBase (version 13.0) was annotated. *Ath*, *Arabidopsis thaliana*; *Osa*, *Oryza sativa*. Black bar = 10 nucleotides (nt).

omal protein encoding genes (ESTs listed in Table 2). In most cases, non-homologous clusters encoded two miRNAs, with the exception of the *Osa-MIR1876-1862d-1884b* cluster. Similarly to the homologous miRNA clusters, this novel *MIRNA* precursor class contained several hairpins, each carrying a single annotated miRNA in the stem (Figure 2b; Figures S2 and S3 in Additional data file 3). A majority of 21- to 22-bp species was found for each predicted miRNA in *Arabidopsis* non-homologous clusters (Genome View browser, *Arabidopsis* Small RNA Project (ASRP) database) [38-40], and we systematically annotated miRNA* to identify *bona fide* miRNAs [41] (Figure S4 in Additional data file 3). However, for *Ath-MIR859*, *Ath-MIR397b* and *Ath-MIR857* loci, no miRNA* could be identified in small RNA databases.

Currently, the *Arabidopsis* genomic regions corresponding to candidate polycistronic non-homologous *MIRNA* genes are represented as independent transcriptional units in the ASRP database (Figure S4 in Additional data file 3). Indeed, no EST comprising tandem miRNAs was available in *Arabidopsis*. ESTs spanning the two *MIRNA* hairpins were reported only for the poplar *Ptc-MIR482-1448* locus (Table 2), indicating that these two miRNAs are indeed co-transcribed. To determine whether *Arabidopsis* non-homologous miRNA clusters are encoded into single polycistronic units, WWwe performed semi-quantitative RT-PCR experiments (Figure 3). Primers designed to amplify precursors spanning the two predicted hairpins revealed expression for *Ath-MIR859-774*, *Ath-MIR850-863*, *Ath-MIR851-771* and *Ath-MIR397b-857* in

Table 2**Summary of targets predicted for non-homologous putative polycistronic clustered miRNAs in *Arabidopsis*, rice and poplar**

Non-homologous MIRNA clusters	ESTs overlapping the cluster	Targets	
		Predicted	Cleavage validated for at least one target [reference]*
<i>Arabidopsis</i>			
Ath-MIR397b-857	No	Laccases (3/1)	Yes [15,49,50]
Ath-MIR859-774	No	F-box proteins (35/5)	Yes [15,48]
Ath-MIR842-846	No	JR/MBP (1/10)	Yes [15]
Ath-MIR851-771	No	EIF2 for miR771 (1) no target for miR851	No [15]
Ath-MIR850-863	EG495879 EG495880 EG514601 DR380439, AV523115†	No target	-
<i>Poplar</i>			
Ptc-MIR482-1448	DB893204 DB893031 BP928209 CX175581	Disease resistance proteins (12/2)	Yes [44]
Ptc-MIR1446b-477a	No	Gibberellin response modulator-like protein (DELLA) (1)	Yes [43,44]
Ptc-MIR1446a-477b	No	Gibberellin response modulator-like protein (DELLA) (1)	Yes [43,44]
<i>Rice</i>			
Osa-MIR1876-1862d-1884b	Os10g0170200‡	No target	-
Osa-MIR1423-1868	Os04g32710‡	No target	-

For predicted targets, the number of targets found for each miRNA of the non-homologous cluster is indicated in parentheses. *Validated using 5' RACE (rapid amplification of cDNA ends) PCR. †ESTs cover the region including only miR863. ‡The consecutive miRNAs are predicted in the intron of a protein coding gene (putative ribosomal protein). Ath, *A. thaliana*; Osa, *O. sativa*; Ptc, *P. trichocarpa*.


seedlings (Figure 3a; Figure S4 in Additional data file 3, with RT-PCR amplified regions indicated as ESTs with green arrows). Cloning and subsequent sequencing of RT-PCR products confirmed the specific amplification of these pri-MIRNAs ([NCBI:GU125419] for Ath-MIR859-774; [NCBI:GU125420] for Ath-MIR850-863; [NCBI:GU125421] for Ath-MIR397b-857; [NCBI:GU125422] for Ath-MIR851-771). However, no amplification was obtained for the Ath-MIR842-846 cluster (even using various primer combinations). Expression of these four polycistronic miRNAs in different *Arabidopsis* organs revealed specific patterns: Ath-MIR859-774 pri-MIRNA was only detectable in roots, Ath-MIR397b-857 in roots and cauline leaves, Ath-MIR851-771 in aerial parts, including flowers, and Ath-MIR850-863 mainly in rosette leaves (Figure 3b).

Overall, these results suggest that clusters comprising functional miRNAs with unrelated sequences exist in plants as single transcriptional units, defining a novel class of plant pri-MIRNA genes.

Polycistronic non-homologous miRNAs regulate related targets

In contrast to animals, *in silico* predictions revealed only a few targets for each plant miRNA based on strong sequence complementarity [42]. Strikingly, all predicted targets corresponding to different miRNAs from the same cluster encode

proteins of the same family (Table 2; complete target list in Additional data file 4; based on the ASRP database for *Arabidopsis*, and on [43-45] for poplar). Indeed, the Ath-MIR397b-857 locus encodes two miRNAs that regulate laccases (three and one gene, respectively), the Ath-MIR842-846 locus encodes two miRNAs that regulate JR/MBP proteins (Jacalin repeat/Myrosinase binding protein; one and ten genes, respectively), and the Ath-MIR859-774 locus encodes two miRNAs that regulate F-box proteins (35 and 5 genes, respectively). More importantly, three of these F-box proteins are likely to be targeted by both Ath-miR859 and Ath-miR774 (Additional data file 4). Similarly, the Ptc-MIR482-1448 locus encodes miRNAs that regulate disease resistance proteins (12 and 2 genes, respectively, based on various gene models [46]), and one of them is probably co-regulated by the two miRNAs (Additional data file 4). Finally, the two Ptc-MIR1446-477 loci encode miRNAs that both target a single 'gibberellin response modulator-like protein' homologous to the *Arabidopsis* RGL1/RGL2 (Repressor of gibberellic acid requiring (GA1)-LIKE; DELLA transcription factors [45]). For the other non-homologous clustered miRNAs, targets were either predicted for only one miRNA of the tandem (for example, a single EIF2 encoding transcript - The *Arabidopsis* Information Resource database entry TAIR:At1g76810 - for Ath-miR771), or no target could be identified (Table 2). Target validation based on 5' RACE (5' rapid amplification of cDNA ends) PCR experiments was determined in *Arabidop-*

**Figure 3**

Expression of *Arabidopsis* polycistronic non-homologous miRNA clusters and selected targets in different organs. **(a)** Detection by RT-PCR analysis of the expression of *Arabidopsis* polycistronic non-homologous clusters as single transcriptional units: Ath-*MIR859-774*, Ath-*MIR397b-857*, Ath-*MIR850-863*, and Ath-*MIR851-771*. Total RNAs from wild-type (Col-0) seedlings were used for DNase I treatment and cDNA synthesis to perform RT-PCR reactions. A control without reverse transcriptase (-RT) was systematically included to check for the absence of genomic DNA. Specificity of PCR amplicons was verified by sequencing. **(b)** Expression analysis by RT-PCR of the different *Arabidopsis* polycistronic non-homologous clusters (Ath-*MIR859-774*, Ath-*MIR397b-857*, Ath-*MIR850-863*, and Ath-*MIR851-771*) in different organs (R, roots; RL, rosette leaves; S, stems; CL, cauline leaves; F, flowers). **(c)** Expression analysis by RT-PCR of selected Ath-*MIR859-774* targets in the same organs (roots, rosette or cauline leaves, stems, flowers) as in (b). The *Arabidopsis* Information Resource database entry TAIR:At3g49510 was used as an Ath-miR859 validated target (by 5' RACE (rapid amplification of cDNA ends) PCR) [15], and TAIR:At3g19890 for Ath-miR774 [48]. At3g18780 encoding an *ACTIN* isoform was used as RNA loading control.

sis (AtPARE database) [47] for Ath-miR859 [TAIR:At3g49510] [15], Ath-miR774 [TAIR:At3g19890] [48], Ath-miR397 ([TAIR:At2g29130], [TAIR:At5g60020] and [TAIR:At2g38080]) [49,50], Ath-miR857 [TAIR:At3g09220], Ath-miR842 [TAIR:At5g38550] and Ath-miR846 [TAIR:At5g49850] [15] (Table 2). In poplar, Ptc-miR477/Ptc-miR1446-mediated cleavage of the DPTF:fgenes4_pg.C_LG_XII000915 target (from the Database of Poplar Transcription Factors) was validated for each miRNA, as well as the JGI-Ptr-v1.1:eugene3.00102261 target for Ptc-miR482 and JGI-Ptr-v1.1:eugene3.01310091 for Ptc-

miR1448 (both from the Joint Genome Institute poplar database) [43,44,46].

These analyses suggest that non-homologous miRNA polycistronic clusters are likely to target transcripts encoding proteins of the same family, suggesting that co-transcription of miRNAs may co-regulate their respective targets.

The polycistronic Ath-*MIR859-774* pri-MIRNA is processed by a DCL1-dependent pathway

To determine the functionality of a non-homologous polycistronic pri-MIRNA *in planta*, the Ath-*MIR859-774* locus was selected. Expression of several Ath-*MIR859-774* predicted targets encoding F-box proteins was analyzed in different organs (Figure 3c). Among validated targets, the Ath-miR774 target TAIR:At3g19890 and the Ath-miR859 target TAIR:At3g49510 exhibited detectable expression in roots that also express Ath-*MIR859-774* pri-MIRNA. This indicates that both partners of this post-transcriptional regulation are present in this organ. We then overexpressed this precursor to analyze the transcriptional regulation of these target genes (Figure 4). Independent lines accumulating the pri-MIRNA transcripts at high levels in wild-type (Col-0) *Arabidopsis* plants were obtained (Figure 4a). Northern blot analyses showed accumulation of both mature approximately 21-bp miRNAs corresponding to Ath-miR859 and Ath-miR774 in comparison to control plants (expressing an empty vector; Figure 4b). In addition, significant down-regulation of both the Ath-miR859 target TAIR:At3g49510 and the Ath-miR774 target TAIR:At3g19890 was observed in these transgenic lines (Figure 4c). These results indicate that both miRNAs from the Ath-*MIR859-774* polycistronic pri-MIRNA can be processed and simultaneously co-regulate the expression of their respective predicted targets.

To determine whether the maturation of this atypical polycistronic MIRNA precursor depends on DCL1, which processes canonical single MIRNA precursors, the Ath-*MIR859-774* overexpression construct was introduced into a *dcl1-9* mutant. This weak DCL1 allele affects miRNA processing but is more viable than the embryonic lethal *dcl1* null alleles [51]. Several independent lines overexpressing Ath-*MIR859-774* did not accumulate Ath-miR859 and Ath-miR774 in aerial tissues of the homozygous *dcl1-9* background (*dcl1-9/dcl1-9*), in contrast to the heterozygous siblings (*dcl1-9/DCL1*); Figure 5a, b). This indicates that the processing of polycistronic MIRNA precursors such as pri-*MIR859-774* requires DCL1. Furthermore, down-regulation of the Ath-miR859 and Ath-miR774 targets was abolished in these *dcl1-9/dcl1-9* transgenic lines (Figure 5c).

Discussion

A comparative genomic analysis of miRNA clustering in three model plants (a monocot, rice, a herbaceous dicot, *Arabidopsis*, and a dicot tree, poplar) led us to identify a novel class of




Figure 4
 Ath-*MIR859-774* encodes a functional polycistronic pri-*MIRNA*. **(a)** Overexpression of Ath-*MIR859-774* pri-*MIRNA* (under the control of a 35S-CaMV promoter) in wild-type Col-0 transgenic lines (seven representative OverExpressing (OE) lines) is shown. Real-time RT-PCR analysis was performed based on equal amounts of total RNAs. A green fluorescent protein encoding transcript present 3' to the overexpressed pri-*MIRNA* was used as a tag to efficiently amplify the overexpressed transcript. Threshold cycles are indicated and error bars indicate standard deviation of values obtained based on two independent cDNA syntheses (technical replicates). **(b)** Detection of mature 21-bp Ath-miR859 and Ath-miR774 by Northern blot analysis in two representative independent transgenic lines (selected from (a)) overexpressing the Ath-*MIR859-774* pri-*MIRNA*. Total RNAs (10 µg/lane) from each sample were blotted and probed with ³²P-labeled standard DNA complementary to *U6* RNA (loading control) or oligonucleotides complementary to mature miRNA sequences. **(c)** Real-time RT-PCR analysis of the relative accumulation of selected Ath-*MIR859-774* validated targets (Ath-miR859 target TAIR:At3g49510 and Ath-miR774 target TAIR:At3g19890) in the two transgenic lines used in (b). The histogram represents amounts of specific PCR amplification products (verified by sequencing and dissociation curve analyses) normalized to reference genes [72] defined using Genorm software [73] (see Materials and methods). Values of target expression in the control line (Col-0) are set to 1. Error bars indicate the standard deviation of values obtained based on three independent cDNA syntheses (technical replicates).

polycistronic *MIRNA* precursors encoding miRNAs with unrelated sequences. These non-homologous miRNA clusters target transcripts encoding proteins of the same family, suggesting that this unusual genomic organization may allow co-regulation of different but related targets.

Most miRNA clusters encode several copies of conserved miRNAs from the same family, that is, miR166, miR169, or miR395. Previous analyses of miR395 clusters in rice and *M. truncatula*, as well as a miR156 cluster in rice, maize, sugar-

cane, sorghum and even a dicot (*Ipomea nil*), have suggested conservation of homologous miRNA clusters in various plant genomes [16,29,30]. Our analysis revealed certain homologous miRNA clusters at syntenic genomic positions, implying a common evolutionary origin across monocot and dicot lineages. Specific miRNA families seem positively selected for expansion and clustering in several genomes. For cultivated species, it has been proposed that this spreading may contribute to advantageous agricultural traits [29,30]. In addition, homologous miRNAs or cluster duplication may lead to the




Figure 5
Ath-MIR859-774 polycistronic pri-MIRNA is processed by a DCL1-dependent pathway. (a) Overexpression of Ath-MIR859-774 pri-MIRNA in *dcl1.9* mutant and in the corresponding *DCL1.9/dcl1.9* heterozygous siblings (three representative OverExpressing (OE) transgenic lines) is shown. Real-time RT-PCR analysis was done on equal amounts of total RNAs. A green fluorescent protein encoding transcript present downstream of the overexpressed pri-MIRNA was used to efficiently detect the overexpressed transcript. Threshold cycles are indicated and error bars indicate standard deviation of values obtained based on two independent cDNA syntheses (technical replicates). (b) Detection of mature 21-bp Ath-miR859 and Ath-miR774 by Northern blot analysis in the three independent transgenic lines (shown in (a)) over-expressing Ath-MIR859-774 pri-MIRNA in *dcl1.9* and in the *DCL1.9/dcl1.9* heterozygous siblings. Total RNAs (10 µg/lane) from each sample were blotted and probed with ³²P-labeled standard DNA complementary to U6 RNA (loading control) or modified oligonucleotides complementary to mature miRNA sequences. (c) Real-time RT-PCR analysis of the relative accumulation of selected Ath-MIR859-774 validated targets (Ath-miR859 target TAIR:At3g49510 and Ath-miR774 target TAIR:At3g19890) in the three transgenic lines (used in (a, b)) overexpressing the corresponding pri-MIRNA in *dcl1.9* and in the related *DCL1.9/dcl1.9* heterozygous background. The histogram represents amounts of specific PCR amplification products (verified by sequencing and dissociation curve analyses) normalized to reference genes [72] defined using Genom software [73] (see Materials and methods). Error bars indicate the standard deviation of values obtained based on three independent cDNA syntheses (technical replicates).

emergence of new spatio-temporal expression patterns through the accommodation of alternative promoter regions [29,52,53].

A combination of tandem duplication of miRNAs as well as segmental duplications of whole clusters has been proposed to explain such genomic organization [29,52]. In animal genomes, miRNAs encoded in the miR17 cluster arose through a complex duplication and loss of individual members as well as duplications of entire clusters [28]. In plant genomes, miR156, miR160, miR162, miR167, miR169, miR171 and miR395 families experienced large expansions

via tandem or segmental duplication events and loss of family members ([29,30,52] and this study). This is in agreement with the detection of two to three miRNAs in most (that is, 70 to 80%) of the clusters in our analysis, similar to protein coding gene clusters [52]. These duplication events may therefore represent a major evolutionary route for birth and death of miRNAs in plants.

Folding of putative transcripts derived from homologous miRNA clusters revealed additional hairpins in the rice *Osa-MIR395h-l* and *Osa-MIR395m-s* clusters, which were not annotated in miRBase. In animal genomes, systematic fold-

ing of genomic regions encoding miRNA clusters has helped to identify additional miRNAs [18]. A recent analysis of rice miRNA clusters has revealed a different genomic organization of upstream sequences corresponding to their promoters [53]. *Osa-MIR156b-c*, *Osa-MIR166k-h*, *Osa-MIR169n-o*, *Osa-MIR172b-806a*, *Osa-MIR395a-g*, *Osa-MIR395h-l*, and *Osa-MIR395m-s* clusters may contain only one promoter and be transcribed as polycistronic units. Interestingly, we found that the *Osa-MIR395t-w* cluster was specific to the rice genome. This cluster has previously been reported as having no predicted promoter [53]; a transposable element identified in its vicinity [29] may be associated with the recent evolution of this *MIRNA* cluster [6,54].

Our results indicate that short range (<1 kb) clustering of 'same DNA strand' miRNAs are highly suggestive of co-transcription as reported in animal genomes [18]. Accidental formation of hairpins followed by loss of miRNAs subsequent to duplication was indeed proposed as a general mechanism for the origin of polycistronic *MIRNA* transcripts in animals [28]. Although the clustered miRNAs characterized were always encoded in independent hairpins, a stem-loop encoding the rice miR159 was recently shown to produce additional approximately 21- to 24-nucleotide small RNAs from the 21 bp next to the miR159 sequence [33]. This unusual case is reminiscent of sequential DCL1-dependent processing of the *Arabidopsis* miR163, and of DCL4-dependent processing of tasiRNAs (trans-acting siRNAs) or young *Arabidopsis* miRNAs, which may correspond to 'proto-miRNAs' [6,55,56]. Our results show that maturation of the *Ath-MIR859-774* polycistronic cluster is mediated by DCL1, but we cannot exclude that other (DCL) enzymatic complexes may contribute to the processing of polycistronic *MIRNA* precursors.

In each of the three plant genomes, we identified several clusters encoding distinct miRNAs, in addition to clusters containing homologous miRNAs. Their low abundance in plant genomes contrasts with animal genomes, where miRNA clusters frequently encode miRNAs from different, although evolutionarily related, families, for example, the miR17 gene cluster [25,28]. These clustered non-homologous *MIRNA* genes are proposed to simultaneously regulate multiple functionally related genes in animals. Indeed, a recent study has demonstrated that two human miRNA clusters regulate various cyclin dependent kinase inhibitors, leading to a coordinated regulation of cell cycle progression [27]. In contrast to animals where hundreds of translational targets are frequently predicted for a single miRNA, plant miRNAs target few transcripts, usually showing an extensive homology with the miRNA leading to its cleavage [42] (Table 2; Additional data file 4). Although the recent identification of translational regulation in plants may affect this view, all known translationally regulated targets presently contain binding sites highly homologous to miRNAs [5]. Interestingly, we show that all predicted targets of the different non-homologous

miRNAs present in a single cluster always corresponded to proteins of the same family.

Ath-MIR859-774 and a representative target of each miRNA were mainly expressed in the roots. However, anti-correlation between the *MIRNA* precursor and target transcript levels was not identified in the different organs tested. This could be due to the fact that several plant miRNAs quantitatively regulate gene expression and a low level of variation in a specific organ could not be detected [6]. Indeed, both miRNA and targets were expressed at low levels in each organ tested (Figure 3; Figure S4 in Additional data file 3). Additionally, spatial expression domains of the miRNAs and their targets may vary in the different cell types constituting an organ, resulting in non-significant differences at the whole organ level or even positive correlations ([6,57] and references therein). Furthermore, post-translational regulations may be superimposed upon post-transcriptional regulations, as in the case of another recently evolved plant miRNA, miR834, initially suspected to be inactive [5]. In the latter case, the absence or near absence of transcriptional anti-correlation between miRNA and target transcripts suggests that post-translational regulation is predominant over mRNA regulation.

Ectopic expression of the *Ath-MIR859-774* pri-*MIRNA* led to the simultaneous down-regulation of distinct F-box transcripts, which are likely to be independently regulated by each miRNA. F-box proteins co-regulated by *Ath-MIR859-774* may participate in specific pathways involving proteasome-dependent degradation of signaling components [58]. *Ptc-MIR1446-477* loci are predicted to target a DELLA-like transcription factor similar to the *Arabidopsis* RGL1/RGL2 proteins involved in gibberellin control of seed germination and floral development [45], and shoot and root development in poplar [59]. The *Ath-MIR397b-857* targets transcripts encoding laccase copper proteins associated with lignin synthesis, metal nutrition and response to abiotic stresses [50,60]. Among the four laccase encoding transcripts targeted by these miRNAs, the knock-out mutant of TAIR:At2g29130 (*AtLAC2*) shows slightly reduced root elongation under osmotic stress. Finally, miRNAs derived from the *Ath-MIR842-846* loci target transcripts encoding related JR/MBP, while *Ptc-MIR482-1448* miRNAs target transcripts encoding disease resistance proteins. Both pathways may affect pathogen defense responses [44,61]. Co-transcription of similar or identical miRNAs has been proposed to have a dosage effect on target expression [29]. Co-expression of different miRNAs may serve to increase the efficiency of the regulatory process. Whereas different miRNAs have been shown to bind a single mRNA target in animal systems to cooperatively control its expression [19,62], only three *Ath-MIR859-774* targets were predicted to be recognized by both miRNAs. This result might be biased due to the restrictive criteria used in plants to predict targets, in contrast to animal genomes [42].

MIRNA genes are proposed to originate from the duplication of a target gene [6,15,56,63]. In the case of polycistronic non-homologous *MIRNA* precursors, we could hypothesize that the duplication of a single target locus may have led to the selection of two divergent 'proto-miRNA' regions targeting other members of the family. An alternative is the duplication of an overlapping region between two clustered target genes, leading to the selection of miRNAs that target both clustered ancestral genes. Indeed, predicted targets of tandem polycistronic non-homologous *MIRNA* precursors are often themselves clustered (Additional data file 4), notably the F-box proteins targeted by *Ath-MIR859-774* and the laccases targeted by *Ath-MIR397b-857* (37 clustered F-box proteins and 4 clustered laccases).

Our results show that plant genomes generally contain less clustered or polycistronic miRNAs than animal genomes. Indeed, approximately 20% of total plant miRNAs are clustered, whereas in animals they represent approximately 50% using a similar criterion (that is, cluster size up to 10 kb) [18]. In animals, the Drosha complex specifically catalyzes maturation of long pri-*MIRNA*s, including the numerous polycistronic clusters, into approximately 70 nucleotide pre-*MIRNA*s hairpins [6]. In plants, however, a Drosha-like enzyme is lacking. We have shown that the processing of at least one *Arabidopsis* polycistronic *MIRNA* is DCL1-dependent, similar to most non-polycistronic *MIRNA* precursors. We can speculate that the occurrence of a single step maturation process of polycistronic precursors in plants may not be functionally equivalent to the two-step process existing in animals.

Conclusions

In contrast to plants, clusters of miRNAs are frequently present in animal genomes. Our comparative genomic analysis in three model plants (rice, poplar and *Arabidopsis*), however, has demonstrated the presence of several clusters containing two to eight miRNA species. Certain ancestral miRNA clusters appeared before the divergence of monocot and dicot lineages, and showed differential expansions in plants. Specific miRNA clusters (such as those coding for miR395, miR169 and miR166) are highly conserved. Interestingly, other clusters comprise functional miRNAs with unrelated sequences (non-homologous miRNAs) and are expressed as single transcriptional units, defining a novel class of plant pri-*MIRNA* genes. These polycistronic non-homologous miRNAs regulate related target genes and are processed by a DCL1-dependent pathway. This mechanism paves the way for using polycistronic *MIRNA* precursors as a new molecular tool in plants to simultaneously express artificial miRNAs [64] that control the expression of different genes.

Materials and methods

Plant genotypes and growth conditions

The wild-type Columbia (Col-0) ecotype of *A. thaliana* was used, as well as a *dcl1-9* mutant backcrossed five times to Col-0 [51,56]. All plants were grown in long day conditions (16-h light/8-h dark photoperiod) at 23°C. Inflorescences, stems and cauline leaves, or rosette leaves were collected from 3-week-old greenhouse-grown plants. Roots were collected from seedlings grown 3 weeks *in vitro* on 1/2 Murashige and Skoog (MS) medium (Sigma, Lyon, France) supplemented with 1% sucrose (Sigma, Lyon, France).

Bioinformatic analysis

Arabidopsis, poplar and rice miRNA sequences (mature and precursor) were downloaded from the microRNA Registry version 13.0 [65]. miRNA coordinates, chromosome locations and DNA strand orientation were retrieved from the microRNA Registry.

MIRNA genes were sorted by their chromosome locations and coordinates to identify miRNA clusters. The distance between two consecutive *MIRNA* loci was calculated by subtracting the start coordinates of the downstream pre-*MIRNA* (that is, hairpin) to the end coordinates of the upstream pre-*MIRNA*. *MIRNA* loci located within a distance of less than 1, 3 or 10 kb were considered to define the best candidates for polycistronic clusters and clusters with stringent or non-stringent criteria, respectively. The DNA strand containing the miRNA sequence was considered in these analyses.

Secondary structures were predicted using the mfold program [66] with default parameters [67] and a window size between 1 and 2 kb depending on *MIRNA* precursors.

Conservation analysis of miRNA clusters between plant genomes

Conservation between selected clustered miRNAs in *Arabidopsis*, poplar, sorghum and rice genomes as well as determination of candidate orthologous regions were determined using Genome VISTA [36,37]. Query sequence (1 to 1.5 kb depending on clusters) was anchored on the reference genome by local alignment matches and then globally aligned to candidate regions in different selected genomes based on the AVID program [68,69]. Alignments were then displayed with the VISTA graphic server [70]. Identified syntenic regions were manually inspected to identify and annotate orthologous miRNA clusters.

Northern blot analysis of small RNA expression

Tissues were frozen in liquid nitrogen, ground to a fine powder with a mortar and pestle, and then homogenized in TRI-Reagent® (Sigma, Lyon, France) supplemented with β -mercaptoethanol. Total RNAs were prepared according to the manufacturer's instructions (Sigma, Lyon, France) with additional steps: samples were extracted with one volume of Tris/HCl-buffered phenol/chloroform (Sigma, Lyon, France),

then with two volumes of chloroform, and finally RNAs were precipitated with three volumes of ice-cold 100% ethanol and one-tenth volume of 3 M sodium acetate (pH 6) in diethylenepycarbonate (DEPC)-treated water. Northern blot analysis of low molecular weight RNAs (10 µg of total RNAs per lane) was carried out on denaturing 15% polyacrylamide (19:1) gels cast in 7 M urea/Tris borate EDTA (TBE) buffer, followed by blotting onto a nylon hybridization membrane (Hybond-NX®, Amersham/Pharmacia, Les Ulis, France) pre-wetted in distilled water. An EDC (1-ethyl-3-[3-dimethylaminopropyl]carbodiimide hydrochloride)-mediated cross-linking step was then performed as described [71]. Blots were hybridized with gamma-ATP ³²P end-labeled oligonucleotides (20 pmoles) complementary to miRNAs, and at the same time with an end-labelled oligonucleotide U6 RNA probe as loading control.

Analysis of gene expression by RT-PCR

Total RNAs were extracted using the total RNA Isolation kit (Macherey-Nagel, Düren Germany). cDNA was synthesized by reverse transcription of 1.5 µg of total RNAs using the SuperScript II Reverse Transcriptase (Invitrogen, Paisley, UK) and (T)16 A/G/C oligonucleotides. Primer pairs used for RT-PCR are listed in Additional data file 5. Specificity of amplification was checked by cloning and sequencing of PCR amplicons, and ESTs corresponding to *Arabidopsis* non-homologous pri-MIRNAs were submitted to GenBank ([NCBI:GU125419] for Ath-MIR859-774; [NCBI:GU125420] for Ath-MIR850-863; [NCBI:GU125421] for Ath-MIR397B-857; [NCBI:GU125422] for Ath-MIR851-771). For Ath-MIR842-846 loci, no amplification was obtained despite testing eight different primer combinations, even on genomic DNA (data not shown). A control without reverse transcriptase was systematically included.

Real-time RT-PCR was performed on an Eppendorf Mastercycler® realplex real-time PCR system (Eppendorf, Hamburg, Germany) using FastStart Universal SYBR Green Master Mix (Rox) from Roche Applied Science (Meylan, France). Technical triplicates were done for each datapoint, and two independent biological replicates (per condition and/or transgenic line) were assayed. Normalization was done with averaged reference genes TAIR:At1g13320, TAIR:At4g26410, and TAIR:At5g15710 [72], which were systematically validated under our experimental conditions using Genorm software [73].

Cloning and transgenic plants

Firstly, pri-MIRNA Ath-MIR859-774 was amplified by RT-PCR from seedling cDNA and cloned into pCR8®/GW/TOPO® TA Cloning® vector (Invitrogen, Paisley, UK). The construct was then transferred to the destination vector pEarlyGate103 [74] using the LR recombination kit (Invitrogen, Paisley, UK). These constructions (based on the 35S-CaMV promoter) were used to transform *A. thaliana* plants by floral dipping [75]. Transgenic plants were selected in T1

generation by spraying seedlings with Basta® solution (120 mg/L glufosinate ammonium; Bayer CropScience, Monheim am Rhein, Germany) successively at 12, 14, and 16 days after germination. Basta-resistant plantlets were then tested for transgene expression by real time RT-PCR as described above. Since amplification across the successive hairpin regions of the Ath-MIR859-774 pri-MIRNA was not efficient and quantitative enough for real time RT-PCR analyses, a GFP mRNA present 3' of the pEarlyGate103 vector cloning site, for which efficient and specific primers were available (Additional data file 5), was used as a 3' tag to analyze transgene expression.

Abbreviations

ASRP: *Arabidopsis* Small RNA Project; Ath: *Arabidopsis thaliana*; DCL1: Dicer-like enzyme 1; EST: expressed sequence tag; JR/MBP: Jacalin repeat/Myrosinase binding protein; miRNA: microRNA; Osa: *Oryza sativa*; pre-MIRNA: miRNA precursor; pri-MIRNA: miRNA primary transcript; Ptc: *Populus trichocarpa*; RGL: Repressor of gibberellic acid requiring (GA1)-LIKE; RISC: RNA-induced silencing complex; TAIR: The *Arabidopsis* Information Resource.

Authors' contributions

FM carried out the molecular genetic studies. AB conceived the study and carried out the bioinformatic analyses. MC drafted the manuscript and participated in its coordination. FF conceived the study, designed experiments and wrote the manuscript. All authors read and approved the final manuscript.

Additional data files

The following additional data are available with the online version of this paper: a table listing the conserved and non-conserved miRNAs in *Arabidopsis*, rice and poplar genomes (Additional data file 1); a table providing a detailed list of clustered miRNAs in *Arabidopsis*, rice and poplar genomes (Additional data file 2); a PDF including Figures S1 to S4 (Additional data file 3); a table providing a detailed list of all targets predicted for *Arabidopsis* non-homologous polycistronic miRNA clusters (Additional data file 4); a table listing primers used in this study (Additional data file 5).

Acknowledgements

We thank Hervé Vaucheret for providing *dcl1-9* seeds in Col-0 background, and for the careful reading of this manuscript. Laëticia Riva-Roveda is acknowledged for her help in genotyping transgenic plants and on initial validation of non-homologous polycistronic MIRNA precursors. Alexis Maizel provided helpful suggestions for the manuscript. FM was funded by a post-doctoral grant from the Spanish Department of Education and Science and from Sevilla University.

References

- Bartel DP: **MicroRNAs: genomics, biogenesis, mechanism, and function.** *Cell* 2004, **116**:281-297.
- Mallory AC, Vaucheret H: **Functions of microRNAs and related small RNAs in plants.** *Nat Genet* 2006, **38**(3):31-36.
- Jones-Rhoades MW, Bartel DP: **Computational identification of plant microRNAs and their targets, including a stress-induced miRNA.** *Mol Cell* 2004, **14**:787-799.
- Xiao C, Rajewsky K: **MicroRNA control in the immune system: basic principles.** *Cell* 2009, **136**:26-36.
- Brodersen P, Sakvarelidze-Achard L, Bruun-Rasmussen M, Dunoyer P, Yamamoto YY, Sieburth L, Voinnet O: **Widespread translational inhibition by plant miRNAs and siRNAs.** *Science* 2008, **320**:1185-1190.
- Voinnet O: **Origin, biogenesis, and activity of plant microRNAs.** *Cell* 2009, **136**:669-687.
- Chapman EJ, Carrington JC: **Specialization and evolution of endogenous small RNA pathways.** *Nat Rev Genet* 2007, **8**:884-896.
- Mallory AC, Elmayan T, Vaucheret H: **MicroRNA maturation and action - the expanding roles of ARGONAUTES.** *Curr Opin Plant Biol* 2008, **11**:560-566.
- Chen X: **MicroRNA metabolism in plants.** *Curr Top Microbiol Immunol* 2008, **320**:117-136.
- Ramachandran V, Chen X: **Small RNA metabolism in Arabidopsis.** *Trends Plant Sci* 2008, **13**:368-374.
- Kim VN, Han J, Siomi MC: **Biogenesis of small RNAs in animals.** *Nat Rev Mol Cell Biol* 2009, **10**:126-139.
- Bollman KM, Aukerman MJ, Park MY, Hunter C, Berardini TZ, Poethig RS: **HASTY, the Arabidopsis ortholog of exportin 5/MSN5, regulates phase change and morphogenesis.** *Development* 2003, **130**:1493-1504.
- Park MY, Wu G, Gonzalez-Sulser A, Vaucheret H, Poethig RS: **Nuclear processing and export of microRNAs in Arabidopsis.** *Proc Natl Acad Sci USA* 2005, **102**:3691-3696.
- Griffiths-Jones S, Saini HK, van Dongen S, Enright AJ: **miRBase: tools for microRNA genomics.** *Nucleic Acids Res* 2009, **36**:D154-D158.
- Fahlgren N, Howell MD, Kasschau KD, Chapman EJ, Sullivan CM, Cumbie JS, Givan SA, Law TF, Grant SR, Dangel JL, Carrington JC: **High-throughput sequencing of Arabidopsis microRNAs: evidence for frequent birth and death of MIRNA genes.** *PLoS ONE* 2007, **2**:e219.
- Sunkar R, Jagadeeswaran G: **In silico identification of conserved microRNAs in large number of diverse plant species.** *BMC Plant Biol* 2008, **8**:37.
- Rodriguez A, Griffiths-Jones S, Ashurst JL, Bradley A: **Identification of mammalian microRNA host genes and transcription units.** *Genome Res* 2004, **14**:1902-1910.
- Altuvia Y, Landgraf P, Lithwick G, Elefant N, Pfeffer S, Aravin A, Brownstein MJ, Tuschl T, Margalit H: **Clustering and conservation patterns of human microRNAs.** *Nucleic Acids Res* 2005, **33**:2697-2706.
- Kim VN, Nam JW: **Genomics of microRNA.** *Trends Genet* 2006, **22**:165-173.
- Lagos-Quintana M, Rauhut R, Lendeckel W, Tuschl T: **Identification of novel genes coding for small expressed RNAs.** *Science* 2001, **294**:853-858.
- Lau NC, Lim LP, Weinstein EG, Bartel DP: **An abundant class of tiny RNAs with probable regulatory roles in Caenorhabditis elegans.** *Science* 2001, **294**:858-862.
- Mourelatos Z, Dostie J, Paushkin S, Sharma A, Charroux B, Abel L, Rappsilber J, Mann M, Dreyfuss G: **miRNPs: a novel class of ribonucleoproteins containing numerous microRNAs.** *Genes Dev* 2002, **16**:720-728.
- Aravin AA, Lagos-Quintana M, Yalcin A, Zavolan M, Marks D, Snyder B, Gaasterland T, Meyer J, Tuschl T: **The small RNA profile during Drosophila melanogaster development.** *Dev Cell* 2003, **5**:337-350.
- Lai EC, Tomancak P, Williams RW, Rubin GM: **Computational identification of Drosophila microRNA genes.** *Genome Biol* 2003, **4**:R42.
- Yu J, Wang F, Yang GH, Wang FL, Ma YN, Du ZW, Zhang JW: **Human microRNA clusters: genomic organization and expression profile in leukemia cell lines.** *Biochem Biophys Res Commun* 2006, **349**:59-68.
- Thatcher EJ, Bond J, Paydar I, Patton JG: **Genomic organization of zebrafish microRNAs.** *BMC Genomics* 2008, **9**:253.
- Kim YK, Yu J, Han TS, Park SY, Namkoong B, Kim DH, Hur K, Yoo MW, Lee HJ, Yang HK, Kim VN: **Functional links between clustered microRNAs: suppression of cell-cycle inhibitors by microRNA clusters in gastric cancer.** *Nucleic Acids Res* 2009, **37**:1672-1681.
- Tanzer A, Stadler PF: **Molecular evolution of a microRNA cluster.** *J Mol Biol* 2004, **339**:327-335.
- Guddeti S, Zhang DC, Li AL, Leseberg CH, Kang H, Li XG, Zhai WX, Johns MA, Mao L: **Molecular evolution of the rice miR395 gene family.** *Cell Res* 2005, **15**:631-638.
- Wang S, Zhu QH, Guo X, Gui Y, Bao J, Helliwell C, Fan L: **Molecular evolution and selection of a gene encoding two tandem microRNAs in rice.** *FEBS Lett* 2007, **581**:4789-4793.
- Chuck G, Cigan AM, Saeteurn K, Hake S: **The heterochronic maize mutant corngrass1 results from overexpression of a tandem microRNA.** *Nat Genet* 2007, **39**:544-549.
- Boualem A, Laporte P, Jovanovic M, Laffont C, Plet J, Combi JP, Niebel A, Crespi M, Frugier F: **MicroRNA166 controls root and nodule development in Medicago truncatula.** *Plant J* 2008, **54**:876-887.
- Lacombe S, Nagasaki H, Santi C, Duval D, Piegu B, Bangratz M, Breitler JC, Guiderdoni E, Brugidou C, Hirsch J, Cao X, Brice C, Panaud O, Karlowski WM, Sato Y, Echeverria M: **Identification of precursor transcripts for 6 novel miRNAs expands the diversity on the genomic organization and expression of miRNA genes in rice.** *BMC Plant Biol* 2008, **8**:123-142.
- Lee Y, Jeon K, Lee JT, Kim S, Kim VN: **MicroRNA maturation: stepwise processing and subcellular localization.** *EMBO J* 2002, **21**:4663-4670.
- Baskerville S, Bartel DP: **Microarray profiling of microRNAs reveals frequent coexpression with neighboring miRNAs and host genes.** *RNA* 2005, **11**:241-247.
- VISTA Plot [<http://genome.lbl.gov/vista/index.shtml>]
- Frazer KA, Pachter L, Poliakov A, Rubin EM, Dubchak I: **VISTA: computational tools for comparative genomics.** *Nucleic Acids Res* 2004, **32**:W273-W279.
- Gustafson AM, Allen E, Givan S, Smith D, Carrington JC, Kasschau KD: **ASRP: the Arabidopsis Small RNA Project database.** *Nucleic Acids Res* 2005, **33**:D637-D640.
- Backman TW, Sullivan CM, Cumbie JS, Miller ZA, Chapman EJ, Fahlgren N, Givan SA, Carrington JC, Kasschau KD: **Update of ASRP: the Arabidopsis Small RNA Project database.** *Nucleic Acids Res* 2008, **36**:D982-D985.
- Arabidopsis Small RNA Project [<http://asrp.cgrb.oregonstate.edu/>]
- Meyers BC, Axtell MJ, Bartel B, Bartel DP, Baulcombe D, Bowman JL, Cao X, Carrington JC, Chen X, Green PJ, Griffiths-Jones S, Jacobsen SE, Mallory AC, Martienssen RA, Poethig RS, Qi Y, Vaucheret H, Voinnet O, Watanabe Y, Weigel D, Zhu JK: **Criteria for annotation of plant MicroRNAs.** *Plant Cell* 2008, **20**:3186-3190.
- Brodersen P, Voinnet O: **Revisiting the principles of microRNA target recognition and mode of action.** *Nat Rev Mol Cell Biol* 2009, **10**:141-148.
- Lu S, Sun YH, Shi R, Clark C, Li L, Chiang VL: **Novel and mechanical stress-responsive MicroRNAs in Populus trichocarpa that are absent from Arabidopsis.** *Plant Cell* 2005, **17**:2186-2203.
- Lu S, Sun YH, Chiang VL: **Stress-responsive microRNAs in Populus.** *Plant J* 2008, **55**:131-151.
- Tyler L, Thomas SG, Hu J, Dill A, Alonso JM, Ecker JR, Sun TP: **Della proteins and gibberellin-regulated seed germination and floral development in Arabidopsis.** *Plant Physiol* 2004, **135**:1008-1019.
- Populus trichocarpa v1.1 [http://genome.jgi-psf.org/Poptr1_1/Poptr1_1.info.html]
- AtPARE Database [http://mpss.udel.edu/at_pare/]
- Lu C, Kulkarni K, Souret FF, MuthuVallippan R, Tej SS, Poethig RS, Henderson IR, Jacobsen SE, Wang W, Green PJ, Meyers BC: **MicroRNAs and other small RNAs enriched in the Arabidopsis RNA-dependent RNA polymerase-2 mutant.** *Genome Res* 2006, **16**:1276-1288.
- Jones-Rhoades MW, Bartel DP: **Computational identification of plant microRNAs and their targets, including a stress-induced miRNA.** *Mol Cell* 2004, **14**:787-799.
- Abdel-Ghany SE, Pilon M: **MicroRNA-mediated systemic down-regulation of copper protein expression in response to low copper availability in Arabidopsis.** *J Biol Chem* 2008, **283**:15932-15945.

51. Schauer SE, Jacobsen SE, Meinke DW, Ray A: **DICER-LIKE1: blind men and elephants in *Arabidopsis* development.** *Trends Plant Sci* 2002, **7**:487-491.
52. Maher C, Stein L, Ware D: **Evolution of *Arabidopsis* microRNA families through duplication events.** *Genome Res* 2006, **16**:510-519.
53. Cui X, Xu SM, Mu DS, Yang ZM: **Genomic analysis of rice microRNA promoters and clusters.** *Gene* 2009, **431**:61-66.
54. Piriyapongsa J, Jordan IK: **Dual coding of siRNAs and miRNAs by plant transposable elements.** *RNA* 2008, **14**:814-821.
55. Vazquez F, Vaucheret H, Rajagopalan R, Lepers C, Gascioli V, Mallory AC, Hilbert JL, Bartel DP, Crete P: **Endogenous trans-acting siRNAs regulate the accumulation of *Arabidopsis* mRNAs.** *Mol Cell* 2004, **16**:69-79.
56. Rajagopalan R, Vaucheret H, Trejo J, Bartel DP: **A diverse and evolutionarily fluid set of microRNAs in *Arabidopsis thaliana*.** *Genes Dev* 2006, **20**:3407-3425.
57. Kawashima CG, Yoshimoto N, Maruyama-Nakashita A, Tsuchiya YN, Saito K, Takahashi H, Dalmay T: **Sulphur starvation induces the expression of microRNA-395 and one of its target genes but in different cell types.** *Plant J* 2009, **57**:313-321.
58. Lechner E, Achard P, Vansiri A, Potuschak T, Genschik P: **F-box proteins everywhere.** *Curr Opin Plant Biol* 2006, **9**:631-638.
59. Busov V, Meilan R, Pearce DW, Rood SB, Ma C, Tschaplinski TJ, Strauss SH: **Transgenic modification of *gai* or *rgll* causes dwarfing and alters gibberellins, root growth, and metabolite profiles in *Populus*.** *Planta* 2006, **224**:288-299.
60. Cai X, Davis EJ, Ballif J, Liang M, Bushman E, Haroldsen V, Torabinejad J, Wu Y: **Mutant identification and characterization of the lacase gene family in *Arabidopsis*.** *J Exp Bot* 2006, **57**:2563-2569.
61. Clay NK, Adio AM, Denoux C, Jander G, Ausubel FM: **Glucosinolate metabolites required for an *Arabidopsis* innate immune response.** *Science* 2009, **323**:95-101.
62. Lewis BP, Shih IH, Jones-Rhoades MW, Bartel DP, Burge CB: **Prediction of mammalian microRNA targets.** *Cell* 2003, **115**:787-798.
63. Allen E, Xie Z, Gustafson AM, Sung GH, Spatafora JW, Carrington JC: **Evolution of microRNA genes by inverted duplication of target gene sequences in *Arabidopsis thaliana*.** *Nat Genet* 2004, **36**:1282-1290.
64. Schwab R, Ossowski S, Riester M, Warthmann N, Weigel D: **Highly specific gene silencing by artificial microRNAs in *Arabidopsis*.** *Plant Cell* 2006, **18**:1121-1133.
65. **microRNA Registry** [<http://www.sanger.ac.uk/Software/Rfam/mirna/>]
66. **mfold Program** [<http://mfold.bioinfo.rpi.edu/cgi-bin/rna-form1.cgi>]
67. Zuker M: **Mfold web server for nucleic acid folding and hybridization prediction.** *Nucleic Acids Res* 2003, **31**:3406-3415.
68. Couronne O, Poliakov A, Bray N, Ishkhanov T, Ryaboy D, Rubin E, Pachter L, Dubchak I: **Strategies and tools for whole-genome alignments.** *Genome Res* 2003, **13**:73-80.
69. Bray N, Dubchak I, Pachter L: **AVID: A global alignment program.** *Genome Res* 2003, **13**:97-102.
70. **VISTA Graphic Server** [<http://genome.lbl.gov/vista/index.shtml>]
71. Pall GS, Codony-Servat C, Byrne J, Ritchie L, Hamilton A: **Carbodiimide-mediated cross-linking of RNA to nylon membranes improves the detection of siRNA, miRNA and piRNA by northern blot.** *Nucleic Acids Res* 2007, **35**:e60.
72. Czechowski T, Stitt M, Altmann T, Udvardi MK, Scheible WR: **Genome-wide identification and testing of superior reference genes for transcript normalization in *Arabidopsis*.** *Plant Physiol* 2005, **139**:5-17.
73. Vandesompele J, De Preter K, Pattyn F, Poppe B, Van Roy N, De Paepe A, Speleman F: **Accurate normalization of real-time quantitative RT-PCR data by geometric averaging of multiple internal control genes.** *Genome Biol* 2002, **3**:-
74. **pEarlyGate Vectors** [<http://www.biology.wustl.edu/pikaard/Vectors%20homepage.html>]
75. Bechtold N, Pelletier G: **In planta *Agrobacterium*-mediated transformation of adult *Arabidopsis thaliana* plants by vacuum infiltration.** *Methods Mol Biol* 1998, **82**:259-266.