

RESEARCH ARTICLE

Open Access



# Identification of *Gossypium hirsutum* long non-coding RNAs (lncRNAs) under salt stress

Fenni Deng, Xiaopei Zhang, Wei Wang, Rui Yuan and Fafu Shen\*

## Abstract

**Background:** Long non-coding RNAs (lncRNAs) represent a class of riboregulators that either directly act in long form or are processed into shorter microRNAs (miRNAs) and small interfering RNAs. Long noncoding RNAs (lncRNAs) are arbitrarily defined as RNA genes larger than 200 nt in length that have no apparent coding potential. lncRNAs have emerged as playing important roles in various biological regulatory processes and are expressed in a more tissue-specific manner than mRNA. Emerging evidence shows that lncRNAs participate in stress-responsive regulation.

**Results:** In this study, in order to develop a comprehensive catalogue of lncRNAs in upland cotton under salt stress, we performed whole-transcriptome strand-specific RNA sequencing for three-leaf stage cotton seedlings treated with salt stress (S\_NaCl) and controls (S\_CK). In total we identified 1117 unique lncRNAs in this study and 44 differentially expressed RNAs were identified as potential non-coding RNAs. For the differentially expressed lncRNAs that were identified as intergenic lncRNAs (lincRNA), we analysed the gene ontology enrichment of *cis* targets and found that *cis* target protein-coding genes were mainly enriched in stress-related categories. Real-time quantitative PCR confirmed that all selected lincRNAs responsive to salt stress. We found lnc\_388 was likely as regulator of Gh\_A09G1182. And lnc\_883 may participate in regulating tolerance to salt stress by modulating the expression of Gh\_D03G0339 MS\_channel. We then predicted the target mimics for miRNA in *Gossypium*. six miRNAs were identified, and the result of RT-qPCR with lncRNA and miRNA suggested that lnc\_973 and lnc\_253 may regulate the expression of ghr-miR399 and ghr-156e as a target mimic under salt stress.

**Conclusions:** We identified 44 lincRNAs that were differentially expressed under salt stress. These lincRNAs may target protein-coding genes via *cis*-acting regulation. We also discovered that specifically-expressed lincRNAs under salt stress may act as endogenous target mimics for conserved miRNAs. These findings extend the current view on lincRNAs as ubiquitous regulators under stress stress.

**Keywords:** lncRNA, Salt stress, *Gossypium hirsutum*, RNA-Seq, RT-qPCR

## Background

Long noncoding RNAs (lncRNAs) are arbitrarily defined as RNA genes larger than 200 bp in length that have no apparent coding potential. lncRNAs have emerged as playing important roles in various biological regulatory processes and have more tissue-specific expression patterns than mRNA [1–4]. It is well known that lncRNA transcripts are transcribed by RNA polymerase II mainly. As polymerase II polyadenylated products, lncRNAs are modified with a 5'-cap and (or) polyadenylated at the 3'-end in animals and plants [5–7]. The

transcription of several lncRNA genes requires specific transcription factors, mediator complexes, histone modification complexes and transcription elongation factor complexes [6, 8–11]. Based on their genomic origins and their location relative to nearby protein-coding genes, lncRNAs can be classified into five groups: (i) sense long non-coding RNA, (ii) natural antisense transcripts (NATs) derived from introns, (iii) long intergenic non-coding (nc) RNAs (lincRNAs), (iv) intronic ncRNAs (incRNAs) and (v) bidirectional long ncRNA [7].

With advances in sequencing and the emergence of new technologies, the discovery process of lncRNA has generally developed in three phases: in the 1980s–1990s, individual lncRNAs, for example, *XIST* and *H19*, were

\* Correspondence: [ffshen@sdau.edu.cn](mailto:ffshen@sdau.edu.cn)

State Key Laboratory of Crop Biology, College of Agronomy, Shandong Agricultural University, Tai'an 271018, Shandong, People's Republic of China



discovered through traditional gene mapping approaches. *XIST* has been identified in mammalian cells, has no significantly conserved open reading frames and does not appear to encode a protein. In addition, *XIST* originates from the X-chromosome inactivation centre and functions as a structural RNA in the nucleus [12]. In the early 2000s, the development of large-scale cDNA sequencing led to the discovery of a surprising number of lncRNA transcripts. Especially in plants, thousands of lncRNAs have been identified, such as in *Arabidopsis thaliana* [13], rice [14, 15] and *Medicago truncatula* [16]. During the mid-2000s, the number of predicted genes in the mammalian genome declined, while the number of detected lncRNA transcripts increased exponentially in several plant species. Microarrays, tiling arrays and next-generation sequencing were used as high-throughput tools for identifying new transcripts [7, 17–19].

In the nucleus, lncRNAs may execute their functions either in close proximity (*cis*-acting) or over a distance (*trans*-acting) in the genome via numerous mechanisms, including activating, gathering or transporting proteins and epigenetic silencing and repression, modifying promoter activities by nucleosome repositioning, epigenetic modification by regulating the level of DNA methylation and histone modifications [3, 17, 18]. Increasing evidence has shown that lncRNAs play a crucial role in growth and development, disease occurrence and genetic and epigenetic regulation in mammals [20, 21].

Recent studies have reported on the function of lncRNAs in plants [18, 19]. Boerner et al. (2012) identified 2473 lncRNAs in *Zea Mays*, classifying small RNA precursors and lncRNAs that are likely to function as longer molecules [22]. In *Arabidopsis*, *COOLAIR* and *COLDAIR* have been identified to regulate the expression of *FLOWERING LOCUS C (FLC)* [18]. A lncRNA that has been identified in *Arabidopsis*, Induced by phosphate starvation 1 (*IPS1*), can bind and sequester miR399 and reduce miR399-mediated cleavage of Phosphate 2 (*PHO2*), which is associated with phosphate uptake [23]. In hybrid rice, long-day-specific male-fertility-associated RNA (*LDMAR*) has been found to regulate photoperiod-sensitive male sterility (*PSMS*) [24]. More recently, more than 2000 lncRNAs were identified in rice during sexual reproduction, including 1624 lincRNAs and 600 long noncoding natural antisense transcripts (lncNATs) [25].

Cotton (*Gossypium hirsutum L.*) is an important economic crop, which produces a natural fibre, provides edible protein for livestock feed and is a source of oil and biofuel. Today, the most extensively cultivated cotton species are the tetraploid *G. hirsutum* and *G. barbadense*. Recently, whole-genome sequencing analyses of diploid (*G. raimondii*; and *G. arboreum*) and tetraploid cotton species (*G. hirsutum* and *G. barbadense*) have

provided valuable reference genomes for cotton, and a number of shotgun sequencing efforts have increased our understanding of the variation in these cotton genomes [26–32]. Upland cotton is a moderate salt-tolerant plant, with a salinity threshold of  $7.7 \text{ dS m}^{-1}$  [33, 34]. Studies on the tolerance of cotton to salt stresses have shown that a saline environment affects cotton growth and development and influences its biological and metabolic pathways. In particular, the growth of seedlings is severely reduced under a high-salinity environment. Salt stress affects photosynthesis and respiration, flowering, fibre quality and ion uptake in cotton, influencing both biological and metabolic pathways [35, 36]. A study of lncRNAs in the fibre development of cotton identified 30,550 lincRNAs and 4718 lncNATs using RNA-seq, and these lncRNAs are thought to regulate the expression of other genes through multiple RNA-mediated mechanisms [37]. However, reports on lncRNAs involved in salt-responsive regulation in *G. hirsutum* are lacking.

## Results

### High-throughput sequencing

In order to develop a comprehensive catalogue of lncRNAs in upland cotton under salt stress, we performed whole-transcriptome strand-specific RNA sequencing for three-leaf stage cotton seedlings treated with salt stress (S\_NaCl) and control (S\_CK), including biological replicates (S\_NaCl1, S\_NaCl2 and S\_CK1, S\_CK2). Sequencing was done on the Illumina HiSeq 4000 platform, and 250 bp paired-end reads were generated. We obtained more than 40,000,000 raw sequence reads by RNA-seq. From the raw reads, we identified more than 98% that were clean reads (Table 1). To estimate the quality of the RNA-seq data, we used Fast QC with a phred-like algorithm to calculate the quality score (Q) of each base pair in the reads. The results were a mean Q-value of 40, showing that the RNA-seq data were highly reliable (Additional file 1: Mean sample quality).

### Identification and characterization of lincRNAs in cotton

All RNA-seq datasets were mapped to the genome of *G. hirsutum* using TopHat in order to reconstruct the cotton transcriptome. According to the percent of reads

**Table 1** RNA-seq data for four samples

Summary	S_CK		S_NaCl	
	S_CK1	S_CK2	S_NaCl1	S_NaCl2
Raw Reads	43,474,829	46,025,954	43,284,887	47,240,877
Clean Reads	43,474,829	45,678,120	42,839,455	46,971,811
Clean Reads Rate (%)	100%	99.24%	98.97%	99.43%
Unique mRNAs	58,471	58,484	58,805	59,080
Unique lncRNAs	1520	1635	1549	1635

mapped to genomic regions, we found approximately 53% clean reads distributed among exonic regions, 41% distributed among intergenic regions and 5.6% distributed across intronic regions in S\_CK1 (Additional file 2: Percent of reads mapped to genome regions).

Next, the transcripts were assembled and annotated using Cufflinks. We identified 58,471, 58,484, 58,805 and 59,080 unique mRNAs from the four cDNA libraries (S\_CK1, S\_CK2, S\_NaCl1, S\_NaCl2), respectively (Table 1). The remaining reads were filtered according to length and coding potential such that transcripts with length < 200 bp were removed and transcripts with a coding potential > 0 were excluded.

Finally we obtained 1520, 1635, 1549, and 1635 unique lincRNAs from the four samples (S\_CK1, S\_CK2, S\_NaCl1, S\_NaCl2), respectively (Table 1). In total we identified 1117 unique lincRNAs in this study. We mapped these lincRNAs to the 13 chromosomes of the *G. hirsutum* genome and found that they were distributed across these chromosomes without a preference of location in either the controls or treated samples. According to the location of these lincRNAs in the cotton genome, we identified 1117 intergenic lincRNAs (lincRNAs). We then assessed the length and expression level of the lincRNA transcripts. The median length of these lincRNAs was 1200 nucleotides (nt), and most were shorter than 2000 nt (Fig. 1a). We then estimated the expression level of each transcript using reads per kilobase of exon per million fragments mapped (RPKM) and found that the lincRNAs in the control and salt treatment groups were expressed at similar levels (Fig. 1b).

However, the overall expression level of lincRNAs was lower than that of mRNAs (Fig. 1b), consistent with previous studies in *G. barbadense*, humans and *Medicago truncatula* [37–39].

Next, we aligned the genomes of *G. hirsutum* [29] and *Arabidopsis* [40] using MultiZ [41] to verify the conservation score (consScore) of each nt using phastCons [42]. We then extracted the consScore of each lincRNA (Additional files 3, 4). Like lincRNAs from rice, we found that the *G. hirsutum* lincRNAs were less conserved than mRNAs (Fig. 2a, b). The lower the conservation level, the lower the conservation score.

We estimated the degree of differential expression between the controls and salt stress samples of the lincRNAs and mRNAs based on the Jensen-Shannon score [39] and found that lincRNAs tended to be more differentially expressed than mRNAs (Fig. 2c). Meanwhile, a strikingly different expression pattern of lincRNAs was found in *Arabidopsis* [19], rice [42], *M. truncatula* [43] and various animals [39], suggesting that these characteristics are conserved for lincRNAs.


### Expression of *G. hirsutum* lincRNAs under salt stress

To identify salt-responsive lincRNAs of *G. hirsutum*, the normalized expression (RPKM) of the lincRNAs was compared between the control and salt treatment groups. We clustered the differentially expressed lincRNAs on the basis of their expression patterns using Cluster3.0. The highly specific expression pattern observed for lincRNAs allowed us to cluster them into two categories, (i) those that were specifically expressed in the controls (for example, lnc\_1045, lnc\_123), and (ii) those that were specifically expressed under salt-treatment (for example, lnc\_26, lnc\_388) (Fig. 3a), suggesting that these lincRNAs may function in response to salt stress. We next compared the levels of the salt-responsive protein coding genes in the control samples (S\_CK1, S\_CK2) and the salt treatment samples (S\_NaCl1, S\_NaCl2) from the two clusters using hierarchical cluster analysis (Fig. 3b). The log<sub>2</sub> ratio values of the salt responsive genes were used for hierarchical cluster analysis with Cluster3.0. Details of the annotations of all differentially expressed protein coding genes shown on the right are provided in Additional file 4. To confirm their expression patterns, we randomly selected 11 lincRNAs and quantified them using real-time quantitative PCR (RT-qPCR), in Fig. 4 and Additional file 5. In Fig. 3a and Additional file 5, the expression patterns of specifically expressed lincRNAs in the sequencing and RT-qPCR results were mostly consistent, although the relative expression levels of all lincRNAs measured by RNA-Seq were greater than those by RT-qPCR. Consequently, four lincRNAs were identified as up-regulated under salt treatment, while two lincRNAs were down-regulated. Additionally, in Fig. 3b and Additional file 6, we found that several salt tolerance-related protein genes were co-expressed with the lincRNAs, for example Gh\_A05G3489, Gh\_A01G0321, Gh\_A01G0639 and Gh\_A11G0366.

### Functional analysis of differentially expressed lincRNAs under salt stress

It has been shown that a number of lincRNAs can regulate the expression of genes in close proximity (*cis*-acting) or at a distance (*trans*-acting). Previous studies in animals and plants showed that lincRNAs are preferentially located in close proximity to protein-coding genes that they regulate [43–45].

To analyse the potential functions of the identified lincRNAs, we selected protein-coding genes that were co-expressed and were spaced less than 20 kb away from the differentially expressed lincRNAs. We analysed the gene ontology (GO) enrichment of these protein-coding genes and found that these protein-coding genes were mainly enriched in stress-related categories, such as “response to stress”, “response to chemical stress”, “biological regulation”, “binging”, “oxidoreductase” and




“transcription regulator” (Fig. 5a). In addition, we detected significant enrichment that on a mean  $P$ -value < 0.05. For example we found GO-term enrichments for biological processes (GO:0006722, triterpenoid metabolic process; GO:0000027, ribosomal large subunit assembly; and GO:0010426, DNA methylation on cytosine within a CHH sequence), cellular component (for example, GO:0016160, amylase activity; GO:0016671, oxidoreductase activity; and GO:0042300, beta-amyrin synthase activity) and molecular function (GO:0010006, Toc complex; GO:0005871, kinesin complex; and GO:0009317, acetyl-CoA carboxylase complex) (Fig. 5b).

These results suggest that the differentially expressed lncRNAs in salt stress may regulate protein-coding genes involved in several important biological processes, such as carbohydrate metabolism, detoxification, energy synthesis, transcription, chromatin modification and post-transcriptional regulation in response to salt stress. lncRNAs can directly regulate the polymerase II transcription machinery in many ways. To illuminate the function of differentially expressed lncRNAs in salt stress and the relationship between lncRNAs and mRNAs which were co-expressed and fall less than

20 kb away from differentially expressed lncRNAs, putative interactive networks were established using Cytoscape (Additional file 7 Network and Fig. 5b). Genes at network nodes were sorted into three groups. First, seven protein-coding genes were involved in oxidation/reduction reactions. These included, for example, the genes encoding peroxidase and thioredoxin. Thioredoxins are small enzymes that participate in redox reactions via the reversible oxidation of an active center disulfide bond. Second, four protein-coding genes were involved in the transport of anions. For example, three genes encoded proteins that are involved in the transport of anions across the cytoplasmic membrane during salt response. Finally, five protein-coding genes were involved in transcription, including Leucine rich repeat, eukaryotic translation initiation factor 5 and RNA recognition motif (Fig. 5b and Addition file 8).

Interestingly, we found that 13 lncRNAs spaced less than 20 kb away from their putative *cis-acting* targets may regulate neighboring protein coding genes by *cis-acting* in Additional file 8. Of which, lnc\_128 was spaced 4736 bp with Gh\_A05G1992 coding eIF-5\_eIF-2B, especially lnc\_883, located approximately 718 bp upstream



**Fig. 2** Conservation of *Gossypium hirsutum* lincRNAs. **a**, The conservation score (consScore) of each nucleotide in the *G. hirsutum* genome; **(b)**, The level of conservation of the lincRNAs. The cumulative distributions of mean phastCons scores derived from the *G. hirsutum* and Arabidopsis whole-genome alignments are shown; **(c)**, The maximal Jensen-Shannon (JS) specificity score distributions for all lincRNAs and mRNAs

of the coding sequence of Gh\_D03G0339 (Fig. 6). Another lincRNA linc\_388 localized at -strand A09, had a 12.7 kb distance with LRR8 (Gh\_A09G1182). To confirm the relationship of lincRNA and neighboring protein coding genes, we selected two lincRNAs we were interested in. Using real-time quantitative PCR (RT-qPCR) lincRNA and their putative *cis-acting* targets were quantified in Fig. 7. We found linc\_388 and putative *cis-acting* targets LRR8 (Gh\_A09G1182) were co-expressed and dramatically up-regulated in salt stress. Meanwhile the expression of linc\_883 and msD3(Gh\_D03G0339) was up-regulated (Fig. 7). These results suggest that linc\_883 may participate in regulating tolerance to salt stress by modulating the expression of Gh\_D03G0339 MS\_channel. And linc\_388 may be involve in salt tolerance by adjusting the expression of Leucine rich repeat8 (Gh\_A09G1182).


#### Cotton lincRNAs as endogenous target mimics for miRNAs

In plants, an important function of lincRNAs is target mimicry, and the miRNA-lincRNA relationship was recently discovered in *Arabidopsis* [46–48]. The first target mimic identified, *IPS1*, was discovered in *Arabidopsis*, in

which it modulates the activity of miR-399 by a complementary mechanism [46]. It is possible that functional miRNA target mimics consist mainly of lincRNAs. In this study, the target mimics of these lincRNAs were predicted by psRNAtarget (<http://plantgrn.noble.org/psRNAtarget>) [48, 49]. In total, six and 30 target mimics were identified in *Gossypium* spp. and in other plants, respectively (Table 2: Analysis of target mimics of lincRNAs for miRNAs, and Fig. 8). Five miRNAs (gra-miR7502, gra-miR8876, gra-miR8693, gra-miR7504d, and ghr-156e) with various functions (i.e. target and target mimicry) were identified (Fig. 8). linc\_361 targets gra-miR7504 and gra-miR8876. For gra-miR7504, the function appears to be more complex, as it is targeted by two lincRNAs (linc\_361 and linc\_828). And linc\_253 was predicted to target ghr-156e. Therefore, linc\_361, linc\_828 and linc\_253 may regulate their opposite miRNA as target mimics and inhibit its function.

In addition, linc\_973 was identified as a target mimic for bdi-miR399a. By sequence alignment, we found that ghr-miR399a and bdi-miR399a differ at the 16th base-pair, which is an A in bdi-miR399a and a G in ghr-






miR399a (Additional file 9). Thus, lnc\_973 may regulate the expression of ghr-miR399 as a target mimic under salt stress.


To investigate the mechanism function of lncRNAs (lnc\_973, lnc\_253), we analysed the expression of lncRNA and corresponding microRNA under salt stress. Interestingly, we found the expression pattern of lnc\_973 and corresponding microRNA ghr-miR399a were identical. lnc\_973 and ghr-miR399a were up-regulated under salt stress, but the target genes of miRNA Gh\_D07G0254 and Gh\_D05G0219 appeared up-regulated. Meanwhile lnc\_253 and ghr-156e shown the similar result in Fig. 9. These evidences suggested lnc\_973 and lnc\_253 may regulate the expression of ghr-miR399 and ghr-156e as a target mimic under salt stress. lncRNA can promote the target of miRNA expression by competing miRNA to downregulate the activity of miRNA.

### Discussion

Salt stress induces osmotic and ionic toxicity and oxidative stress disrupts plant homeostasis [50]. Under stress, the sensor systems in plants are triggered by downstream signalling and transcriptional control cascades, which result in extensive changes in cellular gene expression [51, 52]. In response to salt stress, numerous transcription factors are differentially expressed, such as members of the basic leucine zipper (bZIP), basic helix-loop-helix (bHLH), MYB and NAC families, WRKY and ERF, [53], which play critical roles as transcriptional regulators in plant growth and development [54]. Increasing numbers of functional studies on protein-coding genes and small noncoding RNAs have revealed the high level of complexity of eukaryotic transcriptomes, especially considering the extensive abundance of lncRNAs [55]. lncRNAs are a recently discovered type of molecule with important functions in a wide range of biological processes, including developmental regulation and stress response; nevertheless, the detailed mechanisms involved in these biological processes remain largely unknown [56]. In cotton, several studies have identified small ncRNAs and lncRNAs through small RNA sequencing in fibre development, but no data has been presented for lncRNAs under salt stress. The recent publication of genome sequences and the accumulation of RNA-seq data have allowed the genome-wide identification of lncRNAs. In this study, we identified more than 58,000 mRNAs and 1117 unique lncRNAs by analysing more than 40,000,000 raw sequence reads. The number of lncRNAs identified by sequencing were selected based on strict criteria. Although this may have excluded many lncRNAs, these 1117 unique lncRNAs constitute a reliable set of cotton lncRNAs. According to








**Fig. 6** Structure of lincRNAs and their putative target genes

studies in *A. thaliana* identified an endogenous lincRNA, *IPS1*, which can bind to ath-miR399 with a three-nucleotide bulge between the 10th and 11th positions of ath-miR399 at its 5' end [46]. Such pairing abolished the cleavage effect of ath-miR399 on *IPS1*; thus, *IPS1* serves as a decoy for ath-miR399 to interfere with the binding

of ath-miR399 to its other targets protein gene *PHO2*. This type of inhibitory mechanism of miRNA function is termed target mimicry, and *IPS1* a target mimic of miR399. Subsequently, in mammals, the category of target mimicry lincRNAs was renamed to competing endogenous RNAs (ceRNAs) and was shown to be relevant



**Fig. 7** The analysis of RT-qPCR with lincRNA and their putative target genes. LRR8: Leucine rich repeat8 (Gh\_A09G1182); msD3: MS\_channel (Gh\_D03G0339)




**Table 2** Putative targets and target mimics of lincRNAs for miRNA

miRNA_Acc.	Target_Acc.	Expectation	UPE	miRNA_aligned_fragment	Target_aligned_fragment	Inhibition
ssl-miR395	lnc_1045	3	14.736	GGGAAAUUUUGGGAAACU	AGUUUCCCCAAAAGAUUUUCU	Cleavage
stu-miR166d-5p	lnc_174	3	15.884	AGAAUGUCGUCUGGUUCGAG	UUUGAACUGGGUGACAUUCU	Cleavage
bra-miR9555a-5p	lnc_175	3	13.847	UUCUAAGCUUUACGGGAAAC	GUUUUUUGUGAGGUUUAGAA	Cleavage
cpa-miR8142	lnc_361	3	15.344	UGAGGUAAAGUAGACAGUAAAGUU	GAUUUUUGCUGUUGAUUUUAUCUCA	Translation
gra-miR7504d	lnc_361	2.5	5.165	AGGAAAAAAAUCUGAUUUUGU	GCAAAUCAGUUUUUUUCUU	Translation
gra-miR8766	lnc_361	3	7.582	UUUUUUUGGAAUUAGAAAAGUCGU	GCAUUUUUUUUAAUUC-AAAAUAA	Cleavage
cre-miR1173	lnc_376	3	14.984	AUGGUUGCAAUAGAAUCAU	AUGGUUUGUAUUUUAAACCAU	Cleavage
osa-miR1881	lnc_464	3	16.025	AAUGUUUUGUAGC-GUGGUGGUG	UGCCACCAUUGCUGCAAUAAACAUU	Cleavage
bra-miR9565-3p	lnc_464	3	11.927	CUGAAGCUAGUGAAAGAGAGA	UCUCUCUUUCUCUGUUUUAG	Translation
gra-miR7502b	lnc_612	3	13.131	UUGUUAAAAGUUUCAUCCAU	AUGGGUGAAAUUUUUGAGAA	Cleavage
gra-miR7502c	lnc_612	3	13.131	UUGUUAAAAGUUUCAUCCAU	AUGGGUGAAAUUUUUGAGAA	Cleavage
ath-miR417	lnc_699	3	10.255	GAAGGUAGUGAAUUUGUUCG	UGAACAGAUUCAUUGCUUUU	Cleavage
gma-miR5376	lnc_699	2.5	9.125	UGAAGAUUUGAAGAAUUUGG	CAAAUUCUUCUAAUUUUUCA	Translation
bdi-miR5200c	lnc_723	3	13.74	UGUAGAUACUCUCUAAAGGCU	AGUCUUUAUAGUAUUUACA	Translation
ath-miR5656	lnc_801	3	15.674	ACUGAAGUAGAGAUUGGGUUU	AAAGACAAUUUCUACUUCGGU	Cleavage
mtr-miR5747	lnc_801	3	11.23	AAAAGAAUACUCAUCAUACA	UAUUAGUAUGAGUAUUUGUUU	Cleavage
gra-miR7504d	lnc_828	2.5	14.203	AGGAAAAAAAUCUGAUUUUGUC	GCCAAAUGAGAUUCUUUUUCUU	Translation
tae-miR5049-3p	lnc_828	3	13.657	AAUAUGGAUCGGAGGGAGUA	UGCUCUUUCAGUUCAUUUU	Translation
mtr-miR5285a	lnc_883	3	15.033	UGGGACUUUGGGUAGAAUJAGGC	GUUUAAUUCUGCCCUAAGUCUCU	Translation
mtr-miR5285b	lnc_883	3	15.033	UGGGACUUUGGGUAGAAUJAGGC	GUUUAAUUCUGCCCUAAGUCUCU	Translation
mtr-miR5285c	lnc_883	3	15.033	UGGGACUUUGGGUAGAAUJAGGC	GUUUAAUUCUGCCCUAAGUCUCU	Translation
miRNA_Acc.	Target_Acc.	Expectation	UPE	miRNA_aligned_fragment	Target_aligned_fragment	Inhibition
bcy-miR529	lnc_886	3	15.175	GAAGAAGAGAGAUUGGUAGAG	UUCUCUCAUUUCUUUUUCUUC	Cleavage
ath-miR865-5p	lnc_894	3	12.672	AUGAAUUUGGAUCUAAUUGAG	UUCAAUUAGCUCUAAAGUCAA	Cleavage
gma-miR4393a	lnc_894	3	6.465	UGAGAAAAGGACGGCAGAAAAG	CCUUUUUGCUGUCCAUUUCUCU	Cleavage
gra-miR8693	lnc_955	3	9.809	AGGAUGAAAAUUAUGAUGUAG	UUGUAUUAAUUUUUCAUCUU	Translation
gma-miR1520d	lnc_973	3	13.997	AUCAGAACAUGACACGUGAC	GUUGCGUGUUUUGUUCUGCU	Cleavage
bdi-miR399a	lnc_973	3	13.409	UGCCAAAGGAGAAUACCCUG	CAGGUUAGUUUCCUUUGGCU	Cleavage
gma-miR1520l	lnc_973	3	13.635	AAUCAGAACAUGACACGUGAU	GUUGCGUGUUUUGUUCUGCUU	Cleavage
gma-miR4994-5p	lnc_973	3	12.834	GGUUAGCUCAAGGAUCUCAC	AUGAGUCCUUGAGCUAAUU	Cleavage
ghr-156e	lnc_253	4.5	8.055	UGACAGAAGAGAGUGAGCAC	GUUCUGCUUCUUCUUUGUUA	Cleavage

in many processes [50, 56, 60, 61], implying that these molecules might represent a widespread form of gene regulation. lncRNAs can regulate their corresponding miRNA target genes by having miRNA-binding sites and competing for shared miRNAs. In *Arabidopsis*, after the *IPSI*-target mimic miR399 was identified, Wu et al. [62] predicted endogenous target mimics (eTMs) for 20 conserved miRNAs from intergenic or nc gene-originated regions in *Arabidopsis*, rice, *Populus trichocarpa* and maize, and several *Arabidopsis* eTMs have been shown to be functional [48, 56, 63, 64]. We also predicted that specifically-expressed lincRNAs under salt stress act as endogenous target mimics for conserved miRNAs in

cotton. Four miRNAs (gra-miR7502, gra-miR8876, gra-miR8693 and gra-miR7504d) with different functions (target and target mimicry) were identified. lnc\_973 and lnc\_253 were identified as the target mimic for bdi-miR399a (*Brachypodium distachyon*) and ghr-156e. And the result of RT-qPCR shown miRNA399 is not only expressed in phosphate starvation (Bari Rajendra et al., 2006). Moreover, we proved lnc\_973 and ghr-miR399a were co-expressed and up-regulated under salt stress. Meanwhile lnc\_253 may act as the target mimic for ghr-156e. The importance of lincRNAs in their role as eTMs during plant development and reproduction regulation will continue to emerge in future studies.



**Fig. 8** Putative targets and target mimics of lincRNAs for miRNA. lincRNAs as miRNA targets and target mimics are shown in **a b**; **c**; **d**; **e**; **f**

**Conclusion**


In this study, using RNA sequencing, 1117 unique lincRNAs were identified and 44 lincRNA that were identified as intergenic lincRNAs (lincRNA) differentially expressed under salt stress. Real-time quantitative PCR confirmed that all selected lincRNAs responded to salt stress, which is consistent with RNA-seq. We analysed the gene ontology enrichment of *cis-acting* targets and found that *cis-acting* target protein-coding genes were mainly enriched in stress-related categories. We found linc\_388 likely as regulator of Gh\_A09G1182. And linc\_883 may participate in regulating tolerance to salt stress by modulating the expression of Gh\_D03G0339 MS\_channel. Six miRNAs in *Gossypium* spp. were identified, and linc\_973 and linc\_253 may regulate the expression of ghr-miR399 and ghr-156e as a target mimic under salt stress. In cotton, several studies have identified small noncoding RNAs (ncRNAs) and lincRNAs through small RNA sequencing in fiber development, but no data has been presented for lincRNAs under salt stress. We found these lincRNAs may target protein-coding genes via *cis-acting* regulation. We also discovered that specifically-expressed lincRNAs under salt stress may act as endogenous target mimics for conserved miRNAs

in cotton. These findings extend the current view on lincRNAs as ubiquitous regulators under stress conditions.

**Methods**

**Plant materials and NaCl treatments**

In this study, SN91–11, a salt-tolerant cotton cultivar, was used. This cultivar was obtained by introducing Bluish Dogbane (*Apocynum venetum*) DNA into LM-6, which is a salt-sensitive cotton cultivar, by the pollen tube pathway. The physiological characteristics of SN91–11 have been described in earlier studies [65–67]. Sterilized seeds of SN91–11 were germinated in a mixture of peat and vermiculite at 28 °C. Then, the seedlings were grown under the following conditions: 28 °C/22 °C as day and night temperatures, respectively, under 16 h of light alternating with 8 h of darkness. At the three-leaf stage, the seedlings showing normal growth were randomly divided into two groups; one group was placed into tanks filled with a 250 mM solution of NaCl, and the remaining seedlings were transferred to tanks filled with plain water to serve as controls. After exposure to the two solutions for 24 h, seedlings of the control and



**Fig. 9** The expression results of lincRNA, miRNA and putative target genes of miRNA. RT-qPCR was performed for lincRNA and miRNA under salt stress, UBQ7 expression level was used as the reference gene

treated groups were harvested directly into liquid nitrogen and stored at  $-80^{\circ}\text{C}$  until used for RNA extraction.

#### Construction of lncRNA sequencing library and RNA-sequencing

Total RNA was isolated from each cotton seedling sample using the RNAPrep Pure Plant Kit Polysaccharides & Polyphenolics-rich (TIANGEN Biotech, Beijing, China). TRibosomal RNA was removed using the Epicentre Ribo-Zero Gold Kit (Epicentre, USA). Subsequently, sequencing libraries were generated following manufacturer recommendations with varied index labels by the NEBNext Ultra Directional RNA Library Prep Kit for Illumina (NEB, Ipswich, USA). The libraries were sequenced on the Illumina HiSeq 4000 platform, and 150 bp paired-end reads were generated in.

#### lncRNA identification

We processed raw data by removing the adaptor-polluted reads, removing the low-quality reads and trimming the reads whose number of N bases accounted for more than 5% (quality score,  $Q \geq 30$ ). The reference *G. hirsutum* genome and the annotation files were downloaded from the CottonGen database (<http://www.cottongen.org>). We built the genome index using Bowtie2 v2.2.3 [68], and clean data was mapped to the *G. hirsutum* genome using TopHat v2.0.12 [69]. TopHat calls Bowtie2 for mapping, which makes it more accurate and fast. TopHat specialized software for transcriptome sequencing reads mapping can identify exon-exon junctions by splitting the mapped reads and mapping them to the reference genome again. According to the characteristics of lncRNA, we adopted seven steps to identify lncRNAs from the transcripts of transcriptome assemblies [55]: (1) transcripts with length  $< 200$  bp and exon count  $< 2$  were removed; (2) every transcript with a coverage of  $< 3$  calculated by Cufflinks were selected; (3) the known protein-coding transcripts were removed; (4) transcripts were removed that were known ncRNAs; (5) transcripts were aligned in the Swiss-Prot and Pfam databases to remove those encoding proteins and protein-coding domains; (6) transcripts were eliminated that did not pass the protein-coding-score test using the Coding Potential Calculator (CPC), Coding-Non-Coding Index (CNCI) and Coding Potential Assessment Tool (CPAT) [45, 70]. A transcript was deemed to be noncoding if its protein potential scored less than 0, which meant that the transcript has no capacity of coding a protein; and (7) transcripts were removed that were detected in only one sample.

#### Expression analysis

We estimated the expression of lncRNA and mRNA transcripts using all mapped reads mapping by Cufflinks

[71]. First, all RNA-seq datasets were respectively aligned to the cotton genome using TopHat 2.0 [72]. Then, the transcriptome from each dataset was independently assembled using Cufflinks 2.0. All transcriptomes were merged to produce a final transcriptome using Cuffmerge. After the final transcriptome was generated, the abundance of all transcripts was estimated using Cuffdiff according to the final transcriptome, and a BAM file was produced from the TopHat alignment. The remaining transcripts in this study had biological replicates among each group, so we identified differential gene expression analysis using DESeq v1.16 (based on a negative binomial distribution). A *P*-value was assigned to each gene and adjusted by the Benjamini and Hochberg approach for controlling the false discovery rate. Genes with  $Q \leq 0.05$  and  $|\log_2 \text{ratio}| \geq 1$  were identified as differentially expressed genes.

#### Analysis of lncRNA function

In order to predict the function of salt stress-responsive lncRNAs, the neighbours of lncRNA protein-coding genes were analysed by GO enrichment, and GO terms with  $Q < 0.05$  were considered to be significantly enriched [45]. In accordance with previous investigations, lncRNAs regulate the expression of neighbouring genes through transcriptional activation/repression or epigenetic modification.

#### Target mimicry prediction

Targets were predicted by submitting all of the miRNAs (miRBase Release 21, June 2014) and the discovered lincRNAs to psRNATarget (<http://plantgrn.noble.org/psRNATarget/>) [48, 49], with less than three mismatches and G/U pairs allowed within the lincRNA and miRNA pairing regions. PsRNATarget was used to predict target mimics based on the principles established by Wu et al. [48].

#### Quantitative real-time (RT) PCR

To determine the relative transcript levels of selected lncRNAs and protein-coding genes, RT-qPCR was performed with specific primers (Additional file 10: Primer list for gene-specific primers) according to the manufacturer's instructions for the applied Biosystems 7500 RT-qPCR system (ABI 7500; life technologies Inc., Burlington, ON, Canada) and the SYBR premix ex Taq II system (TaKaRa perfect real time). RNA samples from seedlings of the control and treated plants were collected. Total RNA was isolated from cotton plants using RNAPrep Pure Plant kit Polysaccharides & Polyphenolics-Rich (TIANGEN biotech, Beijing, China), and reverse-transcribed using the PrimeScript RT-qPCR kit (TaKaRa, Dalian, China). The RNA concentrations were quantified by a NanoDrop ND-2000 spectrophotometer. The reverse

transcription reactions were performed using the primer RT-qPCR kit (TaKaRa, Dalian, China) according to the supplier's protocol. The expression profiles of the lncRNAs, miRNA and mRNA were assayed by RT-qPCR. 1 µg of total RNA was used for initiating reverse transcription, and the product was used as template for RT-qPCR using specific primers (primers shown in Additional file 10). Primers were then added to perform PCR. *UBQ7* expression was used as the internal control for RT-qPCR. RT-qPCR was performed as per the manufacturer's instructions (Takara). Briefly, 2 µl of cDNA template was added to 12.5 µl of SYBR premix ex Taq (Tli RNaseH plus), 0.5 µM concentration of each primer, and ddH<sub>2</sub>O to a final volume of 25 µl. The reactions were amplified for 10 s at 95 °C, followed by 40 cycles of 95 °C for 10 s and 59 °C for 30 s. All reactions were performed in triplicate, and controls (no template and no RT) were included for each gene. The  $2^{-\Delta\Delta C_t}$  method was used to calculate relative gene expression values [73–76].

## Additional files

- Additional file 1:** Mean quality distribution of samples in this study. (DOCX 52 kb)
- Additional file 2:** Percent of reads mapped to genome regions in this study. (DOC 111 kb)
- Additional file 3:** Conservation of *Gossypium hirsutum* lincRNAs. (XLS 5268 kb)
- Additional file 4:** The conservation score (consScore) of each nucleotide in the *G. hirsutum* genome. (PDF 3617 kb)
- Additional file 5:** Expression level of lincRNAs. (DOCX 85 kb)
- Additional file 6: Table S1.** Differential protein genes (XLSX 282 kb)
- Additional file 7:** Representatives of predicted interaction networks among lincRNAs and protein-coding RNAs. The triangular and square nodes represent lincRNAs and protein-coding genes. (PDF 41 kb)
- Additional file 8: Table S2.** Analysis of lincRNA and their putative target (XLS 24 kb)
- Additional file 9:** Putative targets and target mimics of lincRNAs Inc\_973. (XLSX 11 kb)
- Additional file 10: Table S3.** Primer list for gene-specific primers. (DOCX 27 kb)

## Abbreviations

ceRNAs: competing endogenous RNAs; CNCI: Coding-Non-Coding Index; CPAT: Coding Potential Assessment Tool; CPC: Coding Potential Calculator; eTMs: endogenous target mimics; lncRNAs: long non-coding RNAs; RPKM: Reads per kilobase of exon per million fragments mapped; RT-qPCR: Real-time quantitative PCR

## Acknowledgments

Thanks to all the research workers in State Key Laboratory of Crop Biology laboratory for their contribution to our study.

## Funding

This research was mainly supported by the China Major Projects for Transgenic Breeding (grant nos. 2015zx08005–002), provided the fund, platform and reference material for our research.

## Availability of data and materials

The datasets generated and during the current study are publicly available in NCBI (<http://plantgrn.noble.org/psRNATarget/>), ID was PRJNA416197.

## Authors' contributions

FND and XPZ carried out the functional analysis, participated in the identification of lncRNAs and drafted the manuscript. WW and RY participated in RT-qPCR and data uploads. FND and XPZ participated in the design of the study and analyzed the data. FFS conceived of the study, and participated in its design and coordination and helped to draft the manuscript. FND, and WW were in charge of the revision of the manuscript. All authors read and approved the final manuscript.

## Ethics approval and consent to participate

SN91–11, a salt-tolerant cotton cultivar, was used in this research. This cultivar was obtained by introducing Bluish Dogbane (*Apocynum venetum*) DNA into LM-6, cultivating by State Key Laboratory of Crop Biology, Shandong Agricultural University.

## Consent for publication

Not applicable.

## Competing interests

All authors declared that they have no competing financial interests.

## Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Received: 23 May 2017 Accepted: 17 January 2018

Published online: 25 January 2018

## References

- Jacob F, Monod J. Genetic regulatory mechanisms in the synthesis of proteins. *J Mol Biol.* 1961;3:318–56.
- Amaral PP, Dinger ME, Mercer TR, Mattick JS. The eukaryotic genome as an RNA machine. *Science.* 2008;319(5871):1787.
- Ponting CP, Oliver PL, Reik W. Evolution and functions of long noncoding RNAs. *Cell.* 2009;136(4):629–41.
- Rinn JL, Chang HY. Genome regulation by long noncoding RNAs. *Annu Rev Biochem.* 2012;81:145–66.
- Rymarquis LA, Kastenmayer JP, Hüttenhofer AG, Green PJ. Diamonds in the rough: mRNA-like non-coding RNAs. *Trends Plant Sci.* 2008;13(7):329–34.
- Guttman M, Amit I, Garber M, French C, Lin MF, Feldser D, Lander ES. Chromatin signature reveals over a thousand highly conserved large non-coding RNAs in mammals. *Nature.* 2009;458(7235):223–7.
- Liu J, Jung C, Xu J, Wang H, Deng S, Bernad L, et al. Genome-wide analysis uncovers regulation of long intergenic noncoding RNAs in *Arabidopsis*. *Plant Cell.* 2012;24(11):4333–45.
- Di C, Yuan J, Wu Y, Li J, Lin H, Hu L, Zhang T, Qi Y, Gerstein MB, Guo Y, Lu ZJ. Characterization of stress-responsive lincRNAs in *Arabidopsis thaliana* by integrating expression, epigenetic and structural features. *Plant J.* 2014;80:848–61.
- Heo JB, Sung S. Vernalization-mediated epigenetic silencing by a long intronic noncoding RNA. *Science.* 2011;331(6013):76–9.
- Kim YJ, Zheng B, Yu Y, Won SY, Mo B, Chen X. The role of mediator in small and long noncoding RNA production in *Arabidopsis thaliana*. *EMBO J.* 2011;30(5):814–22.
- Wang Z-W, Wu Z, Raitskin O, Sun Q, Dean C. Antisense-mediated FLC transcriptional repression requires the P-TEFb transcription elongation factor. *Proc Natl Acad Sci U S A.* 2014;111(20):7468–73.
- Brown CJ, Hendrich BD, Rupert JL, Lafrenière RG, Xing Y, Lawrence J, Willard HF. The human XIST gene: analysis of a 17 kb inactive X-specific RNA that contains conserved repeats and is highly localized within the nucleus. *Cell.* 1992;71(3):527–42.
- Yamada K, Lim J, Dale JM, Chen H, Shinn P, Palm CJ, Ecker JR. Empirical analysis of transcriptional activity in the genome. *Science.* 2003;302(5646):842.
- Osato N, Yamada H, Satoh K, Ooka H, Yamamoto M, Suzuki K, Hayashizaki Y. (2004). Antisense transcripts with rice full-length cDNAs. *Genome Biol.* 5(1), R5-R5.



15. Kouchi H, Takane K-i, So RB, Ladha JK, Reddy PM. Rice ENOD40: isolation and expression analysis in rice and transgenic soybean root nodules. *Plant J.* 1999;18(2):121–9.
16. Campalans A, Kondorosi A, Crespi M. Enod40, a short open reading frame-containing mRNA, induces cytoplasmic localization of a nuclear RNA binding protein in *Medicago Truncatula*. *Plant Cell.* 2004;16(4):1047–59.
17. Kornienko AE, Guenzl PM, Barlow DP, & Pauler FM. (2013). Gene regulation by the act of long non-coding RNA transcription. *BMC Biol.* 11, 59–59.
18. Liu J, Wang H, Chua N-H. Long noncoding RNA transcriptome of plants. *Plant Biotechnol J.* 2015;13(3):319–28.
19. Wang H, Chung PJ, Liu J, Jang I-C, Kean MJ, Xu J, Chua N-H. Genome-wide identification of long noncoding natural antisense transcripts and their responses to light in *Arabidopsis*. *Genome Res.* 2014;24(3):444–53.
20. Neil H, Malabat C, d'Aubenton-Carafa Y, Xu Z, Steinmetz LM, Jacquier A. Widespread bidirectional promoters are the major source of cryptic transcripts in yeast. *Nature.* 2009;457(7232):1038–42.
21. Valencia P, Dias AP, Reed R. Splicing promotes rapid and efficient mRNA export in mammalian cells. *Proc Natl Acad Sci U S A.* 2008;105(9):3386–91.
22. Boerner S, McGinnis KM. Computational identification and functional predictions of long noncoding RNA in *Zea Mays*. *PLoS One.* 2012;7(8):e43047e.
23. Shin H, Shin H-S, Chen R, Harrison MJ. Loss of At4 function impacts phosphate distribution between the roots and the shoots during phosphate starvation. *Plant J.* 2006;45(5):712–26.
24. Ding J, Lu Q, Ouyang Y, Mao H, Zhang P, Yao J, et al. A long noncoding RNA regulates photoperiod-sensitive male sterility, an essential component of hybrid rice. *Proc Natl Acad Sci U S A.* 2012;109(7):2654–9.
25. Zhang Y-C, Liao J-Y, Li Z-Y, Yu Y, Zhang J-P, Li Q-F, et al. Genome-wide screening and functional analysis identify a large number of long noncoding RNAs involved in the sexual reproduction of rice. *Genome Biol.* 2014;15(12):512.
26. Wang K, Wang Z, Li F, Ye W, Wang J, Song G, et al. The draft genome of a diploid cotton *Gossypium Raimondii*. *Nat Genet.* 2012;44(10):1098–103.
27. Paterson AH, Wendel JF, Gundlach H, Guo H, Jenkins J, Jin D., ... Schmutz, J. (2012). Repeated polyploidization of *Gossypium* genomes and the evolution of spinnable cotton fibres. *Nature.* 492(7429), 423–427.
28. Li F, Fan G, Wang K, Sun F, Yuan Y, Song G, et al. Genome sequence of the cultivated cotton *Gossypium Arboreum*. *Nat Genet.* 2014;46(6):567–72.
29. Li F, Fan G, Lu C, Xiao G, Zou C, Kohel RJ, et al. Genome sequence of cultivated upland cotton (*Gossypium Hirsutum* TM-1) provides insights into genome evolution. *Nat Biotechnol.* 2015;33(5):524–30.
30. Zhang T, Hu Y, Jiang W, Fang L, Guan X, Chen J, et al. Sequencing of allotetraploid cotton (*Gossypium Hirsutum* L. acc. TM-1) provides a resource for fiber improvement. *Nat Biotech.* 2015;33(5):531–7.
31. Yuan D, Tang Z, Wang M, Gao W, Tu L, Jin X, et al. The genome sequence of Sea-Island cotton (*Gossypium Barbardense*) provides insights into the allopolyploidization and development of superior spinnable fibres. *Sci Rep.* 2015;5:17662.
32. Page JT, Huynh MD, Liechty ZS, Grupp K, Stelly D, Hulse AM, et al. Insights into the evolution of cotton diploids and Polyploids from whole-genome re-sequencing. *G3: Genes|Genomes|Genetics.* 2013;3(10):1809–18.
33. Ragab R, Malash N, Gawad GA, Arslan A, Ghaibeh A. A holistic generic integrated approach for irrigation, crop and field management: 2. The SALTMED model validation using field data of five growing seasons from Egypt and Syria. *Agric Water Manag.* 2005;78(1–2):89–107.
34. Viswanathan Chinnusamy AJ, Zhu J-K. Understanding and improving salt tolerance in plants. *Crop Sci.* 2005;45:437–48.
35. Sexton PD, Gerard CJ. Emergence force of cotton seedlings as influenced by Salinity1. *Agron J.* 1982;74:699–702.
36. Ashraf M. Salt tolerance of cotton: some new advances. *Crit Rev Plant Sci.* 2002;21(1):1–30.
37. Wang M, Yuan D, Tu L, Gao W, He Y, Hu H, et al. Long noncoding RNAs and their proposed functions in fibre development of cotton (*Gossypium* spp.). *New Phytol.* 2015;207(4):1181–97.
38. Wang T-Z, Liu M, Zhao M-G, Chen R, Zhang W-H. Identification and characterization of long non-coding RNAs involved in osmotic and salt stress in *Medicago Truncatula* using genome-wide high-throughput sequencing. *BMC Plant Biol.* 2015;15(1):1–13. doi:10.1186/s12870-015-0530-5.
39. Cabili MN, Trapnell C, Goff L, Koziol M, Tazon-Vega B, Regev A, Rinn JL. Integrative annotation of human large intergenic noncoding RNAs reveals global properties and specific subclasses. *Genes Dev.* 2011;25:1915–27.
40. Lamesch P, Berardini TZ, Li D, Swarbreck D, Wilks C, Sasidharan R, Muller R, Dreher K, Alexander DL, Garcia-Hernandez M, Karthikeyan AS, Lee CH, Nelson WD, Ploetz L, Singh S, Wensel A, Huala E. The Arabidopsis information resource (TAIR): improved gene annotation and new tools. *Nucleic Acids Res.* 2012;40:D1202–10.
41. Blanchette M, Kent WJ, Riemer C, Elnitski L, Smit AF, Roskin KM, Baertsch R, Rosenbloom K, Clawson H, Green ED, Haussler D, Miller W. Aligning multiple genomic sequences with the threaded blockset aligner. *Genome Res.* 2004;14:708–15.
42. Siepel A, Bejerano G, Pedersen JS, Hinrichs AS, Hou M, Rosenbloom K CH, Spieth J, Hillier LW, Richards S, Weinstock GM, Wilson RK, Gibbs RA, Kent WJ, Miller W, Haussler D. Evolutionarily conserved elements in vertebrate, insect, worm, and yeast genomes. *Genome Res.* 2005;15:1034–50.
43. Hanano S, Goto K. Arabidopsis TERMINAL FLOWER1 is involved in the regulation of flowering time and inflorescence development through transcriptional repression. *Plant Cell.* 2011;23(9):3172–84. doi:10.1105/tpc.111.088641.
44. Rinn JL, Kertesz M, Wang JK, Squazzo SL, Xu X, Bruggmann SA, et al. Functional demarcation of active and silent chromatin domains in human HOX loci by noncoding RNAs. *Cell.* 2007;129(7):1311–23.
45. Pauli A, Valen E, Lin MF, Garber M, Vastenhouw NL, Levin JZ, et al. Systematic identification of long noncoding RNAs expressed during zebrafish embryogenesis. *Genome Res.* 2012;22(3):577–91.
46. Franco-Zorrilla JM, Valli A, Todesco M, et al. Target mimicry provides a new mechanism for regulation of microRNA activity. *Nat Genet.* 2007;39:1033–7.
47. Meng Y, Shao C, Wang H, Jin Y. Target mimics: an embedded layer of microRNA-involved gene regulatory networks in plants. *BMC Genomics.* 2012;13:197.
48. Wu H-J, Wang Z-M, Wang M, Wang X-J. Widespread long noncoding RNAs as endogenous target mimics for microRNAs in plants. *Plant Physiol.* 2013;161:1875–84.
49. Dai X, Zhao PX. psRNATarget: a plant small RNA target analysis server. *Nucleic Acids Res.* 2011;39:W155–9.
50. Shi G, Guo X, Guo J, Liu L, Hua J. Analyzing serial cDNA libraries revealed reactive oxygen species and gibberellins signaling pathways in the salt response of upland cotton (*Gossypium Hirsutum* L.). *Plant Cell Rep.* 2015;34(6):1005–23.
51. Cabello JV, Lodeyro AF, Zurbriggen MD. Novel perspectives for the engineering of abiotic stress tolerance in plants. *Curr Opin Biotechnol.* 2014;26C:62–70.
52. Deinlein U, Stephan AB, Horie T, Luo W, Xu G, Schroeder JI. Plant salt-tolerance mechanisms. *Trends Plant Sci.* 2014;19:371–6.
53. Gollack D, Luking I, Yang O. Plant tolerance to drought and salinity: stress regulating transcription factors and their functional significance in the cellular transcriptional network. *PlantCell Rep.* 2011;30:1383–91.
54. Peng Z, He S, Gong W, et al. Comprehensive analysis of differentially expressed genes and transcriptional regulation induced by salt stress in two contrasting cotton genotypes. *BMC Genomics.* 2014;15:760.
55. Kapusta A, Feschotte C. Volatile evolution of long noncoding RNA repertoires: mechanisms and biological implications. *Trends Genet.* 2014;30:439–52.
56. Kim ED, Sung S. Long noncoding RNA: unveiling hidden layer of gene regulatory networks. *Trends Plant Sci.* 2012;17:16–21.
57. Schmitz K-M, Mayer C, Postepska A, Grummt I. Interaction of noncoding RNA with the rDNA promoter mediates recruitment of DNMT3b and silencing of rRNA genes. *Genes Dev.* 2010;24(20):2264–9.
58. Martianov I, Ramadass A, Serra Barros A, Chow N, Akoulitchev A. Repression of the human dihydrofolate reductase gene by a non-coding interfering transcript. *Nature.* 2007;445(7128):666–70.
59. Jeon Y, Lee JT. YY1 tethers Xist RNA to the inactive X nucleation center. *Cell.* 2011;146(1):119–33.
60. Lee JT. Lessons from X-chromosome inactivation: long ncRNA as guides and tethers to the epigenome. *Genes Dev.* 2009;23:1831–42.
61. Luo Y, Guo Z, Li L. Evolutionary conservation of microRNA regulatory programs in plant flower development. *Dev Biol.* 2013;380:133–44.
62. Wu HJ, Ma YK, Chen T, Wang M, Wang XJ. PsRobot: a web-based plant small RNA meta-analysis toolbox. *Nucleic Acids Res.* 2012;40:W22–8.
63. Guttman M, Rinn JL. Modular regulatory principles of large non-coding RNAs. *Nature.* 2012;482:339–46.



64. Zhang W, Han Z, Guo Q, Liu Y, Zheng Y, et al. Identification of maize long non-coding RNAs responsive to drought stress. *PLoS One*. 2014;9(6):e98958.
65. Chen CX, Yu YJ, Wang HG, Shen FF, Liu FZ. RAPD analysis and physiologic studies on salt-tolerance variation of cotton. *Acta Agron Sin*. 1999;5(5):644–6.
66. Chen CX, Yu YJ, Liu FZ, Shen FF, Wang HG. Genetic analysis of salt-tolerance variant in cotton. *Acta Bot Boreali-Occidental Sin*. 2000;20(2):234–7.
67. Yin Z, Li Y, Yu J, Liu Y, Li C, Han X, Shen F. Difference in miRNA expression profiles between two cotton cultivars with distinct salt sensitivity. *Mol Biol Rep*. 2012;39(4):4961–70.
68. Langmead B, Salzberg SL. Fast gapped-read alignment with bowtie 2. *Nat Methods*. 2012;9(4):357–9.
69. Trapnell C, Pachter L, Salzberg S L. 2009. TopHat: discovering splice junctions with RNA-Seq[J]. *Bioinformatics*,25 (9): 1105-1111. (TopHat).
70. Wang L, Park HJ, Dasari S, Wang S, Kocher J-P, Li W. CPAT: coding-potential assessment tool using an alignment-free logistic regression model. *Nucleic Acids Res*. 2013;41(6):e74–e74.
71. Trapnell C, Roberts A, Goff L, Pertea G, Kim D, Kelley DR, .. Pachter L. (2012). Differential gene and transcript expression analysis of RNA-seq experiments with TopHat and cufflinks. *Nat Protoc*, 7(3), 562–578.
72. Trapnell C, Williams BA, Pertea G, Mortazavi A, Kwan G, van Baren MJ, .. Pachter L. (2010). Transcript assembly and abundance estimation from RNA-Seq reveals thousands of new transcripts and switching among isoforms. *Nat Biotechnol*, 28(5), 511–515.
73. Pfaffl MW. A new mathematical model for relative quantification in real-time RT-PCR. *Nucleic Acids Research*. 2001;29:2002–7.
74. Chang S, Chen W, Yang J. Another formula for calculating the gene change rate in real-time RT-PCR. *Mol Biol Rep*. 2008;36(8):2165–8.
75. Chen C, Ridzon DA, Broomer AJ, Zhou Z, Lee DH, Nguyen JT, .. Guegler KJ (2005). Real-time quantification of microRNAs by stem-loop RT-PCR. *Nucleic Acids Res*, 33(20), e179–e179.
76. Ballantyne MD, McDonald RA, Baker AH. LncRNA/microRNA interactions in the vasculature. *Clin Pharmacol Ther*. 2016;99(5):494–501.

Submit your next manuscript to BioMed Central and we will help you at every step:

- We accept pre-submission inquiries
- Our selector tool helps you to find the most relevant journal
- We provide round the clock customer support
- Convenient online submission
- Thorough peer review
- Inclusion in PubMed and all major indexing services
- Maximum visibility for your research

Submit your manuscript at  
[www.biomedcentral.com/submit](http://www.biomedcentral.com/submit)

