BMC Medical Research Methodology

CORRECTION   **Open Access**

Correction to: Estimation of treatment preference effects in clinical trials when some participants are indifferent to treatment choice

Stephen D. Walter[1*], Robin M. Turner[2], Petra Macaskill[3], Kirsten J. McCaffery[3] and Les Irwig[3]

Correction to: BMC Med Res Methodol
https://doi.org/10.1186/s12874-017-0304-x

In the original publication of this article [1], there is a small calculational error in the illustrative example. The estimated second preference contrast $\Delta\pi'$ should be -3.23, not -0.49 as shown. All the other numerical results in the example are shown correctly. This change does not materially affect the substantive interpretation of the analysis.

Author details
[1]Department of Clinical Epidemiology and Biostatistics, McMaster University, CRL 233, Hamilton, ON L8N 3Z5, Canada. [2]School of Public Health and Community Medicine, University of New South Wales, Sydney, NSW 2052, Australia. [3]Screening and Test Evaluation Program, Sydney School of Public Health, Sydney Medical School, University of Sydney, Sydney, NSW 2006, Australia.

Published online: 14 April 2020

Reference
1.   Walter, et al. Estimation of treatment preference effects in clinical trials when some participants are indifferent to treatment choice. BMC Med Res Methodol. 2017;17:29.