## CORRECTION Open Access



## Correction to: Iron-free and iron-saturated bovine lactoferrin inhibit survivin expression and differentially modulate apoptosis in breast cancer

Jessica A. Gibbons, Jagat R. Kanwar and Rupinder K. Kanwar\*

Correction to: BMC Cancer (2015) 15:425, DOI: 10.1186/s12885-015-1441-4

It has been highlighted that the original article [1] contains an error in Fig. 3 specifically, that it showed an incorrect version of Fig. 3c. This does not affect the Figure legend, results and conclusions of the article. The correct version of Fig. 3 is shown in this Correction article.


Received: 2 July 2018 Accepted: 2 July 2018 Published online: 19 July 2018

## Reference

 Gibbons JA, et al. Iron-free and iron-saturated bovine lactoferrin inhibit surviving expression and differentially modulate apoptosis in breast cancer. BMC Cancer. 2015;15:425. https://doi.org/10.1186/s12885-015-1441-4.

<sup>\*</sup> Correspondence: rupinder.kanwar@deakin.edu.au Nanomedicine - Laboratory for Immunology and Molecular Biomedical Research, Molecular and Medical Research Facility, School of Medicine, Faculty of Health, Deakin University, Geelong, Victoria, Australia





**Fig. 3** Effect on migration and invasion capacity of MDA-MB-231 and MCF-7 cells after treatment with bLf. Migration of MDA-MB-231 and MCF-7 (a) cells after bLf treatments for 24 h represented as a percentage of untreated (1% FBS) control migration. Invasion of MDA-MB-231 and MCF-7 cells (b) after bLf treatments for 24 h represented as a percentage of untreated (1% FBS) control invasion. \* = p < 0.05 compared with 1% FBS group. Representative images (250X magnification) of invaded cells stained with crystal violet (c)