

RESEARCH ARTICLE

Open Access



# Association of socioeconomic and lifestyle-related risk factors with mental health conditions: a cross-sectional study

Miwako Nagasu<sup>1\*</sup> , Kazutaka Kogi<sup>2</sup> and Isamu Yamamoto<sup>3</sup>

## Abstract

**Background:** There is rising public concern over the widening health inequalities in many countries. The aim of this study was to clarify the associations of socioeconomic status (SES)-related variables, such as levels of household disposable income and employment status, and lifestyle factors with mental health conditions among Japanese adults aged 40 to 69.

**Methods:** A cross-sectional study of 3085 participants (1527 males and 1558 females) was undertaken by using a self-administered questionnaire that included the Japanese version of the 12-item General Health Questionnaire (GHQ-12) and questions related to socioeconomic and lifestyle factors.

**Results:** The prevalence of poor mental health conditions, represented by a GHQ-12 score of 4 or more, was 33.4% among males and 40.4% among females. Males whose annual household disposable income was less than 2 million yen had significantly higher GHQ-12 scores than those with an annual household disposable income above 2 million yen. As per binary logistic regression analyses, short sleep duration and the absence of physical exercise were significantly related to poor mental health conditions among both males and females. Among females, a household disposable income of less than 2 million yen could be a risk factor for poor mental health conditions. Age and habitual drinking were inversely associated with poor mental health conditions.

**Conclusions:** Low levels of household disposable income and unhealthy lifestyle factors were significantly associated with mental health conditions. These results suggest the importance of improving unhealthy lifestyle behaviours and developing effective health promotion programmes. In addition, there is a need for social security systems for people from different socioeconomic backgrounds.

**Keywords:** Mental health, Household disposable income, Socioeconomic status, Lifestyle, General Health questionnaire

## Background

The widening health inequalities associated with economic disparities have received significant attention in many countries, since previous studies have found that economic disparities associated with socioeconomic status (SES) may actually contribute to inequalities in health [1, 2]. Such studies have commonly taken account of income levels, employment status, occupations, and differences in educational levels, demonstrating such

socioeconomic factors to be main determinants of both mental and physical health [3–5].

Mortality due to severe mental illnesses is a global public health concern [6]. In particular, severe mental illnesses can lead to suicide [7]. Kawakami et al. reported that the prevalence of mental disorders as assessed with the World Mental Health version of the World Health Organization (WHO) Composite International Diagnostic Interview (WMH-CIDI) and the Diagnostic and Statistical Manual of Mental Disorders 4th edition (DSM-IV) among the Japanese population was 8.8%, of which 17.0% of cases had severe mental disorders [8]. Moreover, Japan has one of the highest suicide rates in the

\* Correspondence: [mnagasu555@gmail.com](mailto:mnagasu555@gmail.com)

<sup>1</sup>Faculty of Economics, Keio University, 〒108-8345 Tokyo-to, Minato-ku, Mita 2-15-45, Tokyo, Japan

Full list of author information is available at the end of the article



world [9]. Qin et al. revealed that mental illnesses and SES-related variables, such as low income and unemployment, were significantly associated with higher suicide risk [10]. Two studies in the UK and Denmark also reported a strong association between current financial difficulties and mental disorders [11]. In addition, poor economic situation has been reported to be significantly associated with life dissatisfaction [12]. Therefore, the associations between mental health conditions, SES, and lifestyle factors have been receiving an increasing amount of attention [13].

The relationship between mental health and lifestyle factors such as sleep duration [14], habitual physical exercise [15, 16], smoking [17–19], and alcohol consumption has been studied. Among the risk factors for poor mental health, short sleep duration was reported to be linearly associated with psychological distress [14]. Low frequency of exercise [15, 16] and high smoking frequency [15] have also been known to significantly contribute to high scores on the 12-item General Health Questionnaire (GHQ-12). Regarding smoking habits, current and former smokers tended to experience more depressive symptoms than never smokers [17]. Taylor et al. reported that those who quit smoking displayed reduced depression, anxiety, and stress and improved mood and quality of life compared with current smokers [19]. Regarding alcohol consumption, it is a well-known cause of many major physical disease outcomes [20]. According to the WHO, alcohol consumption is responsible for approximately 5.9% of deaths worldwide [21]. Collins reported that people with lower SES seemed to bear a disproportionate burden of negative alcohol-related consequences [21]. In general, previous studies have reported linear relationships between mental health and each lifestyle factor. Therefore, this study adjusted for all potential risk factors in a single model to determine risk factors.

It has been reported that people with low SES are more likely to engage in unhealthy behaviours than those with high SES [20, 22]. For example, people with lower SES are more likely to be smokers, experiencing its negative impacts [22]. On the contrary, healthy lifestyle practices may be associated with a reduced risk of mental disorders. Lyu et al. reported that older South Koreans who engage in a greater number of healthy lifestyle practices have a higher overall level of health than those who do not engage in as many healthy lifestyle practices [23]. Identifying lifestyle-related risk factors would, thus, be beneficial in the promotion of improved mental health and might explain the background of income-related health inequalities.

As it has been consistently reported that the prevalence of mental illnesses and lifestyle-related factors differs between males and females, in the present study,

data analysis was stratified by gender [24]. Matud et al. reported that adherence to traditional gender roles had a great impact on psychological well-being [22]. With regard to lifestyle, one specific factor—sleep duration—is a risk factor for depression that has an inherent gender difference; it has been reported that females sleep longer than males. In particular, being pregnant has been associated with longer sleep hours [24]. The gender differences in healthy lifestyles could potentially play confounding roles in the determination of the association between mental health and lifestyle factors, necessitating gender stratification of analyses.

In this study, we analysed a large nationally representative data set and hypothesised that SES and lifestyle-related factors would be significantly associated with mental health conditions, even after controlling for all potential risk factors. Moreover, the associations for both genders were examined separately. Therefore, the objectives of this study were to (1) investigate the prevalence of psychological distress (GHQ-12 score  $\geq 4$ ) among Japanese people aged 40 to 69 and (2) separately identify the associations of SES and lifestyle-related factors with poor mental health conditions among males and females.

## Methods

This study used the data set in 2017 from the Japan Household Panel Survey (JHPS/KHPS) as a cross-sectional data set. This JHPS/KHPS data set has combined two longitudinal panel data set: the JHPS (the Japan Household Panel Survey) and the KHPS (the Keio Household Panel Survey). The data set used, the JHPS/KHPS in 2017, was collected from February to March 2017. The key characteristics of panel studies is that they collect repeated measures from the same participants at different points. The questionnaire for the study in 2017 was distributed to 5030 respondents who participated in the JHPS/KHPS survey in 2016. Of these 5030 respondents, 4626 (92.0%) returned completed questionnaires. In the final analysis of this study, 1541 questionnaires were excluded either because the respondents did not fall within the 40–69 age range or because of incomplete information regarding gender, age, or other important variables. Finally, the data of 3085 respondents were used for analysis (valid response rate: 61.3%). An informed consent form consisting of the aims and details of the study, as well as an assurance regarding anonymity and confidentiality, was sent to all respondents. Those who provided informed consent received the questionnaire.

## Sampling

The JHPS/KHPS respondents, selected through two-stage stratified random sampling. According to the

National Census Survey, Japan is divided into 24 strata, from which 354 research areas were randomly selected. Five to 10 individuals from each research area were selected by using population ratios based on basic resident register information. The KHPS began with 4005 respondents in 2004. In 2007 and 2012, 1400 and 1000 respondents, respectively, were added. As for the JHPS, it began with 4022 respondents in 2009. From 2004 to 2017, a total of 10,458 respondents participated. Over the course of 13 years, the number of respondents declined gradually, and new respondents were added to ensure the survey remained representative of Japan's population distribution.

### **Study variables**

A self-administered questionnaire was used to collect information about socioeconomic factors, lifestyle-related factors, and mental health outcomes.

### **Socioeconomic factors**

The socioeconomic factors included gender, age, employment status, number of persons in the household, and the level of disposable income per household and respondent. Based on age, the participants were divided into three groups: 40–49, 50–59, and 60–69. On the basis of employment status, two groups were created: full-time workers and others (self-employed persons, freelance professionals, part-time workers, and unemployed). The number of persons in the household was categorized as two or more (living with someone) and one (living alone). All respondents provided information regarding disposable income—both the family's and their own—in the last year. This disposable income excluded tax and social insurance fees. The level of disposable income per household was divided into three groups: less than two million yen, two million yen to under six million yen, and six million yen and above. Respondents' level of disposable income was divided into three groups: less than two million yen, two million yen to under five million yen, and five million yen and above.

### **Health outcomes**

The Japanese version of the GHQ-12, used as a screening tool to detect nonpsychotic psychiatric diseases, consists of 12 questions about feelings over the past few weeks. The questions include the following: Have you recently (1) been able to concentrate on whatever you're doing, (2) lost much sleep over worry, (3) felt that you were playing a useful part in things, (4) felt capable of making decisions, (5) felt constantly under strain, (6) felt you couldn't overcome your difficulties, (7) been able to enjoy your normal day-to-day activities, (8) been able to face problems, (9) been feeling unhappy or depressed, (10) been losing confidence in yourself, (11) been

thinking of yourself as a worthless person, and (12) been feeling reasonably happy, all things considered. The scoring system used was as follows: the response categories (1, 2, 3, and 4) were converted into corresponding binary values (0, 0, 1, and 1) to calculate the total score. The sum of the scores indicated the severity of psychological distress. The participants were then divided into two groups: those with high scores (poor mental health conditions:  $\geq 4$  points) and those with low scores (good mental health conditions:  $\leq 3$  points) [25]. Goldberg reported that sensitivity and specificity ranged from 67.0 to 93.5% (median 83.7%) and 59.0 to 93.0% (median 79.0%), respectively, when the cut-off point is between under 3 and 4 points and more as a screening tool of psychological distress [26].

### **Lifestyle factors**

This study included five lifestyle-related questions for assessing lifestyle practices: sleep duration on weekdays, physical exercise frequency, and smoking and drinking habits. About sleep duration on weekdays, the question was "How many hours do you usually sleep on weekdays?" The responses were categorised into three groups: 7 h or more, 6–7 h, and fewer than 6 h. Physical exercise frequency was determined by the question: "How many days per week do you engage in physical exercise that makes you sweat (excluding your work)?" The answers were categorised into three groups: three or more days, two or fewer days, and no exercise. By smoking habit, respondents were categorised as never smokers, ex-smokers, or current smokers. Alcohol consumption was assessed by the question: "How often do you drink alcohol?" The answers were categorised into the following groups: never, two or fewer times/week, and three or more times/week.

### **Statistical methods**

The gender-stratified association of socioeconomic and lifestyle factors with mental health conditions was analysed by Pearson's chi-squared test. The differences in GHQ-12 scores in household disposable income groups stratified by gender were analysed by the Kruskal-Wallis test with the Bonferroni correction and the Student's *t*-test. *P* values less than 0.05 indicated statistical significance. We investigated the adjusted prevalence odds ratios (AORs) and 95% confidence intervals (CI) of high GHQ-12 scores ( $\geq 4$  points) by using binary logistic regression analysis. All factors such as age, number of persons in the household, employment status, annual disposable income per household, and lifestyle-related factors were adjusted for. The data were analysed by gender, annual disposable income per household, and employment status. SPSS 24.0 was used for data analysis.

## Results

This study included 3085 respondents, of which 1527 (49.5%) were males and 1558 (50.5%) were females. The mean age of the males ( $54.9 \pm 8.8$  years) was similar to that of the females ( $54.5 \pm 8.7$  years). Table 1 depicts the gender-specific distribution of socioeconomic, lifestyle, and mental health variables. The results indicated statistically significant gender differences in the ratios of the corresponding categories of all variables except age and disposable household income. The rate of those living alone was 10.5% among males and 7.3% among females. Regarding employment status, 62.8% of the males worked full time, while this figure stood at 21.8% among females. In the context of socioeconomic variables, of those with an annual disposable income of less than two

million yen per respondent, 13.6% were males and 66.2% were females. As for lifestyle-related variables, more females than males had short sleep durations (< 6 h), no exercise, and no smoking and alcohol consumption; these differences were statistically significant. The overall rate of those with a high GHQ-12 score was 36.9%. Stratified by gender, 33.4% of the males and 40.4% of the females, respectively, had high GHQ-12 scores. These gender differences were statistically significant.

Table 2 depicts the differences in GHQ-12 scores by gender; in particular, in terms of stratified disposable income per household and employment status. Male respondents with a household disposable income of less than two million yen had significantly higher GHQ-12 scores than those in the higher-income

**Table 1** Demographic characteristics of the respondents by gender

Variables	Group	Total		Males		Females		P value <sup>1)</sup>
		n	%	n	%	n	%	
Gender	Males	1527	49.5					
	Females	1558	50.5					
Age (years)	40–49	1037	33.6	504	33.0	533	34.2	
	50–59	1018	33.0	490	32.1	528	33.9	n.s
	60–69	1030	33.4	533	34.9	497	31.9	
Number of persons in the household	≥ 2	2742	91.2	1325	89.5	1417	92.7	
	1 person	266	8.8	155	10.5	111	7.3	**
Employment status	Full time	1083	44.5	845	62.8	238	21.8	
	Others	1353	55.5	501	37.2	852	78.2	***
Disposable income per household (JPY)	< 2000 K	287	9.3	129	8.4	158	10.1	
	2000 K–< 6000 K	1479	47.9	746	48.9	733	47.0	n.s.
	≥ 6000 K	1319	42.8	652	42.7	667	42.8	
Disposable income of the respondent (JPY)	< 2000 K	880	37.1	178	13.6	702	66.2	
	2000 K–< 5000 K	805	33.9	515	39.2	290	27.3	***
	≥ 5000 K	689	29.0	620	47.2	69	6.5	
Sleep duration on weekdays (hours)	≥ 7	1241	40.4	656	43.2	585	37.7	
	6–7	1166	38.0	563	37.1	603	38.9	**
	< 6	663	21.6	299	19.7	364	23.5	
Physical exercise (days/week)	≥ 3	429	14.0	225	14.9	204	13.2	
	≤ 2	530	17.3	293	19.4	237	15.3	**
	No exercise	2097	68.6	994	65.7	1103	71.4	
Smoking	Never	1605	52.2	449	29.5	1156	74.5	
	Quit	836	27.2	610	40.1	226	14.6	***
	Sometimes/everyday	633	20.6	463	30.4	170	11.0	
Drinking (times/week)	Never	1168	38.0	391	25.7	777	50.0	
	≤ 2	898	29.2	401	26.4	497	32.0	***
	≥ 3	1008	32.8	729	47.9	279	18.0	
GHQ-12 score	≥ 4 (poor)	1131	36.9	506	33.4	625	40.4	***
	≤ 3	1933	63.1	1010	66.6	923	59.6	

<sup>1)</sup> P value from Pearson's chi-squared test: \*\*  $P < 0.01$ , \*\*\*  $P < 0.001$

**Table 2** Differences in GHQ-12 scores by gender, disposable income per household, and employment status

Disposable income per household						P value <sup>1)</sup>		Employment status				P value <sup>2)</sup>
< 2000 K		2000 K – < 6000 K		≥ 6000 K		2000 K – < 6000 K	≥ 6000 K	Full-time		Others		
mean	SD	mean	SD	mean	SD			mean	SD	mean	SD	
Males												
3.992	4.007	2.878	3.295	2.981	3.402	*	*	3.033	3.360	2.879	3.405	n.s.
Females												
3.790	3.351	3.480	3.340	3.325	3.410	n.s.	n.s.	3.705	3.372	3.579	3.355	n.s.

<sup>1)</sup> The Shapiro-Wilk test was used to assess normality of data distribution for GHQ-12 scores. Comparing GHQ-12 scores between three groups (Ref: Household disposable income < 2000 K). P value from Kruskal-Wallis test with Bonferroni correction: \*P < 0.05

<sup>2)</sup> Mann-Whitney U test

groups. This tendency was not statistically significant among the females.

As Table 3 depicts, household disposable income levels and employment status were associated with mental health conditions and lifestyle factors. Among all respondents, though it was particularly true of males, earning less than two million yen was associated with living alone and not having a full-time job. Regarding lifestyle factors, males with a household disposable income over six million yen had shorter sleep durations than those who earned less than six million yen. About habitual physical exercise, males with a household disposable income of six million yen and above had a higher exercise frequency ( $\geq 3$  days/week) than those with a household disposable income under two million yen. Male respondents with a household disposable income under two million yen were less likely to engage in physical exercise than those in higher-income categories. About smoking habits, both males and females who were categorised into the group with a household disposable income under two million yen were significantly more likely to be smokers than those categorised into the group with a household disposable income of six million yen and above. About alcohol consumption, male respondents with a household disposable income under two million yen had a significantly lower alcohol consumption frequency ( $\geq 3$  times/week) than those with a household disposable income of 6 million yen and above.

Tables 4 and 5 depict the rates of those with high GHQ-12 scores ( $\geq 4$  points), crude odds ratios (OR) and adjusted odds ratios (AOR) by using binary logistic regression analyses. Males who lived alone and had a disposable income of six million yen or above showed significantly higher AORs (AOR 2.548 [95% CI: 1.049–6.189]). About lifestyle-related factors, short sleep duration (< 6 h) had a significantly greater association with poor mental health than sleeping over 6 h per day (males in total: AOR 1.613 [95% CI: 1.171–2.221], males with income 2 million– $\geq 6$  million: AOR 2.199 [95% CI: 1.383–3.498], males without full-time work: AOR 1.837 [95% CI: 0.531–6.363]). Lack of physical exercise was

also significantly associated with poor mental health conditions (males in total: AOR 1.663 [95% CI: 1.122–2.465], males with income  $\geq 6$  million: 1.743 [95% CI: 0.996–3.048], males without full-time work: AOR 5.175 [95% CI: 2.014–13.295]). Therefore, sleeping for fewer than 6 h and a lack of physical exercise could be risk factors for poor mental health among males.

Among females, household disposable income (< 2 million yen: AOR 1.592 [95% CI: 1.000–2.535]) was significantly associated with poor mental health conditions. About lifestyle factors, female respondents with short sleep duration (< 6 h) (females in total: AOR 1.505 [95% CI: 1.077–2.104], full-time work: AOR 2.288 [95% CI: 1.103–4.746]) and without physical exercise (females in total: AOR 1.738 [95% CI: 1.120–2.696], non-full-time work: AOR 1.298 [95% CI: 0.754–2.235]) also had significantly higher rates of those with poor mental health conditions as compared with respondents who slept over 6 h per day and engaged in physical exercise.

However, the results show the existence of an inverse association between age and alcohol consumption and poor mental health. The following AOR levels were statistically significant (females overall aged 50–59: AOR 0.680 [95% CI: 0.511–0.904], aged 60–69: AOR 0.676 [95% CI: 0.479–0.954], females with  $\geq 6$  million aged 50–59: AOR 0.641 [95% CI: 0.422–0.974], aged 60–69: AOR 0.512 [95% CI: 0.283–0.926], working full-time aged 60–69: AOR 0.257 [95% CI: 0.085–0.781], not working full-time aged 50–59: AOR 0.577 [95% CI: 0.328–1.017]). Concerning alcohol consumption, among females with an income under two million yen, there was an inverse association between frequent drinking and mental health conditions ( $\leq 2$  times/week AOR 0.274 [95% CI: 0.100–0.751],  $\geq 3$  times/week: AOR 0.289 [95% CI: 0.092–0.906]).

## Discussion

The results of this study revealed that males with low household disposable income (< 2 million yen) showed psychological distress with the highest GHQ-12 scores among three income levels. Moreover, according to the results of binary logistic regression analyses after





**Table 4** Prevalence and associations of the GHQ-12 score with risk factors among males

Males		Disposable income per household												Employment status										
		< 2000 K				2000 K – < 6000 K				≥ 6000 K				Full time		Others								
		n	%	Crude OR <sup>b)</sup>	95% CI	Adjusted OR <sup>c)</sup>	95% CI	Lower	Upper	Adjusted OR <sup>d)</sup>	95% CI	Lower	Upper	Adjusted OR <sup>e)</sup>	95% CI	Lower	Upper							
Age (years)																								
40–49	167	33.3	1		1			1		1			1			1								
50–59	177	36.3	1.145	0.881	1.488	1.153	0.870	1.528	0.872	0.249	3.049	1.344	0.879	2.055	1.096	0.729	1.647	1.258	0.908	1.742	1.267	0.269	5.970	
60–69	165	30.7	0.890	0.685	1.157	0.908	0.650	1.268	1.101	0.339	3.576	0.781	0.482	1.266	1.191	0.699	2.026	0.845	0.515	1.387	0.391	0.125	1.221	
Number of persons in the household																								
≥ 2	427	32.4	1		1			1		1			1			1		1						
1	68	45.0	<b>1.711</b>	<b>1.217</b>	<b>2.406</b>	1.495	0.992	2.253	1.237	0.465	3.286	1.220	0.702	2.123	<b>2.548</b>	<b>1.049</b>	<b>6.189</b>	1.418	0.830	2.421	1.498	0.566	3.962	
Employment status																								
Full time	284	33.8	1		1			1		1			1			1		NA <sup>f)</sup>				NA <sup>f)</sup>		
Others	156	31.4	0.896	0.706	1.136	0.970	0.732	1.284	0.993	0.380	2.593	0.937	0.626	1.402	1.002	0.640	1.568	NA <sup>f)</sup>				NA <sup>f)</sup>		
Household disposable income (JPY)																								
≥ 6000 K	206	31.9	1		1			NA <sup>f)</sup>		NA <sup>f)</sup>			NA <sup>f)</sup>			NA <sup>f)</sup>		1				1		
2000 K–< 6000 K	245	33.0	1.053	0.840	1.319	1.027	0.793	1.331	NA <sup>f)</sup>	NA <sup>f)</sup>			NA <sup>f)</sup>			NA <sup>f)</sup>		1.177	0.852	1.626	0.682	0.291	1.594	
< 2000 K	55	43.0	<b>1.609</b>	<b>1.093</b>	<b>2.370</b>	1.535	0.918	2.567	NA <sup>f)</sup>	NA <sup>f)</sup>			NA <sup>f)</sup>			NA <sup>f)</sup>		1.693	0.770	3.723	0.714	0.263	1.939	
Sleep duration on weekdays (hours)																								
≥ 7	208	31.9	1		1			1		1			1			1		1				1		
6–7	171	30.5	0.938	0.735	1.197	0.925	0.700	1.221	1.080	0.367	3.181	1.110	0.744	1.655	0.735	0.478	1.132	1.035	0.723	1.481	1.019	0.471	2.203	
< 6	126	42.3	<b>1.564</b>	<b>1.179</b>	<b>2.075</b>	<b>1.613</b>	<b>1.171</b>	<b>2.221</b>	0.985	0.253	3.830	<b>2.199</b>	<b>1.383</b>	<b>3.498</b>	1.233	0.762	1.995	1.460	0.976	2.182	<b>1.837</b>	<b>0.531</b>	<b>6.363</b>	
Physical exercise (days/week)																								
≥ 3	49	21.9	1		1			1		1			1			1		1				1		
≤ 2	86	29.5	1.491	0.995	2.234	1.250	0.795	1.964	1.061	0.147	7.680	1.101	0.530	2.288	1.352	0.728	2.511	1.030	0.575	1.846	7.551	1.885	30.247	
0	367	37.1	<b>2.111</b>	<b>1.499</b>	<b>2.971</b>	<b>1.663</b>	<b>1.122</b>	<b>2.465</b>	1.037	0.182	5.925	1.588	0.870	2.897	<b>1.743</b>	<b>0.996</b>	<b>3.048</b>	1.497	0.885	2.533	<b>5.175</b>	<b>2.014</b>	<b>13.295</b>	
Smoking																								
Never	153	34.1	1		1			1		1			1			1		1				1		
Quit	197	32.6	0.936	0.723	1.213	0.929	0.693	1.246	0.984	0.257	3.770	0.848	0.549	1.311	1.012	0.660	1.553	1.030	0.715	1.485	1.315	0.587	2.945	
Sometimes/ everyday	155	33.6	0.980	0.745	1.290	0.927	0.682	1.261	1.507	0.443	5.128	0.784	0.494	1.244	1.010	0.638	1.597	0.926	0.631	1.357	1.549	0.583	4.112	

**Table 4** Prevalence and associations of the GHQ-12 score with risk factors among males (Continued)

Males		Disposable income per household						Employment status															
		< 2000 K		2000 K – < 6000 K		≥ 6000 K		Full time		Others													
Prevalence (≥ 4) <sup>a)</sup>	Crude OR <sup>b)</sup>	Adjusted OR <sup>d)</sup>	95% CI	Adjusted OR <sup>d)</sup>	95% CI	Adjusted OR <sup>d)</sup>	95% CI	Adjusted OR <sup>e)</sup>	95% CI	Adjusted OR <sup>e)</sup>	95% CI												
n	%	Lower	Upper	Lower	Upper	Lower	Upper	Lower	Upper	Lower	Upper												
Drinking (times/week)																							
0	141	36.2	1	1	1	1	1	1	1	1	1												
≤ 2	122	30.5	0.772	0.574	1.039	0.881	0.632	1.228	1.306	0.364	4.686	0.910	0.573	1.445	0.745	0.437	1.268	0.910	0.602	1.376	0.714	0.261	1.952
≥ 3	241	33.3	0.878	0.678	1.136	1.055	0.780	1.426	1.171	0.369	3.710	1.081	0.701	1.668	1.013	0.631	1.626	1.136	0.771	1.673	0.878	0.404	1.907

<sup>a)</sup> Prevalence of GHQ-12 score ≥ 4

<sup>b)</sup> Crude odds ratio with 95% confidence interval (no adjustment for any variables)

<sup>c)</sup> Adjusted odds ratios with 95% confidence interval (adjusted for age, number of persons in the household, employment status, disposable income per household, sleep duration on weekdays, physical exercise, smoking, and drinking)

<sup>d)</sup> Adjusted odds ratios with 95% confidence interval (adjusted for age, number of persons in the household, employment status, sleep duration on weekdays, physical exercise, smoking, and drinking)

<sup>e)</sup> Adjusted odds ratios with 95% confidence interval (adjusted for age, number of persons in the household, disposable income per household, sleep duration on weekdays, physical exercise, smoking, and drinking)

<sup>f)</sup> NA not applicable

<sup>g)</sup> Bold ratios: statistically significant results



**Table 5** Prevalence and associations of the GHQ-12 score with risk factors among females

Females		Disposable income per household												Employment status										
		< 2000 K				2000 K – < 6000 K				≥ 6000 K				Full time			Others							
		Prevalence (≥ 4) <sup>a)</sup> n	Crude OR <sup>b)</sup>	95% CI	Adjusted OR <sup>c)</sup>	95% CI	Lower	Upper	Adjusted OR <sup>c)</sup>	95% CI	Lower	Upper	Adjusted OR <sup>c)</sup>	95% CI	Lower	Upper	Adjusted OR <sup>e)</sup>	95% CI	Lower	Upper				
Age (Years)																								
40–49	258	48.5	1	1			1					1			1				1					
50–59	204	38.9	<b>0.675</b>	<b>0.680</b>	<b>0.511</b>	<b>0.904</b>	0.830	0.296	2.333	0.682	0.442	1.053	<b>0.422</b>	<b>0.974</b>	0.635	0.352	1.147	<b>0.577</b>	<b>0.328</b>	<b>1.017</b>				
60–69	163	33.2	<b>0.528</b>	<b>0.676</b>	<b>0.479</b>	<b>0.954</b>	0.606	0.218	1.685	0.823	0.503	1.345	<b>0.283</b>	<b>0.926</b>	<b>0.257</b>	<b>0.085</b>	<b>0.781</b>	0.521	0.315	0.862				
Number of persons in the household																								
≥ 2	585	41.6	1	1			1						1		1				1					
1	33	29.7	<b>0.594</b>	<b>0.390</b>	<b>0.905</b>	0.643	0.371	1.113	0.507	0.485	0.201	1.171	1.186	0.321	4.384	1.021	0.311	3.351	0.478	0.187	1.221			
Employment status																								
Full time	106	45.3	1	1			1						1		NA <sup>d)</sup>				NA <sup>d)</sup>					
Others	360	42.4	0.887	0.663	1.187	0.970	0.709	1.327	1.865	0.697	0.422	1.153	1.164	0.747	1.815	NA <sup>d)</sup>			NA <sup>d)</sup>					
Household disposable income (JPY)																								
≥ 6000 K	258	39.0	1	1			NA <sup>d)</sup>						NA <sup>d)</sup>		NA <sup>d)</sup>				1					
2000 K–< 6000 K	296	40.4	1.065	0.859	1.321	1.128	0.862	1.477	NA <sup>d)</sup>				NA <sup>d)</sup>		NA <sup>d)</sup>				1.624	0.499	5.293	0.579	0.277	1.209
< 2000 K	71	45.2	1.290	0.908	1.832	<b>1.592</b>	<b>1.000</b>	<b>2.535</b>	NA <sup>d)</sup>				NA <sup>d)</sup>		NA <sup>d)</sup>				0.923	0.287	2.966	<b>0.567</b>	<b>0.268</b>	<b>1.203</b>
Sleep duration on weekdays (hours)																								
≥ 7	200	34.4	1	1			1						1		1				1					
6–7	247	41.2	<b>1.337</b>	<b>1.056</b>	<b>1.693</b>	1.238	0.920	1.665	1.252	1.166	0.758	1.795	1.632	0.735	3.624	1.546	0.767	3.119	1.357	0.860	2.141			
< 6	176	48.6	<b>1.803</b>	<b>1.379</b>	<b>2.356</b>	<b>1.505</b>	<b>1.077</b>	<b>2.104</b>	1.158	1.568	0.954	2.575	1.767	0.877	3.561	<b>2.288</b>	<b>1.103</b>	<b>4.746</b>	2.632	1.492	4.643			
Physical exercise (days/week)																								
≥ 3	60	29.6	1	1			1						1		1				1					
≤ 2	83	35.0	1.285	0.859	1.921	1.150	0.681	1.941	1.236	0.698	0.311	1.570	1.635	0.736	3.631	2.763	0.585	13.055	1.559	0.738	3.296			
0	479	43.5	<b>1.838</b>	<b>1.329</b>	<b>2.542</b>	<b>1.738</b>	<b>1.120</b>	<b>2.696</b>	1.171	1.804	0.947	3.433	1.755	0.871	3.537	2.364	0.579	9.648	<b>1.298</b>	<b>0.754</b>	<b>2.235</b>			
Smoking																								
Never	457	39.7	1	1			1						1		1				1					
Quit	97	43.3	1.158	0.867	1.547	0.979	0.696	1.376	1.028	0.894	0.542	1.475	1.180	0.694	2.006	1.240	0.591	2.602	1.682	0.834	3.391			
Sometimes/ everyday	68	40.5	1.031	0.741	1.434	0.999	0.668	1.495	0.934	1.163	0.642	2.108	0.864	0.448	1.666	0.978	0.440	2.173	0.712	0.340	1.489			

**Table 5** Prevalence and associations of the GHQ-12 score with risk factors among females (Continued)

Females		Disposable income per household						Employment status														
		< 2000 K		2000 K – < 6000 K		≥ 6000 K		Full time		Others												
Prevalence (≥ 4) <sup>a)</sup>	Crude OR <sup>b)</sup>	Adjusted OR <sup>d)</sup>	95% CI	Adjusted OR <sup>d)</sup>	95% CI	Adjusted OR <sup>d)</sup>	95% CI	Adjusted OR <sup>e)</sup>	95% CI	Adjusted OR <sup>e)</sup>	95% CI											
n	%	Lower	Upper	Lower	Upper	Lower	Upper	Lower	Upper	Lower	Upper											
Drinking (times/week)																						
0	328	42.5	1	1	1	1	1	1	1	1	1											
≤ 2	190	38.4	0.841	0.668	1.060	0.861	0.648	1.143	<b>0.274</b>	<b>0.100</b>	<b>0.751</b>	0.615	1.433	1.085	0.704	1.671	0.996	0.532	1.864	0.621	0.378	1.022
≥ 3	105	37.8	0.820	0.619	1.086	0.738	0.518	1.051	<b>0.289</b>	<b>0.092</b>	<b>0.906</b>	0.431	1.225	0.888	0.511	1.543	0.651	0.282	1.501	1.286	0.721	2.293

<sup>a)</sup> Prevalence of GHQ-12 score ≥ 4

<sup>b)</sup> Crude odds ratio with 95% confidence interval (no adjustment for any variables)

<sup>c)</sup> Adjusted odds ratios with 95% confidence interval (adjusted for age, number of persons in the household, employment status, disposable income per household, sleep duration on weekdays, physical exercise, smoking, and drinking)

<sup>d)</sup> Adjusted odds ratios with 95% confidence interval (adjusted for age, number of persons in the household, employment status, sleep duration on weekdays, physical exercise, smoking, and drinking)

<sup>e)</sup> Adjusted odds ratios with 95% confidence interval (adjusted for age, number of persons in the household, disposable income per household, sleep duration on weekdays, physical exercise, smoking, and drinking)

<sup>f)</sup> NA not applicable

<sup>g)</sup> Bold ratios: statistically significant

controlling for all covariates, the potential risk factors among males and females differed by household disposable income levels and employment status. Low income was a significant risk factor for psychological distress in females. About lifestyle factors, short sleep duration on weekdays and a lack of physical exercise were significant risk factors for psychological distress in both males and females. Among males at the middle level of household disposable income, there was an association between short sleep duration and psychological distress. Among males with a high level of household disposable income, the number of persons in the household and a lack of physical exercise were associated with psychological distress. Among males who did not have full-time jobs, short sleep duration and a lack of physical exercise were potential risk factors for psychological distress. Regarding females, low levels of household disposable income, short sleep duration, and no physical exercise were risk factors for psychological distress. Short sleep duration and a lack of physical exercise were risk factors for full-time female workers and non-full time female workers, respectively. According to these results, potential risk factors for psychological distress differ by gender, disposable income levels, and employment status.

Comparing GHQ-12 scores between the three disposable income levels, males with a disposable income of less than two million yen had the lowest scores ( $3.992 \pm 4.007$ ). Horii et al. reported GHQ-12 scores of  $2.34 \pm 3.14$  in the general male population aged 40–64 [27]. In this study, low-income males had higher GHQ-12 scores than the general population.

Moreover, after controlling for all covariates, females with low income levels had a significant association with poor mental health conditions. In previous studies, lower income group had the highest GHQ scores because financial hardship could be a cause of poor mental health conditions [28, 29]. Two cohort studies in the UK and Denmark also reported a strong association between current economic difficulties and mental disorders [11]. Therefore, both cross-sectional and longitudinal studies have demonstrated a significant association between poor economic situation and psychological disorders. Accordingly, addressing the mechanism linking socio-economic factors with mental health outcomes is a necessity. In particular, enhancing social security systems and organising effective health promotion programmes would be helpful for people from different socioeconomic backgrounds.

After controlling for all variables, lifestyle factors such as shorter sleep duration and lack of physical exercise were significant potential risk factors for poor mental health conditions. Previous studies on adults have also suggested the significant association of short sleep duration [24, 27, 30] with depression [24], anxiety [31, 32],

and poorer GHQ-12 scores [33], as well as the fact that it is a risk factor for later depression [34]. Thus, short sleep duration could be a cause of poor mental health conditions [31]. Our findings suggest that sleeping for more than 7 h per day is critical for preventing psychological distress.

Regarding habitual physical exercise, males with high incomes showed a higher tendency than those with low income. Among both males and females, especially males with a household disposable income of six million yen and above and non-full-time workers, the lack of physical activity could be a potential risk factor for psychological distress. Some studies have reported that physical activity is associated with better mental health outcomes [16, 27, 35]. There is, thus, a need to promote engagement in physical activity to reduce stress and improve mental health.

About alcohol consumption, while both males and females with lower income tended to drink less alcohol than those with higher income, this was especially true of male respondents. This result was consistent with previous findings [36, 37]. The Japanese National Survey reported that male respondents with lower income tended to drink less alcohol than those with high income [36]. The Centers for Disease Control and Prevention also reported that the prevalence of heavy episodic drinking increased with increase in household income [37]. However, people with lower SES tended to experience more negative alcohol-related outcomes than people with higher SES [38]. This implies that while people with lower SES may be less likely to drink alcohol, their health conditions are more negatively affected by unhealthy alcohol drinking habits. More research is necessary to address the long-term associations between drinking habits and health outcomes depending on SES.

As depicted in Table 5, it is noteworthy that one out of two female respondents in the youngest age group (40–49) indicated psychological distress. Some studies have also reported that distress declines with age [27, 29, 39, 40]. The experience of poor mental health may differ between young and old generations, perhaps because of the differences in stressors across the lifespan [29]. In view of the differences in gender roles between Japanese males and females, some specific job-related and family-related stressors should be identified [41, 42]. Moreover, menopause-related symptoms among middle-aged females could be one of the causes of the worst GHQ-12 score levels [43]. Health promotion programmes for middle-aged females would, thus, be required.

Practising two or three kinds of healthy lifestyle behaviours was found to be associated with a reduced risk of psychological distress [23]. As for the results of this and previous studies, healthy lifestyle practices such as getting adequate sleep and exercise could be beneficial for improving mental health conditions [23]. However, it seems that the mechanism connecting SES, lifestyle

factors, and mental health is complex and requires longitudinal investigation [27].

This study was based on a large data set, and thus has some important contributions to make. First, the examination of the association of socioeconomic and lifestyle factors with mental health outcomes among working adults was based on a nationally representative sample of over 3000 respondents. Second, this study found significant associations between socioeconomic and lifestyle factors and mental health outcomes after controlling for relevant factors. The results imply the existence of different mechanisms linking socioeconomic and lifestyle factors with mental health conditions among males and females with different SES.

However, the study has several limitations. First, a part of the panel data set was used in a cross-sectional form. Thus, it is not possible to make causal inferences based on these results. It appears essential to discuss causality running in both directions: poor financial conditions may breed psychological distress, and psychological distress may lead to poorer financial conditions [44]. As a next step, analysing the entire panel data set would be beneficial for identifying causal relationships for further discussion. Second, there is the possibility of selection and information bias due to attrition. Respondents who dropped out of the study might have been more likely to be unhealthy or in unfavourable situations. Finally, as the data on SES and lifestyle factors were self-reported, the possibility of social desirability bias and recall bias cannot be excluded.

## Conclusions

The results indicated that socioeconomic and lifestyle factors were associated with mental health conditions. The relationships between mental health conditions and lifestyles significantly differed by household disposable income levels and employment status. Therefore, it is necessary to minimise the health inequalities associated with disposable income levels by developing a social welfare policy and health promotion programmes to prevent poor mental health conditions. This study revealed the differences in healthy lifestyle behaviours between people with low and high SES. Health promotion programmes are needed to foster healthy practices such as adequate sleep, frequent physical exercise, controlled alcohol consumption, and refraining from smoking in different population groups. There might be a complex mechanism of the association between socioeconomic and health-related factors. Further research with respect to the differential associations of gender, age, and SES with health outcomes is required. Determining the complex mechanisms linking mental health conditions and socioeconomic and lifestyle factors would benefit the development of better preventive mental health programmes and a reduction of the risks associated with health and socioeconomic inequalities in different life stages.

## Abbreviations

AOR: Adjusted odds ratio; CI: Confidence interval; GHQ-12: 12-item General Health Questionnaire; JHPS/KHPS: Japan Household Panel Survey/Keio Household Panel Survey; OR: Odds ratio; WHO: World Health Organization

## Acknowledgements

The authors gratefully acknowledge the time and effort invested by the participants of this study.

## Authors' contributions

MN performed data analysis and contributed to writing and revising the manuscript. KK contributed to writing and revising the manuscript. IY designed the project and contributed to writing and revising the manuscript. All authors read and approved the final manuscript.

## Funding

This study was supported by JSPS KAKENHI Grant Number JP 17H06086. The funders had no role in the design of the study, in collection, analysis, and interpretation of data, or in writing the manuscript.

## Availability of data and materials

The datasets used and analysed in the current study are available on the website of the Panel Data Research Center at Keio University: <https://www.pdrc.keio.ac.jp/en/>. The website explains how to register the request form. The centre may approve within a few days on reasonable request.

## Ethics approval and consent to participate

This study was approved by the Institutional Review Board, Institute for Economic Studies, Keio University (Reference number 15002).

## Consent for publication

Not applicable.

## Competing interests

The authors declare that they have no competing interests.

## Author details

<sup>1</sup>Faculty of Economics, Keio University, 〒108-8345 Tokyo-to, Minato-ku, Mita 2-15-45, Tokyo, Japan. <sup>2</sup>The Ohara Memorial Institute for Science of Labour, Tokyo, Japan. <sup>3</sup>Faculty of Business and Commerce, Keio University, Tokyo, Japan.

Received: 19 September 2018 Accepted: 29 November 2019

Published online: 30 December 2019

## References

- Murakami K, Hashimoto H, Lee JS, Kawakubo K, Mori K, Akabayashi A. Distinct impact of education and income on habitual exercise: a cross-sectional analysis in a rural city in Japan. *Soc Sci Med*. 2011;73(12):1683–8.
- Mukong AK, Van Walbeek C, Ross H. Lifestyle and income-related inequality in health in South Africa. *Inte J Equity Health*. 2017;16:103.
- Marmot M. Social determinants of health inequalities. *Lancet*. 2005;365(9464):1099–104.
- Stringhini S, Sabia S, Shipley M, Brunner E, Nabi H, Kivimäki M, et al. Association of socioeconomic position with health behaviors and mortality. The Whitehall II study. *JAMA*. 2010;303(12):1159–66.
- Hassanzadeh J, Asadi-Lari M, Baghbanian A, Ghaem H, Kassani A, Rezaianzadeh A. Association between social capital, health-related quality of life, and mental health: a structural-equation modeling approach. *Croat Med J*. 2016;57(1):58–65.
- Russ TC, Stamatakis E, Hamer M, Starr JM, Kivimäki M, Batty GD. Association between psychological distress and mortality: individual participant pooled analysis of 10 prospective cohort studies. *BMJ*. 2012;345:e4933.
- Turecki G, Brent DA. Suicide and suicidal behaviour. *Lancet*. 2016;387(10024):1227–39.
- Kawakami N, Takeshima T, Ono Y, Uda H, Hata Y, Nakane Y, et al. Twelve-month prevalence, severity, and treatment of common mental disorders in communities in Japan: preliminary finding from the world mental Health Japan survey 2002–2003. *Psychiatry Clin Neurosci*. 2005;59(4):441–52.
- Wei KC, Chua HC. Suicide in Asia. *Int Rev Psychiatry*. 2008;20(5):434–40.

10. Qin P, Agerbo E, Mortensen PB. Suicide risk in relation to socioeconomic, demographic, psychiatric, and familial factors: a national register-based study of all suicides in Denmark, 1981-1997. *Am J Psychiatry*. 2003;160(4):765-72.
11. Laaksonen E, Martikainen P, Lahelma E, Lallukka T, Rahkonen O, Head J, et al. Socioeconomic circumstances and common mental disorders among Finnish and British public sector employees: evidence from the Helsinki Health study and the Whitehall II study. *Int J Epi*. 2007;36(4):776-86.
12. Rissanen T, Viinamäki H, Honkalampi K, Lehto SM, Hintikka J, Saharinen T, et al. Long term life dissatisfaction and subsequent major depressive disorder and poor mental health. *BMC psychiatry*. 2011;11:140.
13. Hamer M, Stamatakis E, Steptoe A. Psychiatric hospital admissions, behavioral risk factors, and all-cause mortality: the Scottish Health survey. *Arch Inter Med*. 2008;168(22):2474-9.
14. Glozier N, Martiniuk A, Patton G, Ivers R, Li Q, Hickie I, et al. Short sleep duration in prevalent and persistent psychological distress in young adults: the DRIVE study. *Sleep*. 2010;33(9):1139-45.
15. Matsuzaki I, Sagara T, Ohshita Y, Nagase H, Ogino K, Eboshida A, et al. Psychological factors including sense of coherence and some lifestyles are related to General Health Questionnaire-12 (GHQ-12) in elderly workers in Japan. *Environ Health Prev Med*. 2007;12(2):71.
16. Sagara T, Hitomi Y, Kambayashi Y, Hibino Y, Matsuzaki I, Sasahara S, et al. Common risk factors for changes in body weight and psychological well-being in Japanese male middle-aged workers. *Environ Health Prev Med*. 2009;14(6):319-27.
17. Lam TH, Li ZB, Ho SY, Chan WM, Ho KS, Li MP, et al. Smoking and depressive symptoms in Chinese elderly in Hong Kong. *Acta Psychiatr Scand*. 2004;110(3):195-200.
18. Anda RF, Williamson DF, Escobedo LG, Mast EE, Giovino GA, Remington PL. Depression and the dynamics of smoking: a national perspective. *JAMA*. 1990;264(12):1541-5.
19. Taylor G, McNeill A, Girling A, Farley A, Lindson-Hawley N, Aveyard P. Change in mental health after smoking cessation: systematic review and meta-analysis. *BMJ*. 2014;348:g1151.
20. Rehm J, Room R, Graham K, Monteiro M, Gmel G, Sempos CT. The relationship of average volume of alcohol consumption and patterns of drinking to burden of disease: an overview. *Addiction*. 2003;98(9):1209-28.
21. Collins SE. Associations between socioeconomic factors and alcohol outcomes. *Alcohol Res*. 2016;38(1):83-94.
22. Matud MP, López-Curbelo M, Fortes D. Gender and psychological well-being. *Int J Environ Res Public Health*. 2019;16(19):3531.
23. Lyu J, Lee SH, Kim HY. Associations between healthy lifestyles and health outcomes among older Koreans. *Geriatr Gerontol Int*. 2016;16(6):663-9.
24. Krueger PM, Friedman EM. Sleep duration in the United States: a cross-sectional population-based study. *Am J Epi*. 2009;169(9):1052-63.
25. Goldberg DP, Oldehinkel T, Ormel J. Why GHQ threshold varies from one place to another. *Psychol Med*. 1998;28(4):915-21.
26. Goldberg DP, Gater R, Sartorius N, Ustun TB, Piccinelli M, Gureje O, et al. The validity of two versions of the GHQ in the WHO study of mental illness in general health care. *Psychol Med*. 1997;27(1):191-7.
27. Hori D, Tsujiguchi H, Kambayashi Y, Hamagishi T, Kitaoka M, Mitoma J, et al. The associations between lifestyles and mental health using the general Health questionnaire 12-items are different dependently on age and sex: a population-based cross-sectional study in Kanazawa, Japan. *Environ Health Prev Med*. 2016;21(6):410-21.
28. Katikireddi SV, Niedzwiedz CL, Popham F. Employment status and income as potential mediators of educational inequalities in population mental health. *Eur J Pub Health*. 2016;26(5):814-6.
29. Christensen H, Jorm AF, Mackinnon AJ, Korten AE, Jacomb PA, Henderson AS. Age differences in depression and anxiety symptoms: a structural equation modelling analysis of data from a general population sample. *Psychol Med*. 1999;29:325-39.
30. Matamura M, Tochigi M, Usami S, Yonehara H, Fukushima M, Nishida A, et al. Associations between sleep habits and mental health status and suicidality in a longitudinal survey of monozygotic twin adolescents. *J Sleep Res*. 2014;23(3):292-6.
31. Suzuki H, Kaneita Y, Osaki Y, Minowa M, Kanda H, Suzuki K, et al. Clarification of the factor structure of the 12-item general Health questionnaire among Japanese adolescents and associated sleep status. *Psychiatry Res*. 2011; 188(1):138-46.
32. Wong ML, Lau EYY, Wan JHY, Cheung SF, Hui CH, Mok DSY. The interplay between sleep and mood in predicting academic functioning, physical health and psychological health: a longitudinal study. *J Psychosom Res*. 2013;74(4):271-7.
33. Chikawa K, Matsui T, Tsunoda T, Teruya K, Uemura T, Takeda N, et al. The relationships of sleep duration and mental health with electrocardiographic findings: a retrospective-cohort study in Okinawa, Japan. *Environ Health Prev Med*. 2008;13(4):227-33.
34. Szkló-Coxe M, Young T, Peppard PE, Finn LA, Benca RM. Prospective associations of insomnia markers and symptoms with depression. *Am J Epi*. 2010;171(6):709-20.
35. De Mello MT, Lemos VA, Antunes HKM, Bittencourt L, Santos-Silva R, Tufik S. Relationship between physical activity and depression and anxiety symptoms: a population study. *J Affect Disord*. 2013;149(1):241-6.
36. National Survey Health and Nutrition. 2014. In: National Survey Health and Nutrition The Ministry of Health Labour and Welfare Japan 2014. <https://www.mhlw.go.jp/file/04-Houdouhappyou-10904750-Kenkoukyoku-Gantaisakukenkouzhushinka/0000117311.pdf>. Accessed 27 Oct. 2019.
37. Vital signs: Binge drinking prevalence, frequency, and intensity among adults - United States 2010. In: Morbidity and Mortality Weekly Report (MMWR). Centers for Disease Control and Prevention. 2012. <https://www.cdc.gov/mmwr/preview/mmwrhtml/mm6101a4.htm>. Accessed 5 Jul 2018.
38. Lee JO, Herrenkohl TI, Kosterman R, Small CM, Hawkins JD. Educational inequalities in the co-occurrence of mental health and substance use problems, and its adult socio-economic consequences: a longitudinal study of young adults in a community sample. *Public Health*. 2013;127(8):745-53.
39. Jorm AF, Windsor TD, Dear KBG, Anstey KJ, Christensen H, Rodgers B. Age group differences in psychological distress: the role of psychosocial risk factors that vary with age. *Psychol Med*. 2005;35(9):1253-63.
40. Molarius A, Berglund K, Eriksson C, Eriksson HG, Lindén-Boström M, Nordström E, et al. Mental health symptoms in relation to socio-economic conditions and lifestyle factors - a population-based study in Sweden. *BMC Public Health*. 2009;9(1):302.
41. Honda A, Date Y, Abe Y, Aoyagi K, Honda S. Work-related stress, caregiver role, and depressive symptoms among Japanese workers. *Saf Health Work*. 2014;5(1):7-12.
42. Kikuzawa S. Elder care, multiple role involvement, and well-being among middle-aged men and women in Japan. *J Cross-Cult Gerontol*. 2015;30(4): 423-38.
43. Ornat L, Martínez-Deardth R, Chedraui P, Pérez-López FR. Assessment of subjective sleep disturbance and related factors during female mid-life with the Jenkins sleep scale. *Maturitas*. 2014;77(4):344-50.
44. Health at a Glance 2017: OECD Indicators. In OECD 2017 [https://doi.org/10.1787/health\\_glance-2017-en](https://doi.org/10.1787/health_glance-2017-en). Accessed 27 Oct. 2019.

## Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

**Ready to submit your research? Choose BMC and benefit from:**

- fast, convenient online submission
- thorough peer review by experienced researchers in your field
- rapid publication on acceptance
- support for research data, including large and complex data types
- gold Open Access which fosters wider collaboration and increased citations
- maximum visibility for your research: over 100M website views per year

**At BMC, research is always in progress.**

Learn more [biomedcentral.com/submissions](https://biomedcentral.com/submissions)

