

ORIGINAL INVESTIGATION

Open Access



Donor artery stenosis interactions with diastolic blood pressure on coronary collateral flow in type 2 diabetic patients with chronic total occlusion

Ying Shen¹, Zhen Kun Yang¹, Jian Hu¹, Xiao Qun Wang¹, Yang Dai², Su Zhang³, Rui Yan Zhang¹, Lin Lu^{1,2}, Feng Hua Ding^{1*} and Wei Feng Shen^{1,2*}

Abstract

Background: We investigated whether and to what extent stenosis of predominant collateral donor artery (PCDA) affects coronary collateral flow in relation to blood pressure (BP) in type 2 diabetic patients with chronic total occlusion (CTO).

Methods: Collateral flow index (CFI) as derived from intracoronary pressure distal to occluded segment and mean aortic pressure in 220 type 2 diabetic patients and 220 propensity score matched non-diabetic controls undergoing percutaneous coronary intervention for CTO. The severity of PCDA stenosis was graded according to lumen diameter narrowing.

Results: CFI decreased stepwise from mild to severe stenosis of the PCDA and was lower in diabetic patients with moderate or severe PCDA stenosis than in non-diabetic controls (0.36 ± 0.10 vs. 0.45 ± 0.08 , $P < 0.001$; 0.29 ± 0.09 vs. 0.35 ± 0.08 , $P = 0.008$). When the PCDA was mildly stenotic, CFI increased initially along with a reduction in diastolic BP, and decreased when diastolic BP was below 60 mmHg in diabetic patients (0.38 ± 0.16 vs. 0.57 ± 0.09 , $P < 0.001$). In the presence of moderate PCDA stenosis, diabetic patients had significantly lower CFI compared to non-diabetic controls, with a relative reduction of 19.8% at diastolic BP 70–79 mmHg, 28.2% at 60–69 mmHg and 38.2% below 60 mmHg (all $P < 0.05$). A severe PCDA stenosis resulted in a more pronounced decrease in CFI, with a relative reduction of 37.3% for diabetics compared to non-diabetics when diastolic BP was below 60 mmHg ($P = 0.050$).

Conclusions: In the setting of CTO, donor artery stenosis confers greater risk for reduced coronary collateral flow when diastolic BP is decreased. Even a moderate stenosis in the PCDA may be associated with lower collateral flow as diastolic BP decreases below 80 mmHg in type 2 diabetic than in non-diabetic patients.

Keywords: Blood pressure, Collateral circulation, Diabetes, Coronary artery disease, Chronic total occlusion

*Correspondence: ruijindfh@126.com; rjshenweifeng@126.com

¹ Department of Cardiology, Rui Jin Hospital, Shanghai Jiao Tong University School of Medicine, 197 Rui Jin Road II, Shanghai 200025, People's Republic of China

² Institute of Cardiovascular Diseases, Shanghai Jiao Tong University School of Medicine, 197 Rui Jin Road II, Shanghai 200025, People's Republic of China

Full list of author information is available at the end of the article



Background

In patients with stable coronary artery disease, a gradual development of complete coronary obstruction may lead to a sufficient compensation of blood supply via collateral circulation to prevent myocardial damage from ischemic insults [1]. Protection of the jeopardized myocardium by coronary collaterals is clinically relevant, as presence of well-formed collaterals has been associated with reductions in the occurrence and transmural extent of myocardial infarction, and with increased survival [2]. The mechanism of collateral vessel growth is complex in situations where atherosclerosis affects large conductance arteries [3], and even become more complicated by the presence of diabetes mellitus in which multiple biochemical and cellular components are involved [4]. Nevertheless, arteriogenesis with vessel outward remodeling is prevalent, and outweighs much more than angiogenesis because it reduces dramatically collateral resistance to a negligible extent and enables delivery of blood flow to the region at risk [5, 6]. Among numerous factors which could influence coronary collateral flow [4, 7–10], blood pressure (BP), especially diastolic BP, generates the distal pressure within the occluded segment of the coronary artery, which constitutes a physical stimulus for arteriogenesis and promotes collateral formation [11]. Presence of a chronic total occlusion (CTO) is frequently associated with multi-vessel coronary disease and has been considered as a prerequisite for spontaneous collateral recruitment [7, 12]. Collaterals develop due to the pressure gradient from donor to recipient being greater than that of the recipient (often a CTO). Obviously, myocardium distal to the occlusion is almost entirely perfused by retrograde collateral branches from another epicardial coronary artery (i.e., predominant collateral donor artery [PCDA]), and successful recanalization of a chronically occluded lesion has often led to a rapid reduction of pressure-derived recruitable collateral function and an increase in fractional flow reserve of the PCDA [12]. This suggests a potential interaction between coronary collateral flow and donor artery physiology in the setting of chronic coronary total occlusion. However, the effect of donor artery stenosis on coronary collateral flow in relation to BP for patients with diabetes remains unknown, which partly reflects the heterogeneity of study population and semi-quantitative angiographic assessment of coronary collateral circulation in most previous cohort studies [8–10]. In this study, we investigated whether or to what extent combined BP (particularly diastolic BP) and stenosis of the PCDA affects coronary collateral flow in type 2 diabetic and non-diabetic patients with chronic coronary total occlusion. We examined the equilibrium of collateral supply at the time of advancing a microcatheter distal to the occlusion before antegrade flow is


re-established. Collateral flow index (CFI) was derived from intracoronary distal occluded pressure and central aortic pressure taking into account of central venous pressure [7, 13, 14], which has been considered as the most accurate diagnostic tool to assess the capacity of coronary collateral circulation as it correlates closely with clinical signs of myocardial ischemia [15, 16]. For avoiding potential confounding factors, each type 2 diabetic patient was matched to a non-diabetic control for age, sex and risk factors for coronary artery disease.

Methods

Study population

In total, 1147 consecutive patients with stable coronary artery disease and CTO (>3 months) of at least one major epicardial coronary artery between October 2010 and December 2016 were screened from the database of Shanghai Rui Jin Hospital PCI Outcome Program [17, 18]. We excluded 238 patients who were referred for coronary artery bypass grafting (CABG). In the remaining 909 patients undergoing elective PCI, we further excluded those who had a history of PCI within 3 months ($n=42$) or CABG ($n=46$), renal failure requiring hemodialysis ($n=4$), type 1 diabetes ($n=4$), chronic heart failure with NYHA class III or IV ($n=12$), pulmonary heart disease ($n=10$) and malignant tumor or immune system disorders ($n=4$). We also excluded those who had failed PCI for CTO mostly due to inability of guide wire to cross the occluded segment ($n=73$) and those who underwent PCI via a retrograde approach ($n=82$). To reduce the selection bias, we then performed a propensity score matching analysis, resulting in a total number of 440 patients (220 type 2 diabetics and 220 non-diabetics) into the final analyses (Fig. 1).

The duration of CTO was estimated based on information obtained from a previous angiogram, a history of myocardial infarction in the target vessel territory, or the first onset of an abrupt worsening of existing angina. The diagnosis of type 2 diabetes was made according to the criteria of the American diabetes association, including glycosylated hemoglobin (HbA1c) $\geq 6.5\%$, fasting plasma glucose concentration ≥ 7.0 mmol/L, 2-h postprandial glucose concentration ≥ 11.1 mmol/L, or a random plasma glucose ≥ 11.1 mmol/L in a patient with classic symptoms of hyperglycemia or hyperglycemic crisis [19]. Hypertension was defined as systolic BP ≥ 140 mmHg and/or diastolic BP ≥ 90 mmHg, or use of anti-hypertensive agents for controlling BP [20]. Dyslipidemia was defined according to the Third Report of The National Cholesterol Education Program (NCEP) [21]. Stable angina was diagnosed according to the criteria recommended by the American College of Cardiology/American Heart Association [22].



Biochemical investigation

We obtained blood samples at the day of angiography in all patients after an overnight fasting. Assessment of serum levels of creatinine, blood urea nitrogen, uric acid, lipid profiles, glucose, and glycosylated hemoglobin (HbA1c), was made with standard laboratory techniques. Glomerular filtration rate (GFR) was estimated using the chronic kidney disease epidemiology collaboration (CKD-EPI) equation: $GFR_{EPI} (mL/min/1.73 m^2) = 141 \times \min(creatinine/k, 1)^\alpha \times \max(creatinine/k, 1)^{-1.209} \times 0.993^{age} \times 1.018$ [if female], where k is 0.7 for females and 0.9 for males, α is -0.329 for females and -0.411 for males, min indicates the minimum of creatinine/k or 1, and max indicates the maximum of creatinine/k or 1 [23]. Serum high-sensitivity C-reactive protein (hsCRP) level was assayed by ELISA (Biocheck Laboratories, Toledo, OH, USA).

Angiography and analysis

Coronary angiography was performed via femoral or radial access with 6Fr diagnostic catheters. Quantitative angiographic assessment was done independently by two blinded interventional cardiologists according to lesion classification scheme of the American College of Cardiology/American Heart Association [24]. The PCDA was defined as a contra-lateral vessel making the largest collateral contribution [25], and maximal diameter stenosis of the PCDA was classified as mild: <50%, moderate: 50–70% or severe: >70% (GE Centricity VA 1000; GE Healthcare).

Intracoronary pressure and collateral flow measurement

Elective PCI for CTO was performed through an antegrade approach in all patients. After crossing the occluded segment with various types of guide wire, a

microcatheter was advanced distal to the occlusion. Sometimes, a low-profile balloon catheter was passed initially, and then was exchanged for a microcatheter. Meticulous care was taken to ensure complete obstruction of the coronary artery by lack of contrast passage during proximal contrast injection through the guiding catheter while the microcatheter was in place. Intracoronary pressure distal to an occluded segment measured through a microcatheter (Finecross, Terumo Co, Japan) and central aortic pressure determined from a 6Fr guiding catheter were simultaneously recorded using a fluid-filled manometer system (Mac-Lab Hemodynamic Recording System, GE Healthcare, USA). The transducer was kept at the level of mid axillary line, and zero calibrated to atmosphere before measurement [17]. Despite a phasic waveform for distal coronary pressure, significant damping of high frequency pressure components by the small lumen of microcatheter exhibited, thus only the mean distal occluded pressure was used. For measurement of central aortic pressure, the guiding catheter was kept away from the coronary orifice. The pressure-derived coronary collateral flow index (CFI) was calculated as the ratio of $(Pd - CVP)/(Pa - CVP)$, where Pd is mean distal occluded pressure, Pa is mean central aortic pressure; and CVP is the central venous pressure, which was substituted by 5 mmHg, as all patients had no symptoms and signs of heart failure [7, 14, 15].

We have validated the accuracy of intracoronary pressure-derived collateral flow measurement using a microcatheter against the standard pressure wire technique (PressureWire™ Certus, St. Jude Medical, St. Paul, Minnesota) with RadiAnalyzer™ Xpress System (St. Jude Medical, St. Paul, Minnesota). Bland–Altman analysis revealed that in 40 consecutive patients, Pd and CFI determined with a microcatheter correlated significantly with those by a pressure wire ($r=0.751$ and $r=0.679$, both $P<0.001$), with an absolute difference in Pd and CFI between the two techniques of 1.68 mmHg (95% CI -1.42 to 4.77, $P=0.280$) and 0.019 (95% CI -0.02 to 0.053, $P=0.279$), respectively.

Statistical analysis

For patient selection, a propensity score matching analysis was performed in advance with a match tolerance of 0.02 and a ratio of 1:1 for diabetic and non-diabetic patients using a logistic regression model with age, sex and risk factors for coronary artery disease (smoking and history of hypertension or dyslipidemia). Continuous variables are presented as mean \pm standard deviation (SD), and categorical data are summarized as frequencies (percentages). For categorical clinical variables, differences between groups were evaluated with the Chi square test. For continuous variables, the existence of a

normal distribution was evaluated with the Kolmogorov–Smirnov test, and differences among groups were analyzed by one-way analysis of variance (ANOVA) followed by post hoc analysis with the Fisher's least significant difference (LSD) test. Pearson's or Spearman's correlation analysis was run to determine the relationship between CFI and Pd or different levels of aortic BP. Multivariate linear regressions were used to explore the independent determinants for CFI, and the covariates chosen to enter the multivariate analysis model included diabetes and aortic systolic, diastolic and mean BP, respectively, as well as gender, age, body mass index (BMI), risk factors for coronary artery disease (history of hypertension and dyslipidemia and smoking), history of myocardial infarction, severity of coronary artery disease, GFR, total to high-density lipoprotein (HDL) cholesterol ratio, log-transformed hsCRP and left ventricular ejection fraction. All analyses used 2-sided tests with an overall significance level of $\alpha=0.05$, and were performed with the SPSS 20.0 for Windows (SPSS, Inc., Chicago, IL, USA).

Results

Baseline characteristics

Mild, moderate and severe stenosis of the PCDA was observed in 99, 75 and 46 diabetic patients and 132, 60 and 28 non-diabetic patients, respectively. For diabetic and non-diabetic patients, age, proportion of male gender, dyslipidemia and prior myocardial infarction, and serum levels of creatinine and hsCRP increased whereas GFR, left ventricular ejection fraction and proportion of hypertension decreased stepwise from mild to severe stenosis of the PDCA (all $P<0.05$). Multivessel coronary disease was more prevalent in patients with moderate or severe PDCA stenosis. Medications were similar irrespective of stenosis severity of the PDCA (Table 1).

Collateral flow and BP

Despite similar aortic systolic, diastolic and mean BP (Table 1), CFI decreased stepwise from mild to severe stenosis of the PDCA in diabetic and non-diabetic patients (both $P<0.001$), and were significantly lower in diabetic patients than in non-diabetic controls with moderate (0.36 ± 0.10 vs. 0.45 ± 0.08 , $P<0.001$) or severe (0.29 ± 0.09 vs. 0.35 ± 0.08 , $P=0.008$) PCDA stenosis. Furthermore, there was no difference in CFI between diabetic patients with moderate PDCA stenosis and non-diabetic controls with severe PCDA stenosis ($P=0.421$) (Fig. 2).

Effect of medical therapy

Treatments with β -blockers and nitrates were associated with higher CFI in diabetic (0.42 ± 0.12 vs. 0.38 ± 0.13 , $P=0.026$; 0.42 ± 0.12 vs. 0.39 ± 0.12 , $P=0.033$) and

Table 1 Baseline characteristics in patients with stable coronary artery disease and chronic total occlusion with mild, intermediate and severe stenosis of donor artery

Variables	Diabetes (n = 220)			P value	Non-diabetes (n = 220)			P value
	Mild (n = 99)	Moderate (n = 75)	Severe (n = 46)		Mild (n = 132)	Moderate (n = 60)	Severe (n = 28)	
Male, n (%)	94 (94.9)	59 (78.7)	24 (52.2)	<0.001	127 (96.2)	40 (66.7)	14 (50.0)	<0.001
Age, years	57.9 ± 11.7	67.9 ± 10.7	73.7 ± 9.3	<0.001	61.3 ± 12.1	68.0 ± 9.3	74.6 ± 12.4	<0.001
Body mass index, kg/m ²	25.4 ± 2.5	25.5 ± 2.6	25.9 ± 2.1	0.605	25.0 ± 2.5	25.0 ± 1.9	25.7 ± 2.4	0.294
Hypertension, n (%)	102 (77.3)	37 (61.7)	7 (25.0)	<0.001	91 (91.9)	42 (56.0)	11 (23.9)	<0.001
History of dyslipidemia, n (%)	23 (17.4)	25 (41.7)	14 (50.0)	<0.001	2 (2.0)	20 (26.7)	26 (56.5)	<0.001
Smoking, n (%)	27 (27.3)	26 (34.7)	21 (45.7)	0.090	43 (32.6)	18 (30.0)	15 (53.6)	0.072
History of MI, n (%)	19 (19.2)	16 (21.3)	19 (41.3)	0.012	19 (14.4)	12 (20.0)	10 (35.7)	0.030
Severity of CAD, n (%)				<0.001				<0.001
1-vessel	39 (39.4)	2 (2.7)	0 (0.0)	<0.001	51 (38.6)	5 (8.3)	5 (17.9)	<0.001
2-vessel	35 (35.4)	21 (28.0)	17 (37.0)	0.493	55 (41.7)	17 (28.3)	7 (25.0)	0.089
3-vessel	25 (25.2)	52 (69.3)	29 (63.0)	<0.001	26 (19.7)	38 (63.3)	16 (57.1)	<0.001
Multivessel disease	60 (60.6)	73 (97.3)	46 (100.0)	<0.001	81 (61.4)	55 (91.7)	23 (82.1)	<0.001
Stenosis of PCDA, %	37.6 ± 8.5	58.9 ± 5.3	83.4 ± 8.5	<0.001	33.5 ± 10.5	58.9 ± 5.4	82.6 ± 8.4	<0.001
Fast blood glucose, mmol/L	6.3 ± 1.9	6.9 ± 2.2	6.5 ± 2.7	0.214	5.0 ± 0.7	4.7 ± 0.7	5.3 ± 0.6	<0.001
HbA1c, %	7.2 ± 1.2	7.4 ± 1.5	7.1 ± 1.3	0.555	5.7 ± 0.4	5.7 ± 0.4	5.8 ± 0.4	0.980
Triglyceride, mmol/L	1.55 ± 0.72	1.62 ± 0.63	1.68 ± 0.81	0.538	1.42 ± 0.71	1.45 ± 0.83	1.60 ± 0.71	0.521
Total cholesterol, mmol/L	3.75 ± 1.04	4.11 ± 1.12	4.29 ± 1.04	0.009	4.11 ± 1.34	3.89 ± 1.45	4.08 ± 0.88	0.571
HDL cholesterol, mmol/L	0.94 ± 0.20	0.99 ± 0.28	0.99 ± 0.16	0.312	1.01 ± 0.27	0.96 ± 0.19	1.00 ± 0.34	0.469
LDL cholesterol, mmol/L	2.23 ± 0.90	2.47 ± 0.93	2.71 ± 0.87	0.010	2.49 ± 1.05	2.45 ± 1.11	2.40 ± 0.84	0.891
Serum creatinine, μmol/L	85 ± 21	90 ± 30	110 ± 32	<0.001	77 ± 14	94 ± 27	102 ± 29	<0.001
Uric acid, μmol/L	322 ± 79	315 ± 61	331 ± 76	0.515	337 ± 72	319 ± 96	339 ± 69	0.309
GFR, mL/min/1.73 m ²	86.8 ± 18.5	75.5 ± 24.9	55.9 ± 20.6	<0.001	90.3 ± 15.8	70.7 ± 20.2	59.6 ± 21.8	<0.001
hsCRP, mg/L	1.58 (0.66–3.32)	2.46 (1.21–7.55)	4.23 (1.44–17.15)	0.031	0.97 (0.25–3.13)	1.20 (0.27–10.35)	2.66 (0.42–25.82)	<0.001
LVEF, %	60.5 ± 8.3	60.4 ± 9.3	54.7 ± 10.7	0.001	62.3 ± 9.0	61.3 ± 9.5	57.2 ± 12.8	0.042
Medication, n (%)								
ACE inhibitors/ ARBs	53 (53.5)	42 (56.0)	26 (56.5)	0.924	50 (37.9)	20 (33.3)	11 (39.3)	0.798
β blockers	41 (41.4)	30 (40.0)	20 (43.5)	0.931	51 (38.6)	25 (41.7)	15 (53.6)	0.345
Calcium channel blockers	30 (30.3)	23 (30.7)	12 (26.1)	0.845	41 (31.1)	13 (21.7)	8 (28.6)	0.406
Nitrates	57 (57.6)	41 (54.7)	24 (52.2)	0.819	59 (44.7)	32 (53.3)	16 (57.1)	0.339
Statins	69 (69.7)	59 (78.7)	36 (78.3)	0.328	77 (58.3)	33 (55.0)	12 (42.9)	0.325
Anti-diabetic therapy	58 (58.6)	60 (80.0)	40 (87.0)	<0.001	0 (0.0)	0 (0.0)	0 (0.0)	–
Aortic blood pressure, mmHg								
Systolic	150.2 ± 26.2	147.3 ± 23.4	141.2 ± 24.8	0.130	150.2 ± 29.2	147.0 ± 28.2	141.0 ± 17.9	0.260
Diastolic	77.1 ± 9.8	74.9 ± 9.4	73.5 ± 10.8	0.093	78.1 ± 13.2	75.9 ± 9.7	75.6 ± 11.5	0.388
Mean	101.5 ± 13.2	99.0 ± 11.8	96.1 ± 13.5	0.057	101.2 ± 14.7	101 ± 12.9	97.4 ± 7.7	0.407

Table 1 (continued)

Variables	Diabetes (n = 220)			P value	Non-diabetes (n = 220)			P value
	Mild (n = 99)	Moderate (n = 75)	Severe (n = 46)		Mild (n = 132)	Moderate (n = 60)	Severe (n = 28)	
Pd, mmHg	52.1 ± 9.4	39.6 ± 11.0	32.5 ± 10.2	<0.001	53.6 ± 7.1	48.3 ± 8.6	35.6 ± 7.8	<0.001
Collateral flow index	0.49 ± 0.09	0.36 ± 0.10	0.29 ± 0.09	<0.001	0.51 ± 0.08	0.45 ± 0.08	0.35 ± 0.08	<0.001

Data are mean ± SD or median (25th–75th percentiles) or number (%)

ACE angiotensin converting enzyme, ARB angiotensin receptor blocker, CAD coronary artery disease, GFR glomerular filtration rate, HbA1c glycosylated hemoglobin A1c, HDL high-density lipoprotein, hsCRP high-sensitivity C reactive protein, LDL low-density lipoprotein, LVEF left ventricular ejection fraction, MI myocardial infarction, PCDA predominant collateral donor artery, Pd mean pressure distal to occlusion




Fig. 2 Comparison of CFI between diabetic and non-diabetic patients with mild, moderate or severe stenosis of PCDA group. *P < 0.001 vs. mild; #P < 0.001 vs. moderate

non-diabetic (0.49 ± 0.09 vs. 0.46 ± 0.10, P = 0.023; 0.49 ± 0.10 vs. 0.46 ± 0.10, P = 0.019) patients whereas other medications such as angiotensin-converting enzyme inhibitors or angiotensin receptor blockers, calcium channel blockers, statins, oral hypoglycemic agents and insulin did not significantly affect CFI in both groups (all P > 0.05).

Multivariable analysis

Multivariable linear regression models with systolic (per 20 mmHg), diastolic (per 10 mmHg) and mean (per 10 mmHg) BP as well as potential confounding variables such as gender, age, body mass index, history of hypertension and dyslipidemia, smoking, prior myocardial infarction, severity of coronary artery disease, GFR, total-to-HDL cholesterol ratio, log-transferred hsCRP and left ventricular ejection fraction, revealed that after multiple adjustments, CFI correlated negatively with systolic, diastolic, and mean BP in diabetic and non-diabetic patients with mild PCDA stenosis (P ≤ 0.001), was inversely related to diastolic BP in non-diabetic patients with moderate

PCDA stenosis (P = 0.011), but correlated positively with diastolic BP in diabetic patients with severe PCDA stenosis (P = 0.001).

Further analysis showed an interaction between diabetes and diastolic BP in patients with moderate (P interaction = 0.008) and severe (P interaction = 0.032) stenosis of the PCDA (Table 2). When the PCDA was mildly stenotic, CFI was gradually increased along with a reduction in aortic diastolic BP, but it was decreased when diastolic BP was below 60 mmHg in diabetic patients, with a relative reduction of 32.1% compared with non-diabetic controls (0.38 ± 0.16 vs. 0.57 ± 0.09, P < 0.001). In the presence of moderate PCDA stenosis, with decreasing diastolic BP, the difference of CFI between diabetic and non-diabetic patients was gradually increased. When diastolic BP was below 80 mmHg, diabetic patients had a significantly lower CFI compared to non-diabetic controls, with a relative reduction of 19.8% at diastolic BP 70–79 mmHg (0.37 ± 0.10 vs. 0.46 ± 0.07, P < 0.001), 28.2% at 60–69 mmHg (0.33 ± 0.12 vs. 0.46 ± 0.10, P < 0.001) and 38.2% below 60 mmHg (0.28 ± 0.11 vs. 0.46 ± 0.11, P = 0.002), respectively. A severe stenotic lesion in the PCDA led to more pronounced decrease in CFI, with a relative reduction of 37.3% for diabetic patients compared to non-diabetic controls when diastolic BP was below 60 mmHg (0.19 ± 0.07 vs. 0.30 ± 0.12, P = 0.050) (Fig. 3).

Discussion

Our results support the hypothesis that in the setting of CTO, coronary collateral flow is adversely affected by the combined effect of donor artery stenosis severity and aortic diastolic BP. Even a moderate stenosis in the PCDA resulted in lower collateral flow as diastolic BP decreases below 80 mmHg for type 2 diabetic patients compared with non-diabetic patients.

Interactive effects of PDCA stenosis and BP on collateral flow

Physiologically, coronary collateral inflow into the distal vessel of a CTO through a variety of anatomic arteriolar


Table 2 Aortic blood pressure in relation to coronary flow index in diabetic and non-diabetic patients according to stenosis of PCDA

Aortic BP	Mild				Moderate				Severe								
	Diabetes (n = 99)		Non-diabetes (n = 132)		Diabetes (n = 75)		Non-diabetes (n = 60)		Diabetes (n = 46)		Non-diabetes (n = 28)						
	$\beta \pm SE$	P	$\beta \pm SE$	P	$\beta \pm SE$	P	$\beta \pm SE$	P	$\beta \pm SE$	P	$\beta \pm SE$	P					
Systolic	-0.284 ± 0.073	<0.001	-0.352 ± 0.070	<0.001	0.325	0.325	-0.013 ± 0.062	0.840	0.065 ± 0.158	0.682	0.239	0.239	0.154 ± 0.204	0.464	0.025 ± 0.085	0.766	0.651
Diastolic	-0.275 ± 0.079*	0.001	-0.525 ± 0.065	<0.001	0.168	0.168	-0.019 ± 0.066 [#]	0.774	-0.303 ± 0.114	0.011	0.008	0.008	0.317 ± 0.087	0.001	0.199 ± 0.183	0.296	0.032
Mean	-0.334 ± 0.074 ^{&}	<0.001	-0.540 ± 0.061	<0.001	0.081	0.081	-0.018 ± 0.063	0.779	-0.158 ± 0.136	0.253	0.067	0.067	0.604 ± 0.219 ^{&}	0.016	0.147 ± 0.087	0.008	0.669

Values are regression coefficient (β) ± SE, derived from multiple linear regression analyses of coronary flow index with central aortic systolic, diastolic and mean blood pressure, respectively, after adjustment for gender, age, body mass index, current smoking, history of hypertension and dyslipidemia, prior myocardial infarction, severity of coronary artery disease, GFR, total and HDL cholesterol ratio, log-transformed hsCRP and left ventricular ejection fraction. P values for interaction in group of mild, intermediate and severe stenosis of PCDA are given

GFR glomerular filtration rate, HDL high-density lipoprotein, hsCRP high-sensitivity C reactive protein, PCDA predominant collateral donor artery

* P < 0.001, [#] P < 0.01 and [&] P < 0.05 vs. the corresponding β value in non-diabetes



connections is predominantly modulated by driving force for blood flow resulted mainly from pressure gradient across the occluded site [1, 3–5]. Central aortic pressure (especially diastolic BP) generates the distal pressure within the occluded segment, and relatively high intracoronary pressure and tangential fluid shear stress imposed on the collateral endothelium have been suggested to constitute hemodynamic stimuli for arteriogenesis [5–7] and promote collateral blood flow [14, 26, 27]. A collateral donor vessel will have a distal pressure drop with an increasing stenosis and/or lowering of diastolic BP (which contributes more to mean distal occluded pressure as diastole is longer than systole) and hence both are expected to lead to lower distal occluded pressure and CFI in the recipient CTO. In the setting of single vessel occlusion, collateral function is directly affected by systemic arterial pressure [3]. However, for patients with multi-vessel coronary disease, we would expect pressure gradient across a CTO to be reduced when the artery that supplies collateral blood flow exhibits a critical stenosis because of a pressure drop proximal to the origin of the collaterals. This may result in a further reduction in coronary collateral flow particularly when BP is decreased.

The major finding of this study is that coronary collateral flow decreased stepwise as the severity of PCDA stenosis increased, and a moderate PDCA stenosis in type 2 diabetic patients could induce similar extent of collateral flow reduction to that caused by a severe PCDA stenosis in non-diabetic patients. Interestingly, our study showed that at various degrees of PCDA stenosis, type 2 diabetic patients had lower CFI when diastolic BP decreased below 60 mmHg, and even a moderate stenotic lesion in the PCDA was associated with more reduced collateral flow as diastolic BP decreased below 80 mmHg in type 2 diabetic patients compared to non-diabetic controls. These findings highlight that the effect of PCDA stenosis on collateral flow relative to BP may be different between

type 2 diabetic and non-diabetic patients. The reason for this remains unclear, but a likely explanation is the presence of more diffuse coronary atherosclerosis in a diabetic setting [28]. As shown in this study, patients with type 2 diabetes had more severe coronary artery disease as evidenced by a higher percentage of multivessel disease and a greater stenosis severity of the PCDA. It has been suggested that the presence of diffuse atherosclerotic disease in the collateral donor artery would be likely to be associated with a reduced coronary flow reserve [29], and the large increase in coronary flow through collateral donor vessels as a result of the additional flow through the collateral bed could be enough for minor atherosclerotic irregularities to generate sufficient resistance to become flow limiting [30]. Furthermore, data from prior studies which have assessed the microcirculatory function in patients with and without diabetes, have demonstrated that patients with diabetes have substantially adverse functional and structural remodeling of the coronary arterioles and even amongst those diabetic patients without known coronary artery disease, the presence of an abnormal coronary flow reserve is associated with poor outcome, comparable to non-diabetic patients with known coronary disease [31, 32]. Recently, Hinkel et al. [33] reported that diabetic human myocardial explants revealed capillary rarefaction and pericyte loss compared to non-diabetic explants. In a diabetic pig model of hibernating myocardium, hyperglycemia induced microvascular rarefaction in the myocardium even without ischemia, and capillary density further decreased in chronic ischemia hearts. This indicates that type 2 diabetes destabilized microvascular vessels of the heart and may impair the responsiveness of ischemic myocardium to pro-angiogenic factors [33]. It has been possible to determine microvascular resistance using the pressure wire technique in diabetics and non-diabetics, which could give more of an insight why diabetic patients

have a lower CFI in the CTO vessel and are particularly susceptible to low diastolic BP. Previous studies have shown that there exists a pronounced increase of collateral resistance [34] and obliteration of pre-existing blood vessels in diabetes [28], suggesting that diabetic microvascular resistance is higher in the donor vessel and contributes to the obstruction of collateral flow to the CTO. All these vascular changes could contribute to a further reduction in collateral flow for patients with type 2 diabetes.

Clinical implications

Our study findings could partly highlight the importance of an individualized BP lowering strategy [35–37] and a potential influence of stenotic lesions in the collateral donor coronary artery as a revascularization target [38–40]. In patients with multivessel disease, caution should be taken when administering anti-hypertensive therapies, as aggressive reduction in systemic BP (especially diastolic BP) may compromise collateral recruitment and exacerbate myocardial ischemia particularly for those with type 2 diabetes and stenotic lesions in the PCDA. Similarly, if the vasodilatory reserve of the arterioles in the vascular bed supplied by a chronically occluded coronary artery is completely exhausted, whereas that of the PCDA is still preserved, coronary (collateral) steal may result. This phenomenon has been reported to occur in a very high proportion of well collateralized myocardial beds [41] and is most likely to occur in patients with moderate or severe stenosis of the PCDA, as vasodilator-induced increase in flow could cause a pressure drop across the stenotic lesions, thereby lowering collateral perfusion [3, 42]. Consistent with previous findings [7], our present study also demonstrates a positive association between β blockers and CFI. The use of β blockers reduces heart rate, improves fluid shear stress at the endothelial wall, and decrease catecholamine-mediated inflammatory response, favoring coronary collateral flow.

It is now generally accepted that when presented with multivessel disease, we should aim for complete rather than incomplete revascularization [43]. PCI with drug-eluting stent implantation on severe coronary stenotic lesions has become a routine clinical practice, and clinical evidence suggests that recanalization of a CTO as a part of a complete revascularization procedure confers a substantial benefit to survival [39]. Previous studies reported that a hemodynamically ambiguous lesion would not necessarily be of low angiographic complexity, and the need to treat it might alter the long-term outcomes [30]. Our observations on the relationship between PCDA stenosis and coronary collateral flow relative to diastolic BP supports a notion that multiple aspects should be taken into consideration when

planning revascularization procedures, including characteristics of totally occluded lesion, severity of PCDA stenosis, quality of collaterals, and clinical status of patients (diabetes and BP level) [44]. PCI aimed at improving collateral flow could be accomplished by reducing proximal donor artery stenosis, thereby increasing pressure at the collateral takeoff [39]. In patients with moderate PCDA stenosis (especially those with type 2 diabetes), the use of fractional flow reserve to reveal ischemia can help in clinical decision-making [40]. Furthermore, hypotension (especially low diastolic BP) should be avoided during the procedure. Whether such a strategy is particularly useful for type 2 diabetic patients with CTO and moderate PCDA stenosis warrants further investigation.

Limitations

We acknowledge there are limitations worth mentioning. First, this study is cross-sectional for the point of coronary collateral flow investigation, thereby allowing us to detect association, not to determine the difference between diabetic and non-diabetic response of CFI to increasing diastolic pressure and to establish a causative link and to predict clinical outcome. Second, the number of patients in this study is small as compared to previous reports using angiography to assess coronary collateralization in diabetes. However, it was out-weighted by the use of a quantitative assessment of collateral function, whereas the angiographic method is a semi-quantitative grading of collateral contrast filling. Third, the use of a microcatheter to measure pressure distal to the occluded segment may be problematic as the lesion may compress the catheter and cause inaccuracy in the pressure measured. Although a pressure sensewire would have been more accurate, the absolute difference in intracoronary pressure measurement was small using a microcatheter or a pressure wire method. Finally, central venous pressure was not measured, which is not a major concern when measuring coronary fractional flow reserve but could significantly influence the CFI. Collaterals in the whole cohort were quite good (CFI > 0.25) for the majority of patients. This may be because CFI was determined on an assumption of central venous pressure. Obviously, the accuracy of CFI measurement could be significantly affected by minor variation in actual central venous pressure. Likewise, measurement of left ventricular end-diastolic pressure (LVEDP) in these patients was also important as compressive forces of the ventricle can influence collateral support via septal collaterals. It is possible that diabetic patients have stiffer left ventricle and increased LVEDP that impairs collateral support.

Conclusions

In patients with stable coronary artery disease and CTO, donor artery stenosis confers greater risk for reduced coronary collateral flow when diastolic BP is decreased. For type 2 diabetic patients, even a moderate stenosis in the PCDA is associated with more reduced collateral flow as diastolic BP decreases below 80 mmHg compared with non-diabetic patients. These findings may provide clinical insight into the management of patients with coronary artery disease.

Abbreviations

ANOVA: analysis of variance; BMI: body mass index; BP: blood pressure; CABG: coronary artery bypass grafting; CFI: collateral flow index; CKD-EPI: chronic kidney disease epidemiology collaboration; CTO: chronic total occlusion; CVP: central venous pressure; GFR: glomerular filtration rate; HbA1c: glycated hemoglobin; HDL: high-density lipoprotein; hsCRP: high-sensitivity C-reactive protein; LSD: least significant difference; NCEP: national cholesterol education program; NYHA: New York Heart Association; Pa: mean central aortic pressure; PCDA: predominant collateral donor artery; PCI: percutaneous coronary intervention; Pd: mean distal occluded pressure; SD: standard deviation.

Authors' contributions

YS, FHD, WFS wrote the article, substantially contributed to discussion of the content, and edited the manuscript. YS, YZK, HJ, WXQ, DY, ZS performed the experiments and researched data for the article. FHD analyze the data; RYZ, LL substantially contributed to discussion of the content and reviewed the manuscript. All authors read and approved the final manuscript.

Author details

¹ Department of Cardiology, Rui Jin Hospital, Shanghai Jiao Tong University School of Medicine, 197 Rui Jin Road II, Shanghai 200025, People's Republic of China. ² Institute of Cardiovascular Diseases, Shanghai Jiao Tong University School of Medicine, 197 Rui Jin Road II, Shanghai 200025, People's Republic of China. ³ College of Biomedical Engineering, Jiao Tong University, Shanghai 200031, People's Republic of China.

Acknowledgements

Not applicable.

Competing interests

The authors declare that they have no competing interests.

Availability of data and materials

Data generated or analyzed during this study are included in this published article.

Consent for publication

All authors consent this manuscript for publication.

Ethics approval and consent to participate

The study protocol was approved by the Institutional Review Board of Rui Jin Hospital, Shanghai Jiaotong University School of Medicine. Written informed consent was obtained from all patients, and clinical investigation was conducted according to the principle of the Declaration of Helsinki.

Funding

This study was supported in part by the Research Foundation of Chinese National Natural Science (81400327, 81470469, 81400211, 81670451, 81770437, 81770447), Shanghai Science and Technology Committee (14ZR1425800, 14441903002, 15411963700) and Medico-engineering Project (GY2016MS66) and Talent Young Investigators (17XJ11009) of Shanghai Jiao Tong University School of Medicine.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Received: 26 February 2018 Accepted: 26 May 2018

Published online: 01 June 2018

References

- Seiler C, Stoller M, Pitt B, Meier P. The human coronary collateral circulation: development and clinical importance. *Eur Heart J*. 2013;34:2674–82.
- Meier P, Hemingway H, Lansky AJ, Knapp G, Pitt B, Seiler C. The impact of the coronary collateral circulation on mortality: a meta-analysis. *Eur Heart J*. 2012;33:614–21.
- Zimarino M, D'Andrea Matteo M, Waksman R, Epstein SE, De Caterina R. The dynamics of the coronary collateral circulation. *Nat Rev Cardiol*. 2014;11:191–7.
- Shen Y, Ding FH, Dai Y, Wang XQ, Zhang RY, Lu L, Shen WF. Reduced coronary collateralization in type 2 diabetic patients with chronic total occlusion. *Cardiovasc Diabetol*. 2018;17:26.
- Chilian WM, Penn MS, Pung YF, Dong F, Mayorga M, Ohanian V, Logan S, Yin L. Coronary collateral growth—back to the future. *J Mol Cell Cardiol*. 2012;52:905–11.
- Carmeliet P. Mechanisms of angiogenesis and arteriogenesis. *Nat Med*. 2000;6:389–95.
- van der Heoven NW, Teunissen PF, Werner GS, Delewi R, Schirmer SH, Traupe T, van der Laan AM, Tijssen JG, Seiler C, van Royen N. Clinical parameters associated with collateral development in patients with chronic coronary total occlusion. *Heart*. 2013;99:1100–5.
- Shen Y, Lu L, Ding FH, Sun Z, Sun Z, Zhang RY, Zhang Q, Yang ZK, Hu J, Chen QJ, et al. Association of increased serum glycated albumin levels with low coronary collateralization in type 2 diabetic patients with stable angina and chronic total occlusion. *Cardiovasc Diabetol*. 2013;12:165.
- Shen Y, Ding FH, Zhang RY, Zhang Q, Lu L, Shen WF. Association of serum mimecan with angiographic coronary collateralization in patients with stable coronary artery and chronic total occlusion. *Atherosclerosis*. 2016;252:75–81.
- Shen Y, Lu L, Liu ZH, Wu F, Zhu JZ, Sun Z, Zhang RY, Zhang Q, Hu J, Chen QJ, et al. Increased serum level of CTRP1 is associated with low coronary collateralization in stable angina patients with chronic total occlusion. *Int J Cardiol*. 2014;174:203–6.
- Ladwiniee A, Conington MS, Rossington J, Mather AN, Alahmar A, Oliver RM, Nijjer SS, Davies JE, Thackray S, Alamgir F, et al. Collateral donor artery physiology and the influence of a CTO on fractional flow reserve. *Circ Cardiovasc Interv*. 2015;8:e002219.
- Werner GS. The role of coronary collaterals in chronic total occlusion. *Curr Cardiol Rev*. 2014;10:57–64.
- Seiler C, Fleisch M, Garachemani A, Meier B. Coronary collateral quantitation in patients with coronary artery disease using intravascular flow velocity or pressure measurements. *J Am Coll Cardiol*. 1998;32:1272–9.
- Traupe T, Gloekler S, de Marchi SF, Werner GS, Seiler C. Assessment of the human coronary collateral circulation. *Circulation*. 2010;122:1210–20.
- Dervan JP, McKay RG, Baim DS. Assessment of the relationship between distal occluded pressure and angiographically evident collateral flow during coronary angioplasty. *Am Heart J*. 1987;114:491–7.
- Seiler C. Assessment and impact of the human coronary collateral circulation on myocardial ischemia and outcome. *Circ Cardiovasc Interv*. 2013;6:719–28.
- Shen Y, Ding FH, Wu F, Lu L, Zhang RY, Zhang Q, Wu ZG, Shen WF. Association of blood pressure and coronary collateralization in type 2 diabetic and nondiabetic patients with stable angina and chronic total occlusion. *J Hypertens*. 2015;33:621–6.
- Shen Y, Ding FH, Sun JT, Pu LJ, Zhang RY, Zhang Q, Chen QJ, Shen WF, Lu L. Association of elevated apoA-I glycation and reduced HDL-associated paraoxonase1, 3 activity, and their interaction with angiographic severity of coronary artery disease in patients with type 2 diabetes mellitus. *Cardiovasc Diabetol*. 2015;14:52.
- American Diabetes Association. Standards of medical care in diabetes—2014. *Diabetes Care*. 2014;37(suppl 1):S14–80.

20. Miedema MD, Lopez FL, Blaha MJ, Virani SS, Coresh J, Ballantyne CM, Folsom AR. Implications of the eighth joint national committee guidelines for the management of high blood pressure for aging adults: atherosclerosis risk in communities study. *Hypertension*. 2015;66:474–80.
21. Expert Panel on Detection, Evaluation, and Treatment of High Blood Cholesterol in Adults. Executive summary of the third report of the national cholesterol education program (NCEP) expert panel on detection, evaluation, and treatment of high blood cholesterol in adults (adult treatment panel III). *JAMA*. 2001;285:2486–97.
22. Fraker TD Jr, Fihn SD, Gibbons RJ, Abrams J, Chatterjee K, Daley J, Deedwania PC, Douglas JS, Ferguson TB Jr, Gardin JM, et al. 2007 chronic angina focused update of the ACC/AHA 2002 guidelines for the management of patients with chronic stable angina: a report of the American college of cardiology/American heart association task force on practice guidelines writing group to develop the focused update of the 2002 guidelines for the management of patients with chronic stable angina. *J Am Coll Cardiol*. 2007;50:2264–74.
23. Levy AS, Stevens LA, Schmid CH, Zhang YL, Castro AF 3rd, Feldman HI, Kusek JW, Eggers P, Van Lente F, Greene T, Coresh J, CKD-EPI (Chronic Kidney Disease Epidemiology Collaboration). CKD-EPI (chronic kidney disease epidemiology collaboration). A new equation to estimate glomerular filtration rate. *Ann Intern Med*. 2009;150:604–12.
24. Ellis SG, Vandormael MG, Cowley MJ, DiSciascio G, Deligonul U, Topol EJ, Bulle TM. Coronary morphologic and clinical determinants of procedural outcome with angioplasty for multivessel coronary disease: implications for patient selection (multivessel angioplasty prognosis study group). *Circulation*. 1990;82:1193–202.
25. Werner OS, Ferrari M, Heinker S, Kuethel F, Surber R, Richartz BM, Figulla HR. Angiographic assessment of collateral connections in comparison with invasively determined collateral function in chronic coronary occlusion. *Circulation*. 2003;107:1972–7.
26. Wang S, Jing J, Liu CF, Jiang TM, Yang XB, Zhou Y, Chen YD. The relationship between diastolic pressure and coronary collateral circulation in patients with stable angina pectoris and chronic total occlusion. *Am J Hypertens*. 2013;26:630–5.
27. van Royen N, Piek JJ, Buschmann I, Hoefer I, Voskuil M, Schaper W. Stimulation of arteriogenesis: a new concept for the treatment of arterial occlusive disease. *Cardiovasc Res*. 2001;49:543–53.
28. Rocic P. Why is coronary collateral growth impaired in type II diabetes and the metabolic syndrome? *Vascu Pharmacol*. 2012;57:179–86.
29. Gould LL, Johnson NP, Bateman TM, Beanlands RS, Bengel FM, Bober R, Camici PG, Cerqueira MD, Chow BJ, Di Carli MF, et al. Anatomic versus physiologic assessment of coronary artery disease: role of coronary flow reserve, fractional flow reserve, and positron emission tomography imaging in revascularization decision-making. *J Am Coll Cardiol*. 2013;62:1039–53.
30. Ladwiniec A, Hoye A. The hemodynamic effects of collateral donation to a chronic total occlusion: implications for patient management. *Int J Cardiol*. 2015;196:159–66.
31. Werner GS, Ferrari M, Richartz BM, Gastmann O, Figulla HR. Microvascular dysfunction in chronic total coronary occlusions. *Circulation*. 2001;104:1129–34.
32. Kennedy MW, Fabris E, Suryapranata H, Kedhi E. Is ischemia the only factor predicting cardiovascular outcomes in all diabetes mellitus patients? *Cardiovasc Diabetol*. 2017;16:15.
33. Hinkel R, Howe A, Renner S, Ng J, Lee S, Klett K, Kaczmarek V, Moretti A, Laugwitz KL, Skroblin P, et al. Diabetes mellitus–induced microvascular destabilization in the myocardium. *J Am Coll Cardiol*. 2017;69:131–43.
34. De Bruyne B, Hersbach F, Pijls NHJ, Bartunek J, Bech JW, Heyndrickx GR, Gould KL, Wijns W. Abnormal epicardial coronary resistance in patients with diffuse atherosclerosis but ‘normal’ coronary angiography. *Circulation*. 2001;104:2401–6.
35. Naito R, Kasai T. Coronary artery disease in type 2 diabetes mellitus: recent treatment strategies and future perspectives. *World J Cardiol*. 2015;7:119–24.
36. ACCORD Study Group, Cushman WC, Evans GW, Byington RP, Goff DC Jr, Grimm RH Jr, Cutler JA, Simons-Morton DG, Basile JN, Corson MA, Probstfield JL, et al. Effects of intensive blood pressure control in type 2 diabetes mellitus. *N Engl J Med*. 2010;362:1575–85.
37. Rosendorff C, Lackland DT, Allison M, Aronow WS, Black HR, Blumenthal RS, Cannon CP, de Lemos JA, Elliott WJ, Findeiss L, American Heart Association, American College of Cardiology, and American Society of Hypertension, et al. Treatment of hypertension in patients with coronary artery disease: a scientific statement from the American heart association, American college of cardiology, and American society of hypertension. *Hypertension*. 2015;65:1372–407.
38. Valenti R, Migliorini A, Signorini U, Vergara R, Parodi G, Carrabba N, Cerisano G, Antonucci D. Impact of complete revascularization with percutaneous coronary intervention on survival in patients with at least one chronic total occlusion. *Eur Heart J*. 2008;29:2336–42.
39. Khan MF, Wendel CS, Thai HM, Movahed MR. Effects of percutaneous revascularization of chronic total occlusion on clinical outcomes: a meta-analysis comparing successful versus failed percutaneous intervention for chronic total occlusion. *Catheter Cardiovasc Interv*. 2013;82:95–107.
40. Sachdeva R. Effect of revascularization of chronic total occlusion on tandem stenoses in a ‘donor’ artery. *Tex Heart Inst J*. 2014;41:547–50.
41. Demer LL, Gould KL, Goldstein R, Kirkeeide RL. Noninvasive assessment of coronary collaterals in man by PET perfusion imaging. *J Nucl Med*. 1990;31:259–70.
42. Seiler C, Fleisch M, Meier B. Direct intracoronary evidence of collateral steal in humans. *Circulation*. 1997;96:4261–7.
43. Blundhun PK, Wu ZJ, Chen MH. Coronary artery bypass surgery compared with percutaneous coronary interventions in patients with insulin-treated type 2 diabetes mellitus: a systemic review and meta-analysis of 6 randomized controlled trials. *Cardiovasc Diabetol*. 2016;15:2.
44. Kennedy MW, Kaplan E, Hermanides RS, Fabris E, Hemradj V, Koopmans PC, Dambink JH, Marcel Gosselink AT, Van’t Hof AW, Ottervanger JP, et al. Clinical outcomes of deferred revascularisation using fractional flow reserve in patients with and without diabetes mellitus. *Cardiovasc Diabetol*. 2016;15:100.

Ready to submit your research? Choose BMC and benefit from:

- fast, convenient online submission
- thorough peer review by experienced researchers in your field
- rapid publication on acceptance
- support for research data, including large and complex data types
- gold Open Access which fosters wider collaboration and increased citations
- maximum visibility for your research: over 100M website views per year

At BMC, research is always in progress.

Learn more biomedcentral.com/submissions

