

RESEARCH

Open Access



Detection of foci of residual malaria transmission through reactive case detection in Ethiopia

Endalew Zemene^{1*} , Cristian Koepfli², Abebaw Tiruneh¹, Asnakew K. Yeshiwondim³, Dinberu Seyoum⁴, Ming-Chieh Lee², Guiyun Yan^{2†} and Delenasaw Yewhalaw^{1,5†}

Abstract

Background: Sub-microscopic and asymptomatic infections could be bottlenecks to malaria elimination efforts in Ethiopia. This study determined the prevalence of malaria, and individual and household-level factors associated with *Plasmodium* infections obtained following detection of index cases in health facilities in Jimma Zone.

Methods: Index malaria cases were passively detected and tracked in health facilities from June to November 2016. Moreover, family members of the index houses and neighbours located within approximately 200 m from the index houses were also screened for malaria.

Results: A total of 39 index cases initiated the reactive case detection of 726 individuals in 116 households. Overall, the prevalence of malaria using microscopy and PCR was 4.0% and 8.96%, respectively. Seventeen (43.6%) of the index cases were from Doyo Yaya *kebele*, where parasite prevalence was higher. The majority of the malaria cases (90.74%) were asymptomatic. Fever (AOR = 12.68, 95% CI 3.34–48.18) and history of malaria in the preceding 1 year (AOR = 3.62, 95% CI 1.77–7.38) were significant individual-level factors associated with detection of *Plasmodium* infection. Moreover, living in index house (AOR = 2.22, 95% CI 1.16–4.27), house with eave (AOR = 2.28, 95% CI 1.14–4.55), area of residence (AOR = 6.81, 95% CI 2.49–18.63) and family size (AOR = 3.35, 95% CI 1.53–7.33) were main household-level predictors for residual malaria transmission.

Conclusion: The number of index cases per *kebele* may enhance RACD efforts to detect additional malaria cases in low transmission settings. Asymptomatic and sub-microscopic infections were high in the study area, which need new or improved surveillance tools for malaria elimination efforts.

Keywords: Reactive case detection, Malaria, Residual malaria transmission, Low-transmission setting, Ethiopia

Background

The global pattern of malaria epidemiology has changed remarkably over the last decade. Between 2010 and 2015, malaria mortality rates declined by 35% among children under 5 years of age, with 21% fall in incidence among the population at risk [1]. Implementation of key malaria prevention and control measures have played a pivotal role in decreasing morbidity and mortality due to malaria

[2, 3]. Furthermore, concerted control and elimination efforts of malaria in some countries, such as the United Arab Emirates and Sri Lanka, have resulted in malaria-free status in recent years [4, 5]. Despite the achievements gained in the control of malaria, the disease still remains a significant public health problem in many sub-Saharan African countries, including Ethiopia.

The epidemiology of malaria in Ethiopia appears unique, compared to other countries in sub-Saharan Africa, in that both *Plasmodium falciparum* and *Plasmodium vivax* coexist. While almost all cases of malaria are due to the two species, there is high spatiotemporal heterogeneity in the distribution of these parasite species.

*Correspondence: endalew2005@yahoo.com

†Guiyun Yan and Delenasaw Yewhalaw contributed equally to this work

¹ School of Medical Laboratory Sciences, Faculty of Health Sciences, Jimma University, Jimma, Ethiopia

Full list of author information is available at the end of the article



According to the 2015 National Health Sector Development Plan report [6], out of the total microscopy or rapid diagnostic test (RDT) confirmed malaria cases, 63.7% and 36.3% were due to *P. falciparum* and *P. vivax*, respectively. In a recent study done in Jimma in south-western Ethiopia, however, more than three-quarters of the cases were due to *P. vivax* [7]. *Plasmodium ovale* plays a minor role in Ethiopia, and appears to be often misdiagnosed [8].

Over the last decade, during which malaria elimination was put back on the global health agenda, morbidity and mortality due to malaria has remarkably declined in Ethiopia [9, 10]. Besides the sharp decline of malaria including from some of the historically malarious areas of the country [11], no major malaria epidemics, which usually recur every 5- to 8 years, have been reported since 2005 [12]. Implementation and scale-up of the powerful vector control interventions, including indoor residual spraying (IRS) and long-lasting insecticidal nets (LLINs) appear to have played key roles [13]. More than 17 million LLINs have been distributed in 2014/2015 alone, with cumulative number of the nets distributed since 2009 being scaled up to more than 75 million [6]. Access to malaria diagnostics and treatment has also remarkably improved over the last decade, mainly via the innovative health extension programme [14] that operates at community level.

Based on the malaria control achievements gained, and with the help of international partners, Ethiopia has set goals to eliminate malaria by 2030. However, substantial portions of human *Plasmodium* infections are asymptomatic, often remaining undetected by microscopic examination [15]. Asymptomatic infections can serve as reservoirs of infection to the vector mosquitoes [16], potentially sustaining transmission.

To further sustain control of malaria and move towards elimination, adequate detection and prompt treatment of both symptomatic and asymptomatic cases in the community is critical [17]. One of the strategies of addressing malaria cases not presenting to the health care facilities is reactive case detection (RACD) with focal test and treatment methods. Reactive case detection makes use of the spatial clustering trend of malaria carriers particularly in low endemic settings [18, 19]. Hence, in RACD, following passive case detection, household members of the index case and neighbours located at certain distance from the index household are screened. This method has been utilized in several low malaria transmission settings [20, 21], despite lack of established standard approach to the spatial range of neighbouring households to be within the screening radius.

Reactive case detection also allows detection of asymptomatic malaria infections, which play a major role in

sustaining malaria transmission in low-transmission settings [22]. However, active case detection of malaria is not yet fully implemented in the routine health care system in Ethiopia. Thus, this study is aimed at detecting malaria cases using RACD in two health centres in Jimma Zone, south-western Ethiopia.

Methods


Study setting

The study was conducted in catchment *kebeles* (smallest government administrative units in Ethiopia) of Kishe and Nada health centres, located in Shebe Sambo and Omo Nada districts of Jimma Zone, respectively (Fig. 1). Shebe Sambo and Omo Nada districts are located at 415 and 285 kms south west of the capital, Addis Ababa, respectively. The geographical coordinates of Shebe Sambo and Omo Nada are approximately 7°30'14"N, 36°30'44"E and 7°38'00"N, 37°15'05"E, respectively. The inhabitants in both areas mainly depend on subsistence farming, cultivating mainly maize and *teff*. Moreover, in Kishe area rice is cultivated in small scale.

Historically, the catchment areas of both health centres have been malarious [23–25]. As in most parts of Ethiopia, the transmission of malaria in these areas is seasonal. The transmission usually peaks from September to October, following the major rains from June to September, and minor transmission occurs in April and May, following the short rains of February to March. According to the information obtained from both health centres, malaria cases detected in the health facilities have remarkably declined in recent years. A total 43 malaria cases have been registered in Kishe Health Centre in 2016. Of these, 62.8% were due to *P. falciparum*. Kishe Health Centre has been serving a total of 26,843 population in 2016. In the same year, a total of 51 malaria cases were recorded at Nada Health Centre, 49% of which were due to *P. falciparum*. The health centre has been serving a total of 32,264 population in 2016.

Population and sampling

A prospective observational study was conducted for 6 months (June to November 2016) in two health centres and their catchment *kebeles*. Index malaria cases residing within the catchment *kebeles* of the two health centres who did not travel within 2 weeks prior to presenting to the health centres, and diagnosed with malaria at the health centres during the study period were included in the study. The index cases were identified in the health centres based on the routine blood film microscopy by the laboratory staff at each health facility. Following detection of the index cases, household members of the index houses and neighbours within 200 m radius were included in the study.



Data collection

Passive case detection

Febrile patients who sought treatment at Kishe and Nada Health Centres from June 1 to November 31, 2016, were screened for malaria using microscopy by the resident laboratory personnel as a routine practice. Consenting index cases that were microscopy-positive and who agreed that a research team will visit them within 1 week provided their home address. To locate the index household easily, names of three nearby household heads were also recorded.

Reactive case detection

Presence of index cases was communicated to the research team on the same day of presentation to the health centres. The index houses and neighbours were visited within 1 week of detecting the index cases, in most cases, within 3 days. After obtaining consent, family members of the index houses and neighbours within 200 m radius from the index houses were screened for *Plasmodium* infection using RDT. Moreover,

demographic information and some individual and household-level risk factors of malaria were collected using a semi-structured questionnaire. The field data was collected by experienced laboratory technologists.

The individual-level factors assessed included demographic characteristics (age, sex, educational status and occupation), recent travel history, history of malaria infection in the last 1 year, LLIN usage the previous night before the survey and axillary body temperature. Fever was defined in this study as having axillary temperature of ≥ 37.5 °C, which was measured during the survey. The household-level factors assessed included housing conditions such as roof structure, presence of visible hole on the wall, presence of eave and presence of window(s), and presence of animals within the house, family size, total number of LLINs owned during the survey and whether the house was sprayed with insecticide during the last 1 year. In Ethiopia, indoor residual spraying is performed once in a year, usually around July to September (before malaria cases peak) as transmission is mainly seasonal. Apart from the household-level characteristics, an

approximate distance of each house from the index house was estimated, and coordinates of each house was taken using hand-held global positioning system unit (GPS).

All consenting members of the index houses and neighbours were enrolled in this study. Infants less than 6 months of age were not included, with the assumption that they are likely protected from malaria due to passively acquired maternal antibody, and presence of foetal haemoglobin. The household members available during the visit and those found in the nearby farming sites were included in the study. The samples were collected by a team of two laboratory technologists deployed at a time. The data collection team spent, in most cases, all the day around the index houses to maximize coverage of the screened population. However, few individuals who were not available on the day of screening around their houses were not captured, as there was no follow-up to the community members.

Specimen collection and laboratory processing

Finger-prick blood samples were collected from consenting study participants for blood examination by RDT and microscopy. The RDT was done for rapid diagnosis and treatment; hence, RDT-positive individuals were referred to the health centres for confirmation and treatment. Moreover, thick and thin blood smears were prepared for each study participant in the field. After air drying, the thin films were fixed with absolute methanol, and Giemsa stained at Jimma University Medical Parasitology Laboratory on the same day of collection. The slides were examined by two experienced laboratory technologists independently. Approximately 200 high-power fields of the thick blood smears were examined before declaring a microscopy-negative result. The personnel reading the slides were also blinded of the RDT results. Apart from the blood smears, three to four drops of blood were spotted on Whatman 3MM filter paper for further molecular analysis. The blood spots were air-dried and kept individually in air-tight plastic bags and stored at -20°C until DNA extraction.

DNA was extracted from approximately 20 μL whole blood using the QiaCube DNA extraction system as per standard protocols. DNA was eluted in 100 μL buffer, and 4 μL DNA was screened using a multiplex *P. falciparum/P. vivax* qPCR with limit of detection of 1 DNA copy per reaction [26]. Thus, the limit of detection of the qPCR was 1–2 parasites/ μL blood.

Data management and analysis

Household was defined in this study as group of human subjects residing in the same house as family members. Household access to LLINs was considered “sufficient” when the ratio of the total LLINs owned by a household

to the family members is at least 0.5 (assuming that one LLIN covers two individuals), and “not sufficient” when the ratio is less than 0.5 [27]. While a total of 726 individuals participated in this study providing samples for microscopic examination, sufficient DBS sample for PCR was obtained from 603 individuals. Specimens of sufficient quantity could not be obtained from the remaining individuals for PCR following blood film preparation and RDT testing, the results of which were used for immediate care. Hence, the data analysis was based on the PCR-run samples. Asymptomatic malaria infection was considered when an individual who did not experience fever at the time of the survey (axillary body temperature is less than 37.5°C) and no malaria-related symptoms was positive for *Plasmodium* species by PCR.

The collected data were coded, entered into Excel (Microsoft Office 2010) and cleaned. The data were analysed using a statistical software package STATA 12 (StataCorp., TX, USA). Descriptive statistics including frequency, percentages and median were calculated to summarize demographic profile of the study participants. Univariate and multivariate logistic regressions were employed to determine individual and household-level factors associated with malaria infection. Multi-level regression model (mixed-effects logistic regression) was utilized to determine predictors of malaria infection among individual and household-level variables. Variables with significant association with malaria infection by the univariate analysis and those with *p* values less than 0.2 were candidates for the multivariate analyses. Odds ratio and the corresponding 95% confidence intervals were calculated to show the strength of the association. Statistical significance was set at $p < 0.05$ during the analysis.

Results

Socio-demographic characteristics

Thirty-nine passively detected microscopy-positive malaria cases (index cases) initiated the RACD. Most of the index cases (56.4%, $n=22$) were from Nada Health Centre, with the remaining 17 (43.6%) being from Kische Health Centre. Twenty-four (61.5%) and 15 (38.5%) of the index cases were due to *P. falciparum* and *P. vivax* infections, respectively. Following detection of the index cases, a total of 726 individuals residing in 116 households were screened for *Plasmodium* infection. Most of them were female (55.6%, $n=404$), and the median age was 16 years (range 1 to 80 years). Demographic characteristics of the study participants are presented in Table 1.

Table 1 Demographic characteristics of the study participants

Characteristics	Frequency, n (%)
Age group in years	
< 5	144 (19.8)
5–15	221 (30.4)
> 15	361 (49.7)
Sex	
Male	322 (44.4)
Female	404 (55.6)
Site	
Kishe	220 (30.30)
Nada	506 (69.69)
Educational status	
Illiterate	235 (42.6)
Literate	316 (57.4)
Occupation	
Farmers	155 (21.4)
Housewives	143 (19.7)
Students	214 (29.5)
Others	38 (5.2)
House structure	
Iron sheet	496 (68.3)
Thatched	230 (31.7)
Spray status of the house	
Not sprayed	417 (57.4)
Sprayed	309 (42.6)

Prevalence of malaria

The overall prevalence of malaria from the RACD by microscopy and PCR was 4.0% (29/726) and 8.96% (54/603), respectively. Majority of the PCR-detected cases (92.59%, n = 50) were *P. falciparum*, the remaining 4 (7.41%) being *P. vivax*. The prevalence of malaria among

individuals residing in the index houses and neighbours was 17.5% and 6.83%, respectively. The majority (90.74%, n = 49) of malaria cases detected in the RACD were asymptomatic. A relatively larger number of the malaria cases were detected in Doyo Yaya *kebele* (Table 2).

Factors associated with *Plasmodium* infection

Bivariate analysis of individual and household-level factors associated with malaria positivity is shown in Table 3. The analysis revealed that several individual-level factors were significantly associated with *Plasmodium* infection, including age (COR = 1.99, 95% CI 1.06–3.74), self-reported history of malaria infection in the last 1 year before the survey (COR = 3.71, 95% CI 1.94–7.07) and fever (COR = 6.12, 95% CI 1.97–18.98). The study participants were asked if they travelled to other villages in the preceding 2 weeks before the survey. Only 3 (0.5%) reported travel and none of them were positive.

Among the household-level factors assessed in this study, IRS (COR = 6.86, 95% CI 2.69–17.48), living within index house (COR = 2.89, 95% CI 1.60–5.21), area of residence (COR = 6.70, 95% CI 2.63–17.08) and family size (COR = 3.73, 95% CI 1.88–7.38) were significant risk factors of malaria.

Long-lasting insecticidal nets ownership (proportion of households who reported to have at least one LLIN) was 90.9%, but 56.82% of the households having insufficient LLINs. Bivariate analysis showed that individuals living in households with insufficient access to LLINs were at higher risk of *Plasmodium* infection (COR = 2.35, 95% CI 1.18–4.65).

Table 4 shows predictors of *Plasmodium* infection. After adjusting for other variables, fever (AOR = 10.13, 95% CI 2.76–37.16) and history of malaria (AOR = 3.97, 95% CI 1.96–8.04) were the main individual-level predictors of *Plasmodium* infection. Moreover, household-level

Table 2 Residence of the index cases and frequency of individuals screened following the index cases, Jimma, southwestern Ethiopia

Health facility	Kebele	Index cases, n (%)	Examined by PCR, n (%)	Ratio of # screened to index cases	Additional RACD cases, n (%)
Kishe Health Centre	Kishe	7 (17.9)	59 (9.8)	8.4	1 (1.7)
	Halo Sebaka	7 (17.9)	91 (15.1)	13.0	4 (4.1)
	Gasera Kekero	1 (2.6)	6 (1.0)	6.0	0 (0)
	Wala Kela	1 (2.6)	7 (1.2)	7.0	0 (0)
Omo Nada Health Centre	Doyo Yaya	17 (43.6)	375 (62.2)	22.1	49 (13.1)
	Nada	4 (10.3)	18 (3.0)	4.5	0 (0)
	Nada Chala	1 (2.6)	24 (4.0)	24.0	0 (0)
	Nada Sekote	1 (2.6)	23 (3.8)	23.0	0 (0)
Total		39 (100)	603 (100)	15.5	54 (8.96)

Table 3 Bivariate analysis results showing individual and household-level factors associated with *Plasmodium* infection among the study participants, Jimma, south-western Ethiopia

Variables	Examined by PCR, n (%)	Positive, n (%)	p-value	COR (95% CI)
Age (years)				
< 5	117 (19.40)	11 (9.40)	0.336	1.46 (0.68–3.14)
5–15	185 (30.68)	23 (12.43)	0.032	1.99 (1.06–3.74)*
> 15	301 (49.92)	20 (6.64)	Ref	
Sex				
Male	270 (44.78)	24 (8.89)	Ref	
Female	333 (55.22)	30 (9.01)	0.959	1.01 (0.58–1.78)
Malaria in the preceding 1 year				
Yes	72 (11.94)	16 (22.22)	0.001	3.71 (1.94–7.07)*
No	531 (88.06)	38 (7.16)	Ref	
Fever at the time of survey				
Yes	14 (2.32)	5 (35.71)	0.002	6.12 (1.97–18.98)*
No	589 (97.68)	49 (8.32)	Ref	
ITN usage the preceding night				
Yes	243 (40.30)	18 (7.41)	Ref	
No	360 (59.70)	36 (10.00)	0.276	1.39 (0.77–2.51)
Roof structure				
Iron sheet	398 (66.00)	32 (8.04)	Ref	
Thatched	205 (34.00)	22 (10.73)	0.274	1.38 (0.78–2.43)
Visible hole on house wall				
No	418 (69.32)	36 (8.61)	Ref	
Yes	185 (30.68)	18 (9.73)	0.658	1.14 (0.63–2.07)
Presence of window				
No	210 (34.83)	16 (7.62)	Ref	
Yes	393 (65.17)	38 (9.67)	0.402	1.30 (0.71–2.39)
Eave gap				
Absent	450 (74.63)	37 (8.22)	Ref	
Present	153 (25.37)	17 (11.11)	0.281	1.40 (0.76–2.56)
Area of residence				
Doyo Yaya	375 (62.19)	49 (13.07)	0.001	6.70 (2.63–17.08)*
Others**	228 (37.81)	5 (2.19)	Ref	
Spray status of the structure				
Not sprayed	372 (61.69)	49 (13.17)	0.001	6.86 (2.69–17.48)*
Sprayed	231 (38.31)	5 (2.16)	Ref	
Animals sleeping in the house				
No	291 (48.26)	25 (8.59)	Ref	
Yes	312 (51.74)	29 (9.29)	0.762	1.09 (0.62–1.91)
Individual residing in				
Index house	120 (19.90)	21 (17.50)	0.001	2.89 (1.60–5.21)*
Neighbour	483 (80.10)	33 (6.83)	Ref	

Table 3 (continued)

Variables	Examined by PCR, n (%)	Positive, n (%)	p-value	COR (95% CI)
<i>Plasmodium</i> species of the index house				
<i>P. vivax</i>	197 (32.67)	9 (4.57)	Ref	
<i>P. falciparum</i>	406 (67.33)	45 (11.08)	0.011	2.60 (1.25–5.44)*
Family size				
Less than five	279 (46.27)	11 (3.94)	Ref	
Five or more	324 (53.73)	43 (13.27)	0.001	3.73 (1.88–7.38)*
Access to ITN				
Sufficient	217 (35.99)	11 (5.07)	Ref	
Not sufficient	386 (64.01)	43 (11.14)	0.015	2.35 (1.18–4.65)*

COR crude odds ratio, ref reference

* Significant at $p < 0.05$, ** includes all the remaining catchment *kebeles* of the selected health centres**Table 4 Results of mixed-effects logistic regression analysis of individual and household-level factors showing main predictors of *Plasmodium* infection, Jimma, south-western Ethiopia**

Characteristics	Examined by PCR, n (%)	Positive, n (%)	AOR (95% CI)	p-value
Fever at the time of survey				
Yes	14 (2.32)	5 (35.71)	12.68 (3.34–48.18)	0.001*
No	589 (97.68)	49 (8.32)	Ref	
Malaria history in the preceding 1 year				
Yes	72 (11.94)	16 (22.22)	3.62 (1.77–7.38)	0.001*
No	531 (88.06)	38 (7.16)	Ref	
Area of residence				
Doyo Yaya	375 (62.19)	49 (13.07)	6.81 (2.49–18.63)	0.001*
Others**	228 (37.81)	5 (2.19)	Ref	
Individual residing in				
Index house	120 (19.90)	21 (17.50)	2.22 (1.16–4.27)	0.016*
Neighbour	483 (80.10)	33 (6.83)	Ref	
Family size				
< 5	279 (46.27)	11 (3.94)	Ref	
≥ 5	324 (53.73)	43 (13.27)	3.35 (1.53–7.33)	0.003*
Eave				
Absent	450 (74.63)	37 (8.22)	Ref	
Present	153 (25.37)	17 (11.11)	2.28 (1.14–4.55)	0.020*

AOR adjusted odds ratio

* Significant at $p < 0.05$ ** includes all the remaining catchment *kebeles* of the selected health centres

predictors of *Plasmodium* infection include area of residence (AOR=6.81, 95% CI 2.49–18.63), living in index house (AOR=2.22, 95% CI 1.16–4.27), eaves (AOR=2.28, 95% CI 1.14–4.55) and family size (AOR=3.35, 95% CI 1.53–7.33).

Discussion

This study revealed that index cases presenting to health facilities enabled detection of good number of additional asymptomatic malaria in the community. Thus, screening of individuals using PCR following index malaria cases presenting to health facilities is a valuable method for detecting additional malaria cases in low transmission settings. However, it should be noted that PCR-based malaria detection techniques are largely restricted to research settings, and not currently in use by the National Malaria Control Programme in Ethiopia. This calls for evaluation of affordable alternative test methods with better sensitivity compared to microscopy to be utilised in elimination settings.

The overall prevalence of malaria detected using the RACD was 8.96%, more than 90% of the cases being due to *P. falciparum*. This is in contrast to the recent report from Jimma, in which less than a third of the malaria cases were due to *P. falciparum* [7], and the national report (63.7% of the documented malaria cases being caused by *P. falciparum*) [6]. There might be high spatio-temporal variation of *Plasmodium* species in a country, and even among villages within the same district, mainly due to difference in ecological, climatic and socio-economic factors, and interventions carried out to control malaria [28, 29]. Significant clustering of *Plasmodium* infection between *kebeles*, and households was observed in this study. The vast majority of cases were found in one of the eight *kebeles* included in this study, and individuals residing within the index house were at twofold higher risk of *Plasmodium* infections as compared to their neighbours. This has an important implication in prioritizing resources for targeted malaria control and elimination efforts in low transmission setting.

Apart from household-level clustering of the malaria cases, significantly higher prevalence of malaria was documented in Doyo Yaya *kebele* compared to other *kebeles*. Heterogeneity in transmission of malaria is a well-known feature [30]. It appears that the clustering of malaria in this *kebele* is likely due to presence of a village not covered in the preceding IRS operation. Indeed, most of the houses which did not receive IRS within the preceding 12 months of the survey were located in this *kebele*. Malaria control interventions in resource-limited areas should, therefore, target such hotspots that may contribute to sustained transmission, and possibly fuel epidemics in such settings. It is worth noting that screening of

malaria was limited to approximately 200 m from index houses in this study, and thus the overall prevalence in the population was not known. Several studies utilized different screening radii around the index houses, often detecting more cases within the index household and among those residing closer to the index houses [20, 31]. However, the optimum screening radius from the index house that should be included during RACD still remains unclear. It appears that the screening radius is mainly influenced by the local epidemiology of malaria and available resources to implement RACD [32].

Improved housing structure is essential in deterring endophagic/endophilic mosquitoes from reaching the occupants [33], thus possibly reducing vector-human contact. In this study, individuals residing in houses with open eaves were significantly more likely to succumb to malaria compared to those living in houses with closed eaves. Significantly higher odds of malaria in houses with open eaves was also reported previously [34, 35]. Earlier reports around the study area indicate that *Anopheles gambiae* sensu lato (presumably *Anopheles arabiensis*), the primary malaria vector in Ethiopia, to be predominantly endophagic [36], which may explain the observed higher prevalence of malaria in those living in houses with open eaves. Significantly higher number of indoor-resting malaria vectors were also observed in houses with open eaves in a previous study from central Ethiopia [37].

The vast majority of the malaria cases detected in this study were asymptomatic, most of which were sub-microscopic. Reports also show that in low-transmission settings, asymptomatic infections are common and most of the asymptomatic infections are sub-microscopic [22, 38]. As asymptomatic malaria carriers apparently do not seek treatment, they may serve as reservoirs of infection [39], jeopardizing elimination efforts. The challenges asymptomatic malaria infections pose to the malaria elimination efforts is also exacerbated by the diagnostic limitations in detecting the infections [40]. In very low transmission settings, sub-microscopic carriers may contribute up to 50% of human-to-mosquito transmissions [41]. Apart from the possible contribution to onward transmission, persistent asymptomatic infections may be associated with other deleterious health outcomes, such as anaemia [42].

In the current study, it was also found that self-reported history of malaria in the preceding 1 year before the survey was significantly associated with *Plasmodium* infection, after adjusting for the other variables. Individuals with history of malaria were more than three times likely to have *Plasmodium* infection. Relapse with vivax malaria is well known, hence, vivax cases could be due to relapse, even if small number of *P. vivax* cases were detected. It could also be attributed to persistent *P. falciparum*

asymptomatic infections. The duration of persistence of asymptomatic *P. falciparum* parasitaemia is debatable. A recent report from Cambodia shows limited duration of asymptomatic *P. falciparum* parasitaemia in low transmission setting [43], while cases of asymptomatic falciparum malaria persisting more than a decade have been documented elsewhere [44].

In this study, PCR detected 2.2-fold higher *Plasmodium* infection as compared to microscopy. This difference is comparable with other previous studies [22, 45]. As the current routine malaria diagnostic protocol in health facilities in Ethiopia involve blood film microscopy, it is inevitable that sub-microscopic infections remain a huge challenge to the envisaged malaria elimination efforts. Thus, alternative field-friendly and more sensitive diagnostic tools such as highly sensitive RDTs (HS-RDTs) or loop-mediated isothermal amplification (LAMP) need to be evaluated and used for detection of sub-microscopic/asymptomatic malaria.

The difference in the prevalence of malaria was not significantly different among different age groups and between sexes. However, family size was a one of the main predictors. In households with five or more occupants, the risk of infection was threefold, compared to those living in families with fewer than five family members. This could be due to higher number of anopheline mosquitoes related with increased number of household occupants [46].

Long-lasting insecticidal nets were the only personal protection tools utilized by the study participants. However, the difference in prevalence of malaria among households with sufficient number of LLINs was not significant. Similarly, there was no significant difference in LLIN usage the preceding night on malaria risk. These could perhaps be due to poor integrity of the LLINs, rendering only partial protection from mosquito bite, and/or possibly inaccurate information provided on the number of nets owned and their usage. The apparent lack of association of sufficient ownership or usage of the LLINs and *Plasmodium* infection could also be related with biting activity of the anopheline mosquitoes in the area. It was reported in a neighbouring district that the peak biting time of *An. gambiae* s.l. was at the early part of the night, before 21:00 h [47], during which most of the inhabitants likely do not sleep. As the LLIN coverage was high in this study, it may have 'herd effect', in that, those not utilizing the nets may be protected as a result of usage of the rest [48].

Sustained residual transmission of malaria in the areas necessitates assessing vector behavioural characteristics and human activities that may contribute to the on-going transmission. As LLINs and IRS interventions target vectors which feed and rest indoors, malaria elimination

using these interventions alone may not be achieved [49]. Outdoor biting by the vectors as a result of long-term use of the vector control tools, and resistance to insecticides used in IRS and treating the LLINs may contribute to sustained low transmission of malaria [50, 51]. Moreover, night-time human activities may increase malaria risk [52].

The study has the following limitations: First, due to limitation of resources, the RACD was limited to catchment *kebeles* of the two health centres. Hence, passively detected malaria cases which resided out of the catchment of the two selected health centres were not included. Second, this study did not include index cases who might have visited private clinics. However, the areas being predominantly rural, it is expected that most of the residents visit the two public health facilities for treatment of fever. Finally, while this study allowed to identify high number of *Plasmodium* infections around the index cases from one site, no individuals living >200 m from index case houses were screened for infection. Thus, it was not known how large this cluster of transmission was.

Conclusions

The study revealed that substantial number of malaria cases, largely asymptomatic, were detected using RACD. Reactive case detection strategies utilizing malaria diagnostic tools of high sensitivity, therefore, complement the routine passive case detection in low malaria transmission areas, and may be used in malaria elimination programme. Significant household and *kebele* level clustering of the malaria cases were observed. Moreover, history of malaria, houses with eaves and family size were risk factors to *Plasmodium* infection. Further studies incorporating greater distance from index households, and including environmental factors affecting risk of malaria infection are recommended.

Abbreviations

DBS: dried blood spot; DNA: deoxyribonucleic acid; ITNs: insecticide-treated nets; LLINs: long-lasting insecticide-treated nets; PCR: polymerase chain reaction; RACD: reactive case detection; RDT: rapid diagnostic test.

Authors' contributions

EZ and DY: Conceived and designed the study. AT and EZ involved in data collection. CK, AT, AKY, DS and ML involved in data analysis. EZ, CK, DY and GY drafted the manuscript. DY and GY critically reviewed the manuscript. All authors read and approved the final manuscript.

Author details

¹ School of Medical Laboratory Sciences, Faculty of Health Sciences, Jimma University, Jimma, Ethiopia. ² Program in Public Health, College of Health Sciences, University of California at Irvine, Irvine, CA 92697, USA. ³ PATH/MACEPA, Addis Ababa, Ethiopia. ⁴ Department of Statistics, College of Natural Sciences, Jimma University, Jimma, Ethiopia. ⁵ Tropical and Infectious Diseases Research Centre, Jimma University, Jimma, Ethiopia.

Acknowledgements

We thank the study participants for taking part in the study, administration of the study *kebeles*, the two health centres and Jimma University for their support. We are also grateful to the laboratory staff of the health centres and Tropical and Infectious Diseases Research Center (TIDRC), Jimma University, who gave technical support in slide readings.

Competing interests

The authors declare that they have no competing interests.

Availability of data and materials

The datasets used for the current study are available from the corresponding author on reasonable request.

Consent for publication

Not applicable.

Ethics approval and consent to participate

The study protocol was reviewed and approved by the Ethical Review Board (IRB) of College of Health Sciences, Jimma University (Ref No. RPGC/486/06). Permission was obtained from administration of the health facilities prior to data collection. The study participants were provided with full information about the purpose of the study, the specimens to be provided and the procedures followed, before enrolment into the study. Written consent was obtained from each study participant, and from parents/guardians in case of minors. Study participants found RDT-positive for malaria during the field work, and those with fever, regardless of the RDT results, were referred to either health centres for confirmation and treatment.

Funding

The study was supported by Grants from the National Institutes of Health (R01 AI050243, U19 AI129326 and D43 TW001505) and Jimma University. The funders had no role in the design of the study and collection, analysis, and interpretation of data and in writing the manuscript or decision to publish.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Received: 30 May 2018 Accepted: 20 October 2018

Published online: 26 October 2018

References

- WHO. World Malaria Report 2016. Geneva: World Health Organization; 2016. <http://www.who.int/malaria/media/world-malaria-report-2016/en/>.
- Cibulskis RE, Alonso P, Aponte J, Aregawi M, Barrette A, Bergeron L, et al. Malaria: global progress 2000–2015 and future challenges. *Infect Dis Poverty*. 2016;5:61.
- Brown G, Rogerson S. Malaria: global challenges for malaria eradication. *Microbiol Aust*. 2016;37:34–8.
- Meleigy M. The quest to be free of malaria. *Bull World Health Organ*. 2007;85:507–8.
- Lucas G. Malaria-free Sri Lanka. *Sri Lanka J Child Health*. 2017;46:1–2.
- FMOH. Health Sector Development Programme Annual Performance Report EFY 2007 (2014/15). Version 1. Addis Ababa: FMOH; 2015.
- Zhou G, Yewhalaw D, Lo E, Zhong D, Wang X, Degefa T, et al. Analysis of asymptomatic and clinical malaria in urban and suburban settings of southwestern Ethiopia in the context of sustaining malaria control and approaching elimination. *Malar J*. 2016;15:250.
- Alemu A, Fuehrer H-P, Getnet G, Tessema B, Noedl H. *Plasmodium ovale curtisi* and *Plasmodium ovale wallikeri* in North-West Ethiopia. *Malar J*. 2013;12:346.
- Otten M, Aregawi M, Were W, Karema C, Medin A, Jima D, et al. Initial evidence of reduction of malaria cases and deaths in Rwanda and Ethiopia due to rapid scale-up of malaria prevention and treatment. *Malar J*. 2009;8:14.
- Yimer F, Animut A, Erko B, Mamo H. Past five-year trend, current prevalence and household knowledge, attitude and practice of malaria in Abeshge, south-central Ethiopia. *Malar J*. 2015;14:230.
- Toyama Y, Ota M, Molla G, Beyene BB. Sharp decline of malaria cases in the Burie Zuria, Dembia, and Mecha districts, Amhara Region, Ethiopia, 2012–2014: descriptive analysis of surveillance data. *Malar J*. 2016;15:104.
- FMOH. Health Sector Transformation Plan 2015/16–2019/20. Addis Ababa: FMOH; 2015.
- Abeku TA, Helinski ME, Kirby MJ, Kefyalew T, Awano T, Batisso E, et al. Monitoring changes in malaria epidemiology and effectiveness of interventions in Ethiopia and Uganda: beyond Garki Project baseline survey. *Malar J*. 2015;14:337.
- PMI. Malaria operational plan FY 2017. Addis Ababa: PMI; 2017.
- Golassa L, Enweji N, Erko B, Aseffa A, Swedberg G. Detection of a substantial number of sub-microscopic *Plasmodium falciparum* infections by polymerase chain reaction: a potential threat to malaria control and diagnosis in Ethiopia. *Malar J*. 2013;12:352.
- Kiattibutr K, Roobsoong W, Sriwichai P, Saesue T, Rachaphaew N, Suan-somjit C, et al. Infectivity of symptomatic and asymptomatic *Plasmodium vivax* infections to a Southeast Asian vector, *Anopheles dirus*. *Int J Parasitol*. 2017;47:163–70.
- Moonen B, Cohen JM, Snow RW, Slutsker L, Drakeley C, Smith DL, et al. Operational strategies to achieve and maintain malaria elimination. *Lancet*. 2010;376:1592–603.
- Sluydts V, Heng S, Coosemans M, Van Roey K, Gryseels C, Canier L, et al. Spatial clustering and risk factors of malaria infections in Ratanakiri Province, Cambodia. *Malar J*. 2014;13:387.
- Rulisa S, Kateera F, Bizimana JP, Agaba S, Dukuzumuremyi J, Baas L, et al. Malaria prevalence, spatial clustering and risk factors in a low endemic area of Eastern Rwanda: a cross sectional study. *PLoS ONE*. 2013;8:e69443.
- Littrell M, Sow G, Ngom A, Ba M, Mboup B, Dieye Y, et al. Case investigation and reactive case detection for malaria elimination in northern Senegal. *Malar J*. 2013;12:331.
- Smith Gueye C, Sanders KC, Galappaththy GN, Rundi C, Tobgay T, Sovannarothe S, et al. Active case detection for malaria elimination: a survey among Asia Pacific countries. *Malar J*. 2013;12:358.
- Harris I, Sharrock WW, Bain LM, Gray K-A, Bobogare A, Boaz L, et al. A large proportion of asymptomatic Plasmodium infections with low and sub-microscopic parasite densities in the low transmission setting of Temotu Province, Solomon Islands: challenges for malaria diagnostics in an elimination setting. *Malar J*. 2010;9:254.
- Yewhalaw D, Legesse W, Van Bortel W, Gebre-Selassie S, Kloos H, Duchateau L, et al. Malaria and water resource development: the case of Gilgel-Gibe hydroelectric dam in Ethiopia. *Malar J*. 2009;8:21.
- Mekonnen SK, Medhin G, Berhe N, Clouse RM, Aseffa A. Efficacy of artemether–lumefantrine therapy for the treatment of uncomplicated *Plasmodium falciparum* malaria in Southwestern Ethiopia. *Malar J*. 2015;14:317.
- Adera TD. Beliefs and traditional treatment of malaria in Kische settlement area, southwest Ethiopia. *Ethiop Med J*. 2003;41:25–34.
- Rosanas-Urgell A, Mueller D, Betuela I, Barnadas C, Iga J, Zimmerman PA, et al. Comparison of diagnostic methods for the detection and quantification of the four sympatric Plasmodium species in field samples from Papua New Guinea. *Malar J*. 2010;9:361.
- WHO. Methods for maintaining coverage with long-lasting insecticidal nets (LLINs). Geneva: World Health Organization; 2013.
- Kumar D, Andimuthu R, Rajan R, Venkatesan M. Spatial trend, environmental and socioeconomic factors associated with malaria prevalence in Chennai. *Malar J*. 2014;13:14.
- Dhimall M, O'Hara RB, Karki R, Thakur GD, Kuch U, Ahrens B. Spatio-temporal distribution of malaria and its association with climatic factors and vector-control interventions in two high-risk districts of Nepal. *Malar J*. 2014;13:457.
- Mogeni P, Omedo I, Nyundo C, Kamau A, Noor A, Bejon P, et al. Effect of transmission intensity on hotspots and micro-epidemiology of malaria in sub-Saharan Africa. *BMC Med*. 2017;15:121.
- Sturrock HJW, Novotny JM, Kunene S, Dlamini S, Zulu Z, Cohen JM, et al. Reactive case detection for malaria elimination: real-life experience from an ongoing program in Swaziland. *PLoS ONE*. 2013;8:e63830.
- WHO. Disease surveillance for malaria elimination: an operational manual. Geneva: World Health Organization; 2012.

33. Konradsen F, Amerasinghe P, van der Hoek W, Amerasinghe F, Perera D, Piyaratne M. Strong association between house characteristics and malaria vectors in Sri Lanka. *Am J Trop Med Hyg.* 2003;68:177–81.
34. Kirby MJ, Green C, Milligan PM, Sismanidis C, Jasseh M, Conway DJ, et al. Risk factors for house-entry by malaria vectors in a rural town and satellite villages in The Gambia. *Malar J.* 2008;7:2.
35. Ghebreyesus TA, Haile M, Witten KH, Getachew A, Yohannes M, Lindsay SW, et al. Household risk factors for malaria among children in the Ethiopian highlands. *Trans R Soc Trop Med Hyg.* 2000;94:17–21.
36. Lelisa K, Asale A, Taye B, Emanu D, Yewhalaw D. Anopheline mosquitoes behaviour and entomological monitoring in southwestern Ethiopia. *J Vector Borne Dis.* 2017;54:240.
37. Animut A, Balkew M, Lindtjorn B. Impact of housing condition on indoor-biting and indoor-resting *Anopheles arabiensis* density in a highland area, central Ethiopia. *Malar J.* 2013;12:393.
38. Tadesse FG, Pett H, Baidjoe A, Lanke K, Grignard L, Sutherland C, et al. Submicroscopic carriage of *Plasmodium falciparum* and *Plasmodium vivax* in a low endemic area in Ethiopia where no parasitaemia was detected by microscopy or rapid diagnostic test. *Malar J.* 2015;14:303.
39. Ganguly S, Saha P, Guha SK, Biswas A, Das S, Kundu PK, et al. High prevalence of asymptomatic malaria in a tribal population in Eastern India. *J Clin Microbiol.* 2013;51:1439–44.
40. Okell LC, Ghani AC, Lyons E, Drakeley CJ. Submicroscopic infection in *Plasmodium falciparum*—endemic populations: a systematic review and meta-analysis. *J Infect Dis.* 2009;200:1509–17.
41. Okell LC, Bousema T, Griffin JT, Ouedraogo AL, Ghani AC, Drakeley CJ. Factors determining the occurrence of submicroscopic malaria infections and their relevance for control. *Nat Commun.* 2012;3:1237.
42. Chen I, Clarke SE, Gosling R, Hamainza B, Killeen G, Magill A, et al. "Asymptomatic" malaria: a chronic and debilitating infection that should be treated. *PLoS Med.* 2016;13:e1001942.
43. Tripura R, Peto TJ, Chalk J, Lee SJ, Sirithiranont P, Nguon C, et al. Persistent *Plasmodium falciparum* and *Plasmodium vivax* infections in a western Cambodian population: implications for prevention, treatment and elimination strategies. *Malar J.* 2016;15:181.
44. Ashley EA, White NJ. The duration of *Plasmodium falciparum* infections. *Malar J.* 2014;13:500.
45. Manjurano A, Okell L, Lukindo T, Reyburn H, Olomi R, Roper C, et al. Association of sub-microscopic malaria parasite carriage with transmission intensity in north-eastern Tanzania. *Malar J.* 2011;10:370.
46. Kateera F, Mens PF, Hakizimana E, Ingabire CM, Muragijemariya L, Karinda P, et al. Malaria parasite carriage and risk determinants in a rural population: a malariometric survey in Rwanda. *Malar J.* 2015;14:16.
47. Taye B, Lelisa K, Emanu D, Asale A, Yewhalaw D. Seasonal dynamics, longevity, and biting activity of Anopheline mosquitoes in southwestern Ethiopia. *J Insect Sci.* 2016;16:6.
48. Howard SC, Omumbo J, Nevill C, Some ES, Donnelly CA, Snow RW. Evidence for a mass community effect of insecticide-treated bednets on the incidence of malaria on the Kenyan coast. *Trans R Soc Trop Med Hyg.* 2000;94:357–60.
49. Govella NJ, Ferguson H. Why use of interventions targeting outdoor biting mosquitoes will be necessary to achieve malaria elimination. *Front Physiol.* 2012;3:199.
50. Reddy MR, Overgaard HJ, Abaga S, Reddy VP, Caccone A, Kiszewski AE, et al. Outdoor host seeking behaviour of *Anopheles gambiae* mosquitoes following initiation of malaria vector control on Bioko Island, Equatorial Guinea. *Malar J.* 2011;10:184.
51. Haji KA, Khatib BO, Smith S, Ali AS, Devine GJ, Coetzee M, et al. Challenges for malaria elimination in Zanzibar: pyrethroid resistance in malaria vectors and poor performance of long-lasting insecticide nets. *Parasit Vectors.* 2013;6:82.
52. Monroe A, Asamoah O, Lam Y, Koenker H, Psychas P, Lynch M, et al. Outdoor-sleeping and other night-time activities in northern Ghana: implications for residual transmission and malaria prevention. *Malar J.* 2015;14:35.

Ready to submit your research? Choose BMC and benefit from:

- fast, convenient online submission
- thorough peer review by experienced researchers in your field
- rapid publication on acceptance
- support for research data, including large and complex data types
- gold Open Access which fosters wider collaboration and increased citations
- maximum visibility for your research: over 100M website views per year

At BMC, research is always in progress.

Learn more biomedcentral.com/submissions

