

CORRECTION

Open Access



Correction: Longitudinal strain correlates with 6-minute walk distance whereas ejection fraction and diastolic parameters do not

John W. Petersen^{1*}, Natalie Bracewell¹, Kevin M. Schneider², Joshua Latner¹, Shuang Yang³ and Yi Guo³

Correction: *Cardiovasc Ultrasound* 22, 6 (2024).

<https://doi.org/10.1186/s12947-024-00325-z>.

Following publication of the original article [1], the authors identified an error in Table 3. For Long. Strain Inferior LV segments, it was published as correlation estimate of -0.035 with p value 0.792. The correct correlation estimate is -0.323 with p value 0.013).

The correct Table 3 is given below.

The original article has been corrected.

The online version of the original article can be found at <https://doi.org/10.1186/s12947-024-00325-z>.

*Correspondence:

John W. Petersen

john.petersen@medicine.ufl.edu

¹Division of Cardiovascular Medicine, University of Florida, 1600 SW Archer Road, PO Box 100288, Gainesville, FL 32610, USA

²Department of Internal Medicine, University of Florida, Gainesville, FL, USA

³Department of Health Outcomes and Biomedical Informatics, University of Florida, Gainesville, FL, USA



This is a U.S. Government work and not under copyright protection in the US; foreign copyright protection may apply 2024. **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated in a credit line to the data.

Table 3 Correlation of measures of global and regional longitudinal strain with 6-minute walk distance

Correlation with 6-minute walk distance	N	Correlation Estimate ^a	p Value ^a
Global Long. Strain	58	-0.312	0.017
Long. Strain Basal LV segments	58	-0.359	0.006
Long. Strain Mid LV segments	58	-0.209	0.115
Long. Strain Apical LV segments	58	-0.254	0.054
Long. Strain Anteroseptal LV seg.	56	-0.202	0.136
Long. Strain Anterior LV segments	58	-0.321	0.014
Long. Strain Anterolateral LV seg.	58	-0.292	0.026
Long. Strain Inferolateral LV seg.	57	-0.259	0.051
Long. Strain Inferior LV segments	58	-0.323	0.013
Long. Strain Inferoseptal LV seg.	58	-0.252	0.057

Published online: 05 July 2024

References

- 1 Petersen JW, Bracewell N, Schneider KM, et al. Longitudinal strain correlates with 6-minute walk distance whereas ejection fraction and diastolic parameters do not. *Cardiovasc Ultrasound*. 2024;22:6. <https://doi.org/10.1186/s12947-024-00325-z>.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.