

ERRATUM

Open Access



# Erratum to: Physical activity of elderly patients with rheumatoid arthritis and healthy individuals: an actigraphy study

Toshihide Hashimoto<sup>1,2\*</sup>, Kazuhiro Yoshiuchi<sup>2</sup>, Shyuji Inada<sup>2</sup>, Kenji Shirakura<sup>1</sup>, Naoki Wada<sup>1</sup>, Kimihiko Takeuchi<sup>3</sup> and Masatoshi Matsushita<sup>3</sup>

## Erratum

After publication of this article [1], two errors were noticed in Table 1 and also in Table 3.

In Table 1 in the Radiographic evaluation section, one of the characteristics was displayed as:  $\leq$  Larsen IV- no. (%), when it should have been:  $\geq$  Larsen IV- no. (%).

In Table 3, one of the headings was displayed as MCA, when it should have been MAC, to refer to the Mean activity count.

The correct version of both Table 1 (Table 1 here) and Table 3 (Table 2 here) are displayed below. Both Tables have also since been updated in the original article [1].

## Author details

<sup>1</sup>Department of Rehabilitation, Gunma University, 3-39-15 Syowa, Maebashi 371-8511 Gunma, Japan. <sup>2</sup>Department of Stress Sciences and Psychosomatic Medicine, The University of Tokyo, 7-3-1 Hongo, Bukyo-ku 113-8655, Tokyo, Japan. <sup>3</sup>Department of Orthopedics, Isesaki Fukushima Hospital, 556-2 Kashima, Isesaki 372-0015 Gunma, Japan.

Received: 13 October 2015 Accepted: 13 October 2015

Published online: 26 October 2015

## Reference

1. Hashimoto T, Yoshiuchi K, Inada S, Shirakura K, Wada N, Takeuchi K, et al. Physical activity of elderly patients with rheumatoid arthritis and healthy individuals: an actigraphy study. *Bio Psycho Soc Med*. 2015;9:19. doi:10.1186/s13030-015-0046-0.

\* Correspondence: hashimoto@mtj.biglobe.ne.jp

<sup>1</sup>Department of Rehabilitation, Gunma University, 3-39-15 Syowa, Maebashi 371-8511 Gunma, Japan

<sup>2</sup>Department of Stress Sciences and Psychosomatic Medicine, The University of Tokyo, 7-3-1 Hongo, Bukyo-ku 113-8655, Tokyo, Japan

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at  
[www.biomedcentral.com/submit](http://www.biomedcentral.com/submit)



**Table 1** Characteristics of participants

	RA patients (n = 20)	Healthy individuals (n = 20)	p
Age mean (SD)	69.4 (5.1)	69.6 (6.6)	NS
Sex female - no. (%)	16 (80.0)	16 (80.0)	NS
Body mass index mean (SD)	21.4 (3.6)	23.4 (3.4)	NS
Comorbidity one or over - no. (%)	7 (35.0)	8 (40.0)	NS
Employment employed - no. (%)	4 (20.0)	7 (35.0)	NS
Family composition			
with children - no. (%)	9 (45.0)	7 (35.0)	NS
with spouse - no. (%)	9 (45.0)	9 (45.0)	
single - no. (%)	2 (10.0)	4 (20.0)	
Disease duration			
<1y - no. (%)	5 (25.0)	-	
1y-5y - no. (%)	2 (10.0)		
≥5y - no. (%)	13 (65.0)		
Radiographic evaluation			
≤ Larsen III- no. (%)	9 (45.0)	-	
≥ Larsen IV- no. (%)	11 (55.0)		
Operation due to RA			
one or over - no. (%)	5 (25.0)	-	
DAS28-CRP			
mean (SD)	3.59 (1.22)	-	
mild (<2.7) - no. (%)	6 (30.0)		
moderate (2.7-4.1) - no. (%)	9 (45.0)		
high (>4.1) - no. (%)	5 (25.0)		

NS non-significance by Welch two sample t test or Fisher's exact test

**Table 2** Spearman's correlations between subjective measures (SF-36, HAQ-DI) and actigraphic measures

	RA patients (n = 20)			Healthy individuals (n = 20)		
	MAC	PAC	LAR	MAC	PAC	LAR
SF-36						
Physical Functioning	0.38	0.03	-0.45*	0.20	0.35	-0.06
Role-Physical	0.33	0.44	-0.31	0.39	0.55*	-0.26
Bodily Pain	0.35	-0.05	-0.47*	0.43	0.51*	-0.33
General Health	0.26	-0.12	-0.21	-0.28	-0.26	0.22
Perception	0.00	-0.15	-0.08	0.08	0.11	-0.14
Vitality	0.23	0.15	-0.27	0.02	0.00	-0.05
Social Functioning	0.22	0.23	-0.23	0.04	0.06	-0.04
Role-Emotional	0.11	0.04	-0.13	-0.05	0.08	0.05
Mental Health						
HAQ-DI	-0.52*	-0.08	0.58**	-0.05	-0.18	-0.14

SF-36 MOS 36-item short-form health survey, HAQ-DI Health Assessment Questionnaire disability index

MAC Mean activity count, PAC Peak activity count, LAR Low activity ratio

\* $p < 0.05$ , \*\* $p < 0.01$