

ERRATUM

Open Access



Erratum to: Effectiveness of two different doses of rituximab for the treatment of rheumatoid arthritis in an international cohort: data from the CERERRA collaboration

Katerina Chatzidionysiou^{1*}, Elisabeth Lie², Evgeny Nasonov³, Galina Lukina³, Merete Lund Hetland⁴, Ulrik Tarp⁵, Ioan Ancuta⁶, Karel Pavelka⁷, Dan C. Nordström⁸, Cem Gabay⁹, Helene Canhão¹⁰, Matija Tomšič¹¹, Piet L. C. M. van Riel¹², Juan Gomez-Reino¹³, Tore K. Kvien² and Ronald F. van Vollenhoven¹

Erratum

Unfortunately, after publication of this article [1] it was noticed that the author list was incorrectly captured during the production process. The original author list contained the group “Rheumatic Diseases Portuguese Register” however this was included erroneously and instead should only be listed as the affiliation for Helene Canhão. The correct author list can be seen above and the original article has been updated to reflect this change.

Author details

¹Department of Medicine, Karolinska Institute, Unit for Clinical Research Therapy, Inflammatory Diseases (ClinTrid), D1:00, Karolinska Universitetssjukhuset, Stockholm 171 76, Sweden. ²Department of Rheumatology, Diakonhjemmet Hospital, Oslo, Norway. ³ARBITER, Institute of Rheumatology, Moscow, Russia. ⁴DANBIO, Department of Rheumatology, Copenhagen University Hospital at Glostrup, Copenhagen, Denmark. ⁵Department of Rheumatology, Aarhus University Hospital, Aarhus, Denmark. ⁶Cantacuzino Hospital, Bucharest, Romania. ⁷Charles University, Prague, Czech Republic. ⁸ROB-FIN Helsinki University Central Hospital, Helsinki, Finland. ⁹SCQM registry, University Hospital of Geneva, Geneva, Switzerland. ¹⁰Rheumatic Diseases Portuguese Register, Rheumatology Research Unit, Instituto de Medicina Molecular, Lisbon, Portugal. ¹¹University Medical Center, Ljubljana, Slovenia. ¹²Radboud University Nijmegen Medical Centre, Nijmegen, Netherlands. ¹³Hospital Clinico Universidad De Santiago, Santiago, Spain.

Received: 9 June 2016 Accepted: 10 June 2016

Published online: 20 June 2016

Reference

1. Chatzidionysiou K, Lie E, Nasonov E, Lukina G, Hetland ML, Tarp U, et al. Effectiveness of two different doses of rituximab for the treatment of rheumatoid arthritis in an international cohort: data from the CERERRA collaboration. *Arthritis Res Ther*. 2016;18:50. doi:10.1186/s13075-016-0951-z.

* Correspondence: Aikaterini.chatzidionysiou@karolinska.se

¹Department of Medicine, Karolinska Institute, Unit for Clinical Research Therapy, Inflammatory Diseases (ClinTrid), D1:00, Karolinska Universitetssjukhuset, Stockholm 171 76, Sweden

Submit your next manuscript to BioMed Central and we will help you at every step:

- We accept pre-submission inquiries
- Our selector tool helps you to find the most relevant journal
- We provide round the clock customer support
- Convenient online submission
- Thorough peer review
- Inclusion in PubMed and all major indexing services
- Maximum visibility for your research

Submit your manuscript at
www.biomedcentral.com/submit

