

RESEARCH NOTE

Open Access



Biofilm and hemagglutinin formation: a Hallmark for drug resistant uropathogenic *Escherichia coli*

Dawit Gebreegziabihir Hagos^{1*}, Tadele Araya Mezgebo¹, Samuel Berhane² and Araya Abraha Medhanyie³

Abstract

Objective: Urinary tract infection (UTI) is one of the most frequent disease encounters in pregnant mothers, and the most drug resistant, biofilm and hemagglutinin producer Uropathogenic *Escherichia coli* (UPEC) is the major etiologic agent. Therefore, the aim of this study was to assess the association between the antimicrobial resistance, and biofilm and hemagglutinin production of Uropathogenic *Escherichia coli*.

Results: UTI among the study participants was 27.3%; and UPEC was found the major etiologic agent followed by coagulase negative staphylococcus. Risk factors, previous history of catheterization and previous history of UTI were found significantly associated with UTI, recurrent UTI, drug resistance and biofilm formation. Of the tested antibiotics, nitrofurantoin was the most effective drug for UPEC. Nearly 100% of the biofilm producers were resistant to norfloxacin, cotrimoxazole, and gentamicin.

Keywords: UPEC, Biofilm, Drug resistance, UTI

Introduction

UTI is one of the major bacterial infections worldwide and about 50% of women experience at least one episode during their lifetime [1–3]. Nearly 25% [4] and 44% [5] of women develop recurrent UTI within 12 months following acute infection, resulting an estimated prevalence of 150 million cases globally per year [4]. Recurrent UTI contributes significantly to UTI associated morbidity and demands huge resources [4, 5]. For example, in the United States of America (USA), more than 7 million people visit health institutes every year and the costs to manage it reaches approximately over \$1 billion every year [6]. Moreover, UTI is among the highest antibiotics prescribed diseases in the health institutions around the worldwide [7, 8].

Though there is no enough and well documented data in developing, a few studies indicated that UTI is one of the top causes of morbidity [9–11]. A community based

study in Nigeria showed that the prevalence of UTI is about 13%, of which majority of this was due to UPEC [12]. Likewise, in Ethiopia, very limited studies indicated the same trend with other developing countries. A study done at Tikur Anbessa Specialized Teaching Hospital, Addis Ababa, Ethiopia indicated that the overall prevalence of UTI is 23.32%, where *E. coli* was the major isolate [13, 14]. Other studies at Felge Hiwot Referral Hospital, Bahirdar and Hawassa Referral Hospital, also showed that the prevalence of UTI was 30.2% [14] and 48% [15] respectively. Furthermore, the burden of UTI is not only morbidity, mortality and economic impact at individual and nationwide, but it also incurs a huge antibiotic consumption and hence one of the causes for increasing drug resistance to the commonly prescribed drugs [1].

Over 80% [9, 10] of the etiologic agent of UTI is UPEC [16]. Frequently, the patient's own intestinal flora is the source of infection [6, 9]. In addition to the ability of UPEC to firmly attach to the urinary bladder and kidney, biofilm formation is also considered another pathogenic determinant, which allows UPEC to persist for a long time in the urinary tract and interfere with bacterial elimination [17]. Biofilms can define as structured

*Correspondence: dawitg@mu.edu.et

¹ Department of Microbiology and Immunology, College of Health Sciences, Mekelle University, Mekelle, Tigray, Ethiopia
Full list of author information is available at the end of the article



bacterial communities embedded in a self-produced exopolysaccharide matrix adherent to any abiotic or biological surface [12, 18]. Currently, antimicrobial resistance is one of major health threats and it is even more in developing countries, where no strict drug monitoring program. The problem is very much significant in UTI; as the major cause (UPEC) is one of the most drug resistant pathogens.

The microbes have evolved a number of mechanisms to evade antimicrobial therapy and the most important way for UPEC is the ability to form the biofilm [19]. Biofilm endows bacteria with several advantages, such as the acquisition of antibiotic tolerance, expression of several virulence factors and increased resistance against phagocytosis and other host defense mechanisms [18]. Studies comparing biofilm positive versus biofilm negative UPEC strains showed that, drug resistance was significantly higher in vitro biofilm formers than non-former [3, 18]. Other studies also indicated that more UPEC drug resistance was observed in strains which produce hemagglutinin [9]. Therefore, the objective of the study was to assess if the very drug resistant nature of UPEC is associated with biofilm formation and hemagglutinin production.

Main text

Method

This cross-sectional study was conducted to assess the association between antimicrobial resistance and biofilm formation and hemagglutinin production of the most eminent drug resistant Uropathogenic pathogens at Mekelle University, College of Health Sciences, Ayder Comprehensive Specialized hospital, Antenatal Care (ANC) Clinics from December 2017 to August 2018. Using a structured questionnaire-based interview, demographic and clinical data were collected from 323 study participants. Training was given to data collectors on how to collect the midstream urine and data. The midstream urine was collected using wide-mouthed and clean container and transported to Mekelle University, College of Health Sciences, Medical Microbiology laboratory within 1 h of collection and cultured to isolate the etiologic agents. The specimen was cultured on CHROMaga, at 37 °C for 24 h aerobically. The bacterial isolates were further identified by standard biochemical tests [1, 2]. On the UPEC isolates, hemagglutination test was performed by mixing one drop of bacterial suspension with one drop of 3% blood group “O” red blood cells in phosphate-buffered saline (PBS) with and without 3% mannose [4, 6]. Biofilm formation was also tested by Congo red agar (CRA) method. The culture media contain brain heart infusion broth 37 g/L, sucrose 50 g/L, agar 10 g/L and Congo red 8 g/L. Plates were inoculated and incubated aerobically for 24 h at 37 °C [5, 6].

On Muller Hinton agar, Kirby-Bauer disc diffusion assay was carried out to determine the antimicrobial susceptibility profiles [3]. Based on the frequency of drug prescription in the study area, the following antibiotics (Oxoid UK) were included: amoxicillin: 30 mg, gentamicin: 10 mg, cefotaxime: 30 mg, nalidixic acid 30 mg, ciprofloxacin: 5 mg, ofloxacin: 5 mg, norfloxacin: 10 mg, erythromycin: 15 mg, oxacillin: 5 mg, vancomycin: 30 mg, nitrofurantoin: 300 mg and tetracycline: 30 mg. Quality control strains; *S. aureus* (ATCC 25923) and *E. coli* (ATCC 25922) were used as a quality control for culture and susceptibility testing throughout the study. Data analysis was done by SPSS Ver. 20 and descriptive and multivariate analysis was performed. Those variables which were statistically significant, ($p < 0.05$) in bivariate logistic regression were moved to multivariate logistic regression.

Results

A total of 323 pregnant women with symptomatic of UTI were included in this study. Median age of the study subjects were 24 ± 5.4 years old and 297 (91.9%) were urban residents (Table 1). Among the study subjects, 283 (87.6%) were married, and about 176 (54.5%) were house wives. Regarding educational status, about 47 (13.2%) and 55 (17%) of them were illiterate and attended college education (Table 1).

Of the total, bacteriuria was detected in 87 (26.9%) of the study participants. In multivariate analysis previous history of UTI (AOR=7.057, 95% CI (5.269, 18.654), $P=0.002$) and previous history of catheterization (AOR=2.870, 95% CI (1.516, 9.122), $P=0.003$) were significantly associated with UTI. Though the number of patients with previous history of catheterization were small 23 (7.1%), it was found significantly associated with UTI ($P=0.03$). Variables like marital status, residency, education status, other chronic diseases like diabetic mellitus and previous hospitalization were not showed a significant association with UTI (Table 2).

Five bacterial species were isolated and of these with significant bacteriuria, 94.8% were had a single isolate, while the remaining 5.2% were dually infected. Out of the total isolates, *E. coli* was the major isolated, which is followed by coagulase negative staphylococci (CoNS), *S. aureus*, and *K. pneumoniae*. Majority of the isolates were gram negative uropathogens and the resistance rate against Ciprofloxacin, Tetracycline, Trimethoprim-Sulfamethoxazole, Ceftriaxone, Amoxicillin-Clavulanic acid, Norfloxacin, Cotrimoxazole, Cefotaxime, Amikacin, Gentamicin; range from 27.0 to 100%. However, all Gram-negative bacterial isolates were showed relatively low level of resistance to nitrofurantoin (3.8%) and cef-tazidime (17%).

Table 1 Socio-demographic characteristics of the study participants in Ayder Comprehensive Specialized Hospital December 2017–August 2018 (n = 323)

Variables	Frequency	Percent (%)
Age (years)		
15–24	134	41.5
25–34	169	52.3
35–44	20	6.2
Residence		
Urban	297	91.9
Rural	26	8.1
Occupation		
House wife	176	54.5
Merchant	54	16.7
Student	12	3.7
Government employee	81	25.1
Educational level		
Illiterate	47	14.6
Only read and write	26	8.1
Primary school	81	25.1
Secondary school	114	35.2
College and above	55	17.0
Marital status		
Single	15	4.6
Married	283	87.6
Divorced	8	2.5
Widowed	7	2.2
Others	6	3.3

The leading isolate *UPEC*, showed high level of resistance to Ampicilin 13 (33.3%) and tetracycline 9 (23.1%). On contrary, 76.5% of the *UPEC* were relatively sensitive to ceftazidime and norfloxacin each, followed by nitrofurantoin (96.2%). *K. pneumoniae* isolates were 96% resistant to, Gentamicin, Amoxicillin-Clavulanic acid and Trimethoprim-Sulfamethoxazole. Other drugs like Norfloxacin, Amikacin, Nitrofurantoin, Ciprofloxacin, Ceftriaxone, Cefotaxime, Ceftazidime and Ampicillin were resistant to 60.0% of *K. pneumoniae* isolates. All the *UPEC* isolates (n=39) were tested for biofilm formation and hemagglutins production. Of these, hemagglutinin and biofilm formation were detected in 38 (97.4%) and 24 (61.5%) respectively. In this study, 23 (96%) of the biofilm producing *UPEC* isolates were found Multi-drug Resistant (MDR). More interestingly, all biofilm producing *UPEC* were found 100% resistant to antibiotic like norfloxacin, cotrimoxazole, and gentamicin. However, nitrofurantoin was found to be 98% effective to the biofilm producing *UPEC*. On multivariate analysis biofilm and hemagglutins producing *UPEC* isolates were significantly associated with being MDR ($P=0.002$).

Discussion

The prevalence of UTI in our study was 27.3% and this is in agreement with a study done in Addis Ababa 23.3% (25), Nigeria 25.3% [20], Cameroon 23.5% [11], Bar-Dar (30.0%) and lower than the other studies in Dire Dawa 14.4 [21, 22], Tanzania 14.6% [23] and Brazil (15.7%) [24]. On the contrary, our finding was higher from studies done in Bahir Dar 9.5% [21] and Addis Ababa 11.6% [13]. The difference might be due to geographical difference and sample size effect. In addition, the health care system provision, especially the antenatal services are improving from time to time and hence, the time differences might be also the possible explanation for disparities between our result and the reports earlier reports from Ethiopia. In our study, the predominant etiologic agent of UTI was *UPEC* and this is in agreement with almost all studies done around the world [3, 19, 24, 25].

Previous history of UTI and history of catheterization were significantly associated with UTI and that is similar to studies reported from Bahir Dar [26], and Pakistan [27]. The possible justification for this may be, due to empirical treatment and overuse of drugs. However, maternal age, residence, educational level, marital status and occupation are variables which did not show a significant association with UTI and this was in agreement with reports from Bahir Dar [26], Gondar [14], Sudan [28] and Tanzania [23].

As a result of absence of appropriate diagnostic tools for diagnosis and drug susceptibility testing plus extensive use empirical treatment, development of drug resistance is vertically raising [18]. In our case drug resistance is defined as; if an isolate is resistant to two or more different chemical classes, then we considered as MDR isolate. In the present study, 76.7% of the isolates were found MDR and this was higher than a study report from Addis Ababa, Tikur Anbesa Specialized Hospital [25] and lower than the findings from Gondar [14] and Dire Dawa [22] and might be justified by the frequent and inappropriate use of antibiotics.

Adhesion is a crucial and primary step for *UPEC* to establish and cause disease along the urinary tract and this is mediated by type 1 fimbriae. Adhesion to the epithelial cells is very important to resist the host immune response and ascend to the upper parts of the urinary tract. In our study, *UPEC* positive for hemagglutinin formation were significantly associated with MDR ($p=0.001$) and with recurrent UTI ($p=0.01$); and this was in agreement with a study conducted in India [18]. Different studies report that the ability of *UPEC* to cause recurrent UTI is because of their ability to form biofilm on the epithelial layer of urinary tract and inanimate plastic materials. In our study, biofilm formation was detected in about 61.7% and this is in agreement

Table 2 Bivariate and multivariate logistic regression analysis of factors associated with UTI among pregnant women attending ANC clinic of ACSH, December 2017–August 2018

Variables	Culture		COR (95% CI)	P-value	AOR (95% CI)	P-value
	SB negative no. (%)	SB positive no. (%)				
Educational level						
Illiterate	28 (59.6)	19 (40.4)	2.07 (0.23, 7.008)	0.078	3.001 (0.702, 16.903)	0.115
Only write & write	11 (53.4)	10 (47.6)	1.711 (0.513, 7.940)	0.234	3.975 (0.574, 22.876)	0.110
Primary school	55 (73.3)	20 (26.7)	1.561 (0.685, 3.987)	0.209	2.421 (0.692, 9.759)	0.170
Secondary school	69 (75.0)	23 (25.0)	1.602 (0.573, 4.112)	0.363	1.989 (0.766, 7.095)	0.152
Higher education	77 (83.7)	15 (16.3)	1			
History of UTI						
Yes	81 (71.1)	33 (28.9)	9.932 (4.904, 20.114)	0.001	8.015 (4.005, 19.908)	0.001*
No	201 (96.2)	8 (3.8)	1			
History of catheterization						
Yes	86 (83.5)	17 (16.5)	6.003 (3.113, 10.976)	0.002	3.270 (1.316, 8.122)	0.012*
No	214 (97.3)	6 (2.7)	1			
History of hospitalization (last 3 months)						
Yes	62 (78.5)	17 (21.5)	4.038 (1.843, 8.633)	0.032	3.102 (0.535, 11.765)	0.348
No	201 (82.4)	43 (17.6)	1			
History of antibiotic use (last 3 months)						
Yes	37 (61.7)	23 (38.3)	1.227 (2.007, 6.406)	0.057	2.102 (0.724, 3.935)	0.834
No	217 (82.5)	46 (17.5)	1			

SB Significant bacteriuria, ACSH Ayder Comprehensive Specialized Hospital, COR crude odds ratio, AOR adjusted odds ratio

with studies conducted in India (22), and lower than another study conducted in India [29]. The relationship between being drug resistant and biofilm formation was showed a statistically significant relationship ($P=0.02$), and this was in agreement with the studies conducted in India [18]. Moreover, in the present study, biofilm formation was significantly associated with history of catheterization and history of previous UTI, which implies, biofilm could be one of the main factors for UPEC to develop drug resistance. However, UPEC was found uniquely sensitive to nitrofurantoin and hence it could be the best choice of drug for patient suffering from recurrent UTI and for those who failed to respond to the current treatment regimen.

Limitations

The present study did not address the different serotypes of *E. coli*.

Abbreviations

ANC: antenatal Care; CRA: congo red agar; MDR: multi-drug Resistant; PBS: phosphate-Buffered Saline; UTI: urinary tract infection; UPEC: uropathogenic *Escherichia coli*; USA: United States of America.

Acknowledgements

We, authors would like to thank the study participants. Furthermore, I would like to profoundly grateful to Medical Microbiology Laboratory technologists, who helped me during laboratory analysis.

Authors' contributions

DGH: Design the research, perform the research, prepare manuscript and Submit for publication; AAM: Data analysis and Manuscript editing; TAM: sample processing and laboratory analysis; SBG: Consult patient and Participants selection. All authors read and approved the final manuscript.

Funding

This research project is funded by Mekelle University, College of Health Sciences (Grant No. 40,000.00 Eth.Birr) via the small-scale recurrent budget. The funder was not having any role the in study design, data collection, data analysis, interpretation, writing of this manuscripts.

Availability of data and materials

All generated data are included in this article.

Ethics approval and consent to participate

Ethical approval was obtained from the Institutional Review Board of Mekelle University, College of Health Sciences; [Project No. CRPO/CHS/SM/010/09]. Permission to carry out the study was also obtained from the hospital medical director office and all were enrolled after they agreed and signed on the informed consent. We were also prepared an assent form, however all the study participants were greater than 18 years. The minimum age of the study participants were 19 years old. All records of patient were kept confidential and anonymous. Filled out questionnaires were carefully handled and access to results was kept strictly.

Consent to publish

Not applicable.

Competing interests

The authors declare that they have no competing interests.

Author details

¹ Department of Microbiology and Immunology, College of Health Sciences, Mekelle University, Mekelle, Tigray, Ethiopia. ² Department of Internal Medicine, College of Health Sciences, Mekelle University, Mekelle, Tigray, Ethiopia. ³ School of Public Health, College of Health Sciences, Mekelle University, Mekelle, Tigray, Ethiopia.

Received: 18 March 2019 Accepted: 11 June 2019

Published online: 24 June 2019

References

- Köves B (2014) The role of bacterial virulence factors in the clinical course of urinary tract infections. Ph.D. Thesis. Department of Microbiology, Immunology and Glycobiology. Institute of Laboratory Medicine. Lund University, Lund, Sweden
- Sivick KE, Mobley HL. Waging war against uropathogenic *Escherichia coli*: winning back the urinary tract. *Infect Immun*. 2010;78(2):568–85.
- Totsika M, Gomes Moriel D, Idris A, A Rogers B, J Wurpel D, Phan MD, et al. Uropathogenic *Escherichia coli* mediated urinary tract infection. *Curr Drug Targets*. 2012;13(11):1386–99.
- O'Hanley P. Prospects for urinary tract infection vaccines. *Urinary tract infection: molecular pathogenesis and clinical management*. 1996:405–425.
- Hilbert DW, Paulish-Miller TE, Tan CK, Carey AJ, Ulett GC, Mordechai E, et al. Clinical *Escherichia coli* isolates utilize alpha-hemolysin to inhibit in vitro epithelial cytokine production. *Microbes Infect*. 2012;14(7–8):628–38.
- Langermann S, Möllby R, Burlein JE, Palaszynski SR, Gale Auguste C, DeFusco A, et al. Vaccination with fimbriae protects cynomolgus monkeys from colonization and infection by uropathogenic *Escherichia coli*. *J Infect Dis*. 2000;181(2):774–8.
- Wilson ML, Gaido L. Laboratory diagnosis of urinary tract infections in adult patients. *Clin Infect Dis*. 2004;38(8):1150–8.
- Foxman B. Epidemiology of urinary tract infections: incidence, morbidity, and economic costs. *Am J Med*. 2002;113(1):5–13.
- Momtaz H, Karimian A, Madani M, Dehkordi FS, Ranjbar R, Sarshar M, et al. Uropathogenic *Escherichia coli* in Iran: serogroup distributions, virulence factors and antimicrobial resistance properties. *Ann Clin Microbiol Antimicrob*. 2013;12(1):8.
- Cusumano CK, Pinkner JS, Han Z, Greene SE, Ford BA, Crowley JR, et al. Treatment and prevention of urinary tract infection with orally active fimbriae inhibitors. *Sci Transl Med*. 2011;3(109):109ra115–109ra115.
- Gupta K, Hooton TM, Stamm WE. Increasing antimicrobial resistance and the management of uncomplicated community-acquired urinary tract infections. *Ann Intern Med*. 2001;135(1):41–50.
- Blango MG, Mulvey MA. Persistence of uropathogenic *Escherichia coli* in the face of multiple antibiotics. *Antimicrob Agents Chemother*. 2010;54(5):1855–63.
- Kabew G, Abebe T, Miheret A. A retrospective study on prevalence and antimicrobial susceptibility patterns of bacterial isolates from urinary tract infections in Tikur Anbessa Specialized teaching hospital Addis Ababa, Ethiopia, 2011. *Ethiop J Health Dev*. 2013;27(2):111–7.
- Alemu A, Moges F, Shiferaw Y, Tafess K, Kassu A, Anagaw B, et al. Bacterial profile and drug susceptibility pattern of urinary tract infection in pregnant women at University of Gondar Teaching Hospital, Northwest Ethiopia. *BMC Res Notes*. 2012;5(1):197.
- Gizachew M, Kebede M, Merid Y, Sinshaw Y, Tiruneh M, Alemayehu M, et al. *Escherichia coli* isolated from patients suspected for urinary tract infections in Hawassa Referral Hospital, Southern Ethiopia: an institution based cross sectional study. *J Microbiol Res*. 2013;1:9–15.
- Totsika M, Kostakioti M, Hannan TJ, Upton M, Beatson SA, Janetka JW, et al. A fimbriae inhibitor prevents acute bladder infection and treats chronic cystitis caused by multidrug-resistant uropathogenic *Escherichia coli* ST131. *J Infect Dis*. 2013;208(6):921–8.
- Forbes BA, Sahm DF, Weissfeld AS. *Study guide for Bailey and Scott's diagnostic microbiology-e-book*. St Louis: Elsevier Health Sciences; 2016.
- Maheswari UB, Palvai S, Anuradha PR, Kammili N. Hemagglutination and biofilm formation as virulence markers of uropathogenic *Escherichia coli* in acute urinary tract infections and urolithiasis. *Indian J Urol*. 2013;29(4):277.
- Fatima N, Agrawal M, Shukla I, Khan P. Characterization of uropathogenic *E. coli* in relation to virulence factors. *Open Access Scientific reports*. 2012.
- Iregbu K, Nwajioji-Princewill P. Urinary tract infections in a tertiary hospital in Abuja, Nigeria. *Afr J Clin Exp Microbiol*. 2013;14(3):169–73.
- Biadlegne F, Abera B. Antimicrobial resistance of bacterial isolates from urinary tract infections at Felge Hiwot Referral Hospital, Ethiopia. *Ethiop J Health Dev*. 2009;23(3):236–8.
- Derese B, Kedir H, Teklemariam Z, Weldegebreal F, Balakrishnan S. Bacterial profile of urinary tract infection and antimicrobial susceptibility pattern among pregnant women attending at antenatal clinic in Dil Chora Referral Hospital, Dire Dawa, Eastern Ethiopia. *Ther Clin Risk Manage*. 2016;12:251.
- Masinde A, Gumodoka B, Kilonzo A, Mshana S. Prevalence of urinary tract infection among pregnant women at Bugando Medical Centre, Mwanza, Tanzania. *Tanzan J Health Res*. 2009. <https://doi.org/10.4314/thrb.v11i3.47704>.
- de Vasconcelos-Pereira EF, Figueiró-Filho EA, de Oliveira VM, Fernandes ACO, de Moura Fé CS, Coelho LR, et al. Urinary tract infection in high risk pregnant women. *Infection*. 2013;7(25):27–30.
- Assefa A, Asrat D, Woldeamanuel Y, Abdella A, Melesse T. Bacterial profile and drug susceptibility pattern of urinary tract infection in pregnant women at Tikur Anbessa Specialized Hospital Addis Ababa, Ethiopia. *Ethiop Med J*. 2008;46(3):227–35.
- Demilie T, Beyene G, Melaku S, Tsegaye W. Urinary bacterial profile and antibiotic susceptibility pattern among pregnant women in North West Ethiopia. *Ethiop J Health Sci*. 2012;22(2).
- Haider G, Zehra N, Munir AA, Haider A. Risk factors of urinary tract infection in pregnancy. *J Pak Med Assoc*. 2010;60(3):213.
- Hamdan HZ, Ziad AHM, Ali SK, Adam I. Epidemiology of urinary tract infections and antibiotics sensitivity among pregnant women at Khartoum North Hospital. *Ann Clin Microbiol Antimicrob*. 2011;10(1):2.
- Mittal S, Sharma M, Chaudhary U. Biofilm and multidrug resistance in uropathogenic *Escherichia coli*. *Pathog Glob Health*. 2015;109(1):26–9.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.