

RESEARCH

Open Access



Circular RNA BIRC6 depletion promotes osteogenic differentiation of periodontal ligament stem cells via the miR-543/PTEN/PI3K/AKT/mTOR signaling pathway in the inflammatory microenvironment

Xinyuan Zhao¹, Wenjuan Sun², Bing Guo^{3*} and Li Cui^{1,4*} 

Abstract

Background: Periodontal ligament stem cells (PDLSCs) are the ideal seed cells for periodontal tissue regeneration. It is well established that persistent inflammation significantly impairs the osteogenic differentiation capability of PDLSCs. Therefore, maintaining PDLSC osteogenic potential under the inflammatory microenvironment is important for treating bone loss in periodontitis. The aim of our study was to explore the potential role of circular RNA BIRC6 (circBIRC6) in regulating osteogenic differentiation of PDLSCs in the inflammatory conditions.

Methods: Alkaline phosphatase staining, Alizarin Red staining, quantitative real-time polymerase chain reaction, western blotting and immunofluorescence staining were used to evaluate the effects of circBIRC6 on the osteogenic differentiation of PDLSCs. RNA pull-down and luciferase assays were performed to explore the interaction between circBIRC6 and miR-543. Then, the downstream signaling pathway affected by circBIRC6/miR-543 axis was further investigated.

Results: The expression level of circBIRC6 was higher in PDLSCs exposed to inflammatory stimulus and in periodontitis tissues compared to the respective controls. Downregulation of circBIRC6 enhanced the osteogenic potential of PDLSCs under the inflammatory conditions, and upregulation of circBIRC6 led to opposite findings. Mechanistically, we found that circBIRC6 modulated PDLSC osteogenic differentiation through sponging miR-543. More importantly, we have demonstrated that circBIRC6/miR-543 axis regulated the mineralization capacity of PDLSCs via PTEN/PI3K/AKT/mTOR signaling pathway in the inflammatory microenvironment.

Conclusions: In summary, the expression of miR-543 is significantly increased following circBIRC6 downregulation, leading to inhibition of PTEN and subsequently activation of PI3K/AKT/mTOR signaling pathway. Therefore, targeting circBIRC6 might represent a potential therapeutic strategy for improving bone loss in periodontitis.

*Correspondence: guobing@mail.sysu.edu.cn; licui@smu.edu.cn

¹ Stomatological Hospital, Southern Medical University, Guangzhou 510280, China

³ Department of Dentistry, The First Affiliated Hospital, Sun Yat-Sen University, Guangzhou 510080, China

Full list of author information is available at the end of the article



Keywords: circBIRC6, Osteogenic differentiation, Periodontal ligament stem cells, Inflammatory microenvironment, miR-543, PTEN

Background

Periodontitis is a chronic, multifactorial inflammatory disease that is characterized by destruction of bone and supporting soft tissues surrounding the tooth [1, 2]. In addition, accumulating evidence indicates that periodontitis is a potential risk factor for a number of systemic conditions [3]. Unfortunately, the currently available periodontal treatments can only retard the progression of periodontitis with minimal tissue regeneration. Therefore, it is urgently needed to explore and develop effective and safe periodontal therapies.

The goal of periodontal therapy is to fully restore the damaged/lost supporting periodontal tissues. The regeneration of periodontitis-related bone loss is a key challenge in clinical dentistry, and enhancing the osteogenic differentiation capacity of seed cells is of great importance for treating periodontitis. Due to the potential to differentiate into osteoblasts, the capability to modulate the immune and inflammatory milieu and lack of ethical issues, periodontal ligament stem cells (PDLSCs) are becoming attractive candidates for regenerating bone defects in the periodontitis microenvironment [4]. However, increasing evidence has demonstrated that the local periodontitis microenvironment significantly impairs the osteogenic potential of PDLSCs [5, 6]. Thus, maintaining the normal biological function of PDLSCs, especially their osteogenic potential, in the harsh inflammatory microenvironment is a major challenge for periodontal tissue repair and regeneration.

Circular RNAs (circRNAs) are a type of evolutionarily conserved, covalently closed non-coding RNAs generated from backsplicing of exons, introns, or both [7, 8]. CircRNAs are highly abundant in eukaryotic transcriptome and are more stable than the linear RNAs due to the characterized loop structure. CircRNAs have been shown to play a critical role in regulating gene and protein expressions by acting as microRNA (miRNA) sponge, or directly translating proteins, or interacting with proteins to affect downstream signaling pathways, etc. [7]. CircRNAs are important for modulating the osteogenic differentiation capacity of PDLSCs. For instance, the expression profile of circRNAs was found to be significantly changed during PDLSC osteogenic differentiation [9]. Ye et al. [10] reported that circFAT1 depletion suppressed the osteogenic differentiation of PDLSCs in vitro and bone formation in vivo, suggesting that circFAT1 might be a potential

candidate regulator of PDLSC osteogenesis. Similarly, the expression level of CDR1as was significantly increased during osteogenic differentiation of PDLSCs. Both in vitro and in vivo results revealed that targeting CDR1as markedly inhibited the osteogenic potential of PDLSCs [11]. These findings indicate that circRNAs are actively involving in modulating the osteogenic capacity of PDLSCs, and targeted manipulation of circRNA level might be a novel and promising strategy for bone regeneration under the condition of periodontitis.

Circular RNA BIRC6 (circBIRC6) is a recently discovered circRNA and located on chromosome 2. CircBIRC6 is enriched in human embryonic stem cells (hESCs) and plays an important role in maintaining the pluripotent state of hESCs [12]. The expression level of circBirc6 was found to be deregulated in the myocardial infarction mouse hearts [13]. In addition, abnormal expression of circBIRC6 is actively involved in regulation of tumorigenesis [14, 15]. These currently available findings indicate that circBIRC6 is important for maintaining normal cellular function. Our preliminary study revealed that circBIRC6 level was significantly increased in PDLSCs under the in vitro inflammatory microenvironment, suggesting that its aberrant expression might affect the biological functions of PDLSCs. In this study, we aimed to explore the effects of circBIRC6 on the biological functions of PDLSCs under the inflammatory conditions.

Materials and methods

Clinical samples

All clinical samples were obtained from the Stomatological Hospital, Southern Medical University. Written informed consent was obtained from the participants. This study was approved by the Ethics Committee of the Stomatological Hospital, Southern Medical University.

Cell culture

The PDLSCs were isolated from the periodontal ligament tissues of healthy impacted third molars and then maintained in Dulbecco's modified Eagle's medium (DMEM) supplied with 10% fetal bovine serum, 100 U/mL penicillin and 100 µg/mL streptomycin. For the osteogenic differentiation induction, the PDLSCs were cultured with Mesenchymal Stem Cell Osteogenic Differentiation Medium (Cyagen Biosciences, Guangzhou, China). TNF-α (10 ng/mL), IL-1β (10 ng/mL) and *P. gingivalis*-LPS (10 µg/mL) were used to mimic an in vitro inflammatory microenvironment.

Plasmid construction and cell transfection

The circBIRC6 overexpression plasmid was generated by inserting the circBIRC6 sequence into the pLCDH-ciR vector. The shRNAs targeting circBIRC6 were inserted into the lentiviral vector, and shRNA oligonucleotide sequences were designed by GenePharma (Shanghai, China). Mutant circBIRC6 was constructed by changing the interaction residues between circBIRC6 and miR-543. The miR-543 mimic, miR-543 inhibitor and their respective controls were synthesized by GenePharma. The PTEN overexpression plasmid was constructed by inserting full-length PTEN into the vector. The PTEN shRNA lentiviral particles were purchased from Santa Cruz Biotechnology, Inc. (Santa Cruz, CA, USA). Lentiviruses with circBIRC6 overexpression/inhibition and PTEN overexpression were produced as previously described [16]. The PDLSCs were infected with lentiviruses and/or transfected with miRNA mimics/inhibitors using Lipofectamine 3000 (Invitrogen, Carlsbad, CA, USA) according to the manufacturer's instructions.

Quantitative real-time polymerase chain reaction analysis

Total RNA was extracted with the Quick-RNA™ kit (Zymo Research Corp, Irvine, CA, USA). The SuperScript™ III First-Strand Synthesis SuperMix (Invitrogen) was used to synthesize first-strand cDNA. The amplification of cDNA was performed with Light Cycler 480@ SYBR Green I MasterMix (Roche, Applied Science, Indianapolis, IN, USA) on the CFX96 Real-Time PCR detection system (Bio-Rad, Hercules, CA, USA). GAPDH and U6 were used as the internal controls, and the relative gene expression levels were determined by the $2^{-\Delta\Delta CT}$ method. The circBIRC6 primers were designed and provided by GenePharma. The primer sequences used in this study are listed in Additional file 1: Table S1.

Western blotting

Followed by separation with 4–20% tris-glycine gels, the proteins were transferred onto a PVDF membrane using a Trans-Blot Turbo system (Bio-Rad). After soaking the membrane in 5% nonfat milk for 1 h at room temperature, the primary antibodies and HRP-linked secondary antibodies (Proteintech, Chicago, IL, USA) were used to detect the targeted proteins. The primary antibodies for BMP2, OPN, PTEN, PI3K, AKT, p-AKT, mTOR and p-mTOR were from Proteintech. The p-PI3K primary antibody was from Abcam. The primary antibodies against OCN and BSP were from Santa Cruz Biotechnology and Thermo Fisher Scientific, respectively.

Immunofluorescence staining

Briefly, the cells were fixed with paraformaldehyde for 10 min and then permeated with 0.1% Triton X-100 for 10 min. After washing three times with PBS, the samples were blocked with immunostaining blocking buffer (Beyotime, Shanghai, China) for 1 h at room temperature. Then, the cells were incubated with primary antibodies against RUNX2 (Proteintech) and COL1A1 (Proteintech) at 4 °C overnight, and subsequently with a fluorescence-conjugated secondary antibody at room temperature for 1 h. Images were taken with an inverted fluorescence microscope (Leica Microsystems, Wetzlar, Germany).

Alkaline phosphatase (ALP) staining

The cells were fixed with 4% paraformaldehyde for 10 min after 10 days induction of osteogenic differentiation. Then, the samples were incubated in 0.1 M Tris buffer (pH 9.3) containing 0.25% naphthol AS-BI phosphate (Sigma-Aldrich, Saint Louis, MO, USA) and 0.75% Fast Blue BB (Sigma-Aldrich) at 37 °C.

Alizarin Red S (ARS) staining

Briefly, after three-week osteogenic differentiation induction, the fixed cells were stained with Alizarin Red S solution (Cyagen Biosciences) for 20 min at room temperature.

MTT assay

The cells were incubated with MTT solution (Sigma-Aldrich) for 4 h at 37 °C. Then, dimethyl sulfoxide was used to dissolve the sediment. The absorbance was determined with a Synergy HT microplate reader (BioTek Instruments, Winooski, VT, USA) at 570 nm wavelength.

EdU (5-ethynyl-2'-deoxyuridine) assay

The cells were incubated with 10 μM EdU for at 37 °C for 2 h, and then, the fixation and permeabilization of the samples were performed with 3.7% formaldehyde and 0.1% Triton X-100, respectively. The Click-iT™ reaction cocktail (Invitrogen) and Hoechst 33342 were used to stain the cells. Images were taken with an inverted fluorescence microscope (Leica Microsystems).

Flow cytometry

The harvested cells were fixed with ice-cold ethanol overnight. After removing the fixative, the samples were stained with propidium iodide and incubated in dark for 20 min. The cell cycle distribution was detected using

DxFLEX Flow Cytometer (Beckman Coulter, Brea, CA, USA).

RNA immunoprecipitation (RIP)

The RIP assay was performed using a Magna RIP Kit (Millipore, Billerica, MA, USA) according to the manufacturer's instructions. The AGO2 antibody (Abcam) and the negative control IgG antibody were used for the RIP experiments.

Pull-down assay

Briefly, the biotinylated circBIRC6 probe and oligo probe were incubated with Streptavidin-Dyna beads M-280 (Invitrogen). Then, the cell lysates were incubated with beads at 4 °C overnight, and the beads were subsequently washed. The RNA was extracted and evaluated by qRT-PCR.

Luciferase reporter assay

The circBIRC6 wt, circBIRC6 mut, miR-543 mimic, miR-543 inhibitor and miRNA negative controls were transfected into PDLSCs alone or in combination using the Lipofectamine 3000 transfection reagent (Invitrogen). Following 24 h of incubation, the luciferase reporter assay kit (Promega, Madison, WI, USA) was used to determine the relative luciferase activities.

Statistical analysis

All statistical analyses were performed with the GraphPad Prism 9.0 (GraphPad Software, San Diego, CA, USA). Student's t test and one-way ANOVA were used to compare the difference between groups. The correlation between circBIRC6 and miR-543 in periodontitis tissues was evaluated with the Pearson correlation analysis. *P* values less than 0.05 were considered as statistically significant.

Results


The osteogenic differentiation of PDLSCs is impaired in the inflammatory microenvironment

The in vitro inflammatory microenvironment was mimicked by adding TNF- α , IL-1 β and *P. gingivalis*-LPS to the culture medium. Then, the effects of inflammatory conditions on the proliferation and osteogenic differentiation of PDLSCs were evaluated. The MTT assay showed that the OD values changed little between the PDLSCs under the normal condition and those exposed to inflammation (Fig. 1A). Similarly, no significant alteration was found for the percentage of EdU positive cells and cell cycle distribution between the control group and the inflammation group (Fig. 1B, C), indicating that the inflammatory microenvironment has little effects on the proliferation of PDLSCs. For the osteogenic differentiation, the ALP

staining assay showed that the staining intensity was lower in the inflammation group compared to the control group (Fig. 1D). The ARS staining revealed that cells exposure to the inflammatory conditions formed significantly less mineralized nodules than those grown in the normal growth microenvironment (Fig. 1E). In addition, the qRT-PCR and WB demonstrated that the levels of osteogenic differentiation markers OCN, BMP2, BSP and OPN were markedly lower in the inflammation group than in the control group (Fig. 1F, G). Similarly, the immunofluorescence staining showed that lower positive immunofluorescence signals for RUNX2 and COL1A1 were detected in the inflammation group (Fig. 1H). The above results suggest that the in vitro inflammatory microenvironment dramatically reduces the osteogenic differentiation of PDLSCs.

CircBIRC6 depletion promotes osteogenic differentiation of PDLSCs in the inflammatory microenvironment, and vice versa

As shown in Fig. 2A, the expression level of circBIRC6 was significantly increased in the inflammatory microenvironment. In addition, circBIRC6 was overexpressed in periodontitis tissues compared to healthy periodontal tissues (Fig. 2B). The aberrant expression of circBIRC6 both in vitro and in vivo periodontitis models indicates that it might play an important role in regulating the biological functions of PDLSCs under the inflammatory conditions. The shcircBIRC6 significantly reduced the expression level of circBIRC6 in PDLSCs, whereas had little effect on the linear *BIRC6* expression (Fig. 2C). Then, the effects of circBIRC6 downregulation on the proliferation and osteogenic differentiation of PDLSCs under inflammatory conditions were evaluated. The flow cytometry and EdU assays showed that circBIRC6 depletion did not affect the proliferative capacity of PDLSCs (Fig. 2D, E). Interestingly, the ALP staining intensity and the mineralized nodule formation capacity were significantly higher in the shcircBIRC6 groups than in the control group (Fig. 2F, G). The expression levels of OCN, BMP2, BSP and OPN were found consistently upregulated at both mRNA and protein levels in the shcircBIRC6 groups compared to the control group (Fig. 2H, I). Similarly, higher RUNX2 and COL1A1 immunofluorescence staining were observed in the circBIRC6 depletion groups (Fig. 2J). These results support that circBIRC6 might act as a negative regulator of PDLSC osteogenic differentiation in the inflammatory microenvironment. Then, circBIRC6 was overexpressed to explore its effects on the osteogenic differentiation of PDLSCs (Fig. 3A). The ALP and ARS staining assays both demonstrated that circBIRC6 overexpression significantly decreased the osteogenic potential of PDLSCs under the inflammatory conditions (Fig. 3B, C). In





addition, as shown in Fig. 3D, E, and F, the levels of osteogenic differentiation markers were markedly decreased in the circBIRC6-overexpressing group compared to the control group.

CircBIRC6 directly binds to miR-543 and acts as a competitive endogenous RNA

It is well established that circRNAs exert their biological functions through the miRNA sponge mechanism. We first screened the potential miRNAs that might bind the circBIRC6 using the online tool (<https://circinteractome.nia.nih.gov/>). Then, only those that have been reported to be associated with osteogenic differentiation were chosen for further evaluation. Among all tested miRNAs,

miR-543 was the only miRNA that was copiously pulled down by circBIRC6 probe in PDLSCs (Fig. 4A). The RNA immunoprecipitation (RIP) assay showed that circBIRC6 was pulled down in dramatically higher quantity by the anti-AGO2 than anti-IgG. In addition, in the anti-AGO2 pull-down cells, the level of circBIRC6 was significantly higher in the miR-543 mimic group compared to NC mimic group (Fig. 4B). Based on the complementary sequences, the mutant circBIRC6 was constructed by changing the binding sites between circBIRC6 and miR-543. The luciferase reporter assay showed that miR-543 inhibitor promoted circBIRC6 wt reporter activity, while miR-543 mimic suppressed its activity. The luciferase activity of the circBIRC6 mut was not significantly altered





following the miR-543 overexpression or inhibition (Fig. 4C). These findings indicate that circBIRC6 directly binds to miR-543. The correlation between circBIRC6 and miR-543 was further evaluated in periodontitis tissues. The results showed that miR-543 expression was negatively associated with circBIRC6 expression in periodontitis tissues (Fig. 4D).


MiR-543 promotes osteogenic differentiation of PDLSCs under the inflammatory microenvironment

The effects of miR-543 on the osteogenic differentiation of PDLSCs under the inflammatory conditions were then evaluated. The results showed that the ALP staining intensity and the mineralized nodule formation capacity were markedly lower in the miR-543 inhibitor group than in the NC inhibitor group (Fig. 5A, B, and C). In addition, miR-543 inhibitor suppressed the expression of osteogenic differentiation markers in the inflammatory microenvironment (Fig. 5D, E, and F). As shown in Fig. 5G–L, overexpression of miR-543 promoted the

osteogenic differentiation of PDLSCs in the inflammatory conditions. The above results suggest that miR-543 positively regulates the osteogenic potential of PDLSCs in the inflammatory conditions.

CircBIRC6 regulates osteogenic differentiation of PDLSCs by sponging miR-543 in the inflammatory environment

The rescue experiments were performed to verify whether miR-543 was a functional downstream target of circBIRC6. The ALP and ARS staining assays showed that downregulation of circBIRC6 increased the ALP staining intensity and promoted mineralized nodule formation, and this effect was partially suppressed by miR-543 inhibitor (Fig. 6A, B). Similarly, the increased levels of osteogenic differentiation markers induced by shcircBIRC6 were reversed by miR-543 inhibitor (Fig. 6C, D, and E). We then further investigated whether miR-543 overexpression reversed the suppressive effects of circBIRC6 upregulation in osteogenic differentiation. As shown in Fig. 7A–E, the inhibitory effects of circBIRC6



overexpression on osteogenic differentiation of PDLSCs were antagonized by upregulation of miR-543. These results strongly support that miR-543 is functional downstream target of circBIRC6 for regulating osteogenic differentiation under the inflammatory conditions.

CircBIRC6/miR-543 axis regulates PDLSC osteogenic differentiation in the inflammatory conditions through PTEN/PI3K/AKT/mTOR signaling pathway

To further investigate the downstream genes of miR-543 in PDLSCs, we first screened the currently

(See figure on next page.)

Fig. 5 MiR-543 promoted the osteogenic differentiation of PDLSCs under the inflammatory microenvironment. **A** The expression of miR-543 in PDLSCs was analyzed by qRT-PCR after transfection with miR-543 inhibitor and NC inhibitor. **B–F** The osteogenic differentiation capability of PDLSCs transfected with miR-543 inhibitor in the inflammatory conditions was evaluated by ALP staining, ARS staining, qRT-PCR, WB and immunofluorescence staining. **G** The miR-543 level was detected with qRT-PCR after transfection with miR-543 mimic and NC mimic. **H–L** The effects of miR-543 mimic on the osteogenic differentiation of PDLSCs under the inflammatory microenvironment were evaluated by ALP staining, ARS staining, qRT-PCR, WB and immunofluorescence staining. Data are presented as the mean \pm SD (* $P < 0.05$, ** $P < 0.01$, *** $P < 0.001$). Scale bar: 20 μ m (**F** and **L**)




Fig. 5 (See legend on previous page.)





Fig. 6 MiR-543 inhibitor suppressed the osteogenic differentiation capacity of PDLSCs enhanced by circBIRC6 downregulation under the inflammatory microenvironment. **A** ALP staining of PDLSCs with indicated modifications. **B** ARS staining of mineralized nodules formed by PDLSCs with indicated treatments. **C–E** qRT-PCR, WB and immunofluorescence staining were performed to detect the changes in osteogenic differentiation makers in PDLSCs with indicated modifications. Scale bar: 20 μ m. Data are presented as the mean \pm SD (***) P < 0.001

available validated targets. Based on the results of miRWalk (<http://mirwalk.umm.uni-heidelberg.de/>) and miRTarBase (<http://miRTarBase.cuhk>), PTEN was selected as the potential downstream target of further investigation. The WB results showed that shcircBIRC6 suppressed the expression of PTEN, and PTEN level was increased when circBIRC6 was overexpressed (Fig. 8A). Not surprisingly, miR-543 mimic downregulated PTEN expression, and miR-543 inhibitor promoted its expression (Fig. 8B). To further verify the potential role of circBIRC6 in regulating PTEN expression, circBIRC6, circBIRC6 mut, miR-543 mimic and miR-543 inhibitor were transfected into PDLSCs individually or in combination. The results showed that only circBIRC6 and miR-543 inhibitor enhanced the reporter activity of PTEN (Fig. 8C), indicating that circBIRC6/miR-543 axis might be an upstream regulator of PTEN. As shown in Fig. 8D, E, and F,

overexpression of PTEN suppressed the circBIRC6 knockdown-mediated increased osteogenic potential of PDLSCs, and PTEN downregulation partially reversed the suppressive effects of circBIRC6 upregulation in osteogenic differentiation. PTEN is a well-known upstream regulator of PI3K/AKT/mTOR, which plays an essential role in maintaining the osteogenic differentiation capability of PDLSCs. Our WB results showed that shcircBIRC6 promoted the expression levels of p-PI3K, p-AKT and p-mTOR, and this effect was counteracted by PTEN overexpression. Similarly, circBIRC6 upregulation reduced the expression levels of p-PI3K, p-AKT and p-mTOR, and downregulation of PTEN rescued their expression. Figure 8G illustrates the potential molecular mechanisms for circBIRC6 in regulating PDLSC osteogenic differentiation under the inflammatory microenvironment.



Discussion

It is well established that the chronic inflammatory microenvironment attenuates the osteogenic differentiation capacity of MSCs and negatively affects bone repair and regeneration [17, 18]. Alveolar bone defect is one of the common characteristics of periodontitis, and thus, successful bone regeneration under the inflammatory conditions is the key challenge for conquering periodontitis. PDLSCs are the most important seed cells

for periodontal regeneration. Therefore, it is of great importance to investigate how to enhance the defective osteogenic differentiation capability of PDLSCs in the inflammatory microenvironment. In this study, our results have clearly demonstrated that circBIRC6 level is increased in PDLSCs exposed to inflammatory stimulus and in periodontitis tissues. Knockdown of circBIRC6 promotes the osteogenic differentiation of PDLSCs under the inflammatory conditions, and opposite findings are

(See figure on next page.)

Fig. 8 CircBIRC6/miR-543 axis regulated the osteogenic potential of PDLSCs via PTEN/PI3K/AKT/mTOR signaling pathway. **A, B** WB assay of PTEN expression level in PDLSCs with indicated modifications. **C** The relative luciferase activities of PTEN promoter in PDLSCs transfected with circBIRC6 wt, circBIRC6 mut, miR-543 mimic and miR-543 inhibitor alone or in combination. **D–F** WB assays of PTEN, osteogenic differentiation markers, p-PI3K, PI3K, p-AKT, AKT, p-mTOR and mTOR in PDLSCs with indicated treatments. **G** Schematic illustration of the circBIRC6/miR-543/PTEN/PI3K-AKT-mTOR axis in regulating the PDLSC osteogenic differentiation under the inflammatory microenvironment. Data are presented as the mean \pm SD (ns: not significant, * $P < 0.05$, ** $P < 0.01$, *** $P < 0.001$)




Fig. 8 (See legend on previous page.)

observed when circBIRC6 is overexpressed. Mechanistically, circBIRC6 regulates osteogenic differentiation of PDLSCs by sponging miR-543 in the inflammatory environment. More importantly, we have evidenced that circBIRC6/miR-543 axis modulates the mineralization capacity of PDLSCs through PTEN/PI3K/AKT/mTOR signaling pathway.

Recent evidence has demonstrated that circRNAs play an important role in regulating the osteogenic differentiation of MSCs. For instance, circSIPA1L1 was found to promote the osteogenic potential of stem cells from apical papilla by modulating the miR-204-5p/ALPL axis [19]. Similarly, circRNA CDR1as induced the osteogenic differentiation of PDLSCs by sponging miR-7, which further enhanced GDF5 expression and subsequently activated pSmad1/5/8 and p38 MAPK pathway [11]. However, whether circRNAs regulate the osteogenic differentiation of PDLSCs under the inflammatory microenvironment is poorly known. Aberrant expression of circBIRC6 has been reported to be involved in many biological processes such as stem cell pluripotency, myocardial infarction and tumorigenesis [12–14]. To the best of our knowledge, this is the first study showing the downregulation of circBIRC6 enhances the osteogenic differentiation capacity of PDLSCs exposed to the inflammatory conditions, which might provide a novel strategy for promoting bone regeneration in the harsh microenvironment.

Then, we demonstrated that miR-543 was a functional downstream target of circBIRC6 for regulating the osteogenic differentiation of PDLSCs in the inflammatory microenvironment. Consistent with our findings, downregulation of miR-543 suppressed the osteogenic potential of human dental pulp stem cells (DPSCs), and opposite findings were observed when miR-543 was overexpressed [20], indicating that the miR-543 was a positive regulator of DPSC osteogenic differentiation. Similarly, under the normal condition, the expression level of miR-543 was increased during the osteogenic differentiation of PDLSCs. In addition, miR-543 upregulation promoted the osteogenesis of PDLSCs, and vice versa [21]. We speculate that miR-543 is reduced in the inflammatory microenvironment due to the sponging effects of circBIRC6, leading to the defective osteogenic differentiation capability of PDLSCs. In addition to miR-543, whether circBIRC6 sponges other miRNAs for regulating osteogenic differentiation of PDLSCs needs further investigation.

Among the validated downstream targets of miR-543, PTEN is an important regulator for controlling MSC osteogenic differentiation. For instance, PTEN upregulation suppressed the BMP-9 mediated osteogenic potential in MSCs through inhibiting the expression

of Wnt10b [22]. Loss of PTEN in mouse osteoblasts promoted the osteogenic differentiation and reduced apoptosis *in vitro* and *in vivo*, and the enhancing effects were mediated by activating the AKT [23]. PTEN is well-known negative regulator of PI3K/AKT/mTOR signaling pathway, which is crucial for controlling multiple biological processes including MSC osteogenic differentiation [24]. Therefore, following downregulation of circBIRC6, the sponging effects of circBIRC6 on miR-543 are suppressed, leading to the release of miR-543 in the cytosol. The accumulation of miR-543 inhibits the expression of PTEN, which further results in the activation of PI3K/AKT/mTOR signaling pathway. The activated PI3K/AKT/mTOR pathway promotes the osteogenic differentiation of PDLSCs under the inflammatory conditions.

Conclusions

Under the inflammatory conditions, downregulation of circBIRC6 promotes the osteogenic differentiation of PDLSCs by modulating miR-543/ PTEN/PI3K/AKT/ mTOR signaling pathway. These findings have important implications for the design and implementation of circBIRC6 targeting strategies to treat bone loss in periodontitis.

Abbreviations

ALP: Alkaline phosphatase; ARS: Alizarin Red S; circBIRC6: Circular RNA BIRC6; hESCs: Human embryonic stem cells; mTOR: Mammalian target of rapamycin; MSCs: Mesenchymal stem cells; PDLSCs: Periodontal ligament stem cells; PI3K: Phosphatidylinositol 3-kinase; PTEN: Phosphatase and tensin homolog.

Supplementary Information

The online version contains supplementary material available at <https://doi.org/10.1186/s13287-022-03093-7>.

Additional file 1. Sequences of primers and oligos.

Acknowledgements

Not applicable.

Author contributions

LC and BG conceived the study, designed the experiments and supervised the project. XZ, LC and WS performed experiments and acquired and analyzed the data. LC and BG wrote the manuscript. All authors read and approved the final manuscript.

Funding

This work was supported by National Natural Science Foundation of China (81901006), Guangdong Basic and Applied Basic Research Foundation (2020A1515110051) and Scientific Research Talent Cultivation Project of Stomatological Hospital, Southern Medical University (RC202005).

Availability of data and materials

The datasets used and/or analyzed during the current study are available from the corresponding author on reasonable request.

Declarations

Ethics approval and consent to participate

This study was approved by the Ethics Committee of the Stomatological Hospital, Southern Medical University.

Consent for publication

All patients have provided written informed consent for the publication of any associated data and accompanying images.

Competing interests

The authors declare that they have no competing interests.

Author details

¹Stomatological Hospital, Southern Medical University, Guangzhou 510280, China. ²Department of Stomatology, The Third Affiliated Hospital, Sun Yat-Sen University, Guangzhou 510630, China. ³Department of Dentistry, The First Affiliated Hospital, Sun Yat-Sen University, Guangzhou 510080, China. ⁴School of Dentistry, University of California, Los Angeles, Los Angeles, CA 90095, USA.

Received: 28 January 2022 Accepted: 14 May 2022

Published online: 13 August 2022

References

- Huang Y, Zhao X, Cui L, Huang S. Metagenomic and metatranscriptomic insight into oral biofilms in periodontitis and related systemic diseases. *Front Microbiol.* 2021;12:728585.
- Kinane DF, Stathopoulou PG, Papapanou PN. Periodontal diseases. *Nat Rev Dis Prim.* 2017;3:17038.
- Zhao X, Liu Z, Shu D, Xiong Y, He M, Xu S, et al. Association of periodontitis with rheumatoid arthritis and the effect of non-surgical periodontal treatment on disease activity in patients with rheumatoid arthritis. *Med Sci Monit.* 2018;24:5802–10.
- Hu L, Liu Y, Wang S. Stem cell-based tooth and periodontal regeneration. *Oral Dis.* 2018;24:696–705.
- Bao L, Zhang X, Xu Y, Wang M, Song Y, Gu Y, et al. Dysfunction of MiR-148a-NRP1 functional axis suppresses osteogenic differentiation of periodontal ligament stem cells under inflammatory microenvironment. *Cell Reprogram.* 2019;21:314–22.
- Zhou X, Luan X, Chen Z, Francis M, Gopinathan G, Li W, et al. MicroRNA-138 inhibits periodontal progenitor differentiation under inflammatory conditions. *J Dent Res.* 2016;95:230–7.
- Kristensen LS, Andersen MS, Stagsted LVW, Ebbesen KK, Hansen TB, Kjems J. The biogenesis, biology and characterization of circular RNAs. *Nat Rev Genet.* 2019;20:675–91.
- Wang Y, Han D, Zhou T, Zhang J, Liu C, Cao F, et al. Melatonin ameliorates aortic valve calcification via the regulation of circular RNA CircRIC3/miR-204-5p/DPP4 signaling in valvular interstitial cells. *J Pineal Res.* 2020;69:e12666.
- Xie L, Chen J, Ren X, Zhang M, Thuaksuban N, Nuntanarant T, et al. Alteration of circRNA and lncRNA expression profile in exosomes derived from periodontal ligament stem cells undergoing osteogenic differentiation. *Arch Oral Biol.* 2021;121:104984.
- Ye Y, Ke Y, Liu L, Xiao T, Yu J. CircRNA FAT1 regulates osteoblastic differentiation of periodontal ligament stem cells via miR-4781-3p/SMAD5 pathway. *Stem Cells Int.* 2021;2021:5177488.
- Li X, Zheng Y, Zheng Y, Huang Y, Zhang Y, Jia L, et al. Circular RNA CDR1as regulates osteoblastic differentiation of periodontal ligament stem cells via the miR-7/GDF5/SMAD and p38 MAPK signaling pathway. *Stem Cell Res Ther.* 2018;9:232.
- Yu CY, Li TC, Wu YY, Yeh CH, Chiang W, Chuang CY, et al. The circular RNA circBIRC6 participates in the molecular circuitry controlling human pluripotency. *Nat Commun.* 2017;8:1149.
- Garikipati VNS, Verma SK, Cheng Z, Liang D, Truongcao MM, Cimini M, et al. Circular RNA CircFndc3b modulates cardiac repair after myocardial infarction via FUS/VEGF-A axis. *Nat Commun.* 2019;10:4317.
- Yang H, Zhao M, Zhao L, Li P, Duan Y, Li G. CircRNA BIRC6 promotes non-small cell lung cancer cell progression by sponging microRNA-145. *Cell Oncol (Dordr).* 2020;43:477–88.
- Yang G, Wang X, Liu B, Lu Z, Xu Z, Xiu P, et al. circ-BIRC6, a circular RNA, promotes hepatocellular carcinoma progression by targeting the miR-3918/Bcl2 axis. *Cell Cycle.* 2019;18:976–89.
- Cui L, Zhao X, Jin Z, Wang H, Yang SF, Hu S. Melatonin modulates metabolic remodeling in HNSCC by suppressing MTHFD1L-formate axis. *J Pineal Res.* 2021;71:e12767.
- Lacey DC, Simmons PJ, Graves SE, Hamilton JA. Proinflammatory cytokines inhibit osteogenic differentiation from stem cells: implications for bone repair during inflammation. *Osteoarthr Cartil.* 2009;17:735–42.
- Zhou J, Zhang Y, Li L, Fu H, Yang W, Yan F. Human beta-defensin 3-combined gold nanoparticles for enhancement of osteogenic differentiation of human periodontal ligament cells in inflammatory microenvironments. *Int J Nanomed.* 2018;13:555–67.
- Li Y, Bian M, Zhou Z, Wu X, Ge X, Xiao T, et al. Circular RNA SIPA1L1 regulates osteoblastic differentiation of stem cells from apical papilla via miR-204-5p/ALPL pathway. *Stem Cell Res Ther.* 2020;11:461.
- Zhao LD, Xu WC, Cui J, Liang YC, Cheng WQ, Xin BC, et al. Long non-coding RNA maternally expressed gene 3 inhibits osteogenic differentiation of human dental pulp stem cells via microRNA-543/smad ubiquitin regulatory factor 1/runx2-related transcription factor 2 axis. *Arch Oral Biol.* 2020;118:104838.
- Ge Y, Li J, Hao Y, Hu Y, Chen D, Wu B, et al. MicroRNA-543 functions as an osteogenesis promoter in human periodontal ligament-derived stem cells by inhibiting transducer of ERBB2, 2. *J Periodontal Res.* 2018;53:832–41.
- Li FS, Li PP, Li L, Deng Y, Hu Y, He BC. PTEN reduces BMP9-induced osteogenic differentiation through inhibiting Wnt10b in mesenchymal stem cells. *Front Cell Dev Biol.* 2020;8:608544.
- Liu X, Bruxvoort KJ, Zylstra CR, Liu J, Cichowski R, Faugere MC, et al. Lifelong accumulation of bone in mice lacking *Pten* in osteoblasts. *Proc Natl Acad Sci USA.* 2007;104:2259–64.
- Carracedo A, Pandolfi PP. The PTEN-PI3K pathway: of feedbacks and cross-talks. *Oncogene.* 2008;27:5527–41.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Ready to submit your research? Choose BMC and benefit from:

- fast, convenient online submission
- thorough peer review by experienced researchers in your field
- rapid publication on acceptance
- support for research data, including large and complex data types
- gold Open Access which fosters wider collaboration and increased citations
- maximum visibility for your research: over 100M website views per year

At BMC, research is always in progress.

Learn more biomedcentral.com/submissions

