

RESEARCH

Open Access



# On $(p, q)$ -analogue of two parametric Stancu-Beta operators

Mohammad Mursaleen<sup>1\*</sup>, Abdizhahan M Sarsenbi<sup>2,3</sup> and Taqseer Khan<sup>1</sup>

\*Correspondence:

mursaleenm@gmail.com

<sup>1</sup>Department of Mathematics,  
Aligarh Muslim University, Aligarh,  
202002, India

Full list of author information is  
available at the end of the article

## Abstract

Our purpose is to introduce a two-parametric  $(p, q)$ -analogue of the Stancu-Beta operators. We study approximating properties of these operators using the Korovkin approximation theorem and also study a direct theorem. We also obtain the Voronovskaya-type estimate for these operators. Furthermore, we study the weighted approximation results and pointwise estimates for these operators.

**MSC:** 40A30; 41A10; 41A25; 41A36

**Keywords:** Stancu-Beta operators; linear positive operators; Lipschitz function; modulus of continuity; rate of convergence; Voronovskaya-type theorem

## 1 Introduction

The  $q$ -calculus has attracted attention of many researchers because of its applications in various fields such as numerical analysis, computer-aided geometric design, differential equations, and so on. In the field of approximation theory, the application of  $q$ -calculus has been the area of many recent researches.

Lupaş [1] presented the first  $q$ -analogue of the classical Bernstein operators in 1987. He studied the approximation and shape-preserving properties of these operators. Another  $q$ -companion of the classical Bernstein polynomials is due to Phillips [2]. Inspired by this, several authors produced generalizations of well-known positive linear operators based on  $q$ -integers and studied them extensively. For instance, the approximation properties of the Kantorovich-type  $q$ -Bernstein operators [3],  $q$ -BBH operators [4],  $q$ -analogue of generalized Bernstein-Schurer operators [5], weighted statistical approximation by Kantorovich-type  $q$ -Szász-Mirakjan operators [6],  $q$ -Szász-Durrmeyer operators [7], operators constructed by means of  $q$ -Lagrange polynomials and  $A$ -statistical approximation [8], statistical approximation properties of modified  $q$ -Stancu-Beta operators [9], and  $q$ -Bernstein-Schurer-Kantorovich operators [10].

The  $q$ -calculus has led to the discovery of the  $(p, q)$ -calculus. Recently, Mursaleen et al. have used the  $(p, q)$ -calculus in approximation theory. They have applied it to construct a  $(p, q)$ -analogue of the classical Bernstein operators [11], a  $(p, q)$ -analogue of the Bernstein-Stancu operators [12], and a  $(p, q)$ -analogue of the Bernstein-Schurer operators [13] and have studied their approximation properties. Most recently,  $(p, q)$ -analogues of some other operators have been studied in [14–18], and [19].

We now give some basic notions of the  $(p, q)$ -calculus.

The  $(p, q)$ -integer is defined by

$$[n]_{p,q} = \frac{p^n - q^n}{p - q}, \quad n = 0, 1, 2, \dots, 0 < q < p \leq 1.$$

The  $(p, q)$ -companion of the binomial expansion is

$$(ax + by)_{p,q}^n = \sum_{k=0}^n \binom{n}{k}_{p,q} q^{\frac{k(k-1)}{2}} p^{\frac{(n-k)(n-k-1)}{2}} a^{n-k} b^k x^{n-k} y^k,$$

$$(x + y)_{p,q}^n = (x + y)(px + qy)(p^2x + q^2y) \cdots (p^{n-1}x + q^{n-1}y).$$

The  $(p, q)$ -analogues of the binomial coefficients are defined by

$$\binom{n}{k}_{p,q} = \frac{[n]_{p,q}!}{[k]_{p,q}! [n - k]_{p,q}!}.$$

The  $(p, q)$ -analogues of definite integrals of a function  $f$  are defined by

$$\int_0^a f(x) d_{p,q}x = (q - p)a \sum_{k=0}^{\infty} \frac{p^k}{q^{k+1}} f\left(\frac{p^k}{q^{k+1}}a\right) \quad \text{when } \left|\frac{p}{q}\right| < 1$$

and

$$\int_0^a f(x) d_{p,q}x = (p - q)a \sum_{k=0}^{\infty} \frac{q^k}{p^{k+1}} f\left(\frac{q^k}{p^{k+1}}a\right) \quad \text{when } \left|\frac{q}{p}\right| < 1.$$

For  $m, n \in \mathbb{N}$ , the  $(p, q)$ -gamma and the  $(p, q)$ -beta functions are defined by

$$\Gamma_{p,q}(n) = \int_0^{\infty} p^{\frac{n(n-1)}{2}} E_{p,q}(-qx) d_{p,q}x, \quad \Gamma_{p,q}(n + 1) = [n]_{p,q}!$$

and

$$B_{p,q}(m, n) = \int_0^{\infty} \frac{x^{m-1}}{(1 + x)^{m+n}} d_{p,q}x, \tag{1.1}$$

respectively. These two are related by

$$B_{p,q}(m, n) = q^{\frac{2-m(m-1)}{2}} p^{\frac{-m(m-1)}{2}} \frac{\Gamma_{p,q}(n)\Gamma_{p,q}(m)}{\Gamma_{p,q}(m + n)}. \tag{1.2}$$

For  $p = 1$ , all the concepts of the  $(p, q)$ -calculus reduce to those of  $q$ -calculus. The details on  $(p, q)$ -calculus can be found in [20–22].

Stancu [23] introduced the beta operators to approximate the Lebesgue-integrable functions on  $[0, \infty)$  as follows:

$$L_n(f, x) = \frac{1}{B(nx, n + 1)} \int_0^{\infty} \frac{t^{nx}}{(1 + t)^{nx+n+1}} f(t) dt.$$

The  $q$ -companion of the Stancu-Beta operators was given by Aral and Gupta [24] as follows:

$$L_n(f, x) = \frac{K(A, [n]_q x)}{B([n]_q x, [n]_q + 1)} \int_0^{\infty/A} \frac{u^{[n]_q x - 1}}{(1 + u)^{[n]_q x + [n]_q + 1}} f(q^{[n]_q x} u) d_q u.$$

Let  $0 < q < p < 1$ . Mursaleen et al. [25] constructed the  $(p, q)$ -Stancu-Beta operators as follows:

$$L_n^{p,q}(f, x) = \frac{1}{B_{p,q}([n]_{p,q} x, [n]_{p,q} + 1)} \int_0^\infty \frac{u^{[n]_{p,q} x - 1}}{(1 + u)^{[n]_{p,q} x + [n]_{p,q} + 1}} f(p^{[n]_{p,q} x} q^{[n]_{p,q} x} u) d_{p,q} u. \tag{1.3}$$

They investigated the approximating properties and estimated the rate of convergence of these operators. Motivated by this work, we introduce the following sequence of operators:

$$S_{n,p,q}^{\alpha,\beta}(f; x) = \frac{1}{B_{p,q}([n]_{p,q} x, [n]_{p,q} + 1)} \times \int_0^\infty \frac{u^{[n]_{p,q} x - 1}}{(1 + u)^{[n]_{p,q} x + [n]_{p,q} + 1}} f\left(\frac{[n]_{p,q} p^{[n]_{p,q} x} q^{[n]_{p,q} x} u + \alpha}{[n]_{p,q} + \beta}\right) d_{p,q} u, \tag{1.4}$$

where  $0 \leq \alpha \leq \beta$ . We call them two-parametric  $(p, q)$ -Stancu-Beta operators. For  $\alpha = 0 = \beta$ , the operators (1.4) coincide with the operators (1.3). So the latter is a generalization of the former.

### 2 Main results

We shall investigate approximation results for the operators (1.4). We calculate the moments of the operators  $S_{n,p,q}^{\alpha,\beta}(f; x)$  in the following lemma.

**Lemma 2.1** *Let  $S_{n,p,q}^{\alpha,\beta}(f; x)$  be given by (1.4). Then we have the following equalities:*

- (i)  $S_{n,p,q}^{\alpha,\beta}(1; x) = 1,$
- (ii)  $S_{n,p,q}^{\alpha,\beta}(t; x) = \frac{[n]_{p,q}}{([n]_{p,q} + \beta)} x + \frac{\alpha}{([n]_{p,q} + \beta)},$
- (iii)  $S_{n,p,q}^{\alpha,\beta}(t^2; x) = \frac{[n]_{p,q}^3}{pq([n]_{p,q} - 1)([n]_{p,q} + \beta)^2} x^2 + \frac{[n]_{p,q}}{([n]_{p,q} + \beta)^2} \left(\frac{[n]_{p,q}}{pq([n]_{p,q} - 1)} + 2\alpha\right) x + \frac{\alpha^2}{([n]_{p,q} + \beta)^2}.$

*Proof* Using (1.1), (i) is immediate. Further,

$$\begin{aligned} S_{n,p,q}^{\alpha,\beta}(t; x) &= \frac{1}{B_{p,q}([n]_{p,q} x, [n]_{p,q} + 1)} \\ &\times \int_0^\infty \frac{u^{[n]_{p,q} x - 1}}{(1 + u)^{[n]_{p,q} x + [n]_{p,q} + 1}} \left(\frac{[n]_{p,q} p^{[n]_{p,q} x} q^{[n]_{p,q} x} u + \alpha}{([n]_{p,q} + \beta)}\right) d_{p,q} u \\ &= \frac{[n]_{p,q}}{([n]_{p,q} + \beta)} \frac{p^{[n]_{p,q} x} q^{[n]_{p,q} x}}{B_{p,q}([n]_{p,q} x, [n]_{p,q} + 1)} \int_0^\infty \frac{u^{[n]_{p,q} x}}{(1 + u)^{[n]_{p,q} x + [n]_{p,q} + 1}} d_{p,q} u \\ &\quad + \frac{\alpha}{([n]_{p,q} + \beta)} \frac{1}{B_{p,q}([n]_{p,q} x, [n]_{p,q} + 1)} \int_0^\infty \frac{u^{[n]_{p,q} x - 1}}{(1 + u)^{[n]_{p,q} x + [n]_{p,q} + 1}} d_{p,q} u \\ &= \frac{[n]_{p,q}}{([n]_{p,q} + \beta)} L_n^{p,q}(t; x) + \frac{\alpha}{([n]_{p,q} + \beta)} L_n^{p,q}(1; x) \\ &= \frac{[n]_{p,q}}{([n]_{p,q} + \beta)} x + \frac{\alpha}{([n]_{p,q} + \beta)}, \end{aligned}$$

and (ii) is proved;

$$\begin{aligned}
 S_{n,p,q}^{\alpha,\beta}(t^2; x) &= \frac{1}{B_{p,q}([n]_{p,q}x, [n]_{p,q} + 1)} \\
 &\times \int_0^\infty \frac{u^{[n]_{p,q}x-1}}{(1+u)^{[n]_{p,q}x+[n]_{p,q}+1}} \left( \frac{[n]_{p,q}P^{[n]_{p,q}x}q^{[n]_{p,q}x}u + \alpha}{[n]_{p,q} + \beta} \right)^2 d_{p,q}u \\
 &= \frac{[n]_{p,q}^2}{([n]_{p,q} + \beta)^2} \frac{p^{2[n]_{p,q}x}q^{2[n]_{p,q}x}}{B_{p,q}([n]_{p,q}x, [n]_{p,q} + 1)} \int_0^\infty \frac{u^{[n]_{p,q}x+1}}{(1+u)^{[n]_{p,q}x+[n]_{p,q}+1}} d_{p,q}u \\
 &\quad + \frac{2\alpha}{[n]_{p,q}} ([n]_{p,q} + \beta)^2 \frac{q^{[n]_{p,q}x}}{B_{p,q}([n]_{p,q}x, [n]_{p,q} + 1)} \int_0^\infty \frac{u^{[n]_{p,q}x}}{(1+u)^{[n]_{p,q}x+[n]_{p,q}+1}} d_{p,q}u \\
 &\quad + \frac{\alpha^2}{([n]_{p,q} + \beta)^2} \frac{1}{B_{p,q}([n]_{p,q}x, [n]_{p,q} + 1)} \int_0^\infty \frac{u^{[n]_{p,q}x-1}}{(1+u)^{[n]_{p,q}x+[n]_{p,q}+1}} d_{p,q}u \\
 &= \frac{[n]_{p,q}^2}{([n]_{p,q} + \beta)^2} L_n^{p,q}(t^2; x) + \frac{2\alpha [n]_{p,q}}{([n]_{p,q} + \beta)^2} I_n^{p,q}(t; x) + \frac{\alpha^2}{([n]_{p,q} + \beta)^2} L_n^{p,q}(1; x) \\
 &= \frac{[n]_{p,q}^2}{([n]_{p,q} + \beta)^2} \left( \frac{[n]_{p,q}}{pq([n]_{p,q} - 1)} x^2 + \frac{1}{pq([n]_{p,q} - 1)} x \right) \\
 &\quad + \frac{2\alpha [n]_{p,q}}{([n]_{p,q} + \beta)^2} + \frac{\alpha^2}{([n]_{p,q} + \beta)^2} \\
 &= \frac{[n]_{p,q}^3}{pq([n]_{p,q} - 1)([n]_{p,q} + \beta)^2} x^2 + \frac{n}{([n]_{p,q} + \beta)^2} \left( \frac{[n]_{p,q}}{pq([n]_{p,q} - 1)} + 2\alpha \right) x \\
 &\quad + \frac{\alpha^2}{([n]_{p,q} + \beta)^2},
 \end{aligned}$$

which proves (iii).

Hence, the lemma is proved. □

We readily obtain the following lemma.

**Lemma 2.2** *Let  $p, q \in (0, 1)$ . Then, for  $x \in [0, \infty)$ , we have:*

- (i)  $S_{n,p,q}^{\alpha,\beta}((t-x); x) = \frac{\alpha - \beta x}{([n]_{p,q} + \beta)}$ ,
- (ii)  $S_{n,p,q}^{\alpha,\beta}((t-x)^2; x) \leq \left( \frac{[n]_{p,q}}{pq([n]_{p,q} - 1)} - \frac{([n]_{p,q} - \beta)}{([n]_{p,q} + \beta)} \right) x^2 + \frac{1}{pq([n]_{p,q} - 1)} x + \frac{\alpha^2}{([n]_{p,q} + \beta)^2} \leq \frac{2(1+\beta)^2 x^2 + x + \alpha^2}{pq([n]_{p,q} - 1)}$ .

*Proof* We have

$$\begin{aligned}
 S_{n,p,q}^{\alpha,\beta}((t-x); x) &= S_{n,p,q}^{\alpha,\beta}(t; x) - x S_{n,p,q}^{\alpha,\beta}(1; x) \\
 &= \frac{[n]_{p,q}}{([n]_{p,q} + \beta)} x + \frac{\alpha}{([n]_{p,q} + \beta)} - x \\
 &= \left( \frac{[n]_{p,q}}{([n]_{p,q} + \beta)} - 1 \right) x + \frac{\alpha}{([n]_{p,q} + \beta)} \\
 &= \frac{-\beta}{([n]_{p,q} + \beta)} x + \frac{\alpha}{([n]_{p,q} + \beta)} \\
 &= \frac{\alpha - \beta x}{([n]_{p,q} + \beta)},
 \end{aligned}$$

which proves (i). Now

$$\begin{aligned}
 & S_{n,p,q}^{\alpha,\beta}((t-x)^2; x) \\
 &= S_{n,p,q}^{\alpha,\beta}(t^2; x) + x^2 S_{n,p,q}^{\alpha,\beta}(1; x) - 2x S_{n,p,q}^{\alpha,\beta}(t; x) \\
 &= \frac{[n]_{p,q}^3}{pq([n]_{p,q}-1)([n]_{p,q}+\beta)^2} x^2 + \frac{[n]_{p,q}}{([n]_{p,q}+\beta)^2} \left( \frac{[n]_{p,q}}{pq([n]_{p,q}-1)} + 2\alpha \right) x \\
 &\quad + \frac{\alpha^2}{([n]_{p,q}+\beta)^2} - 2x \left( \frac{[n]_{p,q}}{([n]_{p,q}+\beta)} x + \frac{\alpha}{([n]_{p,q}+\beta)} \right) + x^2 \\
 &= \frac{[n]_{p,q}^3}{pq([n]_{p,q}-1)([n]_{p,q}+\beta)^2} - \frac{2[n]_{p,q}}{([n]_{p,q}+\beta)+1} x^2 + \frac{[n]_{p,q}^2}{pq([n]_{p,q}-1)([n]_{p,q}+\beta)^2} \\
 &\quad + \frac{2\alpha[n]_{p,q}}{([n]_{p,q}+\beta)^2} - \frac{2\alpha}{([n]_{p,q}+\beta)} x + \frac{\alpha^2}{([n]_{p,q}+\beta)^2} \\
 &\leq \left( \frac{[n]_{p,q}}{pq([n]_{p,q}-1)} - \frac{([n]_{p,q}-\beta)}{([n]_{p,q}+\beta)} \right) x^2 + \frac{1}{pq([n]_{p,q}-1)} x + \frac{\alpha^2}{([n]_{p,q}+\beta)^2} \\
 &= \frac{\{(p-q)[n]_{p,q}^3 + ([n]_{p,q} + pq[n]_{p,q} - pq)\beta^2 + (2\beta + pq)[n]_{p,q}^2\}x^2 + ([n]_{p,q} + \beta)^2x + pq([n]_{p,q} - 1)\alpha^2}{pq([n]_{p,q} - 1)([n]_{p,q} + \beta)^2} \\
 &= \frac{\{(p^n - q^n)[n]_{p,q}^2 + ([n]_{p,q} + pq[n]_{p,q} - pq)\beta^2 + (2\beta + pq)[n]_{p,q}^2\}x^2 + ([n]_{p,q} + \beta)^2x + pq([n]_{p,q} - 1)\alpha^2}{pq([n]_{p,q} - 1)([n]_{p,q} + \beta)^2} \\
 &\leq \frac{2(\beta^2 + \beta + 1)x^2 + x + \alpha^2}{pq([n]_{p,q} - 1)} \\
 &\leq \frac{2(\beta + 1)^2x^2 + x + \alpha^2}{pq([n]_{p,q} - 1)}
 \end{aligned}$$

which gives (ii). Hence, the lemma is proved. □

Next, we present a direct theorem for the operators  $S_{n,p,q}^{\alpha,\beta}(f; x)$ .

We denote By  $C_B[0, \infty)$ , the space of all real-valued continuous bounded functions  $f$  on the interval  $[0, \infty)$  endowed with the norm

$$\|f\| = \sup_{0 \leq x < \infty} |f(x)|.$$

Let  $\delta > 0$  and  $W^2 = \{h : h', h'' \in C(I), I = [0, \infty)\}$ , then the Peetre  $K$ -functional is defined by

$$K_2(f, \delta) = \inf_{h \in W^2} \{ \|f - h\| + \delta \|h''\| \}.$$

The second-order modulus of continuity  $\omega_2$  of  $f$  is defined as

$$\omega_2(f, \sqrt{\delta}) = \sup_{0 < p < \delta^{\frac{1}{2}}} \sup_{x \in I} |f(x + 2p) - 2f(x + p) + f(x)|.$$

By DeVore-Lorentz theorem (see [26], p.177, Theorem 2.4) there exists a constant  $C > 0$  such that

$$K_2(f, \delta) \leq C\omega_2(f, \sqrt{\delta}). \tag{2.1}$$

Also, by  $\omega(f, \delta)$  we denote the first-order modulus of continuity of  $f \in C(I)$  defined as

$$\omega(f, \delta) = \sup_{0 < p < \delta} \sup_{x \in I} |f(x + p) - f(x)|.$$

We shall use the notation  $v^2(x) = x + x^2$ .

**Theorem 2.3** *Suppose that  $f \in C_B[0, \infty)$  and  $0 < p, q < 1$ . Then for all  $x \in [0, \infty)$  and  $n \geq 2$ , there exists a constant  $C$  such that*

$$|S_{n,p,q}^{\alpha,\beta}(f; x) - f(x)| \leq C\omega_2\left(f, \frac{\delta_n(x)}{\sqrt{pq([n]_{p,q} - 1)}}\right) + \omega\left(f, \frac{\gamma_n(x)}{[n]_{p,q} + \beta}\right),$$

where

$$\delta_n^2(x) = v^2(x) + \frac{2pq\alpha^2}{([n]_{p,q} + \beta)}$$

and

$$\gamma_n^2(x) = (\alpha - \beta x)^2 + [n]_{p,q}([n]_{p,q} + \beta)x^2 + \alpha\beta x.$$

*Proof* Let us define the auxiliary operators

$$S_{n,p,q}^{*\alpha,\beta}(f; x) = S_{n,p,q}^{\alpha,\beta}(f; x) - f\left(\frac{[n]_{p,q}x + \alpha}{[n]_{p,q} + \beta}\right) + f(x). \tag{2.2}$$

By the Lemma 2.1 it is readily seen that these operators are linear and

$$S_{n,p,q}^{*\alpha,\beta}((t - x); x) = 0. \tag{2.3}$$

Suppose that  $g \in W^2$ . By the Taylor expansion we can write

$$g(t) = g(x) + g'(x)(t - x) + \int_x^t (t - u)g''(u) du, \quad t \in [0, \infty).$$

Operating by  $S_{n,p,q}^{*\alpha,\beta}(\cdot; x)$  on both sides of the above and using (2.3), we obtain:

$$\begin{aligned} S_{n,p,q}^{*\alpha,\beta}(g; x) &= g(x) + S_{n,p,q}^{*\alpha,\beta}\left(\int_x^t (t - u)g''(u) du; x\right), \\ S_{n,p,q}^{*\alpha,\beta}(g; x) - g(x) &= S_{n,p,q}^{*\alpha,\beta}\left(\int_x^t (t - u)g''(u) du; x\right), \\ |S_{n,p,q}^{*\alpha,\beta}(g; x) - g(x)| &= \left|S_{n,p,q}^{*\alpha,\beta}\left(\int_x^t (t - u)g''(u) du; x\right)\right|. \end{aligned}$$

Using (2.2) in the right-hand side, we get

$$\begin{aligned} |S_{n,p,q}^{*\alpha,\beta}(g; x) - g(x)| &= \left|S_{n,p,q}^{\alpha,\beta}\left(\int_x^t (t - u)g''(u) du; x\right) \right. \\ &\quad \left. - \int_x^{\frac{[n]_{p,q}x + \alpha}{[n]_{p,q} + \beta}} \left(\frac{[n]_{p,q}x + \alpha}{[n]_{p,q} + \beta} - u\right)g''(u) du\right|. \end{aligned}$$

So we obtain

$$\begin{aligned} & |S_{n,p,q}^{*\alpha,\beta}(g;x) - g(x)| \\ & \leq \left| S_{n,p,q}^{\alpha,\beta} \left( \int_x^t (t-u)g''(u) du; x \right) \right| + \left| \int_x^{\frac{[n]_{p,q}x+\alpha}{[n]_{p,q}+\beta}} \left( \frac{[n]_{p,q}x+\alpha}{[n]_{p,q}+\beta} - u \right) g''(u) du \right| \\ & \leq S_{n,p,q}^{\alpha,\beta} \left( \left| \int_x^t (t-u)g''(u) du \right|; x \right) + \int_x^{\frac{[n]_{p,q}x+\alpha}{[n]_{p,q}+\beta}} \left| \frac{[n]_{p,q}x+\alpha}{[n]_{p,q}+\beta} - u \right| |g''(u)| du. \end{aligned}$$

Using the linearity of the integral operator and the operator  $S_{n,p,q}^{\alpha,\beta}(\cdot;x)$  in the second and first parts of right-hand side, respectively, and using the fact that for all  $x \in [0, \infty)$ ,

$$|g(x)| \leq \|g\|,$$

we obtain

$$|S_{n,p,q}^{*\alpha,\beta}(g;x) - g(x)| \leq \|g''\| S_{n,p,q}^{\alpha,\beta}((t-x)^2;x) + \|g''\| \int_x^{\frac{[n]_{p,q}x+\alpha}{[n]_{p,q}+\beta}} \left| \frac{[n]_{p,q}x+\alpha}{[n]_{p,q}+\beta} - u \right| du. \tag{2.4}$$

In the first part, solving the integral  $\int_x^t |t-u| du$  and using the linearity of the operators  $S_{n,p,q}^{\alpha,\beta}(\cdot;x)$ , we readily see that

$$S_{n,p,q}^{\alpha,\beta} \left( \int_x^t |t-u| du \right) \leq S_{n,p,q}^{\alpha,\beta}((t-x)^2;x),$$

and after some calculations, for the second part of (2.4), we get

$$\begin{aligned} & \int_x^{\frac{[n]_{p,q}x+\alpha}{[n]_{p,q}+\beta}} \left| \frac{[n]_{p,q}x+\alpha}{[n]_{p,q}+\beta} - u \right| du \\ & \leq \frac{([n]_{p,q}x+\alpha)^2 - x([n]_{p,q}x+\alpha)([n]_{p,q}+\beta) + x^2([n]_{p,q}+\beta)^2}{([n]_{p,q}+\beta)^2} \\ & = \frac{(\alpha - \beta x)^2 + [n]_{p,q}x^2([n]_{p,q}+\beta) + \alpha\beta x}{([n]_{p,q}+\beta)^2} \\ & = \left( \frac{\alpha - \beta x}{[n]_{p,q}+\beta} \right)^2 + \frac{[n]_{p,q}}{[n]_{p,q}+\beta} x^2 + \frac{\alpha\beta}{([n]_{p,q}+\beta)^2} x. \end{aligned}$$

So by (2.4), we obtain

$$\begin{aligned} & |S_{n,p,q}^{*\alpha,\beta}(g;x) - g(x)| \\ & \leq \|g''\| \left( S_{n,p,q}^{\alpha,\beta}((t-x)^2;x) + \left( \frac{\alpha - \beta x}{[n]_{p,q}+\beta} \right)^2 + \frac{[n]_{p,q}}{[n]_{p,q}+\beta} x^2 + \frac{\alpha\beta}{([n]_{p,q}+\beta)^2} x \right). \tag{2.5} \end{aligned}$$

Using Lemma 2.2(ii), we obtain

$$\begin{aligned} & S_{n,p,q}^{\alpha,\beta}((t-x)^2;x) + \left( \frac{\alpha - \beta x}{[n]_{p,q}+\beta} \right)^2 + \frac{[n]_{p,q}}{([n]_{p,q}+\beta)} x^2 + \frac{\alpha\beta}{([n]_{p,q}+\beta)^2} x \\ & \leq \left( \frac{[n]_{p,q}}{pq([n]_{p,q}-1)} - \frac{([n]_{p,q}-\beta)}{([n]_{p,q}+\beta)} \right) x^2 + \frac{1}{pq([n]_{p,q}-1)} x + \frac{\alpha^2}{([n]_{p,q}+\beta)^2} \end{aligned}$$

$$\begin{aligned}
 & + \left( \frac{\alpha - \beta x}{[n]_{p,q} + \beta} \right)^2 + \frac{[n]_{p,q}}{([n]_{p,q} + \beta)} x^2 + \frac{\alpha\beta}{([n]_{p,q} + \beta)^2} x \\
 \leq & \frac{(p - q)[n]_{p,q}^3}{pq([n]_{p,q} + \beta)^2([n]_{p,q} - 1)} x^2 + \frac{[n]_{p,q}^2 + 4pq(1 - [n]_{p,q})\alpha\beta}{pq([n]_{p,q} + \beta)^2([n]_{p,q} - 1)} x + \frac{2\alpha^2}{([n]_{p,q} + \beta)^2} \\
 \leq & \frac{(p - q)[n]_{p,q}^3 x^2 + [n]_{p,q}^2 x + 2pq([n]_{p,q} - 1)\alpha^2}{pq([n]_{p,q} + \beta)^2([n]_{p,q} - 1)} \\
 = & \frac{(p^n - q^n)[n]_{p,q}^2 x^2 + [n]_{p,q}^2 x + 2pq([n]_{p,q} - 1)\alpha^2}{pq([n]_{p,q} + \beta)^2([n]_{p,q} - 1)} \\
 \leq & \frac{[n]_{p,q}^2 x^2 + [n]_{p,q}^2 x + 2pq[n]_{p,q}\alpha^2}{pq([n]_{p,q} + \beta)^2([n]_{p,q} - 1)} \\
 \leq & \frac{[n]_{p,q}(1 + x) + 2pq\alpha^2}{pq([n]_{p,q} + \beta)^2([n]_{p,q} - 1)} \\
 \leq & \frac{1}{pq([n]_{p,q} - 1)} \left( v^2(x) + \frac{2pq\alpha^2}{([n]_{p,q} + \beta)} \right) \\
 = & \frac{\delta_n^2(x)}{pq([n]_{p,q} - 1)},
 \end{aligned}$$

where

$$\delta_n^2(x) = v^2(x) + \frac{2pq\alpha^2}{([n]_{p,q} + \beta)}.$$

Therefore, by (2.5) we get

$$|S_{n,p,q}^{*\alpha,\beta}(g; x) - g(x)| \leq \frac{\delta_n^2(x)}{pq([n]_{p,q} - 1)} \|g''\|. \tag{2.6}$$

On the other hand, by (2.2) we have

$$|S_{n,p,q}^{*\alpha,\beta}(f; x)| \leq |S_{n,p,q}^{\alpha,\beta}(f; x)| + 2\|f\| \leq 3\|f\|. \tag{2.7}$$

By (2.2), (2.6), and (2.7), we obtain:

$$\begin{aligned}
 |S_{n,p,q}^{\alpha,\beta}(f; x) - f(x)| & \leq |S_{n,p,q}^{*\alpha,\beta}(f - g; x) - (f - g)(x)| + |S_{n,p,q}^{*\alpha,\beta}(g; x) - g(x)| \\
 & + \left| f \left( \frac{[n]_{p,q}x + \alpha}{[n]_{p,q} + \beta} \right) - f(x) \right| \\
 & \leq 4\|f - g\| + \frac{\delta_n^2(x)}{pq([n]_{p,q} - 1)} \|g''\| \\
 & + \omega \left( f, \frac{\sqrt{(\alpha - \beta x)^2 + [n]_{p,q}([n]_{p,q} + \beta)x^2 + \alpha\beta x}}{[n]_{p,q} + \beta} \right) \\
 & = 4\|f - g\| + \frac{\delta_n^2(x)}{pq([n]_{p,q} - 1)} \|g''\| + \omega \left( f, \frac{\gamma_n(x)}{[n]_{p,q} + \beta} \right), \tag{2.8}
 \end{aligned}$$

where

$$\gamma_n^2(x) = (\alpha - \beta x)^2 + [n]_{p,q}([n]_{p,q} + \beta)x^2 + \alpha\beta x.$$



Taking the infimum over all  $g \in W^2$  on the right-hand side of (2.8), we obtain

$$|S_{n,p,q}^{\alpha,\beta}(f; x) - f(x)| \leq CK_2 \left( f, \frac{\delta_n^2(x)}{pq([n]_{p,q} - 1)} \right) + \omega \left( f, \frac{(\gamma_n)x}{[n]_{p,q} + \beta} \right).$$

Using relation (2.1), for  $p, q \in (0, 1)$ , we get

$$|S_{n,p,q}^{\alpha,\beta}(f; x) - f(x)| \leq C\omega_2 \left( f, \frac{\delta_n(x)}{\sqrt{pq([n]_{p,q} - 1)}} \right) + \omega \left( f, \frac{\gamma_n(x)}{[n]_{p,q} + \beta} \right),$$

and this completes the proof. □

### 3 Rate of approximation

Let  $B_{x^2}[0, \infty)$  denote the set of all functions  $f$  such that  $f(x) \leq M_f(1 + x^2)$ , where  $M_f$  is a constant depending on  $f$ . By  $C_{x^2}[0, \infty)$  we denote the subspace of all continuous functions in the space  $B_{x^2}[0, \infty)$ . Also, we denote by  $C_{x^2}^*[0, \infty)$ , the subspace of all functions  $f \in C_{x^2}[0, \infty)$  for which  $\lim_{x \rightarrow \infty} \frac{f(x)}{1+x^2}$  is finite with

$$\|f\| = \sup_{x \in [0, \infty)} \frac{|f(x)|}{1 + x^2}.$$

For  $a > 0$ , the modulus of continuity of  $f$  over  $[0, a]$  is defined by

$$\omega_a(f, \delta) = \sup_{|t-x| \leq \delta} \sup_{0 \leq x, t \leq a} |f(t) - f(x)|.$$

We have the following proposition.

#### Proposition 3.1

- (i) For  $f \in C_{x^2}[0, \infty)$ , the modulus of continuity  $\omega_a(f, \delta)$ ,  $a > 0$ , approaches to zero.
- (ii) For every  $\delta > 0$ , we have

$$|f(y) - f(x)| \leq \left( 1 + \frac{|y - x|}{\delta} \right) \omega_a(f, \delta)$$

and

$$|f(y) - f(x)| \leq \left( 1 + \frac{(y - x)^2}{\delta^2} \right) \omega_a(f, \delta).$$

In the following theorem, we estimate the rate of convergence of the operators  $S_{n,p,q}^{\alpha,\beta}(f; x)$ .

**Theorem 3.2** Let  $f \in C_{x^2}[0, \infty)$ ,  $p, q \in (0, 1)$ , and let  $\omega_{a+1}(f, \delta)$  be the modulus of continuity on the interval  $[0, 1 + a] \subseteq [0, \infty)$ ,  $a > 0$ . Then, for  $n \geq 2$ , we have

$$\begin{aligned} \|S_{n,p,q}^{\alpha,\beta}(f) - f\|_{C[0,a]} &\leq \frac{4M_f(1 + a^2)(2(1 + \beta)^2 a^2 + a + \alpha^2)}{pq([n]_{p,q} - 1)} \\ &\quad + 2\omega_{1+a} \left( f, \left( \frac{2(1 + \beta)^2 a^2 + a + \alpha^2}{pq([n]_{p,q} - 1)} \right)^{\frac{1}{2}} \right). \end{aligned}$$

*Proof* Let  $x \in [0, a]$  and  $t > a + 1$ . Since  $1 + x < t$ , we have

$$\begin{aligned} |f(t) - f(x)| &\leq M_f(x^2 + t^2 + 2) \leq M_f(2 + 3x^2 + 2(t - x)^2) \\ &\leq M_f(4 + 3x^2)(t - x)^2 \leq 4M_f(1 + a^2)(t - x)^2. \end{aligned} \tag{3.1}$$

For  $\delta > 0, x \in [0, a], t - 1 \leq a$ , by Proposition 3.1 we obtain

$$|f(t) - f(x)| \leq \omega_{1+a}(f, |t - x|) \leq \omega_{1+a}(f, \delta) \left(1 + \frac{1}{\delta} |t - x|\right). \tag{3.2}$$

By (3.1) and (3.2), for  $x \in [0, a]$  and nonnegative  $t$ , we can write

$$|f(t) - f(x)| \leq 4M_f(1 + a^2)(t - x)^2 \omega_{1+a}(f, \delta) \left(1 + \frac{1}{\delta} |t - x|\right). \tag{3.3}$$

Therefore,

$$\begin{aligned} |S_{n,p,q}^{\alpha,\beta}(f; x) - f(x)| &\leq S_{n,p,q}^{\alpha,\beta}(|f(t) - f(x)|; x) \\ &\leq 4M_f(1 + a^2) S_{n,p,q}^{\alpha,\beta}((t - x)^2; x) + \omega_{1+a}(f, \delta) \left(1 + \frac{1}{\delta} (S_{n,p,q}^{\alpha,\beta}((t - x)^2; x))^{\frac{1}{2}}\right). \end{aligned}$$

Hence, using the Lemma 2.2(ii) and the Schwarz inequality, for every  $p, q \in (0, 1)$  and  $x \in [0, a]$ , we obtain

$$\begin{aligned} |S_{n,p,q}^{\alpha,\beta}(f; x) - f(x)| &\leq 4M_f(1 + a^2) \left(\frac{2(1 + \beta)^2 x^2 + x + \alpha^2}{pq([n]_{p,q} - 1)}\right) \\ &\quad + \omega_{1+a}(f, \delta) \left(1 + \frac{1}{\delta} \left(\frac{2(1 + \beta)^2 a^2 + a + \alpha^2}{pq([n]_{p,q} - 1)}\right)^{\frac{1}{2}}\right) \\ &\leq \frac{4M_f(1 + a^2)(2(1 + \beta)^2 a^2 + a + \alpha^2)}{pq([n]_{p,q} - 1)} \\ &\quad + \omega_{1+a} \left(1 + \frac{1}{\delta} \left(\frac{2(1 + \beta)^2 a^2 + a + \alpha^2}{pq([n]_{p,q} - 1)}\right)^{\frac{1}{2}}\right). \end{aligned}$$

By choosing  $\delta^2 = \frac{2(1+\beta)^2 a^2 + a + \alpha^2}{pq([n]_{p,q} - 1)}$  we get the required result. □

#### 4 Weighted approximation

This section is devoted to the study of weighted approximation theorems for the operators (2.2).

**Theorem 4.1** *Suppose that  $p = p_n$  and  $q = q_n$  are two sequences satisfying  $0 < p_n, q_n < 1$  and suppose that  $p_n \rightarrow 1$  and  $q_n \rightarrow 1$  as  $n \rightarrow \infty$ . Then, for each  $f \in C_{x^2}^*[0, \infty)$ , we have*

$$\lim_{n \rightarrow \infty} \|S_{n,p_n,q_n}^{\alpha,\beta}(f) - f\|_{x^2} = 0.$$

*Proof* By the theorem in [27] it suffices to prove that

$$\lim_{n \rightarrow \infty} \|S_{n,p_n,q_n}^{\alpha,\beta}(t^i) - x^i\|_{x^2} = 0 \quad \text{for } i = 0, 1, 2. \tag{4.1}$$

By Lemma 2.1(i)-(ii), the conditions of (4.1) are easily verified for  $i = 0$  and 1. For  $i = 2$ , we can write

$$\begin{aligned} & \|S_{n,p_n,q_n}^{\alpha,\beta}(t^2) - x^2\|_{x^2} \\ &= \sup_{x \in [0,\infty)} \frac{|S_{n,p_n,q_n}^{\alpha,\beta}(t^2) - x^2|}{1 + x^2} \\ &\leq \left( \frac{[n]_{p_n,q_n}^3}{p_n q_n ([n]_{p_n,q_n} - 1) ([n]_{p_n,q_n} + \beta)^2} - 1 \right) \sup_{x \in [0,\infty)} \frac{x^2}{1 + x^2} \\ &\quad + \frac{[n]_{p_n,q_n}^2 + 2p_n q_n [n]_{p_n,q_n} ([n]_{p_n,q_n} - 1)\alpha}{p_n q_n ([n]_{p_n,q_n} - 1) ([n]_{p_n,q_n} + \beta)^2} \sup_{x \in [0,\infty)} \frac{x}{1 + x^2} + \frac{\alpha^2}{([n]_{p_n,q_n} + \beta)^2} \\ &\leq \frac{(p_n^n - q_n^n) [n]_{p_n,q_n}^2 - p_n q_n (2\beta - 1) [n]_{p_n,q_n}^2 - q_n \beta (\beta - 1) [n]_{p_n,q_n} + q_n \beta^2}{p_n q_n ([n]_{p_n,q_n} - 1) ([n]_{p_n,q_n} + \beta)^2} \\ &\quad + \left( \frac{[n]_{p_n,q_n}^2 + 2p_n q_n [n]_{p_n,q_n} ([n]_{p_n,q_n} - 1)\alpha}{p_n q_n ([n]_{p_n,q_n} - 1) ([n]_{p_n,q_n} + \beta)^2} \right) + \frac{\alpha^2}{([n]_{p_n,q_n} + \beta)^2}, \end{aligned}$$

which implies that

$$\lim_{n \rightarrow \infty} \|S_{n,p_n,q_n}^{\alpha,\beta}(t^2, x) - x^2\|_{x^2} = 0.$$

This completes the proof of the theorem. □

**Theorem 4.2** *Let  $p = (p_n)$  and  $q = (q_n)$  be two sequences such that  $0 < p_n, q_n < 1$ , and let  $p_n \rightarrow 1$  and  $q_n \rightarrow 1$  as  $n \rightarrow \infty$ . Then, for each  $f \in C_{x^2}[0, \infty)$  and all  $\alpha > 0$ , we have*

$$\lim_{n \rightarrow \infty} \sup_{x \in [0,\infty)} \frac{|S_{n,p_n,q_n}^{\alpha,\beta}(f; x) - f(x)|}{(1 + x^2)^{1+\alpha^2}} = 0.$$

*Proof* For  $x_0 > 0$  fixed, we have:

$$\begin{aligned} \sup_{x \in [0,\infty)} \frac{|S_{n,p_n,q_n}^{\alpha,\beta}(f; x) - f(x)|}{(1 + x^2)^{1+\alpha^2}} &= \sup_{x \leq x_0} \frac{|S_{n,p_n,q_n}^{\alpha,\beta}(f; x) - f(x)|}{(1 + x^2)^{1+\alpha^2}} + \sup_{x \geq x_0} \frac{|S_{n,p_n,q_n}^{\alpha,\beta}(f; x) - f(x)|}{(1 + x^2)^{1+\alpha^2}} \\ &\leq \|S_{n,p_n,q_n}^{\alpha,\beta}(f) - f\|_{C[0,x_0]} + \|f\|_{x^2} \sup_{x \geq x_0} \frac{|S_{n,p_n,q_n}^{\alpha,\beta}(1 + t^2; x)|}{(1 + x^2)^{1+\alpha^2}} \\ &\quad + \sup_{x \geq x_0} \frac{|f(x)|}{(1 + x^2)^{1+\alpha^2}}. \end{aligned}$$

The first term of this inequality goes to zero by Theorem 3.2. Also, for any fixed  $x_0 > 0$ , it is readily seen from Lemma 2.1 that

$$\sup_{x \geq x_0} \frac{|S_{n,p_n,q_n}^{\alpha,\beta}(1 + t^2; x)|}{(1 + x^2)^{1+\alpha^2}}$$

approaches zero as  $n \rightarrow \infty$ . If we choose  $x_0 > 0$  large enough so that the last part of the last inequality is arbitrarily small, then our theorem is proved.  $\square$

### 5 Voronovskaya-type theorem

This section presents the Voronovskaya-type theorem for the operators  $S_{n,p,q}^{\alpha,\beta}(f;x)$ . We need the following lemma.

**Lemma 5.1** *Suppose that  $p_n, q_n \in (0, 1)$  are such that  $p_n^n \rightarrow a, q_n^n \rightarrow b$  ( $0 \leq a, b < 1$ ) as  $n \rightarrow \infty$ . Then, for every  $x \in [0, \infty)$ , simple computations yield*

$$\begin{aligned} \lim_{n \rightarrow \infty} [n]_{p_n, q_n} S_{n,p_n, q_n}^{\alpha, \beta}((t-x); x) &= \alpha - \beta x, \\ \lim_{n \rightarrow \infty} [n]_{p_n, q_n} S_{n,p_n, q_n}^{\alpha, \beta}((t-x)^2; x) &= (1-a)(1-b)x^2 + x. \end{aligned}$$

**Theorem 5.2** *Assume that  $p_n, q_n \in (0, 1)$  are such that  $p_n^n \rightarrow a, q_n^n \rightarrow b$  ( $0 \leq a, b < 1$ ) as  $n \rightarrow \infty$ . Then, for  $f \in C_{x^2}^*[0, \infty)$  such that  $f', f'' \in C_{x^2}^*[0, \infty)$ , we have*

$$\lim_{n \rightarrow \infty} [n]_{p_n, q_n} (S_{n,p_n, q_n}^{\alpha, \beta}(f; x) - f(x)) = (\alpha - \beta x)f'(x) + \frac{(1-a)(1-b)x^2 + x}{2} f''(x)$$

uniformly on  $[0, A]$  for any  $A > 0$ .

*Proof* Let  $f, f', f'' \in C_{x^2}^*[0, \infty)$  and  $x \in [0, \infty)$ . By the Taylor formula we can write

$$f(t) = f(x) + (t-x)f'(x) + \frac{1}{2}(t-x)^2 f''(x) + r(t;x)(t-x)^2, \tag{5.1}$$

where  $r(t;x)$  is the remainder term,  $r(\cdot;x) \in C_{x^2}^*[0, \infty)$ , and  $\lim_{t \rightarrow x} r(t;x) = 0$ . Operating by  $S_{n,p_n, q_n}^{\alpha, \beta}$  on both sides of (5.1), we get

$$\begin{aligned} [n]_{p_n, q_n} (S_{n,p_n, q_n}^{\alpha, \beta}(f; x) - f(x)) &= [n]_{p_n, q_n} S_{n,p_n, q_n}^{\alpha, \beta}((t-x); x) f'(x) + \frac{1}{2} [n]_{p_n, q_n} S_{n,p_n, q_n}^{\alpha, \beta}((t-x)^2; x) f''(x) \\ &\quad + [n]_{p_n, q_n} S_{n,p_n, q_n}^{\alpha, \beta}(r(\cdot;x)(\cdot-x)^2; x). \end{aligned}$$

It follows from the Cauchy-Schwarz inequality that

$$S_{n,p_n, q_n}^{\alpha, \beta}(r(\cdot;x)(\cdot-x)^2; x) \leq \sqrt{S_{n,p_n, q_n}^{\alpha, \beta}(r^2(\cdot;x); x)} \sqrt{S_{n,p_n, q_n}^{\alpha, \beta}(r((\cdot-x)^4; x))}. \tag{5.2}$$

Note that  $r^2(x;x) = 0$  and  $r^2(\cdot;x) \in C_{x^2}^*[0, \infty)$ . Therefore, it follows that

$$\lim_{n \rightarrow \infty} S_{n,p_n, q_n}^{\alpha, \beta}(r^2(\cdot;x); x) = r^2(x;x) = 0 \tag{5.3}$$

uniformly over  $[0, A]$ .

By Lemma 5.1 and equations (5.2) and (5.3), we obtain

$$\lim_{n \rightarrow \infty} [n]_{p_n, q_n} S_{n,p_n, q_n}^{\alpha, \beta}(r(\cdot;x)(\cdot-x)^2; x) = 0.$$

Thus, we obtain

$$\begin{aligned} & \lim_{n \rightarrow \infty} [n]_{p_n, q_n} (S_{n, p_n, q_n}^{\alpha, \beta}(f; x) - f(x)) \\ &= \lim_{n \rightarrow \infty} \left( [n]_{p_n, q_n} S_{n, p_n, q_n}^{\alpha, \beta}((t-x); x) f'(x) + \frac{1}{2} [n]_{p_n, q_n} S_{n, p_n, q_n}^{\alpha, \beta}((t-x)^2; x) f''(x) \right. \\ & \quad \left. + [n]_{p_n, q_n} S_{n, p_n, q_n}^{\alpha, \beta}(r(\cdot; x)(\cdot-x)^2; x) \right) \\ &= (\alpha - \beta x) f'(x) + \frac{(1-a)(1-b)x^2 + x}{2} f''(x). \end{aligned} \quad \square$$

### 6 Pointwise estimates

In this section, we study pointwise estimates of rate of convergence of the operators  $S_{n, p, q}^{\alpha, \beta}(f; x)$ .

Let  $0 < \nu \leq 1$  and  $E \subset [0, \infty)$ . We say that a function  $f \in C[0, \infty)$  belongs to  $Lip(\nu)$  if

$$|f(t) - f(x)| \leq M_f |t - x|^\nu, \quad t \in [0, \infty), x \in E, \tag{6.1}$$

where  $M_f$  is a constant depending on  $\alpha$  and  $f$  only.

We have the following theorem.

**Theorem 6.1** *Let  $\nu \in (0, 1], f \in Lip(\nu)$ , and  $E \subset [0, \infty)$ . Then, for  $x \in [0, \infty)$ ,*

$$\begin{aligned} & \|S_{n, p, q}^{\alpha, \beta}(f; x) - f(x)\| \\ & \leq M_f \left\{ \left( \left( \frac{[n]_{p, q}}{pq([n]_{p, q} - 1)} - \frac{([n]_{p, q} - \beta)}{([n]_{p, q} + \beta)} \right) x^2 + \frac{1}{pq([n]_{p, q} - 1)} x + \frac{\alpha^2}{([n]_{p, q} + \beta)^2} \right)^{\frac{\nu}{2}} \right. \\ & \quad \left. + 2(d(x, E))^\nu \right\}, \end{aligned}$$

where  $d(x, E)$  denotes the distance of the point  $x$  from the set  $E$ , defined by

$$d(x, E) = \inf\{|x - y| : y \in E\}.$$

*Proof* Taking  $y \in \bar{E}$ , we can write

$$|f(t) - f(x)| \leq |f(t) - f(y)| + |f(y) - f(x)|, \quad x \in [0, \infty).$$

By (6.1) we have

$$\begin{aligned} |S_{n, p, q}^{\alpha, \beta}(f; x) - f(x)| &= |S_{n, p, q}^{\alpha, \beta}(f; x) - S_{n, p, q}^{\alpha, \beta}(f(x); x)| \\ &\leq S_{n, p, q}^{\alpha, \beta}(|f(t) - f(x)|; x) \\ &\leq S_{n, p, q}^{\alpha, \beta}(|f(t) - f(y)|; x) + S_{n, p, q}^{\alpha, \beta}(|f(y) - f(x)|; x) \\ &\leq S_{n, p, q}^{\alpha, \beta}(|f(t) - f(y)|; x) + |f(x) - f(y)| \\ &\leq M_f S_{n, p, q}^{\alpha, \beta}(|t - y|^\nu; x) + |x - y|^\nu \\ &\leq M_f S_{n, p, q}^{\alpha, \beta}(|t - x|^\nu + |x - y|^\nu; x) + |x - y|^\nu \\ &\leq M_f S_{n, p, q}^{\alpha, \beta}(|t - x|^\nu; x) + 2|x - y|^\nu. \end{aligned}$$

Using the Hölder inequality with  $p = \frac{2}{\nu}, q = \frac{2}{2-\nu}$ , we obtain

$$\begin{aligned} & |S_{n,p,q}^{\alpha,\beta}(f; x) - f(x)| \\ & \leq M_f \left\{ (S_{n,p,q}^{\alpha,\beta}(|t - x|^{p\nu}; x))^{\frac{1}{p}} (S_{n,p,q}^{\alpha,\beta}(1^q; x))^{\frac{1}{q}} + 2(d(x, E))^{\nu} \right\} \\ & = M_f \left\{ (S_{n,p,q}^{\alpha,\beta}(|t - x|^2; x))^{\frac{\nu}{2}} + 2(d(x, E))^{\nu} \right\} \\ & = \left\{ \left( \left( \frac{[n]_{p,q}}{pq([n]_{p,q} - 1)} - \frac{([n]_{p,q} - \beta)}{([n]_{p,q} + \beta)} \right) x^2 + \frac{1}{pq([n]_{p,q} - 1)} x + \frac{\alpha^2}{([n]_{p,q} + \beta)^2} \right)^{\frac{\nu}{2}} \right. \\ & \quad \left. + 2(d(x, E))^{\nu} \right\}, \end{aligned}$$

and the theorem is proved. □

We now present a theorem regarding a local direct estimate for the operators  $S_{n,p,q}^{\alpha,\beta}(f; x)$  in terms of the Lipschitz-type maximal function of order  $\nu$  as introduced by Lenze [28]. It is defined by

$$\tilde{\omega}_{\nu}(f; x) = \sup_{y \neq x, y \in [0, \infty)} \frac{|f(y) - f(x)|}{|y - x|^{\nu}}, \quad x \in [0, \infty), \nu \in (0, 1]. \tag{6.2}$$

**Theorem 6.2** *Let  $\nu \in (0, 1]$  and  $f \in C[0, \infty)$ . Then, for each  $x \in [0, \infty)$ , we have*

$$\begin{aligned} & |S_{n,p,q}^{\alpha,\beta}(f; x) - f(x)| \\ & \leq \tilde{\omega}_{\nu}(f; x) \left\{ \left( \frac{[n]_{p,q}}{pq([n]_{p,q} - 1)} - \frac{([n]_{p,q} - \beta)}{([n]_{p,q} + \beta)} \right) x^2 + \frac{1}{pq([n]_{p,q} - 1)} x + \frac{\alpha^2}{([n]_{p,q} + \beta)^2} \right\}^{\frac{\nu}{2}}. \end{aligned}$$

*Proof* By (6.2) we can write

$$|f(t) - f(x)| \leq \tilde{\omega}_{\nu}(f; x) |t - x|^{\nu}$$

and

$$|S_{n,p,q}^{\alpha,\beta}(f; x) - f(x)| \leq S_{n,p,q}^{\alpha,\beta}(|f(t) - f(x)|; x) \leq \tilde{\omega}_{\nu}(f; x) S_{n,p,q}^{\alpha,\beta}(|t - x|^{\nu}; x).$$

Using the Lemma 2.2 and applying the Hölder inequality with  $p = \frac{2}{\nu}, q = \frac{2}{2-\nu}$ , we obtain

$$|S_{n,p,q}^{\alpha,\beta}(f; x) - f(x)| \leq \tilde{\omega}_{\nu}(f; x) S_{n,p,q}^{\alpha,\beta}(|t - x|^{\nu}; x),$$

which proves the theorem. □

**Remark** The further properties of the operators such as convergence properties via summability methods (see, e.g., [29–31]) can be studied.

**7 Conclusions**

In this paper, we have introduced a two-parametric  $(p, q)$ -analogue of the Stancu-Beta operators and studied some approximating properties of these operators. We also obtained the Voronovskaya-type estimate and the weighted approximation results for these operators. Furthermore, we obtained a pointwise estimate for these operators.

**Competing interests**

The authors declare that they have no competing interests.

**Authors' contributions**

All authors of the manuscript have read and agreed to its content and are accountable for all aspects of the accuracy and integrity of the manuscript.

**Author details**

<sup>1</sup>Department of Mathematics, Aligarh Muslim University, Aligarh, 202002, India. <sup>2</sup>Department of Mathematical Methods and Modeling, M. Auezov South-Kazakhstan State University, Tauke khan, 5, Shymkent, Kazakhstan. <sup>3</sup>Institute of Mathematics and Mathematical Modeling, Pushkin 125, Almaty, Kazakhstan.

**Acknowledgements**

Research of the second author is supported by the Grants Nos. 5414/GF4, 0971/GF4 of the Committee of Science of Ministry of Education and Science of the Republic of Kazakhstan.

Received: 21 April 2016 Accepted: 12 July 2016 Published online: 02 August 2016

**References**

- Lupaş, A: A  $q$ -analogue of the Bernstein operator. In: Seminar on Numerical and Statistical Calculus, vol. 87, pp. 85-92. Univ. 'Babeş-Bolyai', Cluj-Napoca (1987)
- Phillips, GM: Bernstein polynomials based on the  $q$ -integers. *Ann. Numer. Math.* **4**, 511-518 (1997)
- Dalmanoglu, Ö: Approximation by Kantorovich type  $q$ -Bernstein operators. In: Proceedings of the 12th WSEAS International Conference on Applied Mathematics, Cairo, Egypt, pp. 113-117 (2007)
- Aral, A, Dođru, O: Bleimann Butzer and Hahn operators based on  $q$ -integers. *J. Inequal. Appl.* **2007**, Article ID 79410 (2007)
- Muraru, CV: Note on  $q$ -Bernstein-Schurer operators. *Stud. Univ. Babeş-Bolyai, Math.* **56**(2), 489-495 (2011)
- Örkücü, M, Dođru, Ö: Weighted statistical approximation by Kantorovich type  $q$ -Szász-Mirakjan operators. *Appl. Math. Comput.* **217**, 7913-7919 (2011)
- Mahmudov, NI: On  $q$ -Szász-Durrmeyer operators. *Cent. Eur. J. Math.* **8**(2), 399-409 (2010)
- Mursaleen, M, Khan, A, Srivastava, HM, Nisar, KS: Operators constructed by means of  $q$ -Lagrange polynomials and  $A$ -statistical approximation. *Appl. Math. Comput.* **219**, 6911-6918 (2013)
- Mursaleen, M, Khan, A: Statistical approximation properties of modified  $q$ -Stancu-Beta operators. *Bull. Malays. Math. Soc.* **36**(3), 683-690 (2013)
- Özarslan, MA, Vedi, T:  $q$ -Bernstein-Schurer-Kantorovich Operators. *J. Inequal. Appl.* **2013**, Article ID 444 (2013)
- Mursaleen, M, Ansari, KJ, Khan, A: On  $(p, q)$ -analogue of Bernstein operators. *Appl. Math. Comput.* **266**, 874-882 (2015) (Erratum: *Appl. Math. Comput.* **278**, 70-71 (2016))
- Mursaleen, M, Ansari, KJ, Khan, A: Some approximation results by  $(p, q)$ -analogue of Bernstein-Stancu operators. *Appl. Math. Comput.* **264**, 392-402 (2015) (Corrigendum: *Appl. Math. Comput.* **269**, 744-746 (2015))
- Mursaleen, M, Nasruzzaman, M, Nurgali, A: Some approximation results on Bernstein-Schurer operators defined by  $(p, q)$ -integers. *J. Inequal. Appl.* **2015**, Article ID 249 (2015)
- Acar, T:  $(p, q)$ -Generalization of Szász-Mirakjan operators. *Math. Methods Appl. Sci.* **39**(10), 2685-2695 (2016)
- Acar, T, Aral, A, Mohiuddine, SA: On Kantorovich modifications of  $(p, q)$ -Baskakov operators. *J. Inequal. Appl.* **2016**, Article ID 98 (2016)
- Acar, T, Aral, A, Mohiuddine, SA: Approximation by bivariate  $(p, q)$ -Bernstein-Kantorovich operators. *Iran. J. Sci. Technol., Trans. A, Sci.* (2016). doi:10.1007/s40995-016-0045-4
- Cai, QB, Zhou, G: On  $(p, q)$ -analogue of Kantorovich type Bernstein-Stancu-Schurer operators. *Appl. Math. Comput.* **276**, 12-20 (2016)
- Mursaleen, M, Alotaibi, A, Ansari, KJ: On a Kantorovich variant of  $(p, q)$ -Szász-Mirakjan operators. *J. Funct. Spaces* **2016**, Article ID 1035253 (2016)
- Mursaleen, M, Khan, F, Khan, A: Approximation by  $(p, q)$ -Lorentz polynomials on a compact disk. *Complex Anal. Oper. Theory* (2016). doi:10.1007/s11785-016-0553-4
- Hounkonnou, MN, Dsir, J, Kyemba, B:  $R(p, q)$ -Calculus: differentiation and integration. *SUT J. Math.* **49**(2), 145-167 (2013)
- Sadjang, PN: On the fundamental theorem of  $(p, q)$ -calculus and some  $(p, q)$ -Taylor formulas. arXiv:1309.3934 [math.QA]
- Sahai, V, Yadav, S: Representations of two parameter quantum algebras and  $p, q$ -special functions. *J. Math. Anal. Appl.* **335**, 268-279 (2007)
- Stancu, DD: On the beta approximating operators of second kind. *Rev. Anal. Numér. Théor. Approx.* **24**, 231-239 (1995)
- Aral, A, Gupta, V: On the  $q$ -analogue of Stancu-Beta operators. *Appl. Math. Lett.* **25**, 67-71 (2012)
- Mursaleen, M, Khan, T: Approximation by Stancu-Beta operators via  $(p, q)$ -calculus. arXiv:1602.06319, submit/1463821
- Devore, RA, Lorentz, GG: *Constructive Approximation*. Springer, Berlin (1993)
- Gadjiev, AD: Theorems of the type of P.P. Korovkin type theorems. *Mat. Zametki* **20**(5), 781-786 (1976) (English translation: *Math. Notes* **20**(5-6), 996-998 (1976))
- Lenze, B: On Lipschitz-type maximal functions and their smoothness spaces. *Indag. Math.* **91**, 53-63 (1988)
- Braha, NL, Srivastava, HM, Mohiuddine, SA: A Korovkin's type approximation theorem for periodic functions via the statistical summability of the type of P.P. Korovkin type theorems. *Appl. Math. Comput.* **228**, 162-169 (2014)
- Edely, OHH, Mohiuddine, SA, Noman, AK: Korovkin type approximation theorems obtained through generalized statistical convergence. *Appl. Math. Lett.* **23**, 1382-1387 (2010)
- Mohiuddine, SA: An application of almost convergence in approximation theorems. *Appl. Math. Lett.* **24**, 1856-1860 (2011)