

RESEARCH

Open Access



# A new conservative nonlinear high-order compact finite difference scheme for the general Rosenau-RLW equation

Huan Wang, Jue Wang\* and Shuguang Li

\*Correspondence:  
wangjue3721@163.com  
School of Science, Harbin  
Engineering University, Harbin,  
150001, China

## Abstract

In this paper, a new conservative high-order compact finite difference scheme is studied for the initial-boundary value problem of the generalized Rosenau-regularized long wave equation. We design new conservative nonlinear fourth-order compact finite difference schemes. It is proved by the discrete energy method that the compact scheme is uniquely solvable; we have the energy conservation and the mass conservation for this approach in discrete Sobolev spaces. The convergence and stability of the difference schemes are obtained, and its numerical convergence order is  $O(\tau^2 + h^4)$  in the  $L^\infty$ -norm. Furthermore, numerical results are given to support the theoretical analysis. Numerical experiment results show that the theory is accurate and the method is efficient and reliable.

**Keywords:** generalized Rosenau-RLW equation; compact finite difference scheme; unique solvability; convergence; stability; conservation

## 1 Introduction

In this paper, we consider the following initial-boundary value problem of the Generalized Rosenau-RLW Regularized Long Wave (RLW) equation (GRRLW):

$$u_t + u_{xxxxt} - u_{xxt} + u_x + (u^p)_x = 0, \quad (x, t) \in \Omega \times (0, T], \quad (1.1)$$

$$u(x_l, t) = u(x_r, t) = 0, \quad u_{xx}(x_l, t) = u_{xx}(x_r, t) = 0, \quad t \in (0, T], \quad (1.2)$$

$$u(x, 0) = u_0(x), \quad x \in \Omega, \quad (1.3)$$

where  $p \geq 2$  is a positive integer,  $\Omega = (x_l, x_r)$  and  $u_0(x)$  are known smooth functions. Let  $H_0^2(\Omega) = \{v(x) \in H^2(\Omega) \mid v(x_l, t) = v(x_r, t) = 0, v_{xx}(x_l, t) = v_{xx}(x_r, t) = 0\}$ . The initial-boundary value problem (1.1)-(1.3) possesses the following conservative quantities:

$$E(t) = \|u\|_{L^2}^2 + \|u_x\|_{L^2}^2 + \|u_{xx}\|_{L^2}^2 = E(0), \quad (1.4)$$

$$Q(t) = \int_{x_l}^{x_r} u(x, t) dx = \int_{x_l}^{x_r} u(x, 0) dx = Q(0). \quad (1.5)$$

For the Schrödinger equation, the Cahn-Hilliard equation, and the Klein-Gordon equation, the existence and uniqueness of numerical solutions were discussed in [1–5], respec-

tively. The convergence and stability of the finite difference schemes were proved in the theory, and their numerical convergence orders are  $O(\tau^2 + h^2)$ . In [6–12], some new finite difference schemes for the initial-boundary value problem of the RLW equation were considered. Two types of conservative finite difference schemes were proposed in [13], which depended on the choice of a parameter. On the basis of the prior estimates as regards the norms, the convergence of the difference solution was proved with order  $O(\tau^2 + h^2)$  in the energy norm in [14, 15]. For the Cahn-Hilliard equation, a three-level linearized high-order compact difference scheme was derived. The unique solvability and unconditional convergence of the difference solution were proved. The convergence order is  $O(\tau^2 + h^4)$  in the maximum norm in [16]. In [17], a new conservative difference scheme for the general Rosenau-RLW equation was proposed. In [18], Pan and Zhang proposed a conservative linearized difference scheme for the general Rosenau-RLW equation which was unconditionally stable and second-order convergent and simulates conservative laws at the same time. In [19], the initial-boundary value problem for the Rosenau-RLW equation was studied. One proposed a three-level linear finite difference scheme, which has the theoretical accuracy of  $O(\tau^2 + h^4)$ .

This paper is organized as follows. In Section 2, a nonlinear and conservative difference scheme for the GRRLW equation is constructed, and the discrete conservative laws of the difference scheme are discussed. The unique solvability of the numerical solutions is also given. In Section 3, the prior error estimates for a fourth-order finite difference approximation of the GRRLW equation are obtained, and the convergence and stability of the difference scheme are proved. Numerical experiments are reported in Section 4.

### 2 Finite difference scheme and conservation law

Let  $h = (x_r - x_l)/J$  be the uniform step size in the spatial direction for positive integer  $J$ . Let  $\tau$  denote the uniform step size in the temporal direction. Denote  $x_j = x_l + jh$  ( $0 \leq j \leq J$ ),  $t^n = n\tau$  ( $0 \leq n \leq N$ ). Let  $U_j^n$  denote the approximation of  $u(x_j, t_n)$ , and let

$$\mathbf{R}_0^J = \{V_j = (V_j)_{j \in \mathbb{Z}} \mid V_0 = V_J = 0\}.$$

As usual, the following notations will be used:

$$\begin{aligned} \delta_x V_i^n &= \frac{V_{i+1}^n - V_i^n}{h}, & \delta_{\bar{x}} V_i^n &= \frac{V_{i+1}^n - V_{i-1}^n}{2h}, & \delta_{\bar{x}} V_i^n &= \frac{V_{i+2}^n - V_{i-2}^n}{4h}, \\ \delta_x^2 V_i^n &= \frac{V_{i+1}^n - 2V_i^n + V_{i-1}^n}{h^2}, & \delta_x^4 V_i^n &= \delta_x^2(\delta_x^2 V_i^n), & \partial_t V_i^n &= \frac{V_i^{n+1} - V_i^n}{\tau}, \\ V_i^{n+\frac{1}{2}} &= \frac{V_i^{n+1} + V_i^n}{2}, & \mathcal{A}_1 V_i &= \left(1 + \frac{h^2}{12} \delta_x^2\right) V_i, & \mathcal{A}_2 V_i &= \left(1 + \frac{h^2}{6} \delta_x^2\right) V_i. \end{aligned}$$

We now introduce the discrete  $L^2$ -inner product and the associated norm

$$(U, V)_h = h \sum_{i=1}^{J-1} U_i V_i, \quad U, V \in \mathbf{R}_0^J, \quad \|V\|_h = (V, V)_h^{\frac{1}{2}}.$$

The discrete  $H^m$ -seminorm  $|\cdot|_{m,h}$ , the  $H^m$ -norm  $\|\cdot\|_{m,h}$  and the  $L^\infty$ -norm  $\|\cdot\|_{\infty,h}$  are defined, respectively, as

$$|V|_{m,h} = \left(h \sum_{i=0}^{J-m} |\delta_x^m V_i|^2\right)^{\frac{1}{2}}, \quad \|V\|_{m,h} = \left(\sum_{s=0}^m |V|_{s,h}^2\right)^{\frac{1}{2}}, \quad \|V\|_{\infty,h} = \max_{0 \leq i \leq J} |V_i|,$$

where the  $\delta_x^m$  ( $m \geq 1$ ) denote the  $m$ th-order forward difference quotient operators in the  $x$  direction. It is convenient to let  $L_h^2(\Omega_h)$  and  $H_h^m(\Omega_h)$  ( $m \geq 1$ ) denote the normed vector space, respectively, as

$$L_h^2(\Omega_h) := \{\mathbf{R}_0^J, \|\cdot\|_h\}, \quad H_h^m(\Omega_h) := \{\mathbf{R}_0^J, \|\cdot\|_{m,h}\},$$

where  $\Omega_h = \{x_j = x_l + jh \mid 0 < j < J\}$ .

For the discretization of the first-order derivatives  $u_x$ , the second-order derivatives  $u_{xx}$  and the fourth-order derivatives  $u_{xxxx}$  of the function  $u(x)$ , we have the following formulas:

$$\begin{aligned} \mathcal{A}_1 u_{xx}(x_i) &= \delta_x^2 u(x_i) + O(h^4) \quad \Rightarrow \quad u_{xx}(x_i) = \mathcal{A}_1^{-1} \delta_x^2 u(x_i) + O(h^4), \\ \mathcal{A}_2 u_x(x_i) &= \delta_{\hat{x}} u(x_i) + O(h^4) \quad \Rightarrow \quad u_x(x_i) = \mathcal{A}_2^{-1} \delta_{\hat{x}} u(x_i) + O(h^4), \\ \mathcal{A}_2 u_{xxxx}(x_i) &= \delta_x^4 u(x_i) + O(h^4) \quad \Rightarrow \quad u_{xxxx}(x_i) = \mathcal{A}_2^{-1} \delta_x^4 u(x_i) + O(h^4), \end{aligned}$$

omitting the small terms  $O(h^4)$ , we obtain the approximation of  $u_{xx}$ ,  $u_x$ , and  $u_{xxxx}$  as

$$\begin{aligned} \mathcal{A}_1 u_{xx}(x_i) &\approx \delta_x^2 U_i \quad \Rightarrow \quad u_{xx}(x_i) \approx \mathcal{A}_1^{-1} \delta_x^2 U_i, \\ \mathcal{A}_2 u_x(x_i) &\approx \delta_{\hat{x}} U_i \quad \Rightarrow \quad u_x(x_i) \approx \mathcal{A}_2^{-1} \delta_{\hat{x}} U_i, \\ \mathcal{A}_2 u_{xxxx}(x_i) &\approx \delta_x^4 U_i \quad \Rightarrow \quad u_{xxxx}(x_i) \approx \mathcal{A}_2^{-1} \delta_x^4 U_i, \end{aligned}$$

where  $U_i$  is the approximation of  $u(x_i)$ . The corresponding matrix form is

$$\begin{aligned} M_1(\Pi_h u_{xx}) &\approx \delta_x^2 U \quad \Rightarrow \quad \Pi_h u_{xx} \approx M_1^{-1} \delta_x^2 U, \\ M_2(\Pi_h u_x) &\approx \delta_{\hat{x}} U \quad \Rightarrow \quad \Pi_h u_x \approx M_2^{-1} \delta_{\hat{x}} U, \\ M_2(\Pi_h u_{xxxx}) &\approx \delta_x^4 U \quad \Rightarrow \quad \Pi_h u_{xxxx} \approx M_2^{-1} \delta_x^4 U, \end{aligned}$$

where

$$\begin{aligned} U &= (U_1, U_2, \dots, U_{J-1}), \\ \Pi_h u_x &= (u_x(x_1), u_x(x_2), \dots, u_x(x_{J-1})), \\ \Pi_h u_{xx} &= (u_{xx}(x_1), u_{xx}(x_2), \dots, u_{xx}(x_{J-1})), \\ \Pi_h u_{xxxx} &= (u_{xxxx}(x_1), u_{xxxx}(x_2), \dots, u_{xxxx}(x_{J-1})), \end{aligned}$$

and

$$\begin{aligned} M_1 &= \frac{1}{12} \begin{pmatrix} 10 & 1 & 0 & \dots & 0 & 0 \\ 1 & 10 & 1 & \dots & 0 & 0 \\ \vdots & \vdots & \vdots & \dots & \vdots & \vdots \\ 0 & 0 & 0 & \dots & 1 & 10 \end{pmatrix}_{(J-1) \times (J-1)}, \\ M_2 &= \frac{1}{6} \begin{pmatrix} 4 & 1 & 0 & \dots & 0 & 0 \\ 1 & 4 & 1 & \dots & 0 & 0 \\ \vdots & \vdots & \vdots & \dots & \vdots & \vdots \\ 0 & 0 & 0 & \dots & 1 & 4 \end{pmatrix}_{(J-1) \times (J-1)}. \end{aligned}$$

Imposing the compact difference scheme of the GRRLW equations (1.1)-(1.3) gives

$$\begin{aligned} & \partial_t U_i^n + \mathcal{A}_2^{-1} \delta_x^4 \partial_t U_i^n + \mathcal{A}_2^{-1} \delta_{\bar{x}} U_i^{n+\frac{1}{2}} - \mathcal{A}_1^{-1} \delta_x^2 \partial_t U_i^n \\ & + \frac{4p}{3(p+1)} \left[ (U_i^{n+\frac{1}{2}})^{p-1} \delta_{\bar{x}} U_i^{n+\frac{1}{2}} + \delta_{\bar{x}} (U_i^{n+\frac{1}{2}})^p \right] \\ & - \frac{p}{3(p+1)} \left[ (U_i^{n+\frac{1}{2}})^{p-1} \delta_{\bar{x}} U_i^{n+\frac{1}{2}} + \delta_{\bar{x}} (U_i^{n+\frac{1}{2}})^p \right] = 0, \\ & 1 \leq i \leq J-1, 0 \leq n \leq N-1, \end{aligned} \tag{2.1}$$

$$U_0^n = U_J^n = 0, \quad \delta_x^2 U_0^n = \delta_x^2 U_J^n = 0, \quad 0 \leq n \leq N, \tag{2.2}$$

$$U_i^0 = u_0(x_i), \quad 0 \leq i \leq J. \tag{2.3}$$

**Lemma 2.1** [20] *The eigenvalues of the matrices  $M_1, M_2$  are, respectively, in the following forms:*

$$\lambda_{M_1,k} = \frac{1}{6} \left( 5 + \cos \frac{k\pi}{J} \right), \quad \lambda_{M_2,k} = \frac{1}{3} \left( 2 + \cos \frac{k\pi}{J} \right), \quad k = 1, 2, \dots, J-1.$$

For the real symmetric positive definite matrices  $M_1, M_2$ , we let  $H_1 = M_1^{-1}$  and  $H_2 = M_2^{-1}$ . Then  $H_1, H_2$  are also real symmetric positive definite matrices. Now, we introduce the following discrete norm:

$$\| |V| \|_{1,h} = [(H_1 \delta_x V, \delta_x V)]^{\frac{1}{2}}, \quad \| |V| \|_{2,h} = [(H_2 \delta_x^2 V, \delta_x^2 V)]^{\frac{1}{2}}, \quad V \in \mathbf{R}_0^J. \tag{2.4}$$

**Lemma 2.2** *The discrete norms  $\| | \cdot \|_{l,h}$  and  $| \cdot |_{l,h}$  ( $l = 1, 2$ ) are equivalent. In fact, for any grid function  $V \in \mathbf{R}_0^J$ , we have*

$$c_1 |V|_{1,h} \leq \| |V| \|_{1,h} \leq c_2 |V|_{1,h}, \quad c_1 |V|_{2,h} \leq \| |V| \|_{2,h} \leq c_3 |V|_{2,h}, \tag{2.5}$$

where  $c_1 = 1, c_2 = \sqrt{\frac{3}{2}}, c_3 = \sqrt{3}$ .

*Proof* It follows from Lemma 2.1 that the eigenvalues of  $H_1$  and  $H_2$  satisfy

$$1 \leq \lambda_{H_1,k} \leq \frac{3}{2}, \quad 1 \leq \lambda_{H_2,k} \leq 3, \quad k = 1, 2, \dots, J-1.$$

these give the spectral radius  $\rho(H_1) \leq \frac{3}{2}, \rho(H_2) \leq 3$ , and consequently

$$1 \leq \|H_1\| = \|\rho(H_1)\| \leq \frac{3}{2}, \quad 1 \leq \|H_2\| = \|\rho(H_2)\| \leq 3. \tag{2.6}$$

Thus we have

$$\begin{aligned} |V|_{1,h}^2 & \leq (H_1 \delta_x V, \delta_x V)_h \leq \|H_1\| (\delta_x V, \delta_x V)_h \leq \frac{3}{2} |V|_{1,h}^2, \\ |V|_{2,h}^2 & \leq (H_2 \delta_x^2 V, \delta_x^2 V)_h \leq \|H_2\| (\delta_x^2 V, \delta_x^2 V)_h \leq 3 |V|_{2,h}^2. \end{aligned} \tag{2.7}$$

□

**Lemma 2.3** [17] For  $U, V \in \mathbf{R}_0^J$ , we have

$$(\delta_{\hat{x}}U, V)_h = -(U, \delta_{\hat{x}}V)_h, \quad (\delta_{\hat{x}}U, V)_h = -(U, \delta_{\hat{x}}V)_h.$$

**Lemma 2.4** [21] For any discrete function  $V \in \mathbf{R}_0^J$ , we have interpolation formulas as follows:

$$\|V\|_{k,h} \leq K_0 \|V\|_{\frac{k}{n,h}}^{\frac{k}{n}} \|V\|_h^{1-\frac{k}{n}}, \tag{2.8}$$

for  $0 \leq k \leq n$ , and

$$\|V\|_{\infty,h} \leq K \|V\|_{\frac{1}{n,h}}^{\frac{1}{n}} \|V\|_h^{1-\frac{1}{n}}, \tag{2.9}$$

for  $n \geq 1$ , where  $K_0$  and  $K$  are constants independent of  $h$  and  $V$ .

**Lemma 2.5** [21] For  $V \in H_h^1(\Omega_h)$ , we have

$$\|V\|_h^2 \leq K_1 |V|_{1,h}^2,$$

where  $K_1$  is a constant independent of  $h$  and  $V$ .

**Lemma 2.6** [22] For  $V \in H_h^2(\Omega_h)$ , we have

$$|V|_{1,h}^2 \leq K_2 |V|_{2,h}^2,$$

where  $K_2$  is a constant independent of  $h$  and  $V$ .

**Lemma 2.7** [23] Let  $(H, (\cdot, \cdot)_h)$  be a finite-dimensional inner product space,  $\|\cdot\|_h$  be the associated norm, and  $g : H \rightarrow H$  be continuous. Assume, moreover, that  $\exists \alpha > 0, \forall z \in H, \|z\|_h = \alpha, (g(z), z) \geq 0$ . Then there exists a  $z^* \in H$  such that  $g(z^*) = 0$  and  $\|z^*\|_h \leq \alpha$ .

**Lemma 2.8** [19] Suppose that the discrete function  $\{\omega^n \mid n = 0, 1, 2, \dots, N; N\tau = T\}$  satisfies the inequality

$$\omega^n - \omega^{n-1} \leq A\tau\omega^n + B\tau\omega^{n-1} + C_n\tau,$$

where  $A, B$ , and  $C_n$  are nonnegative constants. Then

$$\max_{1 \leq n \leq N} |\omega^n| \leq \left( \omega^0 + \tau \sum_{l=1}^N C_l \right) e^{2(A+B)T},$$

where  $\tau$  is sufficiently small, such that  $(A + B)\tau \leq \frac{N-1}{2N}$  ( $N > 1$ ).

The matrix form of the difference scheme (2.1)-(2.3) can be written as

$$\begin{aligned} & \partial_t U^n + H_2 \delta_x^4 \partial_t U^n + H_2 \delta_{\bar{x}} U^{n+\frac{1}{2}} - H_1 \delta_x^2 \partial_t U^n \\ & + \frac{4p}{3(p+1)} [(U^{n+\frac{1}{2}})^p \delta_{\bar{x}} U^{n+\frac{1}{2}} + \delta_{\bar{x}} (U_i^{n+\frac{1}{2}})^p] \\ & - \frac{p}{3(p+1)} [(U^{n+\frac{1}{2}})^{p-1} \delta_{\bar{x}} U^{n+\frac{1}{2}} + \delta_{\bar{x}} (U^{n+\frac{1}{2}})^p] = 0, \quad 0 \leq n \leq N-1, \end{aligned} \tag{2.10}$$

$$U^n|_{\partial\Omega_h} = 0, \quad \delta_x^2 U^n|_{\partial\Omega_h} = 0, \quad 0 \leq n \leq N, \tag{2.11}$$

$$U_i^0 = u_0(x_i), \quad 0 \leq i \leq J. \tag{2.12}$$

Let  $Z_h^0 = \{V_j = (V_j)_{j \in \mathbb{Z}} \mid V_0 = V_J = 0, \delta_x^2 V_0 = \delta_x^2 V_J = 0\}$ , obviously, the solution  $U^n \in Z_h^0$  of the difference scheme (2.1)-(2.3), then there are the following lemmas:

**Theorem 2.9** *Assume  $u_0 \in H_0^2(\Omega)$ , then the finite difference scheme (2.1)-(2.3) is conservative for the discrete energy and the discrete mass, i.e.*

$$E^n = \|U^n\|_h^2 + \|U^n\|_{1,h}^2 + \|U^n\|_{2,h}^2 = \dots = E^0 \tag{2.13}$$

and

$$Q^n = h \sum_{j=1}^{J-1} U_j^n = Q^{n-1} = \dots = Q^0.$$

*Proof* Taking the inner product of (2.10) with  $2U^{n+\frac{1}{2}}$ , we obtain

$$\begin{aligned} & (\partial_t U^n, 2U^{n+\frac{1}{2}})_h + (H_2 \delta_x^4 \partial_t U^n, 2U^{n+\frac{1}{2}})_h + (H_2 \delta_{\bar{x}} U^{n+\frac{1}{2}}, 2U^{n+\frac{1}{2}})_h \\ & - (H_1 \delta_x^2 \partial_t U^n, 2U^{n+\frac{1}{2}})_h + \frac{4p}{3(p+1)} (((U^{n+\frac{1}{2}})^{p-1} \delta_{\bar{x}} U^{n+\frac{1}{2}} + \delta_{\bar{x}} (U^{n+\frac{1}{2}})^p), 2U^{n+\frac{1}{2}})_h \\ & - \frac{p}{3(p+1)} (((U^{n+\frac{1}{2}})^{p-1} \delta_{\bar{x}} U^{n+\frac{1}{2}} + \delta_{\bar{x}} (U^{n+\frac{1}{2}})^p), 2U^{n+\frac{1}{2}})_h = 0, \end{aligned} \tag{2.14}$$

letting

$$\begin{aligned} \phi(U, U) &= \frac{4p}{3(p+1)} (U^{p-1} \delta_{\bar{x}} U + \delta_{\bar{x}} (U^p)), \\ \psi(U, U) &= \frac{p}{3(p+1)} (U^{p-1} \delta_{\bar{x}} U + \delta_{\bar{x}} (U^p)), \end{aligned}$$

from Lemma 2.3, we have

$$\begin{aligned} (\phi(U, U), U)_h &= \frac{4p}{3(p+1)} (U^{p-1} \delta_{\bar{x}} U + \delta_{\bar{x}} (U^p), U)_h \\ &= \frac{4p}{3(p+1)} [(U^{p-1} \delta_{\bar{x}} U, U)_h + (\delta_{\bar{x}} (U^p), U)_h] \\ &= \frac{4p}{3(p+1)} [(\delta_{\bar{x}} U, U^p)_h - (\delta_{\bar{x}} U, U^p)_h] = 0 \end{aligned} \tag{2.15}$$

and

$$\begin{aligned}
 (\psi(U, U), U)_h &= \frac{p}{3(p+1)} (U^{p-1} \delta_{\bar{x}} U + \delta_{\bar{x}}(U^p), U)_h \\
 &= \frac{p}{3(p+1)} [(U^{p-1} \delta_{\bar{x}} U, U)_h + (\delta_{\bar{x}}(U^p), U)_h] \\
 &= \frac{p}{3(p+1)} [(\delta_{\bar{x}} U, U^p)_h - (\delta_{\bar{x}} U, U^p)_h] = 0.
 \end{aligned}
 \tag{2.16}$$

Thus from (2.14)-(2.16), we can obtain

$$(\|U^{n+1}\|_h^2 - \|U^n\|_h^2) + (\|U^{n+1}\|_{1,h}^2 - \|U^n\|_{1,h}^2) + (\|U^{n+1}\|_{2,h}^2 - \|U^n\|_{2,h}^2) = 0.
 \tag{2.17}$$

Let  $E^n$  denote the following discrete energy:

$$E^n = \|U^n\|_h^2 + \|U^n\|_{1,h}^2 + \|U^n\|_{2,h}^2,
 \tag{2.18}$$

then from (2.17), we get

$$E^n = E^{n-1} = \dots = E^0.$$

Multiplying (2.1) with  $h$ , according to the boundary condition (2.2), summing for  $j$  from 1 to  $J - 1$ , we obtain

$$h \sum_{j=1}^{J-1} (U_j^{n+1} - U_j^n) = 0,$$

letting

$$Q^n = h \sum_{j=1}^{J-1} U_j^n,$$

then we have

$$Q^n = Q^{n-1} = \dots = Q^0.$$

This completes the proof. □

**Lemma 2.10** *Assume  $u_0 \in H_0^2(\Omega)$ , then there is the estimation for the solution of the difference scheme (2.1)-(2.3)*

$$\|U^n\|_{1,h} \leq \sqrt{\frac{(2K_2 + 1)E^0}{K_2 + 1}}, \quad \|U^n\|_{\infty,h} \leq K \sqrt{\frac{(2K_2 + 1)E^0}{K_2 + 1}}.$$

*Proof* From Lemma 2.2, Lemma 2.6, and Theorem 2.9, we have

$$\left(\frac{K_2 + 1}{K_2}\right) |U^n|_{1,h}^2 + \|U^n\|_h^2 \leq \|U^n\|_h^2 + |U^n|_{1,h}^2 + |U^n|_{2,h}^2 \leq E^0, \quad n \geq 0.$$

Hence, we can get

$$\|U^n\|_{1,h} = \sqrt{\|U^n\|_h^2 + |U^n|_{1,h}^2} \leq \sqrt{E^0 + \frac{K_2 E^0}{K_2 + 1}} = \sqrt{\frac{(2K_2 + 1)E^0}{K_2 + 1}}.$$

It follows from Lemma 2.4 that

$$\|U^n\|_{\infty,h} \leq K \|U^n\|_{1,h} \leq K \sqrt{\frac{(2K_2 + 1)E^0}{K_2 + 1}}.$$

This completes the proof. □

**Lemma 2.11** For  $V \in \mathbf{Z}_h^0$ , we have

$$\|\delta_{\bar{x}} V\|_h^2 \leq \|\delta_{\hat{x}} V\|_h^2 \leq \|\delta_x V\|_h^2.$$

*Proof* From the definition of  $\|\cdot\|_h$ , we have

$$\begin{aligned} \|\delta_{\bar{x}} V\|_h^2 &= h \sum_{j=2}^{J-2} (\delta_{\bar{x}} V_j)^2 = \frac{h}{4} \sum_{j=2}^{J-2} (\delta_{\bar{x}} V_{j+1} + \delta_{\bar{x}} V_{j-1})^2 \\ &= \frac{h}{4} \sum_{j=2}^{J-2} ((\delta_{\bar{x}} V_{j+1})^2 + (\delta_{\bar{x}} V_{j-1})^2 + 2(\delta_{\bar{x}} V_{j+1})(\delta_{\bar{x}} V_{j-1})) \leq \|\delta_{\bar{x}} V\|_h^2 \end{aligned}$$

and

$$\begin{aligned} \|\delta_{\hat{x}} V\|_h^2 &= h \sum_{j=1}^{J-1} (\delta_{\hat{x}} V_j)^2 = \frac{h}{4} \sum_{j=1}^{J-1} (\delta_x V_j + \delta_x V_{j-1})^2 \\ &= \frac{h}{4} \sum_{j=1}^{J-1} ((\delta_x V_j)^2 + (\delta_x V_{j-1})^2 + 2(\delta_x V_j)(\delta_x V_{j-1})) \leq \|\delta_x V\|_h^2. \end{aligned}$$

The proof is completed. □

**Theorem 2.12** The difference scheme (2.1)-(2.3) is uniquely solvable.

*Proof* For a fixed  $n$ , (2.10) can be written as

$$\begin{aligned} U^{n+\frac{1}{2}} - U^n + H_2 \delta_x^4 (U^{n+\frac{1}{2}} - U^n) + \frac{\tau}{2} H_2 \delta_{\bar{x}} U^{n+\frac{1}{2}} - H_1 \delta_x^2 (U^{n+\frac{1}{2}} - U^n) \\ + \frac{\tau}{2} \phi(U^{n+\frac{1}{2}}, U^{n+\frac{1}{2}}) - \frac{\tau}{2} \psi(U^{n+\frac{1}{2}}, U^{n+\frac{1}{2}}) = 0, \end{aligned} \tag{2.19}$$

we define  $F$  on  $\mathbf{Z}_h^0$  as follows:

$$\begin{aligned} F(\xi) &= \xi - U^n + H_2 \delta_x^4 \xi - H_2 \delta_x^4 U^n + \frac{\tau}{2} H_2 \delta_{\bar{x}} \xi \\ &\quad - H_1 \delta_x^2 \xi + H_1 \delta_x^2 U^n + \frac{\tau}{2} \phi(\xi, \xi) - \frac{\tau}{2} \psi(\xi, \xi), \end{aligned} \tag{2.20}$$



obviously,  $F$  is continuous. Computing the inner product of (2.20) with  $\xi$  and considering  $(\phi(\xi, \xi), \xi)_h = 0$ ,  $(\psi(\xi, \xi), \xi)_h = 0$  and  $(H_2\delta_x^2\xi, \xi)_h = 0$ , we obtain

$$\begin{aligned} (F(\xi), \xi)_h &= \|\xi\|_h^2 - (U^n, \xi)_h + \|\xi\|_{2,h}^2 - (H_2\delta_x^2U^n, \delta_x^2\xi)_h + \|\xi\|_{1,h}^2 + (H_1\delta_x^2U^n, \xi)_h \\ &\geq \|\xi\|_h^2 - \frac{1}{2}(\|\xi\|_h^2 + \|U^n\|_h^2) + \|\xi\|_{2,h}^2 + \|\xi\|_{1,h}^2 \\ &\quad - (H_2\delta_x^2U^n, \delta_x^2\xi)_h + (H_1\delta_x^2U^n, \xi)_h \\ &\geq \frac{1}{2}(\|\xi\|_h^2 - \|U^n\|_h^2) + \|\xi\|_{1,h}^2 + \|\xi\|_{2,h}^2 - \frac{1}{2}(\|\xi\|_{2,h}^2 + \|U^n\|_{2,h}^2) \\ &\quad - \frac{1}{2}(\|\xi\|_{1,h}^2 + \|U^n\|_{1,h}^2) \\ &= \frac{1}{2}(\|\xi\|_h^2 - \|U^n\|_h^2) + \frac{1}{2}\|\xi\|_{2,h}^2 + \frac{1}{2}\|\xi\|_{1,h}^2 - \frac{1}{2}\|U^n\|_{2,h}^2 - \frac{1}{2}\|U^n\|_{1,h}^2 \\ &\geq \frac{1}{2}\|\xi\|_h^2 - \frac{1}{2}(\|U^n\|_h^2 + \|U^n\|_{1,h}^2 + \|U^n\|_{2,h}^2). \end{aligned}$$

Hence, for all  $\xi \in \mathbf{Z}_h^0$ , let  $\|\xi\|_h^2 = \|U^n\|_h^2 + \|U^n\|_{1,h}^2 + \|U^n\|_{2,h}^2 + 1$ , then there exists  $(F(\xi), \xi)_h > 0$ . It follows from Lemma 2.7 that there exists a  $\xi^* \in \mathbf{Z}_h^0$  which satisfies  $F(\xi^*) = 0$ . Let  $U^{n+1} = 2\xi^* - U^n$ , then it can be proved that  $U^{n+1} \in \mathbf{Z}_h^0$  is the solution of scheme (2.1)-(2.3).

Next, we will give the uniqueness of the difference solution. Assume that  $U^n$  and  $V^n$  satisfy scheme (2.1)-(2.3), letting  $w^n = V^n - U^n$ , we have

$$\begin{aligned} &\partial_t w^n + H_2\delta_x^4\delta_t w^n + H_2\delta_x w^{n+\frac{1}{2}} - H_1\delta_x^2\partial_t w^n \\ &\quad + [\phi(V^{n+\frac{1}{2}}, V^{n+\frac{1}{2}}) - \phi(U^{n+\frac{1}{2}}, U^{n+\frac{1}{2}})] - [\psi(V^{n+\frac{1}{2}}, V^{n+\frac{1}{2}}) - \psi(U^{n+\frac{1}{2}}, U^{n+\frac{1}{2}})] \\ &= 0. \end{aligned} \tag{2.21}$$

Computing the inner product of (2.21) with  $2w^{n+\frac{1}{2}}$ , we have

$$\begin{aligned} 0 &= (\|w^{n+1}\|_h^2 - \|w^n\|_h^2) + (\|w^{n+1}\|_{2,h}^2 - \|w^n\|_{2,h}^2) + (\|w^{n+1}\|_{1,h}^2 - \|w^n\|_{1,h}^2) \\ &\quad + 2\tau(\phi(V^{n+\frac{1}{2}}, V^{n+\frac{1}{2}}) - \phi(U^{n+\frac{1}{2}}, U^{n+\frac{1}{2}}), w^{n+\frac{1}{2}})_h \\ &\quad - 2\tau(\psi(V^{n+\frac{1}{2}}, V^{n+\frac{1}{2}}) - \psi(U^{n+\frac{1}{2}}, U^{n+\frac{1}{2}}), w^{n+\frac{1}{2}})_h, \end{aligned} \tag{2.22}$$

by Lemma 2.10, we can estimate (2.22) as follows:

$$\begin{aligned} &(\phi(V^{n+\frac{1}{2}}, V^{n+\frac{1}{2}}) - \phi(U^{n+\frac{1}{2}}, U^{n+\frac{1}{2}}), w^{n+\frac{1}{2}})_h \\ &= \frac{4ph}{3(p+1)} \sum_{i=1}^{J-1} [(V_i^{n+\frac{1}{2}})^{p-1} \delta_x V_i^{n+\frac{1}{2}} - (U_i^{n+\frac{1}{2}})^{p-1} \delta_x U_i^{n+\frac{1}{2}}] w_i^{n+\frac{1}{2}} \\ &\quad + \frac{4ph}{3(p+1)} \sum_{i=1}^{J-1} [\delta_x (V_i^{n+\frac{1}{2}})^p - \delta_x (U_i^{n+\frac{1}{2}})^p] w_i^{n+\frac{1}{2}} \\ &\leq \frac{2p}{3(p+1)} \max\{K_3^{p-1}, (p-1)K_3^{p-1}\} (\|w^{n+1}\|_{1,h}^2 + \|w^n\|_{1,h}^2), \end{aligned} \tag{2.23}$$

$$\begin{aligned}
 & (\psi(V^{n+\frac{1}{2}}, V^{n+\frac{1}{2}}) - \psi(U^{n+\frac{1}{2}}, U^{n+\frac{1}{2}}), w^{n+\frac{1}{2}})_h \\
 &= \frac{ph}{3(p+1)} \sum_{i=2}^{J-2} [(V_i^{n+\frac{1}{2}})^{p-1} \delta_{\bar{x}} V_i^{n+\frac{1}{2}} - (U_i^{n+\frac{1}{2}})^{p-1} \delta_{\bar{x}} U_i^{n+\frac{1}{2}}] w_i^{n+\frac{1}{2}} \\
 & \quad + \frac{ph}{3(p+1)} \sum_{i=2}^{J-2} [\delta_{\bar{x}} (V_i^{n+\frac{1}{2}})^p - \delta_{\bar{x}} (U_i^{n+\frac{1}{2}})^p] w_i^{n+\frac{1}{2}} \\
 & \leq \frac{p}{6(p+1)} \max\{K_3^{p-1}, (p-1)K_3^{p-1}\} (\|w^{n+1}\|_{1,h}^2 + \|w^n\|_{1,h}^2), \tag{2.24}
 \end{aligned}$$

where  $K_3 = K \sqrt{\frac{(2K_2+1)E^0}{K_2+1}}$ . Substituting (2.23) and (2.24) into (2.22), from Lemma 2.2, we obtain

$$\begin{aligned}
 & (\|w^{n+1}\|_h^2 + \|w^{n+1}\|_{1,h}^2 + \|w^{n+1}\|_{2,h}^2) - (\|w^n\|_h^2 + \|w^n\|_{1,h}^2 + \|w^n\|_{2,h}^2) \\
 & \leq \frac{5p\tau}{3(p+1)} \max\{K_3^{p-1}, (p-1)K_3^{p-1}\} (\|w^{n+1}\|_{1,h}^2 + \|w^n\|_{1,h}^2) \\
 & \leq K_4 \tau (\|w^{n+1}\|_{1,h}^2 + \|w^n\|_{1,h}^2) \\
 & \leq K_4 \tau (\|w^{n+1}\|_h^2 + \|w^{n+1}\|_{1,h}^2 + \|w^{n+1}\|_{2,h}^2) \\
 & \quad + K_4 \tau (\|w^n\|_h^2 + \|w^n\|_{1,h}^2 + \|w^n\|_{2,h}^2), \tag{2.25}
 \end{aligned}$$

where  $K_4 = \frac{5p}{3(p+1)} \max\{K_3^{p-1}, (p-1)K_3^{p-1}\}$ .

Choosing small enough  $\tau$ , we obtain by Lemma 2.8

$$\|w^n\|_h^2 + \|w^n\|_{1,h}^2 + \|w^n\|_{2,h}^2 = 0. \tag{2.26}$$

This completes the proof. □

### 3 Convergence and stability of the difference solution

In this section, we will consider the convergence and stability of the finite difference scheme (2.1)-(2.3). Assume that the solution  $u(x, t)$  of (1.1)-(1.3) is sufficiently smooth. We define the net function  $u_i^n = u(x_i, t_n)$  and the truncation errors as follows:

$$\begin{aligned}
 & \partial_t u_i^n + \mathcal{A}_2^{-1} \delta_x^4 \partial_t u_i^n + \mathcal{A}_2^{-1} \delta_{\bar{x}} u_i^{n+\frac{1}{2}} - \mathcal{A}_1^{-1} \delta_x^2 \partial_t u_i^n \\
 & \quad + \frac{4p}{3(p+1)} [(u_i^{n+\frac{1}{2}})^{p-1} \delta_{\bar{x}} u_i^{n+\frac{1}{2}} + \delta_{\bar{x}} (u_i^{n+\frac{1}{2}})^p] \\
 & \quad - \frac{p}{3(p+1)} [(u_i^{n+\frac{1}{2}})^{p-1} \delta_{\bar{x}} u_i^{n+\frac{1}{2}} + \delta_{\bar{x}} (u_i^{n+\frac{1}{2}})^p] = r_i^n, \\
 & 1 \leq i \leq J-1, 0 \leq n \leq N-1, \tag{3.1}
 \end{aligned}$$

$$u_0^n = u_J^n = 0, \quad \delta_x^2 u_0^n = \delta_x^2 u_J^n = 0, \quad 0 \leq n \leq N, \tag{3.2}$$

$$u_i^0 = u_0(x_i), 0 \leq i \leq J. \tag{3.3}$$

Suppose that  $u_0 \in H_0^2(\Omega)$  and  $u(x, t) \in C^{6,4}$ , then from Taylor’s expansion, the truncation errors of scheme (3.1) satisfy

$$|r_i^n| = O(\tau^2 + h^4), \quad \text{as } \tau \rightarrow 0, h \rightarrow 0. \tag{3.4}$$

**Theorem 3.1** *Suppose that  $u_0 \in H_0^2(\Omega)$  and  $u(x, t) \in C^{6,4}$ , then the solution of the difference scheme (2.1)-(2.3) converges to the solution of the problem (1.1)-(1.3) with order  $O(\tau^2 + h^4)$  by the  $L^\infty$  norm.*

*Proof* Subtracting (2.1)-(2.3) from (3.1)-(3.3) letting  $e_i^n = u_i^n - U_i^n$ , we obtain

$$\begin{aligned} \partial_t e^n + H_2 \delta_x^4 \delta_t e^n + H_2 \delta_{\bar{x}}^2 e^{n+\frac{1}{2}} - H_1 \delta_x^2 \partial_t e^n + [\phi(u^{n+\frac{1}{2}}, u^{n+\frac{1}{2}}) - \phi(U^{n+\frac{1}{2}}, U^{n+\frac{1}{2}})] \\ - [\psi(u^{n+\frac{1}{2}}, u^{n+\frac{1}{2}}) - \psi(U^{n+\frac{1}{2}}, U^{n+\frac{1}{2}})] = r^n, \quad 0 \leq n \leq N-1, \end{aligned} \tag{3.5}$$

$$e^n|_{\partial\Omega_h} = 0, \quad \delta_x^2 e^n|_{\partial\Omega_h} = 0, \quad 0 \leq n \leq N, \tag{3.6}$$

$$e_i^0 = 0, \quad 0 \leq i \leq J. \tag{3.7}$$

Computing the inner product of (3.5) with  $2e^{n+\frac{1}{2}}$ , we have

$$\begin{aligned} 2\tau (r^n, e^{n+\frac{1}{2}})_h \\ = (\|e^{n+1}\|_h^2 - \|e^n\|_h^2) + (\|e^{n+1}\|_{1,h}^2 - \|e^n\|_{1,h}^2) + (\|e^{n+1}\|_{2,h}^2 - \|e^n\|_{2,h}^2) \\ + 2\tau (\phi(u^{n+\frac{1}{2}}, u^{n+\frac{1}{2}}) - \phi(U^{n+\frac{1}{2}}, U^{n+\frac{1}{2}}), e^{n+\frac{1}{2}})_h \\ - 2\tau (\psi(u^{n+\frac{1}{2}}, u^{n+\frac{1}{2}}) - \psi(U^{n+\frac{1}{2}}, U^{n+\frac{1}{2}}), e^{n+\frac{1}{2}})_h. \end{aligned} \tag{3.8}$$

Similarly to the proof of Theorem 2.12, we have

$$\begin{aligned} (\|e^{n+1}\|_h^2 + \|e^{n+1}\|_{1,h}^2 + \|e^{n+1}\|_{2,h}^2) - (\|e^n\|_h^2 + \|e^n\|_{1,h}^2 + \|e^n\|_{2,h}^2) \\ \leq \tau \|r^n\|_h^2 + K_5 \tau (\|e^{n+1}\|_h^2 + \|e^{n+1}\|_{1,h}^2 + \|e^{n+1}\|_{2,h}^2) \\ + K_5 \tau (\|e^n\|_h^2 + \|e^n\|_{1,h}^2 + \|e^n\|_{2,h}^2), \end{aligned} \tag{3.9}$$

where  $K_5 = K_4 + \frac{1}{2}$ . Let  $B^n = \|e^n\|_h^2 + \|e^n\|_{1,h}^2 + \|e^n\|_{2,h}^2$ , then (3.9) can be rewritten as

$$B^{n+1} - B^n \leq \tau \|r^n\|_h^2 + \tau K_5 (B^{n+1} + B^n). \tag{3.10}$$

Choosing small enough  $\tau$ , from Lemma 2.8, we obtain

$$B^n \leq C(B^0 + (\tau^2 + h^4)^2). \tag{3.11}$$

From the discrete initial conditions, we know that

$$B^0 \leq O(\tau^2 + h^4)^2. \tag{3.12}$$

Then we have

$$\|e^n\|_h \leq O(\tau^2 + h^4), \quad \|e^n\|_{1,h} \leq O(\tau^2 + h^4), \quad \|e^n\|_{2,h} \leq O(\tau^2 + h^4). \tag{3.13}$$

**Table 1** The errors of numerical solutions at  $t = 60$  with  $\tau = h$  for  $p = 2$

$h$	$\ u^n - U^n\ _{\infty, h}$	$\ u^n - U^n\ _{\infty, h}$ [17]
0.4	$3.5235 \times 10^{-3}$	$1.9587 \times 10^{-2}$
0.2	$8.0413 \times 10^{-4}$	$4.9838 \times 10^{-3}$
0.1	$1.9123 \times 10^{-4}$	$1.2520 \times 10^{-3}$
0.05	$4.6595 \times 10^{-5}$	$3.1346 \times 10^{-4}$

**Table 2** The errors of numerical solutions at  $t = 60$  with  $\tau = h$  for  $p = 3$

$h$	$\ u^n - U^n\ _{\infty, h}$	$\ u^n - U^n\ _{\infty, h}$ [17]
0.4	$5.2629 \times 10^{-3}$	$4.2510 \times 10^{-2}$
0.2	$1.4684 \times 10^{-3}$	$1.0804 \times 10^{-2}$
0.1	$3.8492 \times 10^{-4}$	$2.7090 \times 10^{-3}$
0.05	$9.8927 \times 10^{-5}$	$6.7722 \times 10^{-4}$

**Table 3** The errors of numerical solutions at  $t = 60$  with  $\tau = h$  for  $p = 6$

$h$	$\ u^n - U^n\ _{\infty, h}$	$\ u^n - U^n\ _{\infty, h}$ [17]
0.4	$3.1535 \times 10^{-2}$	$6.3539 \times 10^{-2}$
0.2	$7.4328 \times 10^{-3}$	$1.6496 \times 10^{-2}$
0.1	$1.8246 \times 10^{-3}$	$4.1593 \times 10^{-3}$
0.05	$4.5437 \times 10^{-4}$	$1.0409 \times 10^{-3}$

**Table 4** The maximum norm errors and spatial convergence order with fixed time step  $\tau = \frac{1}{1,000}$

$J$	$p = 2$		$p = 3$		$p = 6$	
	$\ u^n - U^n\ _{\infty, h}$	order1	$\ u^n - U^n\ _{\infty, h}$	order1	$\ u^n - U^n\ _{\infty, h}$	order1
125	$9.7465 \times 10^{-4}$	–	$2.9741 \times 10^{-3}$	–	$4.2113 \times 10^{-3}$	–
250	$7.0687 \times 10^{-5}$	3.7854	$2.3162 \times 10^{-4}$	3.6826	$3.5045 \times 10^{-4}$	3.5870
500	$4.5932 \times 10^{-6}$	3.9439	$1.5359 \times 10^{-5}$	3.9164	$2.6008 \times 10^{-5}$	3.7522
1,000	$2.9062 \times 10^{-7}$	3.9823	$9.9102 \times 10^{-7}$	3.9540	$1.6700 \times 10^{-6}$	3.9610

**Table 5** The maximum norm errors and temporal convergence order with the fixed space step  $h = 0.1$

$N$	$p = 2$		$p = 3$		$p = 6$	
	$\ u^n - U^n\ _{\infty, h}$	order2	$\ u^n - U^n\ _{\infty, h}$	order2	$\ u^n - U^n\ _{\infty, h}$	order2
10	$3.0491 \times 10^{-5}$	–	$6.7612 \times 10^{-5}$	–	$8.8887 \times 10^{-5}$	–
20	$7.3366 \times 10^{-6}$	2.0552	$1.6812 \times 10^{-5}$	2.0078	$2.2979 \times 10^{-5}$	1.9517
40	$1.8011 \times 10^{-6}$	2.0262	$4.2029 \times 10^{-6}$	2.0000	$5.8571 \times 10^{-6}$	1.9721
80	$4.4617 \times 10^{-7}$	2.0132	$1.0439 \times 10^{-6}$	2.0094	$1.4822 \times 10^{-6}$	1.9824

By Lemma 2.2, we obtain



$$|e^n|_{1, h} \leq O(\tau^2 + h^4), \quad |e^n|_{2, h} \leq O(\tau^2 + h^4). \tag{3.14}$$

It follows from Lemma 2.4 that

$$\|e^n\|_{\infty, h} \leq O(\tau^2 + h^4). \tag{3.15}$$

This completes the proof. □

We can similarly prove the stability of the difference solution.





**Theorem 3.2** *Under the conditions of Theorem 3.1, the solution of conservative finite difference scheme (2.1)-(2.3) is stable by the  $L^\infty$  norm.*

#### 4 Numerical experiments

In this section, numerical results are provided to demonstrate the accuracy and efficiency of the compact scheme (2.1)-(2.3). The exact solution of the system (1.1)-(1.3) is

$$u(x, t) = \exp\left(\frac{\ln \frac{(p+1)(3p+1)(p+3)}{2(p^2+3)(p^2+4p+7)}}{p-1}\right) \operatorname{sech}^{\frac{4}{p-1}}\left(\left(\frac{p-1}{\sqrt{4p^2+8p+20}}\right)(x-ct)\right), \tag{4.1}$$

where  $c = \frac{p^4+4p^3+14p^2+20p+25}{p^4+4p^3+10p^2+12p+21}$  is the wave velocity. In order to compare with the literature [17], we choose  $x_l = -30, x_r = 120$ , and consider three cases:  $p = 2, p = 3$  and  $p = 6$  in Tables 1, 2, and 3, respectively. Tables 1, 2, and 3 give the errors in the sense of the  $L^\infty$ -





norm of the numerical solutions under various steps of  $\tau = h = 0.4, 0.2, 0.1, 0.05$  at  $t = 60$  for  $p = 2, 3$  and  $6$ .

Denote

$$\text{order1} = \log_2 \frac{E(\tau, 2h)}{E(\tau, h)}, \quad \text{order2} = \log_2 \frac{E(2\tau, h)}{E(\tau, h)},$$

where  $E(\tau, h) = \|u^n - U^n\|_{\infty, h}$ . First, we test the spatial errors and convergence orders by letting  $h$  vary and fixing the time step size  $\tau$  sufficiently small to avoid contamination of the temporal. Table 4 shows the numerical results when  $\tau = \frac{1}{1,000}, h = \frac{150}{125}, h = \frac{150}{250}, h = \frac{150}{500}$ , and  $h = \frac{150}{1,000}$ . It can be seen from Table 4 that the convergence order of the compact difference scheme (2.1)-(2.3) is about 4 with respect to the spatial step size.

We further test the temporal errors and convergence orders. Fix  $h = 0.1$ , a value small enough so that the spatial error is negligible as compared with the temporal error. Take



$\tau = \frac{1}{10}, \frac{1}{20}, \frac{1}{40}, \frac{1}{80}$ , respectively. Table 5 shows that the convergence order of the compact difference scheme (2.1)-(2.3) with respect to the temporal variable is about 2.

Figures 1, 2, and 3 plot the conservative law of discrete energy  $E^n$ , computed by scheme (2.1)-(2.3) with  $\tau = 0.5, h = 0.2$  for  $p = 2, 3$  and 6. Figures 4, 5, and 6 plot the exact solutions at  $t = 0$  and the numerical solutions computed by scheme (2.1)-(2.3) with  $\tau = 0.1, h = 0.2$  at  $t = 30, 60$ , which also show the accuracy of scheme (2.1)-(2.3).

**Competing interests**

The authors declare that they have no competing interests.

**Authors' contributions**

All authors contributed equally to the writing of this paper. All authors read and approved the final manuscript.

**Acknowledgements**

Supported by the National Natural Science Foundation of China (Grant No. 11401183) and the Fundamental Research Funds for the Central Universities.

**References**

1. Zhang, F, Pérez-García, VM, Vázquez, L: Numerical simulation of nonlinear Schrödinger systems: a new conservative scheme. *Appl. Math. Comput.* **71**(2-3), 165-177 (1995)
2. Choo, SM, Chung, SK: Conservative nonlinear difference scheme for the Cahn-Hilliard equation. *Comput. Math. Appl.* **36**(7), 31-39 (1998)
3. Choo, SM, Chung, SK, Kim, KI: Conservative nonlinear difference scheme for the Cahn-Hilliard equation. II. *Comput. Math. Appl.* **39**(1-2), 229-243 (2000)
4. Li, S, Vu-Quoc, L: Finite difference calculus invariant structure of a class of algorithms for the nonlinear Klein-Gordon equation. *SIAM J. Numer. Anal.* **32**(6), 1839-1875 (1995)
5. Wong, YS, Chang, Q, Gong, L: An initial-boundary value problem of a nonlinear Klein-Gordon equation. *Appl. Math. Comput.* **84**(1), 77-93 (1997)
6. Zhang, L: A finite difference scheme for generalized regularized long-wave equation. *Appl. Math. Comput.* **168**, 962-972 (2005)
7. Hu, J, Hu, B, Hu, Y: C-N difference schemes for dissipative symmetric regularized long wave equations with damping term. *Math. Probl. Eng.* **2011**, 651642 (2011)
8. Pan, X, Zhang, L: A new finite difference scheme for the Rosenau-Burgers equation. *Appl. Math. Comput.* **218**, 8917-8924 (2012)
9. Xue, G, Zhang, L: A new finite difference scheme for generalized Rosenau-Burgers equation. *Appl. Math. Comput.* **222**, 490-496 (2013)
10. Hu, J, Hu, B, Xu, Y: Average implicit linear difference scheme for generalized Rosenau-Burgers equation. *Appl. Math. Comput.* **217**, 7557-7563 (2011)
11. Hu, B, Xu, Y, Hu, J: Crank-Nicolson finite difference scheme for the Rosenau-Burgers equation. *Appl. Math. Comput.* **204**, 311-316 (2008)
12. Hu, J, Zheng, K: Two conservative difference schemes for the generalized Rosenau equation. *Bound. Value Probl.* **2010**, 543503 (2010)
13. Shao, X, Xue, G, Li, C: A conservative weighted finite difference scheme for regularized long wave equation. *Appl. Math. Comput.* **219**, 9202-9209 (2013)
14. Zhang, L: Convergence of a conservative difference scheme for a class of Klein-Gordon-Schrödinger equations in one space dimension. *Appl. Math. Comput.* **163**, 343-355 (2005)
15. Omrani, K, Abidi, F, Achouric, T, Khiaria, N: A new conservative finite difference scheme for the Rosenau equation. *Appl. Math. Comput.* **201**, 35-43 (2008)
16. Juan, L, Zhong, S, Xuan, Z: A three level linearized compact difference scheme for the Cahn-Hilliard equation. *Sci. China Math.* **55**(4), 805-826 (2012)
17. Zuo, JM, Zhang, YM, Zhang, TD, Chang, F: A new conservative difference scheme for the general Rosenau-RLW equation. *Bound. Value Probl.* **2010**, 516260 (2010)
18. Pan, XT, Zhang, LM: Numerical simulation for general Rosenau-RLW equation: an average linearized conservative scheme. *Math. Probl. Eng.* **2012**, 517818 (2012)
19. Hu, JS, Wang, YL: A high-accuracy linear conservative difference scheme for Rosenau-RLW equation. *Math. Probl. Eng.* **2013**, 870291 (2013)
20. Wang, TC: Optimal point-wise error estimate of a compact difference scheme for the Klein-Gordon-Schrödinger equation. *J. Math. Anal. Appl.* **412**, 155-167 (2014)
21. Zhou, YL: *Applications of Discrete Functional Analysis to the Finite Difference Method*. International Academic Publishers, Beijing (1990)
22. Guo, B: *The Differential Method of Partial Differential Equations*. Science Publishers, Beijing (1988)
23. Wang, TC, Guo, BL: Analysis of some finite difference schemes for two-dimensional Ginzburg-Landau equation. *Numer. Methods Partial Differ. Equ.* **27**(5), 1340-1363 (2011)

Submit your manuscript to a SpringerOpen<sup>®</sup> journal and benefit from:

- Convenient online submission
- Rigorous peer review
- Immediate publication on acceptance
- Open access: articles freely available online
- High visibility within the field
- Retaining the copyright to your article

---

Submit your next manuscript at ► [springeropen.com](http://springeropen.com)

---