

RESEARCH

Open Access



# Existence of solutions of fractional boundary value problems with $p$ -Laplacian operator

Nazim I Mahmudov and Sinem Unul\*

\*Correspondence:  
sinem.unul@emu.edu.tr  
Eastern Mediterranean University,  
Gazimagusa, TRNC, Mersin 10,  
Turkey

## Abstract

In this paper, the existence of the solutions of the fractional differential equation with  $p$ -Laplacian operator and integral conditions is discussed. By Green's functions and the fixed point theorems, we state and prove the existence and uniqueness results of the problem. Two examples are given to illustrate the results.

**Keywords:** existence and uniqueness; fractional calculus;  $p$ -Laplacian

## 1 Introduction

Differential equations are useful in modern physics, engineering, and in various fields of science. In these days, the theory on existence and uniqueness of boundary value problems of linear and/or nonlinear fractional equations has attracted the attention of many authors. There are comprehensive studies in this area. At the same time, it is known that the  $p$ -Laplacian operator is also used in analyzing mechanics, physics and dynamic systems, and the related fields of mathematical modeling. However, there are few studies of the existence and uniqueness of boundary conditions of fractional differential equations with the  $p$ -Laplacian operator, see [1–27] and the references therein.

Zhang *et al.* [4] studied the eigenvalue problem for a class of singular  $p$ -Laplacian fractional differential equations involving a Riemann-Stieltjes integral boundary condition:

$$\begin{aligned} -D_t^\beta (\phi_p(D_t^\alpha x))(t) &= \lambda f(t, x(t)), \quad t \in (0, 1), \\ x(0) &= 0, \quad D_t^\alpha x(0) = 0, \\ x(1) &= \int_0^1 x(s) dA(s), \end{aligned}$$

where  $D_t^\beta$  and  $D_t^\alpha$  are standard Riemann-Liouville derivatives with  $1 < \alpha \leq 2$ ,  $0 < \beta \leq 1$ ,  $A$  is a function of the bounded variation, and  $\int_0^1 x(s) dA(s)$  is the standard Riemann-Stieltjes integral. In their study, the results are based on upper and lower solution methods and the Schauder fixed point theorem.

In [5], Su *et al.* studied the existence criteria of non-negative solutions of nonlinear  $p$ -Laplacian fractional differential equations with first order derivative,

$$\begin{cases} \varphi_p({}^c D^\alpha u(t)) = \varphi_p(\lambda) f(t, u(t), u'(t)), & \text{for } t \in (0, 1), \\ k_0 u(0) - k_1 u(1) = 0, \\ m_0 u(0) - m_1 u(1) = 0, \\ x^{(r)}(0) = 0, & r = 2, 3, \dots, [\alpha], \end{cases}$$

where  $\varphi_p$  is  $p$ -Laplacian operator, i.e.  $\varphi_p(s) = |s|^{p-2}s, p > 1$ , and  $\varphi_p^{-1} = \varphi_q, \frac{1}{p} + \frac{1}{q} = 1, {}^c D^\alpha$  is the Caputo derivative and we have the function  $f(t, u, u') : [0, 1] \times [0, \infty) \times (-\infty, +\infty) \rightarrow [0, \infty)$  which satisfies the Carathéodory type conditions. Moreover, the nonlinear alternative of Leray-Schauder type and Banach fixed point theorems are used.

Han *et al.* [6] studied nonlinear fractional differential equations with  $p$ -Laplacian operator and boundary value conditions,

$$\begin{aligned} D_{0+}^\alpha (\varphi_p(D_{0+}^\alpha u(t))) + a(t)f(u) &= 0, & \text{for } 0 < t < 1, \\ u(0) &= \gamma u(\xi) + \lambda, \\ \varphi_p(D_{0+}^\alpha u(0)) &= (\varphi_p(D_{0+}^\alpha u(1)))' = (\varphi_p(D_{0+}^\alpha u(0))), \end{aligned}$$

where  $0 < \alpha \leq 1, 2 < \beta \leq 3$ , and  $D_{0+}^\alpha, D_{0+}^\beta$  are Caputo fractional derivatives,  $\varphi_p(s) = |s|^{p-2}s, p > 1$ , and  $\varphi_p^{-1} = \varphi_q, \frac{1}{p} + \frac{1}{q} = 1$ , and the parameters are  $0 \leq \gamma < 1, 0 \leq \xi \leq 1, \lambda > 0$ . The continuous functions  $a : (0, 1) \rightarrow [0, \infty)$  and  $f : [0, \infty) \rightarrow [0, \infty)$  are given. The Green's function properties and the Schauder fixed point theorem are used.

In [2], Liu *et al.* studied the solvability of the Caputo fractional differential equation with boundary value conditions involving the  $p$ -Laplacian operator. The existence and uniqueness of the problem is found by the Banach fixed point theorem. The problem is given in the following:

$$(\varphi_p(D_{0+}^\alpha x(t)))' = f(t, x(t)), \quad \text{for } t \in (0, 1),$$

with boundary value conditions

$$\begin{aligned} x(0) &= r_0 x(1), \\ x'(0) &= r_1 x'(1), \\ x^{(i)}(0) &= 0, \end{aligned}$$

where  $i = 2, 3, \dots, [\alpha] - 1$ . Here,  $\varphi_p$  is the  $p$ -Laplacian operator and  $D_{0+}^\alpha$  is the Caputo fractional derivative,  $1 < \alpha \in R$ , and the nonlinear function  $f \in C([0, 1] \times R, R)$  is given.

In [7], Lu *et al.* studied the existence of nonnegative solutions of a nonlinear fractional boundary value problem with the  $p$ -Laplacian operator:

$$\begin{aligned} D_{0+}^\beta (\varphi_p(D_{0+}^\alpha u(t))) &= f(t, u(t)), & \text{for } 0 < t < 1, \\ u(0) = u'(0) = u'(1) &= 0, \\ D_{0+}^\alpha u(0) = D_{0+}^\alpha u(1) &= 0, \end{aligned}$$

where  $2 < \alpha \leq 3, 1 < \beta \leq 2$ , and  $D_{0+}^\alpha, D_{0+}^\beta$  are the standard Riemann-Liouville fractional derivatives. Green's functions, the Guo-Krasnoselskii theorem, and the Leggett-Williams fixed point theorems are used.

In [1], Wang and Xiang used upper and lower solutions method to find the existence results of at least one non-negative solution of the  $p$ -Laplacian fractional boundary value problem, which is given in the following:

$$\begin{aligned}
 D_{0+}^{\gamma}(\phi_p(D_{0+}^{\alpha}u(t))) &= f(t, u(t)), \quad \text{for } 0 < t < 1, \\
 u(0) = 0, \quad u'(1) &= au(\xi), \\
 D_{0+}^{\alpha}u(0) = 0, \quad D_{0+}^{\alpha}u(1) &= bD_{0+}^{\alpha}u(\eta),
 \end{aligned}$$

where  $1 < \alpha, \gamma \leq 2, 0 \leq a, b \leq 1, 0 < \xi, \eta < 1$ , and also  $D_{0+}^{\alpha}, D_{0+}^{\gamma}$  are Riemann-Liouville fractional operators.

In this paper, we focus on the existence of solutions of the fractional differential equation

$$D_{0+}^{\beta}\phi_p(D_{0+}^{\alpha}u(t)) = f(t, u(t), D_{0+}^{\gamma}u(t)), \tag{1}$$

with the  $p$ -Laplacian operator and integral boundary conditions,

$$\begin{aligned}
 u(0) + \mu_1u(1) &= \sigma_1 \int_0^1 g(s, u(s)) ds, \\
 u'(0) + \mu_2u'(1) &= \sigma_2 \int_0^1 h(s, u(s)) ds, \\
 D_{0+}^{\alpha}u(0) &= 0, \\
 D_{0+}^{\alpha}u(1) &= \nu D_{0+}^{\alpha}u(\eta),
 \end{aligned} \tag{2}$$

where  $D_{0+}^{\alpha}, D_{0+}^{\beta}$  are for the Caputo fractional differential equation with  $1 < \alpha \leq 2, 1 < \beta \leq 2$ ,  $\nu, \mu_i, \sigma_i (i = 1, 2)$  are non-negative parameters.  $f, g, h$  are continuous functions. By the Green's functions and fixed point theorems, we state and prove the existence and uniqueness results of the solutions. Two examples are given to illustrate the results.

### 2 Preliminaries

The basic definitions are given in the following.

**Definition 1** The Riemann-Liouville fractional integral of order  $\alpha > 0$  for a function  $f : (0, +\infty) \rightarrow R$  is defined as

$$I_{0+}^{\alpha}f(t) = \frac{1}{\Gamma(\alpha)} \int_0^t (t-s)^{\alpha-1}f(s) ds,$$

provided that the right hand side of the integral is pointwise defined on  $(0, +\infty)$  and  $\Gamma$  is the gamma function.

**Definition 2** The Caputo derivative of order  $\alpha > 0$  for a function  $f : (0, +\infty) \rightarrow R$  is written as

$$D_{0+}^{\alpha}f(t) = \frac{1}{\Gamma(n-\alpha)} \int_0^t (t-s)^{n-\alpha-1}f^{(n)}(s) ds,$$

where  $n = [\alpha] + 1, [\alpha]$  is the integral part of  $\alpha$ .

**Lemma 3** Let  $u \in C(0,1) \cap L^1(0,1)$  with the fractional derivative of order  $\alpha > 0$  that belongs to  $C(0,1) \cap L^1(0,1)$ . Then

$$I_{0+}^\alpha D_{0+}^\alpha u(t) = u(t) + c_1 t^{\alpha-1} + c_2 t^{\alpha-2} + \dots + c_n t^{\alpha-n},$$

for  $c_i \in R$  ( $i = 1, 2, \dots, n$ ), where  $n$  is the smallest integer greater than or equal to  $\alpha$ .

**Lemma 4** Let  $\alpha > 0$ . Then the differential equation  $D_{0+}^\alpha f(t) = 0$  has solutions

$$f(t) = k_0 + k_1 t + k_2 t^2 + \dots + k_{n-1} t^{n-1}$$

and

$$I_{0+}^\alpha D_{0+}^\alpha f(t) = f(t) + k_0 + k_1 t + k_2 t^2 + \dots + k_{n-1} t^{n-1},$$

where  $k_i \in R$  and  $i = 1, 2, \dots, n = [\alpha] + 1$ .

The Caputo fractional derivative of order  $n - 1 < \alpha < n$  for  $t^\gamma$  is given by

$$D_{0+}^\alpha t^\gamma = \begin{cases} \frac{\Gamma(\gamma+1)}{\Gamma(\gamma-\alpha+1)} t^{\gamma-\alpha}, & \gamma \in N \text{ and } \gamma \geq n \text{ or } \gamma \notin N \text{ and } \gamma > n - 1, \\ 0, & \gamma \in \{0, 1, \dots, n - 1\}. \end{cases} \tag{3}$$

Also, for brevity, we set

$$\begin{aligned} \omega_1 &= \frac{\sigma_1}{1 + \mu_1} - \frac{\sigma_2 \mu_1}{(1 + \mu_1)(1 + \mu_2)}, & \omega_2 &= \frac{\sigma_2}{1 + \mu_2}, \\ c_1(\eta) &= \frac{\nu^{p-1} \eta^{\beta-1}}{(1 - \nu^{p-1} \eta) \Gamma(\beta + 1)}, & L &= c t^{\beta-1} c_1(\eta). \end{aligned}$$

We use the following properties of the  $p$ -Laplacian operator:  $\phi_p(u) = |u|^{p-2} u$ ,  $p > 1$ , and  $\phi_p^{-1} = \phi_q$ ,  $\frac{1}{p} + \frac{1}{q} = 1$ .

(L1) If  $1 < p < 2$ ,  $uv > 0$ ,  $|u|, |v| \geq r > 0$ , then

$$|\phi_p(u) - \phi_p(v)| \leq (p - 1)r^{p-2} |u - v|.$$

(L2) If  $p > 2$ ,  $|u|, |v| \leq R$  then

$$|\phi_p(u) - \phi_p(v)| \leq (p - 1)R^{p-2} |u - v|.$$

We define two Green's functions  $G(t, s)$  and  $H(t, s)$ ,

$$G(t, s) = \begin{cases} \frac{(t-\tau)^{\alpha-1}}{\Gamma(\alpha)} - \left( \frac{\mu_1(1+\mu_2)+t\mu_2(1+\mu_1)}{(1+\mu_1)(1+\mu_2)} \right) \frac{(1-\tau)^{\alpha-1}}{\Gamma(\alpha)} \\ \quad + \frac{\mu_1\mu_2(1-\tau)^{\alpha-2}}{(1+\mu_1)(1+\mu_2)\Gamma(\alpha-1)}, & t \geq \tau, \\ -\left( \frac{\mu_1(1+\mu_2)+t\mu_2(1+\mu_1)}{(1+\mu_1)(1+\mu_2)} \right) \frac{(1-\tau)^{\alpha-1}}{\Gamma(\alpha)} \\ \quad + \frac{\mu_1\mu_2(1-\tau)^{\alpha-2}}{(1+\mu_1)(1+\mu_2)\Gamma(\alpha-1)}, & t \leq \tau, \end{cases}$$

and

$$H(t, s) = \begin{cases} \frac{[(t-s)]^{\beta-1}}{\Gamma(\beta)} - \frac{t(1-s)^{\beta-1}}{(1-\nu^{p-1}\eta)\Gamma(\beta)}, & 0 \leq s \leq t \leq 1, \eta \leq s, \\ \frac{[(t-s)]^{\beta-1}}{\Gamma(\beta)} - \frac{t(1-s)^{\beta-1}}{(1-\nu^{p-1}\eta)\Gamma(\beta)} + \frac{t\nu^{p-1}(\eta-s)^{\beta-1}}{(1-\nu^{p-1}\eta)\Gamma(\beta)}, & 0 \leq s \leq t \leq 1, \eta \geq s, \\ \frac{-t(1-s)^{\beta-1}}{(1-\nu^{p-1}\eta)\Gamma(\beta)}, & 0 \leq t \leq s \leq 1, \eta \leq s, \\ \frac{-t(1-s)^{\beta-1}}{(1-\nu^{p-1}\eta)\Gamma(\beta)} + \frac{t\nu^{p-1}(\eta-s)^{\beta-1}}{(1-\nu^{p-1}\eta)\Gamma(\beta)}, & 0 \leq t \leq s \leq 1, \eta \geq s. \end{cases}$$

**Lemma 5** *Let  $f, g, h \in C(0, 1)$ , and with  $1 < \alpha \leq 2$  we have the following fractional boundary value problem:*

$$D_{0+}^{\beta} \phi_p(D_{0+}^{\alpha} u(t)) = f(t), \tag{4}$$

$$\begin{cases} u(0) + \mu_1 u(1) = \sigma_1 \int_0^1 g(s) ds, \\ u'(0) + \mu_2 u'(1) = \sigma_2 \int_0^1 h(s) ds, \end{cases} \tag{5}$$

$$D_{0+}^{\alpha} u(0) = 0, \tag{6}$$

$$D_{0+}^{\alpha} u(1) = \nu D_{0+}^{\alpha} u(\eta),$$

it has a unique solution which is given by

$$(\mathcal{T}u)(t) = \int_0^t G(t, s) \phi_q \left( \int_0^1 H(t, \tau) f(\tau) d\tau \right) ds + \omega_1 + \omega_2 t,$$

with

$$\omega_1 = \frac{\sigma_1}{1 + \mu_1} - \frac{\sigma_2 \mu_1}{(1 + \mu_1)(1 + \mu_2)} \quad \text{and} \quad \omega_2 = \frac{\sigma_2}{1 + \mu_2}.$$

*Proof* By applying  $I_{0+}^{\beta}$  to both sides of (4), we get

$$\begin{aligned} \phi_p(D_{0+}^{\alpha} u(t)) &= \int_0^t \frac{(t-s)^{\beta-1}}{\Gamma(\beta)} f(s) ds - b_1 - b_2 t, \quad b_1, b_2 \in \mathbb{R}, \\ D_{0+}^{\alpha} u(t) &= \phi_q \left( \int_0^t \frac{(t-s)^{\beta-1}}{\Gamma(\beta)} f(s) ds - b_1 - b_2 t \right). \end{aligned}$$

Using the boundary conditions  $D_{0+}^{\alpha} u(0) = 0$  and  $D_{0+}^{\alpha} u(1) = \nu D_{0+}^{\alpha} u(\eta)$ , we have

$$\phi_q(-b_1) = 0 \implies b_1 = 0,$$

and secondly,

$$\begin{aligned} \phi_q \left( \int_0^1 \frac{(1-s)^{\beta-1}}{\Gamma(\beta)} f(s) ds - b_2 \right) &= \nu \phi_q \left( \int_0^{\eta} \frac{(\eta-s)^{\beta-1}}{\Gamma(\beta)} f(s) ds - b_2 \eta \right) \\ &= \phi_q \left( \nu^{\frac{1}{q-1}} \left( \int_0^{\eta} \frac{(\eta-s)^{\beta-1}}{\Gamma(\beta)} f(s) ds - b_2 \eta \right) \right). \end{aligned}$$

Moreover, since  $\phi_p$  is one-to-one,

$$I_{0+}^{\beta} f(1) - b_2 = \nu^{p-1} (I_{0+}^{\beta} f(\eta) - b_2 \eta) = \nu^{p-1} I_{0+}^{\beta} f(\eta) - \nu^{p-1} b_2 \eta,$$

$$I_{0+}^\beta f(1) - \nu^{p-1} I_{0+}^\beta f(\eta) = (1 - \nu^{p-1} \eta) b_2.$$

Then

$$\begin{aligned} b_2 &= \frac{1}{(1 - \nu^{p-1} \eta)} I_{0+}^\beta f(1) - \frac{\nu^{p-1}}{(1 - \nu^{p-1} \eta)} I_{0+}^\beta f(\eta) \\ &= \frac{1}{(1 - \nu^{p-1} \eta)} \int_0^1 \frac{(1-s)^{\beta-1}}{\Gamma(\beta)} f(s) ds - \frac{\nu^{p-1}}{(1 - \nu^{p-1} \eta)} \int_0^\eta \frac{(\eta-s)^{\beta-1}}{\Gamma(\beta)} f(s) ds. \end{aligned}$$

Since  $\phi_p(D_{0+}^\alpha u(t)) = I_{0+}^\beta f(t) - b_1 - b_2 t$ ,

$$\begin{aligned} \phi_p(D_{0+}^\alpha u(t)) &= \int_0^t \frac{(t-s)^{\beta-1}}{\Gamma(\beta)} f(s, u(s)) ds - \frac{t}{(1 - \nu^{p-1} \eta)} \int_0^1 \frac{(1-s)^{\beta-1}}{\Gamma(\beta)} f(s, u(s)) ds \\ &\quad + \frac{t \nu^{p-1}}{(1 - \nu^{p-1} \eta)} \int_0^\eta \frac{(\eta-s)^{\beta-1}}{\Gamma(\beta)} f(s, u(s)) ds \\ &= \int_0^1 H(t, s) f(s) ds, \\ D_{0+}^\alpha u(t) &= \phi_q \left( \int_0^1 H(t, s) f(s) ds \right), \\ u(t) &= \int_0^t \frac{(t-\tau)^{\alpha-1}}{\Gamma(\alpha)} \phi_q \left( \int_0^1 H(\tau, s) f(s) ds \right) d\tau - c_1 - c_2 t. \end{aligned} \tag{7}$$

By the boundary conditions (5), we get

$$\begin{aligned} -c_1 + \mu_1 \left( \int_0^1 \frac{(1-\tau)^{\alpha-1}}{\Gamma(\alpha)} \phi_p \left( \int_0^1 H(\tau, s) f(s) ds \right) d\tau - c_1 - c_2 \right) &= \sigma_1 \int_0^1 g(s) ds, \\ \mu_1 \int_0^1 \frac{(1-\tau)^{\alpha-1}}{\Gamma(\alpha)} \phi_p \left( \int_0^1 H(\tau, s) f(s) ds \right) d\tau - c_2 \mu_1 - \sigma_1 \int_0^1 g(s) ds &= c_1 (1 + \mu_1), \\ c_1 &= \frac{\mu_1}{(1 + \mu_1)} \int_0^1 \frac{(1-\tau)^{\alpha-1}}{\Gamma(\alpha)} \phi_p \left( \int_0^1 H(\tau, s) f(s) ds \right) d\tau - c_2 \frac{\mu_1}{(1 + \mu_1)} \\ &\quad - \frac{\sigma_1}{(1 + \mu_1)} \int_0^1 g(s) ds, \\ c_2 &= \frac{\mu_2}{(1 + \mu_2)} \int_0^1 \frac{(1-\tau)^{\alpha-2}}{\Gamma(\alpha-1)} \phi_p \left( \int_0^1 H(\tau, s) f(s) ds \right) d\tau - \frac{\sigma_2}{(1 + \mu_2)} \int_0^1 h(s) ds. \end{aligned} \tag{8}$$

Inserting  $c_2$  into (8), we get the values of  $c_1$ , and inserting  $c_1$  and  $c_2$  into (7), we have

$$\begin{aligned} u(t) &= \int_0^t \frac{(t-\tau)^{\alpha-1}}{\Gamma(\alpha)} \phi_p \left( \int_0^1 H(\tau, s) f(s, u(s)) ds \right) d\tau \\ &\quad - \left( \frac{\mu_1(1 + \mu_2) + t \mu_2(1 + \mu_1)}{(1 + \mu_1)(1 + \mu_2)} \right) \int_0^1 \frac{(1-\tau)^{\alpha-1}}{\Gamma(\alpha)} \phi_p \left( \int_0^1 H(\tau, s) f(s, u(s)) ds \right) d\tau \\ &\quad + \frac{\mu_1 \mu_2}{(1 + \mu_1)(1 + \mu_2)} \int_0^1 \frac{(1-\tau)^{\alpha-2}}{\Gamma(\alpha-1)} \phi_p \left( \int_0^1 H(\tau, s) f(s, u(s)) ds \right) d\tau \\ &\quad + \frac{\sigma_1}{1 + \mu_1} \int_0^1 g(s, u(s)) ds - \left( \frac{\sigma_2 \mu_1 - t \sigma_2(1 + \mu_1)}{(1 + \mu_1)(1 + \mu_2)} \right) \int_0^1 h(s, u(s)) ds. \end{aligned} \quad \square$$

**Lemma 6** *The functions  $G(t, s)$  and  $H(t, s)$  are continuous on  $[0, 1] \times [0, 1]$  and  $H(t, s)$  satisfies the following properties:*

- (1)  $H(t, s) \leq 0$ , for  $t, s \in [0, 1]$ ,
- (2)  $H(t, s) \geq H(s, s)$ , for  $t, s \in [0, 1]$ ,
- (3) *the Green's function  $H(t, s)$  satisfies the following condition:*

$$0 \leq \int_0^1 |H(t, s)| ds \leq \frac{B(\beta, \beta)}{(1 - \nu^{p-1}\eta)\Gamma(\beta)},$$

where  $B$  is the Beta function.

*Proof* The proofs of properties (1)-(2) are given in [1]. Thus we will prove property (3) for any  $t, s \in [0, 1]$ . The Green's function  $H(t, s)$  is negative. Therefore,

$$0 \leq \int_0^1 |H(t, s)| ds \leq \int_0^1 |H(s, s)| ds \leq \frac{B(\beta, \beta)}{(1 - \nu^{p-1}\eta)\Gamma(\beta)}. \quad \square$$

### 3 Existence and uniqueness results

In this section, we state and prove existence and uniqueness results of the fractional BVP (1)-(2) by using the Banach fixed point theorem. Our study concerns the space

$$C_\gamma([0, 1], R) = \{u \in C([0, 1], R), D_{0+}^\gamma u \in C([0, 1], R)\},$$

which is shown in the form

$$\|u\|_\gamma = \|u\|_c + \|D_{0+}^\gamma u\|_c,$$

where  $\|\cdot\|_c$  is the sup norm in  $C([0, 1], R)$ .

The following notations will be used throughout this paper:

$$\begin{aligned} \Delta_1 &= \frac{1}{\Gamma(\alpha + 1)} \left[ 1 + \frac{|\mu_1||1 + \mu_2| + |\mu_2||1 + \mu_1|}{|1 + \mu_1||1 + \mu_2|} \right] + \frac{1}{\Gamma(\alpha)} \left[ \frac{|\mu_1||\mu_2|}{|1 + \mu_1||1 + \mu_2|} \right], \\ \Delta_2 &= \frac{1}{\Gamma(\alpha - \gamma + 1)} \left[ 1 + \frac{|\mu_2|}{\Gamma(2 - \gamma)|1 + \mu_2|} \right], \\ \Delta_g &= \frac{|\sigma_1|}{|1 + \mu_1|}, \\ \Delta_{h_1} &= \frac{|\sigma_2||\mu_1 + |1 + \mu_1||}{|1 + \mu_1||1 + \mu_2|}, \quad \Delta_{h_2} = \frac{|\sigma_2|}{\Gamma(2 - \gamma)|1 + \mu_2|}. \end{aligned}$$

To state and prove our first result, we pose the following conditions:

- (A1) The function  $f : [0, 1] \times R \times R \rightarrow R$  is jointly continuous.
- (A2) There exists a function  $l_f \in L^{\frac{1}{\gamma}}([0, 1], R^+)$  such that

$$|f(t, u_1, u_2) - f(t, v_1, v_2)| \leq l_f(t)(|u_1 - v_1| + |u_2 - v_2|),$$

for all  $(t, u_1, u_2), (t, v_1, v_2) \in [0, 1] \times R \times R$ .

(A3) The functions  $g$  and  $h$  are jointly continuous and there exists  $l_g, l_h \in L^1([0, 1], R^+)$  such that

$$|g(t, u) - g(t, v)| \leq l_g(t)|u - v|$$

and

$$|h(t, u) - h(t, v)| \leq l_h(t)|u - v|,$$

for each  $(t, u), (t, v) \in [0, 1] \times R$ .

Next, we define an operator,  $\mathcal{T}_0$  which is  $\mathcal{T}_0 : C[0, 1] \rightarrow C[0, 1]$  as follows:

$$\mathcal{T}_0 x(t) = \phi_q \left( \int_0^1 H(t, s) f(s, x(s), D_{0+}^\gamma x(s)) ds \right).$$

**Lemma 7** Assume (A1)-(A3) hold and  $q > 2$ . There exists a constant  $l_{\mathcal{T}_0} > 0$  such that

$$|\mathcal{T}_0 u(t) - \mathcal{T}_0 v(t)| \leq l_{\mathcal{T}_0} \|u - v\|_\gamma,$$

for all  $u, v \in B_r$ . We have

$$l_{\mathcal{T}_0} = (q - 1)L_H^{q-2} \|l_f\|_\infty \int_0^1 |H(s, s)| ds \leq (q - 1)L_H^{q-2} \|l_f\|_\infty \frac{B(\beta, \beta)}{(1 - \nu^{p-1}\eta)\Gamma(\beta)}.$$

*Proof* If  $p > 2$  and  $t > 0$  we have the following estimation:

$$\begin{aligned} \left| \int_0^1 H(t, s) f(s, u(s), D_{0+}^\gamma u(s)) ds \right| &\leq \int_0^1 |H(t, s)| |f(s, u(s), D_{0+}^\gamma u(s))| ds \\ &\leq \int_0^1 |H(t, s)| l_f(s) (|u(s)| + |D_{0+}^\gamma u(s)| + |f(s, 0, 0)|) ds \\ &\leq (\|l_f\|_\infty \|u\|_\gamma + M) \int_0^1 |H(s, s)| ds \\ &\leq (\|l_f\|_\infty r + M) \int_0^1 |H(s, s)| ds \\ &= L_H, \end{aligned}$$

where  $M = \max_{s \in [0, 1]} |f(s, 0, 0)|$ . Now using the property (L2), we get the desired inequality,

$$\begin{aligned} &|(\mathcal{T}_0 u)(t) - (\mathcal{T}_0 v)(t)| \\ &= \left| \phi_q \left( \int_0^1 H(t, s) f(s, u(s), D_{0+}^\gamma u(s)) ds \right) - \phi_q \left( \int_0^1 H(t, s) f(s, v(s), D_{0+}^\gamma v(s)) ds \right) \right| \\ &\leq (q - 1)L_H^{q-2} \left| \int_0^1 H(t, s) (f(s, u(s), D_{0+}^\gamma u(s)) - f(s, v(s), D_{0+}^\gamma v(s))) ds \right| \\ &\leq (q - 1)L_H^{q-2} \|l_f\|_\infty \|u - v\|_\gamma \int_0^1 |H(s, s)| ds \end{aligned}$$



$$\begin{aligned} &\leq (q-1)L_H^{q-2}\|l_f\|_\infty \frac{B(\beta, \beta)}{(1-\nu^{p-1}\eta)\Gamma(\beta)}\|u-v\|_\gamma \\ &= l_{\mathcal{T}_0}\|u-v\|_\gamma. \end{aligned} \quad \square$$

**Theorem 8** Assume (A1)-(A3) hold. If

$$\left\{ l_{\mathcal{T}_0} \left( \sum_{i=1}^2 \Delta_i \right) + \Delta_g \|l_g\|_1 + \left( \sum_{i=1}^2 \Delta h_i \right) \|l_h\|_1 \right\} < 1, \tag{9}$$

then our BVP (1)-(2) has a unique solution on  $[0, 1]$ .

*Proof* Let us define the operator  $\mathcal{T} : C_\gamma([0, 1], R) \rightarrow C_\gamma([0, 1], R)$  to transform our BVP (1)-(2) into a fixed point problem,

$$\begin{aligned} (\mathcal{T}u)(t) &= \int_0^t \frac{(t-s)^{\alpha-1}}{\Gamma(\alpha)} \mathcal{T}_0(f(s, u(s), D_{0+}^\gamma u(s))) ds \\ &\quad - \frac{\mu_1}{(1+\mu_1)} \int_0^1 \frac{(1-s)^{\alpha-1}}{\Gamma(\alpha)} \mathcal{T}_0(f(s, u(s), D_{0+}^\gamma u(s))) ds \\ &\quad + \frac{\mu_1\mu_2}{(1+\mu_1)(1+\mu_2)} \int_0^1 \frac{(1-s)^{\alpha-2}}{\Gamma(\alpha-1)} \mathcal{T}_0(f(s, u(s), D_{0+}^\gamma u(s))) ds \\ &\quad - \frac{\mu_2 t}{(1+\mu_2)} \int_0^1 \frac{(1-s)^{\alpha-1}}{\Gamma(\alpha)} \mathcal{T}_0(f(s, u(s), D_{0+}^\gamma u(s))) ds \\ &\quad + \frac{\sigma_1}{(1+\mu_1)} \int_0^1 g(s, u(s)) ds - \frac{\sigma_2(\mu_1 - (1+\mu_1)t)}{(1+\mu_2)(1+\mu_1)} \int_0^1 h(s, u(s)) ds. \end{aligned} \tag{10}$$

Taking the  $\gamma$ th fractional derivative, we get

$$\begin{aligned} D_{0+}^\gamma(\mathcal{T}u)(t) &= \int_0^t \frac{(t-s)^{\alpha-\gamma-1}}{\Gamma(\alpha-\gamma)} \mathcal{T}_0(f(s, u(s), D_{0+}^\gamma u(s))) ds \\ &\quad - \frac{\mu_2}{(1+\mu_2)} \frac{t^{1-\gamma}}{\Gamma(2-\gamma)} \int_0^1 \frac{(1-s)^{\alpha-\gamma-1}}{\Gamma(\alpha-\gamma)} \mathcal{T}_0(f(s, u(s), D_{0+}^\gamma u(s))) ds \\ &\quad + \frac{\sigma_2}{(1+\mu_2)} \frac{t^{1-\gamma}}{\Gamma(2-\gamma)} \int_0^1 h(s, u(s)) ds \end{aligned} \tag{11}$$

for  $t \in [0, 1]$ . Since  $f, g, h$  are continuous, the expression (10) and (11) are well defined. Clearly, the fixed point of the operator  $\mathcal{T}$  is the solution of the problem (1)-(2). To show the existence and uniqueness of the solution, the Banach fixed point theorem is used and then we show  $\mathcal{T}$  is contraction. We have

$$\begin{aligned} &|(\mathcal{T}u)(t) - (\mathcal{T}v)(t)| \\ &\leq \int_0^t \frac{(t-s)^{\alpha-1}}{\Gamma(\alpha)} l_{\mathcal{T}_0} \|u-v\|_\gamma ds \\ &\quad + \frac{|\mu_1|}{|1+\mu_1|} \int_0^1 \frac{(1-s)^{\alpha-1}}{\Gamma(\alpha)} l_{\mathcal{T}_0} \|u-v\|_\gamma ds \end{aligned}$$

$$\begin{aligned}
 & + \frac{|\mu_1||\mu_2|}{|1 + \mu_1||1 + \mu_2|} \int_0^1 \frac{(1-s)^{\alpha-2}}{\Gamma(\alpha-1)} l_{\mathcal{T}_0} \|u-v\|_\gamma ds \\
 & + \frac{|\mu_2|}{|1 + \mu_2|} \int_0^1 \frac{(1-s)^{\alpha-1}}{\Gamma(\alpha)} l_{\mathcal{T}_0} \|u-v\|_\gamma ds \\
 & + \frac{|\sigma_1|}{|1 + \mu_1|} \int_0^1 l_g(s) (|u(s)-v(s)| + |D_{0+}^\gamma u(s) - D_{0+}^\gamma v(s)|) ds \\
 & + \frac{|\sigma_2||\mu_1 + |1 + \mu_1||}{|1 + \mu_2||1 + \mu_1|} \int_0^1 l_h(s) (|u(s)-v(s)| + |D_{0+}^\gamma u(s) - D_{0+}^\gamma v(s)|) ds \\
 \leq & \left\{ l_{\mathcal{T}_0} \left( \int_0^t \frac{(t-s)^{\alpha-1}}{\Gamma(\alpha)} ds + \frac{|\mu_1||1 + \mu_2| + |\mu_2||1 + \mu_1|}{|1 + \mu_1||1 + \mu_2|} \int_0^1 \frac{(1-s)^{\alpha-1}}{\Gamma(\alpha)} ds \right. \right. \\
 & + \frac{|\mu_1||\mu_2|}{|1 + \mu_1||1 + \mu_2|} \int_0^1 \frac{(1-s)^{\alpha-2}}{\Gamma(\alpha-1)} ds \Big) \\
 & + \frac{|\sigma_1|}{|1 + \mu_1|} \int_0^1 l_g(s) ds + \frac{|\sigma_2||\mu_1 + |1 + \mu_1||}{|1 + \mu_2||1 + \mu_1|} \int_0^1 l_h(s) ds \Big\} \|u-v\|_\gamma \\
 \leq & \left\{ l_{\mathcal{T}_0} \left( \frac{1}{\Gamma(\alpha+1)} + \frac{|\mu_1||1 + \mu_2| + |\mu_2||1 + \mu_1|}{\Gamma(\alpha+1)|1 + \mu_1||1 + \mu_2|} + \frac{|\mu_1||\mu_2|}{\Gamma(\alpha)|1 + \mu_1||1 + \mu_2|} \right) \right. \\
 & \left. + \frac{|\sigma_1|}{|1 + \mu_1|} \|l_g\|_1 + \frac{|\sigma_2||\mu_1 + |1 + \mu_1||}{|1 + \mu_2||1 + \mu_1|} \|l_h\|_1 \right\} \|u-v\|_\gamma. \tag{12}
 \end{aligned}$$

By using the Hölder inequality, we get

$$\begin{aligned}
 |\mathcal{T}u(t) - \mathcal{T}v(t)| & \leq \{ l_{\mathcal{T}_0} \Delta_1 + \Delta_g \|l_g\|_1 + \Delta_{h_1} \|l_h\|_1 \} \|u-v\|_\gamma, \tag{13} \\
 |D_{0+}^\gamma (\mathcal{T}u)(t) - D_{0+}^\gamma (\mathcal{T}v)(t)| & \\
 \leq & \int_0^t \frac{(t-s)^{\alpha-\gamma-1}}{\Gamma(\alpha-\gamma)} l_{\mathcal{T}_0} \|u-v\|_\gamma ds \\
 & + \frac{|\mu_2|}{|1 + \mu_2|} \frac{t^{1-\gamma}}{\Gamma(2-\gamma)} \int_0^1 \frac{(1-s)^{\alpha-\gamma-1}}{\Gamma(\alpha-\gamma)} l_{\mathcal{T}_0} \|u-v\|_\gamma ds \\
 & + \frac{|\sigma_2|}{|1 + \mu_2|} \frac{t^{1-\gamma}}{\Gamma(2-\gamma)} \int_0^1 l_h(s) (|u(s)-v(s)| + |D_{0+}^\gamma u(s) - D_{0+}^\gamma v(s)|) ds \\
 \leq & \left\{ \frac{l_{\mathcal{T}_0}}{\Gamma(\alpha-\gamma)} \int_0^t (t-s)^{\alpha-\gamma-1} ds \right. \\
 & + \frac{l_{\mathcal{T}_0} t^{1-\gamma} |\mu_2|}{\Gamma(\alpha-\gamma)\Gamma(2-\gamma)|1 + \mu_2|} \int_0^1 (1-s)^{\alpha-\gamma-1} ds \\
 & \left. + \frac{|\sigma_2| t^{1-\gamma}}{|1 + \mu_2|\Gamma(2-\gamma)} \int_0^1 l_h(s) ds \right\} \|u-v\|_\gamma \\
 \leq & \left\{ l_{\mathcal{T}_0} \left( \frac{1}{\Gamma(\alpha-\gamma+1)} + \frac{|\mu_2|}{\Gamma(\alpha-\gamma+1)\Gamma(2-\gamma)|1 + \mu_2|} \right) \right. \\
 & \left. + \frac{|\sigma_2|}{|1 + \mu_2|\Gamma(2-\gamma)} \int_0^1 l_h(s) ds \right\} \|u-v\|_\gamma \\
 \leq & \left\{ \frac{l_{\mathcal{T}_0}}{\Gamma(\alpha-\gamma+1)} \left( 1 + \frac{|\mu_2|}{\Gamma(2-\gamma)|1 + \mu_2|} \right) \right. \\
 & \left. + \frac{|\sigma_2|}{|1 + \mu_2|\Gamma(2-\gamma)} \|l_h\|_1 \right\} \|u-v\|_\gamma. \tag{14}
 \end{aligned}$$

Similarly,

$$|D_{0+}^\gamma (\mathcal{T}u(t)) - D_{0+}^\gamma (\mathcal{T}v(t))| \leq \{l_{\mathcal{T}_0} \Delta_2 + \Delta_{h_2} \|l_h\|_1\} \|u - v\|_\gamma. \tag{15}$$

With the help of (13)-(15), we find that

$$\begin{aligned} & \|Tu - Tv\|_\gamma \\ & \leq \{l_{\mathcal{T}_0}(\Delta_1 + \Delta_2) + \Delta_g \|l_g\|_1 + (\Delta_{h_1} + \Delta_{h_2}) \|l_h\|_1\} \|u - v\|_\gamma \\ & = \left\{ l_{\mathcal{T}_0} \left( \sum_{i=1}^2 \Delta_i \right) + \Delta_g \|l_g\|_1 + \left( \sum_{i=1}^2 \Delta_{h_i} \right) \|l_h\|_1 \right\} \|u - v\|_\gamma. \end{aligned}$$

Thus  $\mathcal{T}$  is a contraction mapping by condition (9). By the Banach fixed point theorem,  $\mathcal{T}$  has a fixed point which is the solution of the BVP. □

### 4 Existence results

**Theorem 9** *Assume:*

(iv) *There exist non-decreasing functions  $\varphi : [0, \infty) \times [0, \infty) \rightarrow [0, \infty)$  and  $\psi_i : [0, \infty) \rightarrow [0, \infty)$ ,  $i = 1, 2$ , and the functions  $l_f \in L^{\frac{1}{\alpha}}([0, 1], R^+)$  and  $l_g, l_h \in L^1([0, 1], R^+)$  such that*

$$\begin{aligned} |f(t, u, v)| & \leq l_f(t) \varphi(|u| + |v|), \\ |g(t, u)| & \leq l_g(t) \psi_1(|u|), \\ |h(t, u)| & \leq l_h(t) \psi_2(|u|), \end{aligned}$$

for all  $t \in [0, 1]$  and  $u, v \in R$ .

(v) *There exists a constant  $\mathcal{N} > 0$  such that*

$$\left[ \frac{\mathcal{N}}{\varphi(\|u\|_\gamma) l_{\mathcal{T}_0} \sum_{i=1}^2 \Delta_i + \psi_1(\|u\|_\gamma) \|l_g\|_1 \Delta_g + \psi_2(\|u\|_\gamma) \|l_h\|_1 \sum_{i=1}^2 \Delta_{h_i}} \right] > 1. \tag{16}$$

Thus problem (1)-(2) has at least one solution on  $[0, 1]$ .

*Proof* Let  $B_r = \{u \in C_\gamma([0, 1], R) : \|u\|_\gamma \leq r\}$ .

*Step 1:* Let the operator  $\mathcal{T} : C_\gamma([0, 1], R) \rightarrow C_\gamma([0, 1], R)$  be given in (10) which defines  $B_r$  to be a bounded set. For all  $u \in B_r$ , we get

$$\begin{aligned} & |(\mathcal{T}u)(t)| \\ & \leq \frac{\varphi(r)}{\Gamma(\alpha)} l_{\mathcal{T}_0} \int_0^t (t-s)^{\alpha-1} ds \\ & \quad + \frac{|\mu_1| |1 + \mu_2| + |\mu_2| |1 + \mu_1|}{|1 + \mu_1| |1 + \mu_2|} \frac{\varphi(r)}{\Gamma(\alpha)} l_{\mathcal{T}_0} \int_0^1 (1-s)^{\alpha-1} ds \\ & \quad + \frac{|\mu_1| |\mu_2|}{|1 + \mu_1| |1 + \mu_2|} \frac{\varphi(r)}{\Gamma(\alpha - 1)} l_{\mathcal{T}_0} \int_0^1 (1-s)^{\alpha-2} ds \\ & \quad + \frac{|\sigma_1|}{|1 + \mu_1|} \psi_1(r) \int_0^1 |l_g(s)| ds + \frac{|\sigma_2| |\mu_1 + |1 + \mu_1||}{|1 + \mu_2| |1 + \mu_1|} \psi_2(r) \int_0^1 |l_h(s)| ds \end{aligned}$$

and

$$\begin{aligned}
 & |D_{0+}^{\gamma}(\mathcal{T}u)(t)| \\
 & \leq \frac{\varphi(r)}{\Gamma(\alpha - \gamma)} l_{\mathcal{T}_0} \int_0^t (t - s)^{\alpha - \gamma - 1} ds \\
 & \quad + \frac{|\mu_2|}{|1 + \mu_2|} \frac{t^{1 - \gamma}}{\Gamma(2 - \gamma)} \frac{\varphi(r)}{\Gamma(\alpha - \gamma)} l_{\mathcal{T}_0} \int_0^1 (1 - s)^{\alpha - \gamma - 1} ds \\
 & \quad + \frac{|\sigma_2|}{|1 + \mu_2|} \frac{t^{1 - \gamma}}{\Gamma(2 - \gamma)} \psi_2(r) \int_0^1 |l_h(s)| ds.
 \end{aligned}$$

By the Hölder inequality,

$$\begin{aligned}
 & |(\mathcal{T}u)(t)| \\
 & \leq \frac{\varphi(r)l_{\mathcal{T}_0}}{\Gamma(\alpha + 1)} + \frac{(|\mu_1||1 + \mu_2| + |\mu_2||1 + \mu_1|)\varphi(r)l_{\mathcal{T}_0}}{|1 + \mu_1||1 + \mu_2|\Gamma(\alpha + 1)} + \frac{|\mu_1||\mu_2|\varphi(r)l_{\mathcal{T}_0}}{|1 + \mu_1||1 + \mu_2|\Gamma(\alpha)} \\
 & \quad + \frac{|\sigma_1|\psi_1(r)\|l_g\|_1}{|1 + \mu_1|} + \frac{|\sigma_2||\mu_1 + |1 + \mu_1||\psi_2(r)\|l_h\|_1}{|1 + \mu_2||1 + \mu_1|} \\
 & \leq \varphi(r)l_{\mathcal{T}_0} \Delta_1 + \Delta_g \psi_1(r)\|l_g\|_1 + \Delta_{h_1} \psi_2(r)\|l_h\|_1, \\
 & |D_{0+}^{\gamma}(\mathcal{T}u)(t)| \\
 & \leq \frac{\varphi(r)l_{\mathcal{T}_0}}{\Gamma(\alpha - \gamma + 1)} + \frac{|\mu_2|\varphi(r)l_{\mathcal{T}_0}}{|1 + \mu_2|\Gamma(2 - \gamma)\Gamma(\alpha - \gamma + 1)} + \frac{|\sigma_2|\psi_2(r)\|l_h\|_1}{|1 + \mu_2|\Gamma(2 - \gamma)} \\
 & \leq \varphi(r)l_{\mathcal{T}_0} \Delta_2 + \Delta_{h_2} \psi_2(r)\|l_h\|_1.
 \end{aligned}$$

Therefore,

$$\|(\mathcal{T}u)\|_{\gamma} \leq \varphi(r)l_{\mathcal{T}_0}(\Delta_1 + \Delta_2) + \Delta_g \psi_1(r)\|l_g\|_1 + (\Delta_{h_1} + \Delta_{h_2})\psi_2(r)\|l_h\|_1.$$

Step 2: The families  $\{(\mathcal{T}u) : u \in B_r\}$  and  $\{D_{0+}^{\gamma}(\mathcal{T}u) : u \in B_r\}$  are equicontinuous. For  $t_1 < t_2$ , we get

$$\begin{aligned}
 & |(\mathcal{T}u)(t_2) - (\mathcal{T}u)(t_1)| \\
 & \leq \frac{\varphi(r)l_{\mathcal{T}_0}}{\Gamma(\alpha)} \left[ \int_0^{t_1} ((t_1 - s)^{\alpha - 1} + (t_2 - s)^{\alpha - 1}) ds - \int_{t_1}^{t_2} (t_2 - s)^{\alpha - 1} ds \right] \\
 & \quad + \frac{|\mu_2||t_2 - t_1|}{|1 + \mu_2|} \frac{\varphi(r)l_{\mathcal{T}_0}}{\Gamma(\alpha)} \int_0^{t_1} (1 - s)^{\alpha - 1} ds \\
 & \quad + \frac{|\sigma_2||1 + \mu_1||t_2 - t_1|\psi_2(r)}{|\mu_2||1 + \mu_1|} \int_0^1 |l_h(s)| ds \rightarrow 0 \quad \text{as } t_2 \rightarrow t_1.
 \end{aligned}$$

Similarly,

$$\begin{aligned}
 & |D_{0+}^{\gamma}(\mathcal{T}u)(t_2) - D_{0+}^{\gamma}(\mathcal{T}u)(t_1)| \\
 & \leq \frac{\varphi(r)l_{\mathcal{T}_0}}{\Gamma(\alpha - \gamma)} \left[ \int_0^{t_1} ((t_1 - s)^{\alpha - \gamma - 1} + (t_2 - s)^{\alpha - \gamma - 1}) ds - \int_{t_1}^{t_2} (t_2 - s)^{\alpha - \gamma - 1} ds \right]
 \end{aligned}$$

$$\begin{aligned}
 &+ \frac{\varphi(r)l_{\mathcal{T}_0}|\mu_2|t_2^{1-\gamma} - t_1^{1-\gamma}}{\Gamma(\alpha-\gamma)|1 + \mu_2|\Gamma(2-\gamma)} \int_0^1 (1-s)^{\alpha-\gamma-1} ds \\
 &+ \frac{|\sigma_2|t_2^{1-\gamma} - t_1^{1-\gamma}|\psi_2(r)}{|1 + \mu_2|\Gamma(2-\gamma)} \int_0^1 |l_h(s)| ds \rightarrow 0 \quad \text{as } t_2 \rightarrow t_1.
 \end{aligned}$$

Thus  $\{(\mathcal{T}u) : u \in B_r\}$  and  $\{D_{0+}^\gamma(\mathcal{T}u) : u \in B_r\}$  are equicontinuous and relatively compact in  $C([0, 1], R)$  by the Arzela-Ascoli theorem. Therefore  $\mathcal{T}(B_r)$  is a relatively compact subset of  $C_\gamma([0, 1], R)$  and the operator  $\mathcal{T}$  is compact.

*Step 3:* Let  $u = \lambda(\mathcal{T}u)$  and  $u = \lambda(D_{0+}^\gamma(\mathcal{T}u))$  for  $0 < \lambda < 1$ . For each  $t \in [0, 1]$ , define  $\overline{\mathcal{M}} = \{\|u\|_\gamma \in C_\gamma([0, 1], R), \|u\|_\gamma < \mathcal{N}\}$ . Then we get

$$\begin{aligned}
 \|u\|_c &= \|\lambda(\mathcal{T}u)\|_c \\
 &\leq \varphi(\|u\|_\gamma)l_{\mathcal{T}_0}\Delta_1 + \Delta_g\psi_1(\|u\|_\gamma)\|l_g\|_1 + \Delta_{h_1}\psi_2(\|u\|_\gamma)\|l_h\|_1, \\
 \|u\|_c &= \|\lambda(D_{0+}^\gamma(\mathcal{T}u))\|_c \\
 &\leq \varphi(\|u\|_\gamma)l_{\mathcal{T}_0}\Delta_2 + \Delta_{h_2}\psi_2(\|u\|_\gamma)\|l_h\|_1.
 \end{aligned}$$

Thus

$$\|u\|_\gamma \leq \varphi(\|u\|_\gamma)l_{\mathcal{T}_0} \sum_{i=1}^2 \Delta_i + \psi_1(\|u\|_\gamma)\|l_g\|_1\Delta_g + \psi_2(\|u\|_\gamma)\|l_h\|_1 \sum_{i=1}^2 \Delta_{h_i}.$$

That means

$$\frac{\|u\|_\gamma}{\varphi(\|u\|_\gamma)l_{\mathcal{T}_0} \sum_{i=1}^2 \Delta_i + \psi_1(\|u\|_\gamma)\|l_g\|_1\Delta_g + \psi_2(\|u\|_\gamma)\|l_h\|_1 \sum_{i=1}^2 \Delta_{h_i}} \leq 1.$$

For a non-negative  $\mathcal{N}$  and  $\|u\|_\gamma < \mathcal{N}$ , the operator  $\mathcal{T}$  which is defined in  $\overline{\mathcal{M}}$  to be  $C_\gamma([0, 1], R)$  is continuous and compact. Therefore  $\mathcal{T}$  has a fixed point in  $\overline{\mathcal{M}}$ . □

### 5 Examples

**Example 10** Consider the following boundary value problem of a fractional differential equation:

$$\begin{cases}
 D_{0+}^{\frac{3}{2}}(\phi_p D_{0+}^{\frac{3}{2}} u)(t) = l_f\left(\frac{|u(t)|}{|u(t)|+1} + \frac{|D_{0+}^{\frac{3}{2}} u(t)|}{|D_{0+}^{\frac{3}{2}} u(t)|+1}\right), \\
 u(0) + 0.1u(1) = 0.01 \int_0^1 \frac{u(s)}{(1+s)^2} ds, \\
 u'(0) + 0.1u'(1) = 0.01 \int_0^1 \left(\frac{e^s u(s)}{1+2e^s} + \frac{1}{2}\right) ds.
 \end{cases} \tag{17}$$

Here

$$\begin{aligned}
 \alpha, \beta &= 1.5, & \mu_1, \mu_2 &= 0.1, & \sigma_1, \sigma_2 &= 0.01, \\
 \nu, \eta &= 0.3, & \tau &= 0.4, & \gamma &= 0.01,
 \end{aligned}$$

and

$$f(t, u, \nu) = \frac{|u|}{|u| + 1} + \frac{|\nu|}{|\nu| + 1},$$

$$g(t, u) = \frac{u}{(1+s)^2}, \quad h(t, u) = \frac{e^s u}{(1+2e^s)} + \frac{1}{2}.$$

Since  $0.88 < \Gamma(1.5) < 0.89$ ,  $\Gamma(2) = 1$ ,  $\Gamma(2.5) = 1$ , we find

$$\begin{aligned} \Delta_1 &= 0.89, & \Delta_2 &= 0.82, & \Delta_g &= 0.009, \\ \Delta_{h_1} &= 0.0099, & \Delta_{h_2} &= 0.009, & l_g &= l_h = 1, \end{aligned}$$

with simple calculations. Therefore

$$\begin{aligned} &\{l_{\mathcal{T}_0}(\Delta_1 + \Delta_2) + 2\Delta_g \|l_g\|_1 + (\Delta_{h_1} + \Delta_{h_2}) \|l_h\|_1\} \\ &< 1.73l_{\mathcal{T}_0} + 0.04 \\ &< 1. \end{aligned}$$

Then we can choose

$$l_{\mathcal{T}_0} < 0.562.$$

Thus all assumptions of Theorem 8 satisfied. Therefore the problem has a unique solution on  $[0, 1]$ .

**Example 11** Consider the following boundary value problem of fractional differential equation:

$$\begin{cases} D_{0+}^{\frac{3}{2}}(\phi_p D_{0+}^{\frac{3}{2}} u)(t) = \frac{|u(t)|^3}{9(|u(t)|^3+3)} + \frac{|\sin D_{0+}^{\frac{3}{2}} u(t)|}{9(\sin D_{0+}^{\frac{3}{2}} u(t)+1)} + \frac{1}{12}, \\ u(0) + 0.1u(1) = 0, 01 \int_0^1 \frac{u(s)}{3(1+s)^2} ds, \\ u'(0) + 0.1u'(1) = 0, 01 \int_0^1 \frac{e^s u(s)}{3(1+e^s)^2} ds, \\ D_{0+}^{\frac{3}{2}} u(0) = 0, \\ D_{0+}^{\frac{3}{2}} u(1) = 0, 3D_{0+}^{\frac{3}{2}} u(0, 3), \end{cases} \tag{18}$$

where  $f$  is given by

$$f(t, u, v) = \frac{|u|^3}{9(|u|^3+3)} + \frac{|\sin v|}{9(\sin v+1)} + \frac{1}{12}.$$

We have

$$|f(t, u, v)| \leq \frac{|u|^3}{9(|u|^3+3)} + \frac{|\sin v|}{9(\sin v+1)} + \frac{1}{12}, \quad u \in R.$$

Here

$$\begin{aligned} \alpha, \beta &= 1.5, & \mu_1, \mu_2 &= 0.1, & \sigma_1, \sigma_2 &= 0.01, \\ \nu, \eta &= 0.3, & \tau &= 0.4, & \gamma &= 0.01, \\ \Delta_1 &= 0.89, & \Delta_2 &= 0.82, & \Delta_g &= 0.009, \end{aligned}$$

$$\Delta_{h_1} = 0.0099, \quad \Delta_{h_2} = 0.009, \quad l_g = l_h = 0.1,$$

and  $g(t, u) = \frac{u(s)}{3(1+s)^2}$ ,  $h(t, u) = \frac{e^s u(s)}{3(1+e^s)^2}$ ,  $\varphi(\mathcal{N}) = \psi_1(\mathcal{N}) = \psi_2(\mathcal{N}) = \mathcal{N}$ . If

$$\frac{\mathcal{N}}{\varphi(\mathcal{N})(0.561)(0.89 + 0.82) + \psi_1(\mathcal{N})(0.1)(0.009) + \psi_2(\mathcal{N})(0.1)(0.0099 + 0.009)} > 1,$$

$$\frac{\mathcal{N}}{\mathcal{N}(0.96) + \mathcal{N}(0.0009) + \mathcal{N}(0.0019)} > 1,$$

$$\frac{\mathcal{N}}{0.9628\mathcal{N}} > 1,$$

$$1.04 > 1,$$

then (16) is satisfied. Then there exists at least one solution of the BVP on  $[0, 1]$ .

**Competing interests**

The authors declare that they have no competing interests.

**Authors' contributions**

All authors contributed equally to the manuscript and typed, read, and approved the final manuscript.

**Acknowledgements**

The authors would like to thank the anonymous referees for their valuable comments and suggestions.

Received: 19 March 2015 Accepted: 25 May 2015 Published online: 14 June 2015

**References**

1. Wang, J, Xiang, H: Upper and lower solutions method for a class of singular fractional boundary value problems with  $p$ -Laplacian operator. *Abstr. Appl. Anal.* **2010**, 971824 (2010)
2. Liu, X, Jia, M, Xiang, X: On the solvability of a fractional differential equation model involving the  $p$ -Laplacian operator. *Comput. Math. Appl.* **64**, 3267-32777 (2012)
3. Ahmad, B: Nonlinear fractional differential equations with anti-periodic type fractional boundary conditions. *Differ. Equ. Dyn. Syst.* **21**(4), 387-401 (2013)
4. Zhang, X, Liu, L, Wiwatanapataphee, B, Wu, Y: The eigenvalue for a class of singular  $p$ -Laplacian fractional differential equations involving the Riemann-Stieltjes integral boundary condition. *Appl. Math. Comput.* **235**, 412-422 (2014)
5. Su, Y, Li, Q, Liu, X: Existence criteria for positive solutions of  $p$ -Laplacian fractional differential equations with derivative terms. *Adv. Differ. Equ.* **2013**, 119 (2013)
6. Han, Z, Lu, H, Sun, S, Yang, D: Positive solutions to boundary-value problems of  $p$ -Laplacian fractional differential equations with a parameter in the boundary. *Electron. J. Differ. Equ.* **2012**, 213 (2012)
7. Han, Z, Lu, H, Sun, S, Liu, J: Existence on positive solutions for boundary value problems of nonlinear fractional differential equations with  $p$ -Laplacian. *Adv. Differ. Equ.* **2013**, 30 (2013). doi:10.1186/1687-1847-2013-30
8. Zhang, J, Liangand, S: Existence and uniqueness of positive solutions for integral boundary problems of nonlinear fractional differential equations with  $p$ -Laplacian operator. *Rocky Mt. J. Math.* **44**(3), 953-974 (2014)
9. Khan, RA, Khan, A, Samad, A, Khan, H: On existence of solutions for fractional differential equation with  $p$ -Laplacian operator. *J. Fract. Calc. Appl.* **5**(2), 28-37 (2014)
10. Wang, L, Zhou, Z, Zhou, H: Positive solutions for singular  $p$ -Laplacian fractional differential system with integral boundary conditions. *Abstr. Appl. Anal.* **2014**, 984875 (2014)
11. Wei, L, Duan, L, Agarwal, RP: Existence and uniqueness of the solution to integro-differential equation involving the generalized  $p$ -Laplacian operator with mixed boundary conditions. *J. Math.* **33**, 1009-1018 (2013)
12. Wei, L, Agarwal, RP, Wong, PJY: Study on integro-differential equation with generalized  $p$ -Laplacian operator. *Bound. Value Probl.* **2012**, 131 (2012)
13. Wei, L, Agarwal, RP: Discussion on the existence of solutions to nonlinear boundary value problems with generalized  $p$ -Laplacian operator. *Acta Math. Sci. Ser. A Chin. Ed.* **32**(1), 201-211 (2012)
14. Wei, L, Zhou, H, Agarwal, RP: Existence of solutions to nonlinear Neumann boundary value problems with  $p$ -Laplacian operator and iterative construction. *Acta Math. Appl. Sinica (Engl. Ser.)* **27**, 463-470 (2011)
15. Wei, L, Agarwal, RP, Wong, PJY: Existence of solutions to nonlinear parabolic boundary value problems with generalized  $p$ -Laplacian operator. *Adv. Math. Sci. Appl.* **20**(2), 423-445 (2010)
16. Wei, L, Agarwal, RP: Existence of solutions to nonlinear Neumann boundary value problems with generalized  $p$ -Laplacian operator. *Comput. Math. Appl.* **56**(2), 530-541 (2008)
17. Agarwal, RP, O'Regan, D, Papageorgiun, NS: On the existence of two nontrivial solutions of periodic problems with operators of  $p$ -Laplacian type. *Differ. Equ.* **43**(2), 157-163 (2007)
18. Jiang, DQ, O'Regan, D, Agarwal, RP: A generalized upper and lower solution method for singular discrete boundary value problems for the one-dimensional  $p$ -Laplacian. *J. Appl. Anal.* **11**(1), 35-47 (2005)
19. Aktuğlu, H, Özarslan, MA: Solvability of differential equations of order  $2 < \alpha \leq 3$  involving the  $p$ -Laplacian operator with boundary conditions. *Adv. Differ. Equ.* **2013**, 358 (2013)

20. Kong, X, Wang, D, Li, H: Existence of unique positive solution to a two-point boundary value problem of fractional-order switched system with  $p$ -Laplacian operator. *J. Fract. Calc. Appl.* **5**(2), 9-16 (2014)
21. Khan, RA, Khan, A, Samad, A, Khan, H: On existence of the solution for fractional differential equation with  $p$ -Laplacian operator. *J. Fract. Calc. Appl.* **5**(2), 28-37 (2014)
22. Hu, Z, Liu, W, Liu, J: Existence of solutions for a coupled system of fractional  $p$ -Laplacian equations at resonance. *Adv. Differ. Equ.* **2013**, 312 (2013)
23. Chang, X, Qiao, Y: Existence of periodic solutions for a class of  $p$ -Laplacian operators. *Bound. Value Probl.* **2013**, 96 (2013)
24. Chen, T, Liu, W: Solvability of some boundary value problems for fractional  $p$ -Laplacian equation. *Abstr. Appl. Anal.* **2013**, 432509 (2013)
25. Chen, T, Liu, W, Liu, J: Existence of solutions for some boundary value problems of fractional  $p$ -Laplacian equation at resonance. *Bull. Belg. Math. Soc. Simon Stevin* **20**, 503-517 (2013)
26. Liu, Y, Qian, T: Existence of solutions of periodic-type boundary value problems for multi-term fractional differential equations. *Math. Methods Appl. Sci.* (2012). doi:10.1002/mma.2746
27. Tang, X, Yang, C, Liu, Q: Existence of the solutions of two-point boundary value problems for fractional  $p$ -Laplacian differential equations at resonance. *J. Appl. Math. Comput.* **41**, 119-131 (2013). doi:10.1007/s12190-012-0598-0

**Submit your manuscript to a SpringerOpen<sup>®</sup> journal and benefit from:**

- ▶ Convenient online submission
- ▶ Rigorous peer review
- ▶ Immediate publication on acceptance
- ▶ Open access: articles freely available online
- ▶ High visibility within the field
- ▶ Retaining the copyright to your article

---

Submit your next manuscript at ▶ [springeropen.com](http://springeropen.com)

---