

RESEARCH

Open Access



Studies on a $2n$ th-order p -Laplacian differential equation with singularity

Yun Xin and Shan Zhao*

*Correspondence:
chengdanxin2008@126.com
College of Computer Science and
Technology, Henan Polytechnic
University, Jiaozuo, 454000, China

Abstract

In this paper, we consider the $2n$ th-order p -Laplacian differential equation with singularity

$$(\varphi_p(x(t))^{(n)})^{(n)} + f(x(t))x'(t) + g(t, x(t - \sigma)) = e(t).$$

By applications of coincidence degree theory and some analysis techniques, sufficient conditions for the existence of positive periodic solutions are established.

MSC: 34C25; 34K13; 34K40

Keywords: positive periodic solution; p -Laplacian; $2n$ th-order; singularity

1 Introduction

Generally speaking, differential equations with singularities have been considered from the very beginning of the discipline. The main reason is that singular forces are ubiquitous in applications, gravitational and electromagnetic forces being the most obvious examples. In 1979, Taliaferro [1] discussed the model equation with singularity

$$y'' + \frac{q(t)}{y^\alpha} = 0, \quad 0 < t < 1, \quad (1.1)$$

subject to

$$y(0) = 0 = y(1),$$

and obtained the existence of a solution for the problem. Here $\alpha > 0$, $q \in C(0, 1)$ with $q > 0$ on $(0, 1)$ and $\int_0^1 t(1-t)q(t) dt < \infty$. We call it the equation with the strong force condition if $\alpha \geq 1$ and we call it the equation with the weak force condition if $0 < \alpha < 1$.

Ding's work has attracted the attention of many specialists in differential equations. More recently, topological degree theory [2–4], the Schauder fixed point theorem [5, 6], the Krasnoselskii fixed point theorem in a cone [7–9], the Poincaré-Birkhoff twist theorem [10–12], and the Leray-Schauder alternative principle [13–15] have been employed to investigate the existence of positive periodic solutions of singular second-order, third-order and fourth-order differential equations. In 1996, using coincidence degree theory, Zhang

[2] considered the existence of T -periodic solutions for the scalar Liénard equation

$$x''(t) + f(x(t))x'(t) + g(t, x(t)) = 0,$$

when g becomes unbounded as $x \rightarrow 0^+$. The main emphasis was on the repulsive case, *i.e.* when $g(t, x) \rightarrow +\infty$, as $x \rightarrow 0^+$. In 2007, Torres [5] studied singular forced semilinear differential equation

$$x'' + a(t)x' = f(t, x) + e(t). \tag{1.2}$$

By the Schauder fixed point theorem, the author has shown that the additional assumption of a weak singularity enabled new criteria for the existence of periodic solutions. Afterwards, Wang [3] investigated the existence and multiplicity of positive periodic solutions of the singular systems (1.2) by the Krasnoselskii fixed point theorem. The conditions he presented to guarantee the existence of positive periodic solutions are beautiful. Recently, Cheng and Ren [14] discussed a kind of fourth-order singular differential equation,

$$x^{(4)}(t) + ax'''(t) + bx''(t) + cx'(t) + dx(t) = f(t, x(t)) + e(t). \tag{1.3}$$

By application of Green's function and some fixed point theorems, *i.e.*, the Leray-Schauder alternative principle and Schauder's fixed point theorem, the authors established two existence results of positive periodic solutions for nonlinear fourth-order singular differential equation.

Motivated by [2, 3, 5, 14], in this paper, we consider the high-order p -Laplacian differential equation with singularity

$$(\varphi_p(x(t)))^{(n)} + f(x(t))x'(t) + g(t, x(t - \sigma)) = e(t), \tag{1.4}$$

where $p \geq 2$, $\varphi_p(x) = |x|^{p-2}x$ for $x \neq 0$, and $\varphi_p(0) = 0$; g is continuous function defined on \mathbb{R}^2 and periodic in t with $g(t, \cdot) = g(t + T, \cdot)$, g has a singularity at $x = 0$; σ is a constant and $0 \leq \sigma < T$; $e : \mathbb{R} \rightarrow \mathbb{R}$ are continuous periodic functions with $e(t + T) \equiv e(t)$ and $\int_0^T e(t) dt = 0$. T is a positive constant; n is positive integer.

The paper is organized as follows. In Section 2, we introduce some technical tools and present all the auxiliary results; in Section 3, by applying coincidence degree theory and some new inequalities, we obtain sufficient conditions for the existence of positive periodic solutions for (1.4), an example is also given to illustrate our results. Our new results generalize in several aspects some recent results contained in [2, 3, 5].

2 Lemmas

For the sake of convenience, throughout this paper we will adopt the following notation:

$$\begin{aligned} |u|_\infty &= \max_{t \in [0, T]} |u(t)|, & |u|_0 &= \min_{t \in [0, T]} |u(t)|, \\ |u|_p &= \left(\int_0^T |u|^p dt \right)^{\frac{1}{p}}, & \bar{h} &= \frac{1}{T} \int_0^T h(t) dt. \end{aligned}$$

Let X and Y be real Banach spaces and $L : D(L) \subset X \rightarrow Y$ be a Fredholm operator with index zero, here $D(L)$ denotes the domain of L . This means that $\text{Im} L$ is closed in Y and $\dim \text{Ker} L = \dim(Y/\text{Im} L) < +\infty$. Consider supplementary subspaces X_1, Y_1 of X, Y , respectively, such that $X = \text{Ker} L \oplus X_1, Y = \text{Im} L \oplus Y_1$. Let $P : X \rightarrow \text{Ker} L$ and $Q : Y \rightarrow Y_1$ denote the natural projections. Clearly, $\text{Ker} L \cap (D(L) \cap X_1) = \{0\}$ and so the restriction $L_P := L|_{D(L) \cap X_1}$ is invertible. Let K denote the inverse of L_P .

Let Ω be an open bounded subset of X with $D(L) \cap \Omega \neq \emptyset$. A map $N : \overline{\Omega} \rightarrow Y$ is said to be L -compact in $\overline{\Omega}$ if $QN(\overline{\Omega})$ is bounded and the operator $K(I - Q)N : \overline{\Omega} \rightarrow X$ is compact.

Lemma 2.1 (Gaines and Mawhin [16]) *Suppose that X and Y are two Banach spaces, and $L : D(L) \subset X \rightarrow Y$ is a Fredholm operator with index zero. Let $\Omega \subset X$ be an open bounded set and $N : \overline{\Omega} \rightarrow Y$ be L -compact on $\overline{\Omega}$. Assume that the following conditions hold:*

- (1) $Lx \neq \lambda Nx, \forall x \in \partial\Omega \cap D(L), \lambda \in (0, 1)$;
- (2) $Nx \notin \text{Im} L, \forall x \in \partial\Omega \cap \text{Ker} L$;
- (3) $\deg\{JQN, \Omega \cap \text{Ker} L, 0\} \neq 0$, where $J : \text{Im} Q \rightarrow \text{Ker} L$ is an isomorphism.

Then the equation $Lx = Nx$ has a solution in $\overline{\Omega} \cap D(L)$.

Lemma 2.2 ([17]) *If $\omega \in C^1(\mathbb{R}, \mathbb{R})$ and $\omega(0) = \omega(T) = 0$, then*

$$\int_0^T |\omega(t)|^p dt \leq \left(\frac{T}{\pi_p}\right)^p \int_0^T |\omega'(t)|^p dt,$$

where $1 \leq p < \infty, \pi_p = 2 \int_0^{(p-1)/p} \frac{ds}{(1-s^p)^{1/p}} = \frac{2\pi(p-1)^{1/p}}{p \sin(\pi/p)}$.

Lemma 2.3 *If $x(t) \in C^n(\mathbb{R}, \mathbb{R})$ and $x^{(j)}(t + T) = x^{(j)}(t), j = 0, 1, 2, \dots, n - 1$, then*

$$\int_0^T |x^{(i)}(t)|^p dt \leq \left(\frac{T}{\pi_p}\right)^{p(n-i)} \int_0^T |x^{(n)}(t)|^p dt, \quad i = 1, 2, \dots, n - 1,$$

where $\frac{1}{p} + \frac{1}{q} = 1, p \geq 2$.

Proof From $x^{(i-1)}(0) = x^{(i-1)}(T)$, there is a point $t_i \in [0, T]$ such that $x^{(i)}(t_i) = 0$. Let $\omega_i(t) = x^{(i)}(t + t_i)$, and then $\omega_i(0) = \omega_i(T) = 0$. From $x^{(i)}(0) = x^{(i)}(T)$, there is a point $t_{i+1} \in [0, T]$ such that $x^{(i+1)}(t_{i+1}) = 0$. Let $\omega_{i+1}(t) = x^{(i+1)}(t + t_{i+1})$, and then $\omega_{i+1}(0) = \omega_{i+1}(T) = 0$. Continuing this way we get from $x^{(n-i)}(0) = x^{(n-i)}(T)$ a point $t_{n-i+1} \in [0, T]$ such that $x^{(n)}(t_{n-i+1}) = 0$. Let $\omega_{n-i}(t) = x^{(n-i+1)}(t + t_{n-i+1})$, and then $\omega_{n-i}(0) = \omega_{n-i}(T) = 0$. From Lemma 2.2, we have

$$\begin{aligned} \int_0^T |x^{(i)}(t)|^p dt &= \int_0^T |\omega_i(t)|^p dt \\ &\leq \left(\frac{T}{\pi_p}\right)^p \int_0^T |\omega_i'(t)|^p dt \\ &= \left(\frac{T}{\pi_p}\right)^p \int_0^T |x^{(i+1)}(t)|^p dt \\ &= \left(\frac{T}{\pi_p}\right)^p \int_0^T |\omega_{i+1}(t)|^p dt \\ &\leq \left(\frac{T}{\pi_p}\right)^{2p} \int_0^T |\omega_{i+1}'(t)|^p dt \end{aligned}$$

$$\begin{aligned}
 & \dots \\
 & \leq \left(\frac{T}{\pi_p}\right)^{p(n-i)} \int_0^T |\omega'_{n-i-1}(t)|^p dt \\
 & = \left(\frac{T}{\pi_p}\right)^{p(n-i)} \int_0^T |x^{(n)}(t)|^p dt.
 \end{aligned} \tag{2.1}$$

□

In order to apply coincidence degree theorem, we rewrite (1.4) in the form

$$\begin{cases} x_1^{(n)}(t) = \varphi_q(x_2(t)), \\ x_2^{(n)}(t) = -f(x_1(t))x_1'(t) - g(t, x_1(t - \sigma)) + e(t), \end{cases} \tag{2.2}$$

where $\frac{1}{p} + \frac{1}{q} = 1$. Clearly, if $x(t) = (x_1(t), x_2(t))^T$ is a T -periodic solution to (2.2), then $x_1(t)$ must be a T -periodic solution to (1.4). Thus, the problem of finding a T -periodic solution for (1.4) reduces to finding one for (2.2).

Now, set $X = \{x = (x_1(t), x_2(t)) \in C(\mathbb{R}, \mathbb{R}^2) : x(t + T) \equiv x(t)\}$ with the norm $|x|_\infty = \max\{|x_1|_\infty, |x_2|_\infty\}$; $Y = \{x = (x_1(t), x_2(t)) \in C^1(\mathbb{R}, \mathbb{R}^2) : x(t + T) \equiv x(t)\}$ with the norm $\|x\| = \max\{|x|_\infty, |x'|_\infty\}$. Clearly, X and Y are both Banach spaces. Meanwhile, define

$$L : D(L) = \{x \in C^{2n}(\mathbb{R}, \mathbb{R}^2) : x(t + T) = x(t), t \in \mathbb{R}\} \subset X \rightarrow Y$$

by

$$(Lx)(t) = \begin{pmatrix} x_1^{(n)}(t) \\ x_2^{(n)}(t) \end{pmatrix}$$

and $N : X \rightarrow Y$ by

$$(Nx)(t) = \begin{pmatrix} \varphi_q(x_2(t)) \\ -f(x_1)x_1'(t) - g(t, x_1(t - \sigma)) + e(t) \end{pmatrix}. \tag{2.3}$$

Then (2.2) can be converted into the abstract equation $Lx = Nx$. From the definition of L , one can easily see that

$$\text{Ker } L \cong \mathbb{R}^2, \quad \text{Im } L = \left\{ y \in Y : \int_0^T \begin{pmatrix} y_1(s) \\ y_2(s) \end{pmatrix} ds = \begin{pmatrix} 0 \\ 0 \end{pmatrix} \right\}.$$

So L is a Fredholm operator with index zero. Let $P : X \rightarrow \text{Ker } L$ and $Q : Y \rightarrow \text{Im } Q \subset \mathbb{R}^2$ be defined by

$$Px = \begin{pmatrix} x_1(0) \\ x_2(0) \end{pmatrix}; \quad Qy = \frac{1}{T} \int_0^T \begin{pmatrix} y_1(s) \\ y_2(s) \end{pmatrix} ds,$$

then $\text{Im } P = \text{Ker } L$, $\text{Ker } Q = \text{Im } L$. Setting $L_P = L|_{D(L) \cap \text{Ker } P}$ and $L_P^{-1} : \text{Im } L \rightarrow D(L)$ denoting the inverse of L_P , then

$$[L_P^{-1}y](t) = \begin{pmatrix} (Gy_1)(t) \\ (Gy_2)(t) \end{pmatrix},$$

$$\begin{aligned}
 [Gy_1](t) &= \sum_{i=1}^{n-1} \frac{1}{i!} x_1^{(i)}(0)t^i + \frac{1}{(n-1)!} \int_0^t (t-s)^{n-1} y_1(s) ds, \\
 [Gy_2](t) &= \sum_{i=1}^{n-1} \frac{1}{i!} x_2^{(i)}(0)t^i + \frac{1}{(n-1)!} \int_0^t (t-s)^{n-1} y_2(s) ds,
 \end{aligned}
 \tag{2.4}$$

where $x_j^{(i)}(0), i = 1, 2, \dots, n - 1$ and $j = 1, 2$, are defined by the following:

$$E_1 Z = B, \quad \text{where } E_1 = \begin{pmatrix} 1 & 0 & 0 & \cdots & 0 & 0 \\ c_1 & 1 & 0 & \cdots & 0 & 0 \\ c_2 & c_1 & 1 & \cdots & 0 & 0 \\ \cdots & & & & & \\ c_{n-3} & c_{n-4} & c_{n-5} & \cdots & 1 & 0 \\ c_{n-2} & c_{n-3} & c_{n-4} & \cdots & c_1 & 0 \end{pmatrix}_{(n-1) \times (n-1)}.$$

$Z = (x_1^{(n-1)}(0), \dots, x_1''(0), x_1'(0))^\top, B = (b_1, b_2, \dots, b_{n-1})^\top, b_i = -\frac{1}{i!T} \int_0^T (T-s)^i y_1(s) ds$, and $c_k = \frac{T^k}{(k+1)!}, k = 1, 2, \dots, n - 2$.

From (2.3) and (2.4), it is clearly that QN and $K(I - Q)N$ are continuous, $QN(\bar{\Omega})$ is bounded and then $K(I - Q)N(\bar{\Omega})$ is compact for any open bounded $\Omega \subset X$, which means N is L -compact on $\bar{\Omega}$.

3 Existence of positive periodic solutions for (1.1)

Assume that

$$\psi(t) = \limsup_{x \rightarrow +\infty} \frac{g(t, x)}{x^{p-1}}, \tag{3.1}$$

exists uniformly a.e. $t \in [0, T]$, i.e., for any $\varepsilon > 0$ there is $g_\varepsilon \in L^2(0, T)$ such that

$$g(t, x) \leq (\psi(t) + \varepsilon)x^{p-1} + g_\varepsilon(t) \tag{3.2}$$

for all $x > 0$ and a.e. $t \in [0, T]$. Moreover, $\psi \in C(\mathbb{R}, \mathbb{R})$ and $\psi(t + T) = \psi(t)$.

For the sake of convenience, we list the following assumptions which will be used repeatedly in the sequel:

(H₁) There exist constants $0 < D_1 < D_2$ such that if x is a positive continuous T -periodic function satisfying

$$\int_0^T g(t, x(t)) dt = 0,$$

then

$$D_1 \leq x(\tau) \leq D_2$$

for some $\tau \in [0, T]$.

(H₂) $\bar{g}(x) < 0$ for all $x \in (0, D_1)$, and $\bar{g}(x) > 0$ for all $x > D_2$.

(H₃) $g(t, x) = g_0(x) + g_1(t, x)$, where $g_0 \in C((0, \infty); \mathbb{R})$ and $g_1 : [0, T] \times [0, \infty) \rightarrow \mathbb{R}$ is an L^2 -Carathéodory function, i.e. it is measurable in the first variable and continuous in the

second variable, and for any $b > 0$ there is $h_b \in L^2(0, T; \mathbb{R}_+)$ such that

$$|g_1(t, x)| \leq h_b(t), \quad \text{a.e. } t \in [0, T], \forall 0 \leq x \leq b.$$

$$(H_4) \int_0^1 g_0(x) dx = -\infty.$$

Theorem 3.1 *Assume that conditions (H₁)-(H₄) hold. If $|\psi|_\infty \frac{T^{\frac{p}{q}+1}}{2^{p-1}} (\frac{T}{\pi_p})^{p(n-1)} < 1$, then (1.4) has at least a positive T -periodic solution.*

Proof Consider the equation

$$Lx = \lambda Nx, \quad \lambda \in (0, 1).$$

Set $\Omega_1 = \{x : Lx = \lambda Nx, \lambda \in (0, 1)\}$. If $x(t) = (x_1(t), x_2(t))^T \in \Omega_1$, then

$$\begin{cases} x_1^{(n)}(t) = \lambda \varphi_q(x_2(t)), \\ x_2^{(n)}(t) = -\lambda f(x_1(t))x_1'(t) - \lambda g(t, x_1(t - \sigma)) + \lambda e(t). \end{cases} \tag{3.3}$$

Substituting $x_2(t) = \lambda^{1-p} \varphi_p[x_1^{(n)}(t)]$ into the second equation of (3.3)

$$(\varphi_p(x_1^{(n)}(t)))^{(n)} + \lambda^p f(x_1(t))x_1'(t) + \lambda^p g(t, x_1(t - \sigma)) = \lambda^p e(t). \tag{3.4}$$

Integrating both sides of (3.4) from 0 to T , we have

$$\int_0^T g(t, x_1(t - \sigma)) dt = 0. \tag{3.5}$$

In view of (H₁), there exist positive constants D_1, D_2 , and $\xi \in [0, T]$ such that

$$D_1 \leq |x_1(\xi)| \leq D_2.$$

Then we have

$$|x_1(t)| = \left| x_1(\xi) + \int_\xi^t x_1'(s) ds \right| \leq D_2 + \int_\xi^t |x_1'(s)| ds, \quad t \in [\xi, \xi + T],$$

and

$$|x_1(t)| = |x_1(t - T)| = \left| x_1(\xi) - \int_{t-T}^\xi x_1'(s) ds \right| \leq D_2 + \int_{t-T}^\xi |x_1'(s)| ds, \quad t \in [\xi, \xi + T].$$

Combing the above two inequalities, we obtain

$$\begin{aligned} |x_1|_\infty &= \max_{t \in [0, T]} |x_1(t)| = \max_{t \in [\xi, \xi + T]} |x_1(t)| \\ &\leq \max_{t \in [\xi, \xi + T]} \left\{ D_2 + \frac{1}{2} \left(\int_\xi^t |x_1'(s)| ds + \int_{t-T}^\xi |x_1'(s)| ds \right) \right\} \\ &\leq D_2 + \frac{1}{2} \int_0^T |x_1'(s)| ds. \end{aligned} \tag{3.6}$$

Multiplying both sides of (3.4) by $x_1(t)$ and integrating over interval $[0, T]$, we get

$$\begin{aligned} & \int_0^T (\varphi_p(x_1^{(n)}(t)))^{(n)} x_1(t) dt + \lambda^p \int_0^T f(x_1(t))x_1'(t)x_1(t) dt + \lambda^p \int_0^T g(t, x_1(t - \sigma))x_1(t) dt \\ & = \lambda^p \int_0^T e(t)x_1(t) dt. \end{aligned} \tag{3.7}$$

Substituting $\int_0^T (\varphi_p(x_1^{(n)}(t)))^{(n)} x_1(t) dt = (-1)^n \int_0^T |x_1^{(n)}(t)|^p dt$, $\int_0^T f(x_1(t))x_1'(t)x_1(t) dt = 0$ into (3.7), we have

$$(-1)^n \int_0^T |x_1^{(n)}(t)|^p dt = -\lambda^p \int_0^T g(t, x_1(t - \sigma))x_1(t) dt + \lambda^p \int_0^T e(t)x_1(t) dt.$$

Namely,

$$\begin{aligned} \int_0^T |x_1^{(n)}(t)|^p dt & \leq \int_0^T |g(t, x_1(t - \sigma))||x_1(t)| dt + \int_0^T |e(t)||x_1(t)| dt \\ & \leq |x_1|_\infty \int_0^T |g(t, x_1(t - \sigma))| dt + |x_1|_\infty |e|_\infty T. \end{aligned} \tag{3.8}$$

Write

$$I_+ = \{t \in [0, T] : g(t, x_1(t - \sigma)) \geq 0\}; \quad I_- = \{t \in [0, T] : g(t, x_1(t - \sigma)) \leq 0\}.$$

Then we get from (3.2) and (3.5)

$$\begin{aligned} \int_0^T |g(t, x_1(t - \sigma))| dt & = \int_{I_+} g(t, x_1(t - \sigma)) dt - \int_{I_-} g(t, x_1(t - \sigma)) dt \\ & = 2 \int_{I_+} g(t, x_1(t - \sigma)) dt \\ & \leq 2 \int_{I_+} ((\psi(t) + \varepsilon)x_1^{p-1}(t - \sigma) + g_\varepsilon(t)) dt \\ & \leq 2(|\psi|_\infty + \varepsilon) \int_0^T |x_1(t)|^{p-1} dt + 2 \int_0^T |g_\varepsilon(t)| dt. \end{aligned} \tag{3.9}$$

Substituting (3.9) into (3.8), we have

$$\begin{aligned} \int_0^T |x_1^{(n)}(t)|^p dt & \leq 2|x_1|_\infty (|\psi|_\infty + \varepsilon) \int_0^T |x_1(t)|^{p-1} dt \\ & \quad + |x_1|_\infty \left(2 \int_0^T |g_\varepsilon(t)| dt + |e|_\infty T \right) \\ & \leq 2(|\psi|_\infty + \varepsilon) T |x_1|_\infty^p + |x_1|_\infty \left(2T^{\frac{1}{2}} \left(\int_0^T |g_\varepsilon(t)|^2 dt \right)^{\frac{1}{2}} + |e|_\infty T \right) \\ & = 2(|\psi|_\infty + \varepsilon) T |x_1|_\infty^p + |x_1|_\infty (2T^{\frac{1}{2}} |g_\varepsilon|_2 + |e|_\infty T). \end{aligned} \tag{3.10}$$

From (3.6) and Lemma 2.3, we have

$$\begin{aligned}
 |x_1|_\infty &\leq D_2 + \frac{1}{2} \int_0^T |x_1'(t)| dt \leq D_2 + \frac{T^{\frac{1}{q}}}{2} \left(\int_0^T |x_1'(t)|^p dt \right)^{\frac{1}{p}} \\
 &\leq D_2 + \frac{T^{\frac{1}{q}}}{2} \left(\frac{T}{\pi_p} \right)^{n-1} \left(\int_0^T |x_1^{(n)}(t)|^p dt \right)^{\frac{1}{p}}.
 \end{aligned}
 \tag{3.11}$$

Substituting (3.11) into (3.10), we have

$$\begin{aligned}
 &\int_0^T |x_1^{(n)}(t)|^p dt \\
 &\leq 2(|\psi|_\infty + \varepsilon) T \left(D_2 + \frac{T^{\frac{1}{q}}}{2} \left(\frac{T}{\pi_p} \right)^{n-1} \left(\int_0^T |x_1^{(n)}(t)|^p dt \right)^{\frac{1}{p}} \right)^p \\
 &\quad + \left(D_2 + \frac{T^{\frac{1}{q}}}{2} \left(\frac{T}{\pi_p} \right)^{n-1} \left(\int_0^T |x_1^{(n)}(t)|^p dt \right)^{\frac{1}{p}} \right) (2T^{\frac{1}{2}} |g_\varepsilon|_2 + |e|_\infty T) \\
 &= 2(|\psi|_\infty + \varepsilon) T \left(\frac{T^{\frac{p}{q}}}{2^p} \left(\frac{T}{\pi_p} \right)^{p(n-1)} \int_0^T |x_1^{(n)}(t)|^p dt + p D_2 \frac{T^{\frac{p-1}{q}}}{2^{p-1}} \left(\frac{T}{\pi_p} \right)^{(p-1)(n-1)} \right. \\
 &\quad \cdot \left. \left(\int_0^T |x_1^{(n)}(t)| dt \right)^{\frac{p-1}{p}} + \dots + p D_2^{p-1} \frac{T^{\frac{1}{q}}}{2} \left(\frac{T}{\pi_p} \right)^{n-1} \left(\int_0^T |x_1'(t)|^p dt \right)^{\frac{1}{p}} + D_2^p \right) \\
 &\quad + \left(D_2 + \frac{T^{\frac{1}{q}}}{2} \left(\frac{T}{\pi_p} \right)^{n-1} \left(\int_0^T |x_1^{(n)}(t)|^p dt \right)^{\frac{1}{p}} \right) (2T^{\frac{1}{2}} |g_\varepsilon|_2 + |e|_\infty T) \\
 &= (|\psi|_\infty + \varepsilon) \frac{T^{\frac{p}{q}+1}}{2^{p-1}} \left(\frac{T}{\pi_p} \right)^{p(n-1)} \int_0^T |x_1^{(n)}|^p dt \\
 &\quad + (|\psi|_\infty + \varepsilon) p D_2 \frac{T^{\frac{p-1}{q}+1}}{2^{p-2}} \left(\frac{T}{\pi_p} \right)^{(p-1)(n-1)} \left(\int_0^T |x_1^{(n)}(t)|^p dt \right)^{\frac{p-1}{p}} + \dots \\
 &\quad + (2(|\psi|_\infty + \varepsilon) T p D_2^{p-1} + 2T^{\frac{1}{2}} |g_\varepsilon|_2 + |e|_\infty T) \frac{T^{\frac{1}{q}}}{2} \left(\frac{T}{\pi_p} \right)^{n-1} \\
 &\quad \cdot \left(\int_0^T |x_1^{(n)}(t)|^p dt \right)^{\frac{1}{p}} + 2(|\psi|_\infty + \varepsilon) T D_2^p + D_2 (2T^{\frac{1}{2}} |g_\varepsilon|_2 + |e|_\infty T).
 \end{aligned}$$

Since ε sufficiently small, we know that $|\psi|_\infty \frac{T^{\frac{p}{q}+1}}{2^{p-1}} \left(\frac{T}{\pi_p} \right)^{p(n-1)} < 1$. So, it is easy to see that there exists a positive constant M'_1 such that

$$\int_0^T |x_1^{(n)}(t)|^p dt \leq M'_1.$$

From (3.11), we have

$$\begin{aligned}
 |x_1|_\infty &\leq D_2 + \frac{T^{\frac{1}{q}}}{2} \left(\frac{T}{\pi_p} \right)^{n-1} \left(\int_0^T |x_1^{(n)}(t)|^p dt \right)^{\frac{1}{p}} \\
 &\leq D_2 + \frac{T^{\frac{1}{q}}}{2} \left(\frac{T}{\pi_p} \right)^{n-1} (M'_1)^{\frac{1}{p}} := M_1.
 \end{aligned}
 \tag{3.12}$$

Since $x_1(0) = x_1(T)$, there exists a point $\eta_1 \in [0, T]$ such that $x_1'(\eta_1) = 0$. From Lemma 2.3, we can easily get

$$\begin{aligned} |x_1'|_\infty &\leq \frac{1}{2} \int_0^T |x_1''(t)| dt \\ &\leq \frac{T^{\frac{1}{q}}}{2} \left(\int_0^T |x_1''(t)|^p dt \right)^{\frac{1}{p}} \\ &\leq \frac{T^{\frac{1}{q}}}{2} \left(\frac{T}{\pi_p} \right)^{(n-2)} \left(\int_0^T |x_1^{(n)}|^p \right)^{\frac{1}{p}} \\ &\leq \frac{T^{\frac{1}{q}}}{2} \left(\frac{T}{\pi_p} \right)^{(n-2)} (M_1')^{\frac{1}{p}} := M_2. \end{aligned} \tag{3.13}$$

On the other hand, form $x_2^{(n-2)}(0) = x_2^{(n-2)}(T)$, there exists a point $\eta_2 \in [0, T]$ such that $x_2^{(n-1)}(\eta_2) = 0$, from the second equation of (3.3) and (3.9), we have

$$\begin{aligned} |x_2^{(n-1)}|_\infty &\leq \frac{1}{2} \max \left| \int_0^T x_2^{(n)}(t) dt \right| \\ &\leq \frac{\lambda}{2} \int_0^T |-f(x_1(t))x_1'(t) - g(t, x_1(t), x_1(t - \sigma)) + e(t)| dt \\ &\leq \frac{\lambda}{2} (|f|_{M_1} TM_2 + 2(|\psi|_\infty + \varepsilon) TM_1^{p-1} + 2\sqrt{T}|g_\varepsilon|_2 + T|e|_\infty) := \lambda M_{n-1}, \end{aligned}$$

where $|f|_{M_1} = \max_{0 < x_1(t) \leq M_1} |f(x_1(t))|$. Since $x_2(0) = x_2(T)$, there exists a point $\eta_3 \in [0, T]$ such that $x_2'(\eta_3) = 0$. From the Wirtinger inequality (see [18], Lemma 2.4), we can easily get

$$\begin{aligned} |x_2'|_\infty &\leq \frac{1}{2} \int_0^T |x_2''(t)| dt \leq \frac{T^{\frac{1}{2}}}{2} \left(\int_0^T |x_2''(t)|^2 dt \right)^{\frac{1}{2}} \\ &\leq \frac{T}{2} \left(\frac{T}{2\pi} \right)^{(n-3)} |x_2^{(n-1)}|_\infty \\ &\leq \frac{T}{2} \left(\frac{T}{2\pi} \right)^{(n-3)} (\lambda M_{n-1}) := \lambda M_3. \end{aligned} \tag{3.14}$$

By the first equation of (3.3), we have

$$\int_0^T |x_2(t)|^{q-2} x_2(t) dt = 0,$$

which implies that there is a constant $\eta_4 \in [0, T]$ such that $x_2(\eta_4) = 0$, so

$$|x_2|_\infty \leq \frac{1}{2} \int_0^T |x_2'(t)| dt \leq \frac{T}{2} |x_2'|_\infty \leq \frac{\lambda T}{2} M_3 := \lambda M_4. \tag{3.15}$$

Next, it follows from (3.4) that

$$(\varphi_p(x_1^{(n)}(t + \sigma)))^{(n)} + \lambda^p (f(x_1(t + \sigma))x_1'(t + \sigma) + g(t + \sigma, x_1(t))) = \lambda^p e(t + \sigma). \tag{3.16}$$

Namely,

$$\begin{aligned} & (\varphi_p(x_1^{(n)}(t + \sigma)))^{(n)} + \lambda^p f(x_1(t + \sigma))x_1'(t + \sigma) + \lambda^p (g_0(x_1(t)) + g_1(t + \sigma, x_1(t))) \\ & = \lambda^p e(t + \sigma). \end{aligned} \tag{3.17}$$

Multiplying both sides of (3.17) by $x_1'(t)$, we get

$$\begin{aligned} & (\varphi_p(x_1^{(n)}(t + \sigma)))^{(n)} x_1'(t) + \lambda^p f(x_1(t + \sigma))x_1'(t + \sigma)x_1'(t) \\ & \quad + \lambda^p g_0(x_1(t))x_1'(t) + \lambda^p g_1(t + \sigma, x_1(t))x_1'(t) \\ & = \lambda^p e(t + \sigma)x_1'(t). \end{aligned} \tag{3.18}$$

Let $\tau \in [0, T]$, for any $\tau \leq t \leq T$, we integrate (3.18) on $[\tau, t]$ and get

$$\begin{aligned} & \lambda^p \int_{x_1(\tau)}^{x_1(t)} g_0(u) du \\ & = \lambda^p \int_{\tau}^t g_0(x_1(s))x_1'(s) ds \\ & = - \int_{\tau}^t (\varphi_p(x_1^{(n)}(s + \sigma)))^{(n)} x_1'(s) ds - \lambda^p \int_{\tau}^t f(x_1(s + \sigma))x_1'(s + \sigma)x_1'(s) ds \\ & \quad - \lambda^p \int_{\tau}^t g_1(s + \sigma, x_1(s))x_1'(s) ds + \lambda^p \int_{\tau}^t e(s + \sigma)x_1'(s) ds. \end{aligned} \tag{3.19}$$

By (3.12), (3.13), and (3.16), we have

$$\begin{aligned} & \left| \int_{\tau}^t (\varphi_p(x_1^{(n)}(s + \sigma)))^{(n)} x_1'(s) ds \right| \\ & \leq \int_{\tau}^t |(\varphi_p(x_1^{(n)}(s + \sigma)))^{(n)}| |x_1'(s)| ds \\ & \leq |x_1'|_{\infty} \int_0^T |(\varphi_p(x_1^{(n)}(t + \sigma)))^{(n)}| dt \\ & \leq \lambda^p |x_1'|_{\infty} \left(\int_0^T |f(x_1(t))| |x_1'(t)| dt + \int_0^T |g(t, x_1(t - \sigma))| dt + \int_0^T |e(t)| dt \right) \\ & \leq \lambda^p M_2 (|f|_{M_1} M_2 + 2(|\psi|_{\infty} + \varepsilon) T M_1^{p-1} + 2T^{\frac{1}{2}} |g_{\varepsilon}^+|_2 + T |e|_{\infty}). \end{aligned}$$

Also we have

$$\begin{aligned} & \left| \int_{\tau}^t f(x_1(s + \sigma))x_1'(s + \sigma)x_1'(s) ds \right| \leq |f|_{M_1} M_2^2 T, \\ & \left| \int_{\tau}^t g(s + \sigma, x_1(s))x_1'(s) ds \right| \leq |x_1'|_{\infty} \int_0^T |g(t, x(t - \sigma))| dt \leq M_2 \sqrt{T} |g_{M_1}|_2, \end{aligned}$$

where $g_{M_1} = \max_{0 \leq x \leq M_1} |g_1(t, x)| \in L^2(0, T)$ is as in (H₃).

$$\left| \int_{\tau}^t e(t + \sigma)x_1'(t) dt \right| \leq M_2 T |e|_{\infty}.$$

From these inequalities we can derive from (3.19) that

$$\left| \int_{x_1(\tau)}^{x_1(t)} g_0(u) du \right| \leq M'_5, \tag{3.20}$$

for some constant M'_5 which is independent on λ, x , and t . In view of the strong force condition (H_4) , we know that there exists a constant $M_5 > 0$ such that

$$x_1(t) \geq M_5, \quad \forall t \in [\tau, T]. \tag{3.21}$$

The case $t \in [0, \tau]$ can be treated similarly.

From (3.12), (3.13), (3.14), (3.15), and (3.21), we get

$$\Omega = \{x = (x_1, x_2)^\top : E_1 \leq |x_1|_\infty \leq E_2, |x'_1|_\infty \leq E_3, |x_2|_\infty \leq E_4 \text{ and } |x'_2|_\infty \leq E_5, \forall t \in [0, T]\},$$

where $0 < E_1 < \min(M_5, D_1)$, $E_2 > \max(M_1, D_2)$, $E_3 > M_2$, $E_4 > M_4$, and $E_5 > M_3$. $\Omega_2 = \{x : x \in \partial\Omega \cap \text{Ker } L\}$, then $\forall x \in \partial\Omega \cap \text{Ker } L$

$$QNx = \frac{1}{T} \int_0^T \begin{pmatrix} \varphi_q(x_2(t)) \\ -f(x_1(t))x'_1(t) - g(t, x_1(t - \sigma)) + e(t) \end{pmatrix} dt.$$

If $QNx = 0$, then $x_2(t) = 0$, $x_1 = E_2$ or $-E_2$. But if $x_1(t) = E_2$, we know

$$0 = \int_0^T \{g(t, E_2) - e(t)\} dt.$$

From assumption (H_2) , we have $x_1(t) \leq D_2 \leq E_2$, which yields a contradiction. Similarly if $x_1 = -E_2$. We also have $QNx \neq 0$, i.e., $\forall x \in \partial\Omega \cap \text{Ker } L, x \notin \text{Im } L$, so conditions (1) and (2) of Lemma 2.1 are both satisfied. Define the isomorphism $J : \text{Im } Q \rightarrow \text{Ker } L$ as follows:

$$J(x_1, x_2)^\top = (x_2, -x_1)^\top.$$

Let $H(\mu, x) = -\mu x + (1 - \mu)JQNx$, $(\mu, x) \in [0, 1] \times \Omega$, then $\forall (\mu, x) \in (0, 1) \times (\partial\Omega \cap \text{Ker } L)$,

$$H(\mu, x) = \begin{pmatrix} -\mu x_1 - \frac{1-\mu}{T} \int_0^T [g(t, x_1) - e(t)] dt \\ -\mu x_2 - (1 - \mu)|x_2|^{q-2}x_2 \end{pmatrix}.$$

We have $\int_0^T e(t) dt = 0$. So, we can get

$$H(\mu, x) = \begin{pmatrix} -\mu x_1 - \frac{1-\mu}{T} \int_0^T g(t, x_1) dt \\ -\mu x_2 - (1 - \mu)|x_2|^{q-2}x_2 \end{pmatrix}, \quad \forall (\mu, x) \in (0, 1) \times (\partial\Omega \cap \text{Ker } L).$$

From (H_2) , it is obvious that $x^\top H(\mu, x) < 0$, $\forall (\mu, x) \in (0, 1) \times (\partial\Omega \cap \text{Ker } L)$. Hence

$$\begin{aligned} \deg\{JQN, \Omega \cap \text{Ker } L, 0\} &= \deg\{H(0, x), \Omega \cap \text{Ker } L, 0\} \\ &= \deg\{H(1, x), \Omega \cap \text{Ker } L, 0\} \\ &= \deg\{I, \Omega \cap \text{Ker } L, 0\} \neq 0. \end{aligned}$$

So condition (3) of Lemma 2.1 is satisfied. By applying Lemma 2.1, we conclude that the equation $Lx = Nx$ has a solution $x = (x_1, x_2)^T$ on $\bar{\Omega} \cap D(L)$, i.e., (1.4) has a positive T -periodic solution $x_1(t)$. □

Example 3.1 Consider the high-order p -Laplacian differential equation with singularity

$$(\varphi_p(x(t)'''))'''' + f(x(t))x'(t) + \frac{1}{6}(\sin 2t + 3)x^3(t - \sigma) - \frac{1}{x^\kappa(t - \sigma)} = \cos 2t, \tag{3.22}$$

where $\kappa \geq 1$ and $p = 4$, f is continuous function, σ is a constant, and $0 \leq \sigma < T$.

It is clear that $T = \pi$, $n = 3$, $g(t, x) = \frac{1}{6}(\sin 2t + 3)x^3(t - \sigma) - \frac{1}{x^\kappa(t - \sigma)}$, $\psi(t) = \frac{1}{6}(\sin 2t + 3)$, $|\psi|_\infty = \frac{2}{3}$. It is obvious that (H₁)-(H₄) hold. Now we consider the assumption condition

$$\begin{aligned} |\psi|_\infty & \frac{T^{\frac{p}{q}+1}}{2^{p-1}} \left(\frac{T}{\pi_p}\right)^{p(n-1)} \\ & = |\psi|_\infty \frac{T^{\frac{p}{q}+1}}{2^{p-1}} \left(\frac{T}{\frac{2\pi(p-1)^{1/p}}{p \sin(\pi/p)}}\right)^{p(n-1)} \\ & = \frac{2}{3} \cdot \frac{\pi^{\frac{4}{3}}}{2^3} \left(\frac{\pi}{\frac{2\pi(4-1)^{1/4}}{4 \sin \pi/4}}\right)^8 \\ & = \frac{4\pi^{\frac{4}{3}}}{27} < 1. \end{aligned}$$

So by Theorem 3.1, we know (3.22) has at least one positive π -periodic solution.

Competing interests

The authors declare that they have no competing interests.

Authors' contributions

YX and SZ worked together in the derivation of the mathematical results. Both authors read and approved the final manuscript.

Acknowledgements

YX and SZ would like to thank the referee for invaluable comments and insightful suggestions. This work was supported by Natural Science Foundation of China (No. 11326124) and the Fundamental Research Funds for the Universities of Henan Province (NSFRF140142).

Received: 18 September 2015 Accepted: 20 November 2015 Published online: 26 January 2016

References

1. Taliaferro, S: A nonlinear singular boundary value problem. *Nonlinear Anal. TMA* **3**, 897-904 (1979)
2. Zhang, MR: Periodic solutions of linear and quasilinear neutral functional differential equations. *J. Math. Anal. Appl.* **189**, 378-392 (1995)
3. Wang, ZH: Periodic solutions of Liénard equation with a singularity and a deviating argument. *Nonlinear Anal., Real World Appl.* **16**, 227-234 (2014)
4. Cheng, ZB: Existence of positive periodic solutions for third-order differential equation with strong singularity. *Adv. Differ. Equ.* **2014**, 162 (2014)
5. Torres, P: Weak singularities may help periodic solutions to exist. *J. Differ. Equ.* **232**, 277-284 (2007)
6. Li, X, Zhang, ZH: Periodic solutions for damped differential equations with a weak repulsive singularity. *Nonlinear Anal. TMA* **70**, 2395-2399 (2009)
7. Chu, JF, Torres, P, Zhang, MR: Periodic solution of second order non-autonomous singular dynamical systems. *J. Differ. Equ.* **239**, 196-212 (2007)
8. Wang, HY: Positive periodic solutions of singular systems with a parameter. *J. Differ. Equ.* **249**, 2986-3002 (2010)
9. Chu, JF, Zhou, ZC: Positive solutions for singular non-linear third-order periodic boundary value problems. *Nonlinear Anal. TMA* **64**, 1528-1542 (2006)
10. Cheng, ZB, Ren, JL: Periodic and subharmonic solutions for Duffing equation with singularity. *Discrete Contin. Dyn. Syst., Ser. A* **32**, 1557-1574 (2012)

11. Fonda, A, Manásevich, R: Subharmonics solutions for some second order differential equations with singularities. *SIAM J. Math. Anal.* **24**, 1294-1311 (1993)
12. Xia, J, Wang, ZH: Existence and multiplicity of periodic solutions for the Duffing equation with singularity. *Proc. R. Soc. Edinb., Sect. A* **137**, 625-645 (2007)
13. Cheng, ZB, Ren, JL: Studies on a damped differential equation with repulsive singularity. *Math. Methods Appl. Sci.* **36**, 983-992 (2013)
14. Cheng, ZB, Ren, JL: Multiplicity results of positive solutions for four-order nonlinear differential equation with singularity. *Math. Methods Appl. Sci.* (2015). doi:10.1002/mma.3481
15. Ren, JL, Cheng, ZB, Chen, YL: Existence results of periodic solutions for third-order nonlinear singular differential equation. *Math. Nachr.* **286**, 1022-1042 (2013)
16. Gaines, RE, Mawhin, JL: *Coincidence Degree and Nonlinear Differential Equation*. Springer, Berlin (1977)
17. Zhang, MR: Nonuniform nonresonance at the first eigenvalue of the p -Laplacian. *Nonlinear Anal. TMA* **29**, 41-51 (1996)
18. Torres, P, Cheng, ZB, Ren, JL: Non-degeneracy and uniqueness of periodic solutions for $2n$ -order differential equations. *Discrete Contin. Dyn. Syst.* **33**, 2155-2168 (2013)

Submit your manuscript to a SpringerOpen[®] journal and benefit from:

- ▶ Convenient online submission
- ▶ Rigorous peer review
- ▶ Immediate publication on acceptance
- ▶ Open access: articles freely available online
- ▶ High visibility within the field
- ▶ Retaining the copyright to your article

Submit your next manuscript at ▶ springeropen.com
