



RESEARCH ARTICLE

Open Access

Paraoxonase-1 gene Q192R and L55M polymorphisms and risk of cardiovascular disease in Egyptian patients with type 2 diabetes mellitus

Dalia El-Lebedy*, Mona Kafoury, Dalia Abd-El Haleem, Alshaymaa Ibrahim, Eman Awadallah and Ingy Ashmawy

Abstract

Background: Increased oxidative stress or an impaired antioxidant defense mechanism may play a crucial role in the onset and progression of atherosclerosis. Recently, Paraoxonase –1 (PON1) which accounts for most of the antioxidant effect of high density lipoprotein (HDL) cholesterol has been presented as a potential therapeutic agent against atherosclerosis development. Allele frequencies for PON1 gene that influence enzyme concentration as well as activity differ greatly among ethnic groups and data from several studies showed ethnic variations in the interpretation of cardiovascular disease (CVD) associated with PON1 polymorphisms. In this work, we investigated PON1 Q192R and L55M polymorphisms in Egyptian patients with type 2 diabetes mellitus (T2DM) and its association with CVD.

Methods: The study included 184 subjects classified into 3 groups; T2DM, T2DM + CVD, and healthy controls. PON1 polymorphisms were genotyped by real-time PCR and PON1 concentration was assayed in serum by ELISA (enzyme linked immunosorbent assay).

Results: Genotype and allele frequencies of Q192R were significantly different between controls and diabetic patients. Frequency of QQ genotype was significantly higher in healthy controls, while QR and RR genotypes were significantly higher in diabetic patients ($p = 0.02$). Frequency of 55LL and LM genotypes were significantly higher in patients than in controls ($p = 0.009$). Q192R polymorphism associated with CVD in our diabetic patients ($p = 0.01$) and with low serum PON1 concentration ($p = 0.04$). Multiple logistic regression analysis revealed significant correlations between 192R and other independent CVD risk factors.

Conclusion: PON1 192R and 55 L alleles are associated with T2DM. Q192R polymorphism is associated with CVD and lower serum enzyme concentration and might represents a novel risk factor for CVD in Egyptian patients with T2DM.

Keywords: Paraoxonase-1 gene, Cardiovascular disease, Type 2 diabetes mellitus

Introduction

Diabetes mellitus (DM) is a chronic life-threatening disease and is the second leading cause of death worldwide with at least 1 in 10 deaths among adults between 35 and 64 years old is attributable to diabetes [1]. Cardiovascular disease (CVD) is the most common cause of morbidity and mortality among diabetic patients. People with diabetes are 2 to 4 times more likely to develop coronary artery disease (CAD), 5 times more likely to develop peripheral vascular disease (PVD), and 2 to 6 times more likely to develop myocardial infarction (MI) or stroke than

people without diabetes [2]. Several factors such as dyslipidemia, obesity, smoking, exercise, alcohol intake, oxidative stress and genetic variants have been identified as risk factors for CVD in type 2 diabetes (T2DM) [3].

Recently, dysfunctional HDL or proinflammatory HDL was indicated as a proatherogenic factor [4]. The protective function of HDL against atherogenesis could be partly explained by its main constituent, PON1 (paraoxonase-1) [5]. PON1 is a 355 amino acid glycoprotein which is synthesized in the liver and secreted into the blood where it associates with HDL and accounts for most of its antioxidant effect to prevent oxidation of LDL [6,7].

Increased oxidative stress or an impaired antioxidant defense mechanism may play a crucial role in the onset

* Correspondence: d_lebedy@yahoo.co.uk

Department of Clinical and Chemical Pathology, Medical Research Division, National Research Center, Al-Bohouth Street, Cairo 12311, Egypt

and progression of atherosclerosis [8]. Since reduced PON1 concentration and/or activity might have a deleterious effect on the protective function of HDL [9]. PON1 has been the focus of intensive research and has been reported as an independent risk factor for CVD [10]. The most convincing data to link PON1 with heart disease comes from transgenic mouse studies. HDL from PON1 knockout mice have an increased capacity to oxidize LDL and are more prone to develop atherosclerosis than their wild-type siblings as a consequence of increased oxidative stress [11].

Two single nucleotide polymorphisms in the PON1 coding region affect the anti-atherogenic properties and have been associated with risk for CVD [12], a substitution of glutamine (Q) for arginine (R) at position 192 (Q192R) and a substitution of leucine (L) for methionine (M) at position 55 (L55M) [13-16]. The Q192R polymorphism alters the enzyme's ability to protect LDL from oxidation in vitro with 192Q is the most protective, while L55M polymorphism affects the enzyme concentration with 55 M is associated with lower serum concentration [17].

Data from several studies showed ethnic variations in the interpretation of CVD associated with PON1 polymorphisms [18-21]. The aim of this work is to investigate PON1 Q192R and L55M polymorphisms in Egyptian patients with type 2 diabetes mellitus (T2DM) and its association with CVD.

Materials and methods

Subjects

The studied subjects were recruited from the Outpatients Clinic of the National Research Center and National Diabetes & Endocrinology Institute, Cairo, Egypt. Data of family and medical history, smoking habits and physical activity was obtained by questionnaire. Cigarette smokers were considered if they had once smoked even if they were no longer smokers, and physical activity was defined as exercise for 2-3 days/week for at least 30 minutes. Clinical examination including systolic blood pressure (SBP) and diastolic blood pressure (DBP) measurements was applied. Anthropometric measurements (weight and height) were collected and used for BMI calculation according to the standard formula $BMI = \text{weight (kg)} / [\text{height (m)}]^2$. Hypertension was defined as blood pressure above 140/90 mmHg or taking antihypertensive drugs. According to the criteria of American Diabetes Association Classification 2010 [22], subjects were classified into 3 groups:

Normal healthy control group

Included 50 subjects with fasting plasma glucose (FPG) < 100 mg/dL. Exclusion criteria were hyperlipidemia, hypertension, family history of any form of CVD, diabetes mellitus, hepatic and renal diseases, endocrine disease, metabolic disorders, autoimmune diseases and those under medication.

T2DM patients without CVD

Included 68 subjects fulfilled the diabetes mellitus diagnostic criteria of FPG \geq 126 mg/dL or were under antidiabetic treatment (oral and/or insulin) with no history or signs of any CVD. Exclusion criteria included renal disease, hepatic disease, endocrine disease, other metabolic disorders and autoimmune diseases.

T2DM complicated with CVD

Included 66 subjects diagnosed to have diabetes with FPG \geq 126 mg/dL or under antidiabetic treatment and complicated with any of the vascular disease e.g. ischemic heart disease (IHD), macroangiopathy and/or cerebrovascular disease. Exclusion criteria included renal disease, hepatic disease, endocrine disease, other metabolic disorders and autoimmune diseases.

Informed consent was obtained from all subjects and the study protocol was approved by the Ethics Committee of the National Research Center.

Methods

Lipid analysis and biochemical markers

Venous blood samples were collected from all subjects after 12 hours of overnight fast. Total cholesterol (TC), Triglycerides (TG), high density lipoprotein (HDL-C), glucose and glycosylated hemoglobin (HbA1c) were quantified using an automated clinical chemistry analyzer Roche Diagnostics (Germany). Low density lipoprotein (LDL-C) level was calculated using Friedewald formula [23].

Genotyping of PON1 Q192R and L55M polymorphisms

Genomic DNA was isolated from whole peripheral blood using QIAamp DNA extraction kit (Qiagen Hilden, Germany, Cat no. 51304) according to the manufacturer's protocol.

L55M (rs854560) and Q192R (rs662) polymorphisms of the PON1 gene were determined by Taqman-based allelic discrimination assay using the following primers and probes: For L55M: 5'-TTTCTGTTCTCTTTTCTG GCAGAAA-3' forward and 5'-GAAAACACTCACAG AGCTAATGAAAGC-3' reverse; 55 M: 5'-FAM-CCAT TAGGCAGTATCTCCATGTCTTCAGAGCC-3' and 55 L: 5'-VIC-TCCATTAGGCAGTATCTCCAAGTCTTCAG AGCC-3'. For Q192R: 5'-GGACCTGAGCACTTTTA TGGCA-3' forward and 5'-GACAACATACGACCAC GCTAAACC-3' reverse; 192Q: 5'-FAM-TTCTTGACC CCTACTTACAATCCTGGGAGATGT-3' and 192R: 5'-VIC-CTTGACCCCTACTTACGATCCTGGGAGATGT-3'.

All primers and probes were designed by Applied Biosystems. Genotyping assays were performed on ABI 7500 Real Time PCR (Applied Biosystems). The genotyping success rates were greater than 95% for all SNPs. For the genotyping quality control, 10% of samples were randomly

selected and measured in duplicates and the concordance rate was 100%.

Assay of PON1 enzyme concentration in serum

Serum PON1 concentration was measured by enzyme linked immunosorbent assay (ELISA) using Human Paraoxonase-1 (PON1) ELISA kit Cat# 95462 (Glory Science Co., Ltd, TX, USA).

Statistical analysis

Data were analyzed using SPSS version 16.0 for Windows (Chicago, IL, USA). Chi-square was used to test the difference in genotypes frequency between the groups. Means of serum PON1 concentration according to different genotypes were compared using analysis of variance (ANOVA) and t-tests. Logistic regression analysis and odds ratios were calculated to test the association of genotypes and CVD, keeping wild type as the reference. The association of Q192R polymorphism with other CVD risk factors was tested by multiple logistic regression analysis, assessing independence from potential confounders. Variables such as age, male gender, smoking, BMI, SBP, TC, HDL, LDL and TG were included into the logistic model. P value less than 0.05 was considered significant.

Results

Our study included 184 subjects classified into 68 patients with T2DM, 66 patients with T2DM+CVD, and 50 control

subjects. Our CVD patients were as follows: 45 (68%) had ischemic heart disease (IHD), 7 (11%) had cerebrovascular disease, 8 (12%) had macroangiopathy, 4 (6%) had combined IHD and cerebrovascular, and 2 (3%) had combined macroangiopathy and cerebrovascular disease.

The general characteristics and biochemical variables of the study population

Demographic, clinical and biochemical data of different groups is shown in Table 1. A significant age difference was found between controls and CVD group ($p < 0.01$). Also a significant sex difference was found between controls and CVD group ($p = 0.01$) due to the high prevalence of males among our CVD patients. Diabetic patients had lower levels of physical activity ($p = 0.01$) as compared to controls and patients with CVD had lower levels of physical activity when compared to patients without CVD ($p = 0.005$). There was a significant association between CVD and the duration of diabetes ($p = 0.004$). Other clinical data such as BMI, SBP and DBP were significantly higher in both groups when compared to controls. CVD patients had higher frequency of hypertension ($p = 0.01$) as compared with patients without CVD.

The mean of serum PON1 enzyme concentration was significantly lower in patient groups compared to controls ($p = 0.006$). No significant difference was found between diabetic patients with CVD and those without CVD ($p > 0.05$).

Table 1 General characteristics and biochemical variables of the study population

Variable	Controls (n=50)	T2DM (n=68)	T2DM+CVD (n=66)
Age (Years)	51.24 7.62	51.75 6.00	58.20 7.12**
Sex (male/female)	27/23	38/30	50/16**
BMI (kg/m ²)	23.21 4.62	27.59 4.90*	29.81 5.55**
SBP (mmHg)	121.90 15.29	131.67 20.60*	144.14 21.16**
DBP (mmHg)	81.19 8.93	88.75 16.05*	90.14 9.27**
Hypertension (%)	-	33.3	94.3
Smokers (%)	9.5	11.1	14.3
Physical activity (%)	57.1	58.3	25.7**
Diabetes duration (Years)	-	8.50 7.41	13.00 6.97
Glucose (mg/dL)	84.90 9.61	146.83 58.31*	164.00 68.27**
HbA1c (%)	5.42 0.61	6.30 1.19*	6.61 1.37**
Triglyceride (mg/dL)	117.57 31.28	143.61 73.14*	169.00 66.43**
TC (mg/dL)	186.86 28.68	193.92 50.59*	206.71 44.98**
LDL-C (mg/dL)	111.95 15.95	129.22 42.05*	141.60 45.04**
HDL-C (mg/dL)	55.71 11.24	49.93 11.26*	41.97 12.79**
Serum PON1 (nmol/L)	18.87 17.04	7.82 5.44*	9.22 14.94**

Significance was confirmed by Bonferroni test.

*Significant p in comparison between controls and T2DM, **Significant p in comparison between controls and T2DM+CVD, Significant p in comparison between T2DM and T2DM+CVD.

BMI: body mass index; SBP: systolic blood pressure; DBP: diastolic blood pressure; HbA1c: hemoglobin A1c; TC: total cholesterol; LDL-C: low density lipoprotein cholesterol; HDL-C: high density lipoprotein cholesterol; VLDL-C: very low density lipoprotein cholesterol; PON1: paraoxonase-1.

PON1 polymorphisms genotype and allele distribution in patients vs. controls

Analysis of genotypes and alleles distribution of Q192R and L55M polymorphisms in T2DM, with and without CVD, and control subjects is shown in Table 2. Data revealed that the genotype and allele frequencies of Q192R were significantly different between controls and diabetic patients. Genotype 192QQ was detected in 66% of controls vs. 44.8% of diabetic patients, while QR and RR genotypes were more frequent in diabetic patients than in controls (44% and 11.2% vs. 24% and 10%, respectively), $p = 0.02$. Allele Q was found in 78% of controls vs. 66.8% of patients, while allele R was more frequent in patients (33.2%) than in controls (22%); $p = 0.03$.

As regards L55M, the frequency of genotype MM was higher in controls (58%) than in patients (33.5%), while the frequency of genotypes LM and LL were higher in patients than in controls (49.3% and 17.2% vs. 28% and 14%, respectively); $p = 0.009$. Allele M was found in 72% of controls vs. 58% of patients, while Allele L was more frequent in patients than in controls (42% vs. 28%); $p = 0.015$.

PON1 polymorphisms genotype and allele distribution in T2DM patients with and without CVD

Analysis of genotypes and alleles distribution of Q192R and L55M polymorphisms in T2DM and T2DM + CVD patients is shown in Table 3. Results showed a significant difference between the two groups with QQ genotype more frequent in T2DM patients than in T2DM + CVD patients (57.4% vs. 32%), while the frequency of QR and RR genotypes were significantly increased in T2DM + CVD (54.4% and 13.6% vs. 33.8% and 8.8%, respectively), $p = 0.01$. Allele Q was detected in 74% of patients without CVD vs. 59% of patients with CVD, while allele R was

more frequent in patients with CVD (41%) than in patients without CVD (26%); $p = 0.008$. No statistical significant difference was found between the two groups as regards L55M genotype distribution ($p = 0.9$) or alleles frequency ($p = 0.8$).

Effects of Q192R and L55M polymorphisms on PON1 serum enzyme concentration

Means of serum PON1 concentration according to different genotypes are shown in Table 4. Q192R polymorphism associated with low serum PON1 concentrations, and lowest concentrations were observed in QR and RR genotypes ($p = 0.04$). L55M did not associate with significant difference in enzyme concentration. Though patients with 55 L allele had higher enzyme concentrations, yet that was of no statistical significance ($p = 0.18$).

Association of Q192R with CVD in T2DM patients

Multiple logistic regression analysis was used to investigate potential correlations of 192R allele with other cardiovascular risk factors: age, male gender, smoking, BMI, SBP, TC, HDL, LDL and TG. We demonstrated statistical significant correlations between 192R and age (ORs 1.39, 95% CI 1.1 1.73, $p = 0.04$), male gender (ORs 0.94, 95% CI 0.45 0.86, $p = 0.016$), SBP (ORs 1.92, 95% CI 1.38 2.78, $p = 0.007$), TC (ORs 0.97, 95% CI 0.36 0.86, $p = 0.016$), HDL (ORs 2.21, 95% CI 1.38 2.79, $p = 0.001$) and LDL (ORs 3.20, 95% CI 2.31 6.58, $p = 0.001$) and TG (ORs 0.54, 95% CI 0.36 0.86, $p = 0.012$). No significant correlations were detected with smoking (ORs 0.34, 95% CI 0.29 0.79, $p = 0.23$) or BMI (ORs 0.54, 95% CI 0.24 0.63, $p = 0.12$).

Table 2 Genotypes and alleles frequency of PON1 Q192R and L55M polymorphisms in patients and control groups

	Control N=50	Patients N=134	OR (95% CI)	P value
Q192R				
QQ	33 (66%)	60 (44.8%)	Reference	0.02
QR	12 (24%)	59 (44%)	2.1 (1.2 4.8)	
RR	5 (10%)	15 (11.2%)	3.2 (1.3 6.6)	
L55M				
LL	7 (14%)	23 (17.2%)	Reference	0.009
LM	14 (28%)	66 (49.3%)	2.1 (1.4 5.5)	
MM	29 (58%)	45 (33.5%)	2.8 (1.7-6.2)	
Allele	2n=100	2n=268		
192Q	78 (78%)	179 (66.8%)	Reference	0.03
192R	22 (22%)	89 (33.2%)	1.9 (1.2 3.2)	
55 L	28 (28%)	112 (42%)	Reference	0.01
55 M	72 (72%)	156 (58%)	2.0 (1.1 3.5)	

Table 3 Genotypes and alleles frequency of PON1 Q192R and L55M polymorphisms in T2DM and T2DM+CVD patient groups

	T2DM N = 68	T2DM+CVD N=66	OR (95% CI)	P value
Q192R				
QQ	39 (57.4%)	21 (32%)	Reference	0.01
QR	23 (33.8%)	36 (54.4%)	1.7 (1.0 3.8)	
RR	6 (8.8%)	9 (13.6%)	2.1 (1.3 4.6)	
L55M				
LL	11 (35%)	12 (41%)	Reference	0.9
LM	34 (47%)	32 (45 %)	1.08 (0.8 1.4)	
MM	23 (18%)	22 (14%)	0.99 (0.7 1.2)	
Allele	2n=136	2n=132		
192Q	101 (74%)	78 (59%)	Reference	0.008
192R	35 (26%)	54 (41%)	1.3 (1.0 2.7)	
55 L	56 (59%)	56 (64%)	Reference	0.8
55 M	80 (41%)	76 (36%)	0.9 (0.8 1.1)	

Table 4 Means of serum PON1 concentration in different genotypes of Q192R and L55M polymorphisms

	Serum PON-1 meanSD (nmol/L)	P value
192 QQ	12.97 15.75	0.04
192QR+192RR	8.95 10.2	
55MM	9.07 6.22	0.18
55LM+LL	11.80 15.77	

Discussion

Oxidative modification of LDL is believed to be a major triggering event in the initiation and progression of atherosclerosis and cardiovascular events [24]. Recently, PON1 became the focus of intense research after the identification of its antioxidant properties, particularly its capacity to protect LDL from oxidative damage.

In the present study, we investigated the association of PON1 Q192R and L55M polymorphisms with the CVD in T2DM in Egyptian patients. We found an association of 192R and 55 L alleles with T2DM. Serum PON1 enzyme concentration was significantly reduced in our diabetic patients compared to controls and lowest enzyme concentrations were associated with the 192R allele. As regards the CVD, a significant association with 192R allele was demonstrated. Serum concentration of PON1 was not significantly different between diabetic patients with CVD and those without CVD. Allele 55 L was previously reported to be associated with higher concentrations of the enzyme [25], but lower enzyme activity [18]. L55 isoform was found to be more stable and resistant to proteolysis [26]. This could explain the relatively higher serum enzyme levels found in our L allele carrier patients.

Our findings support the previously reported hypothesis that reduced PON1 concentration and/or activity might play a role in the pathogenesis of T2DM itself. It is suggested that oxidative stress induced by reduced PON1 concentration and/or activity results in reducing glucose uptake from blood by muscle cells and develops into insulin resistance. Results from previous studies in which decreased PON1 activity and concentration has been linked with impaired glucose tolerance [27,28] and increased insulin resistance in healthy subjects [29,30] also support this hypothesis. Reduced PON1 concentration and activity in Type I and Type II diabetic patients has been reported in many studies [31-36].

Logistic regression analysis with other independent risk factors revealed an independent correlation of 192R with CVD risk. The Q192R polymorphism in the PON1 gene hampers the ability of PON1 to inhibit LDL oxidation, suggesting that the 192R allele is less effective than 192Q in preventing LDL oxidation [37,38]. Thus, carriers of the R allele are more susceptible to develop CVD than carriers of the Q allele. Randa et al. [39] reported that individuals with PON1 RR genotype have 9-fold risks to develop

CAD in Egyptians while those with the PON1 QR genotype have 4-fold risks. Also, the risk factors for CAD (diabetes, hypertension, dyslipidemia) were significantly associated with PON1 RR genotype.

Lower PON1 activity was associated with carotid arteriosclerosis and cerebral atherosclerosis in stroke patients [13], and was reported as a risk factor for CVD in systemic lupus patients [14] and CAD in type 2 diabetes in North-West Indians [15]. Also, an association between PON1 polymorphisms and CAD risk was demonstrated in Thai population, where the frequencies of 192R allele and 192RR genotype, as well as, 55 M allele and 55LM genotype were significantly higher in CAD patients [16].

In a study by Ito et al. [40] involving the PON1-192R allele and the environmental risk factors, PON1-192R allele was reported as the most predictive independent risk factor for coronary spasm followed by cigarette smoking. Also, plasma levels of TBARS (Thiobarbituric acid-reactive substances), markers of oxidative stress, were higher in RR than in QQ genotypes suggesting that PON1-192R allele may play an important role in the genesis of coronary spasm, probably by attenuating the suppression of oxidative stress.

In a recent study; including PON1 genotyping, activity and lipid profile and their association with significant coronary stenosis (SCS) in Tunisian population, PON1 activity was lower in patients with SCS than in controls. Low activity was associated with 192R and 55 L alleles. In the presence of diabetes, PON1-192RR genotype associated with an increased risk of SCS while 55MM genotype associated with lower risk [18]. In contrast, in a study on 589 patients (419 Caucasian, 120 South Asian, 50 other) to determine factors which modulate serum PON1 in type 2 diabetes, PON1 activity associated negatively with insulin resistance, triglycerides and 55 M allele; and positively with 192R allele. While PON1 concentration associated negatively with Caucasian ethnicity and duration of diabetes, and positively with 192R allele [19].

Altuner et al. declared that RR genotype associated with higher PON activity than QQ or QR genotypes, while LL genotype associated with higher PON activity than MM genotype in Turkish population [20]. In Asian Indians, CAD was associated PON1 192R allele. Haplotype LQ (55 L and 192Q) showed protective effect, while haplotype MR (55 M and 192R) was associated with increased risk of CAD [21]. A recent meta-analysis of 88 studies comprising 24,702 CAD cases and 38, 232 controls revealed a significant association of Q192R polymorphism with the disease risk [41]. The controversial data in the interpretation of CVD association with PON1 polymorphism might be attributed to ethnic variation, sample size, patients recruiting criteria, gene gene and gene environment interactions [42-45].

Conclusion

Our results indicate that PON1 192R and 55 L alleles are associated with T2DM, and suggest that they may play a role in the pathogenesis of the disease. Q192R is associated with CVD and lower serum enzyme concentration and might represent a novel risk factor for CVD in Egyptian patients with T2DM.

Competing interests

The authors declare that they have no competing interests.

Authors contributions

DE made the study design, contributed to selection of patients, participated in the acquisition, analysis and interpretation of data and wrote the drafting manuscript. DE, MK, DA, AI and IA participated in the genetic analysis. DE, MK, DA, AI and EA participated in the laboratory investigations. DE and MK performed the ELISA assay. All authors revised the manuscript and approved the final version.

Acknowledgements

This work was funded by the National Research Center (Research Project No. 10010309).

Received: 24 October 2014 Accepted: 14 December 2014

Published online: 20 December 2014

References

1. Diabetes Action Now. *An Initiative of the World Health Organization and the International Diabetes Federation* 2004 [http://www.who.int/diabetes/actionnow/en/DANbooklet.pdf]
2. Garber JA: **Attenuating CV, risk factors in patients with diabetes: clinical evidence to clinical practice.** *Diabetes Obesity Metab* 2002, **4**(suppl 1):5-12.
3. Chaudhary R, Likidilid A, Thavatchai Peerapatdit T, Damras Tresukosol D, Srisuma S, Ratanamaneechat S, Sriatanasathavorn C: **Apolipoprotein E gene polymorphism: effects on plasma lipids and risk of type 2 diabetes and coronary artery disease.** *Cardiovasc Diabetol* 2012, **11**:36-47.
4. Sahin I: **HDL-C as a new therapeutic target in the treatment of dyslipidemia.** *Med Sci* 2012, **1**(2):141-149.
5. Guo ZG, Li C, Zhong JK, Tu Y, Xie D: **Laboratory investigation of dysfunctional HDL.** *Chem Phys Lipids* 2012, **165**(1):32-37.
6. Quillen EE, Rainwater DL, Dyer TD, Carless MA, Curran JE, Johnson MP, Gring HH, Cole SA, Rutherford S, Maccluer JW, Moses EK, Blangero J, Almasly L, Mahaney MC: **Novel associations of nonstructural Loci with paraoxonase activity.** *J Lipids* 2012:189681.
7. Narshi CB, Giles IP, Rahman A: **The endothelium: an interface between autoimmunity and atherosclerosis in systemic lupus erythematosus?** *Lupus* 2011, **20**(1):5-13.
8. Tsimikas S, Brilakis ES, Miller ER, McConnell JP, Lennon RJ, Kornman KS, Witztum JL, Berger PB: **Oxidized phospholipids, Lp(a) lipoprotein, and coronary artery disease.** *N Eng J Med* 2005, **353**:46-57.
9. Gugliucci A, Kotani K, Kimura S: **Paraoxonase 1 in chronic kidney failure.** *J Lipids* 2012:726048.
10. Fuhrman B: **Regulation of hepatic paraoxonase-1 expression.** *J Lipids* 2012: 684010.
11. Andersen HR, Wohlfahrt-Veje C, Dalgard C, Christiansen L, Main KM, Nellesmann C, Murata K, Jensen TK, Skakkebk NE, Grandjean P: **Paraoxonase 1 polymorphism and prenatal pesticide exposure associated with adverse cardiovascular risk profiles at school age.** *PLoS ONE* 2012, **7**(5):e36830.
12. Seo D, Goldschmidt-Clermont P: **Paraoxonase gene family and atherosclerosis.** *Curr Atheroscler Rep* 2009, **11**:182-187.
13. Shenhar-Tsarfaty S, Waisskopf N, Ofek K, Shopin L, Usher S, Berliner S, Shapira I, Bornstein NM, Ritov Y, Soreq H, Ben Assayag E: **Atherosclerosis and arteriosclerosis parameters in stroke patients associate with paraoxonase polymorphism and esterase activities.** *Eur J Neurol* 2013, **20**(6):891-898.
14. Dasgupta S, Demirci FY, Dressen AS, Kao AH, Rhew EY, Ramsey-Goldman R, Manzi S, Kammerer CM, Kamboh MI: **Association analysis of PON genetic variants with serum paraoxonase activity and systemic lupus erythematosus.** *BMC Med Genet* 2011, **12**:7.
15. Gupta N, Binu KB, Singh S, Maturu NV, Sharma YP, Bhansali A, Gill KD: **Low serum PON1 activity: an independent risk factor for coronary artery disease in North-West Indian type 2 diabetics.** *Gene* 2012, **498**(1):13-19.
16. Likidilid A, Akrawinhawong K, Poldee S, Sriatanasathavorn C: **Paraoxonase 1 polymorphisms as the risk factor of coronary heart disease in a Thai population.** *Acta Cardiol* 2010, **65**(6):681-691.
17. Deakin SP, James RW: **The importance of high-density lipoproteins for paraoxonase-1 secretion, stability, and activity.** *Free Radic Biol Med* 2004, **37**:1986-1994.
18. Rejeb J, Omezzine A, Rebhi L, Boumaiza I, Mabrouk H, Rhif H, Rejeb NB, Nabili N, Douki W, Abdelaziz AB, Boughzala E, Bouslama A: **Association of PON1 and PON2 polymorphisms with PON1 activity and significant coronary stenosis in a Tunisian population.** *Biochem Genet* 2013, **51**(1-2):76-91.
19. Mackness B, Marsillach J, Elkeles RS, Godstrand IF, Feher MD, Rubens MB, Flather MD, Humphries SE, Cooper J, Mackness M: **Paraoxonase-1 is not associated with coronary artery calcification in type 2 diabetes: results from the PREDICT study.** *Dis Markers* 2012, **33**(2):101-112.
20. Altuner D, Ates I, Suzen SH, Koc GV, Aral Y, Karakaya A: **The relationship of PON1 QR 192 and LM 55 polymorphisms with serum paraoxonase activities of Turkish diabetic patients.** *Toxicol Ind Health* 2011, **27**(10):873-878.
21. Ahmad I, Narang R, Venkatraman A, Das N: **Two- and three-locus haplotypes of the paraoxonase (PON1) gene are associated with coronary artery disease in Asian Indians.** *Gene* 2012, **506**(1):242-247.
22. American Diabetes Association: **Diagnosis and classification of diabetes mellitus.** *Diabetes Care* 2010, **33**(Suppl. 1):S62-S69.
23. Friedewald WT, Levy RI, Fredrickson DS: **Estimation of the concentration of lowdensity lipoprotein cholesterol in plasma without use of the preparative ultracentrifuge.** *Clin Chem* 1972, **18**:499-502.
24. Hassan MA, Al-Attas OS, Hussain T, Al-Daghri NM, Alokail MS, Mohammed AK, Vinodson B: **The Q192R polymorphism of the paraoxonase 1 gene is a risk factor for coronary artery disease in Saudi subjects.** *Mol Cell Biochem* 2013, **380**:121-128.
25. Blatter Garin M-C, James RW, Dussoix P, Blanch H, Passa P, Froguel P, Ruiz J: **Paraoxonase polymorphism Met-Leu54 is associated with modified serum concentrations of the enzyme a possible link between the paraoxonase gene and increased risk of cardiovascular disease in diabetes.** *J Clin Invest* 1997, **99**(1):62-66.
26. Leviev I, Deakin S, James RW: **Decreased stability of the M54 isoform of paraoxonase as a contributory factor to variations in human serum paraoxonase concentrations.** *J Lipid Res* 2001, **42**:528-535.
27. Leviev I, Kalix B, Brulhart-Meynet MC, James RW: **The paraoxonase PON1 promoter polymorphism C-107T is associated with increased serum glucose concentrations in non-diabetic patients.** *Diabetologia* 2001, **44**:1177-1183.
28. Deakin S, Leviev I, Nicaud V, Brulhart-Meynet MC, Tiret L, James RW: **Paraoxonase-1 L55M polymorphism is associated with an abnormal oral glucose tolerance test and differentiates high risk coronary disease families.** *J Clin Endocrinol Metab* 2002, **87**:1268-1273.
29. Barbieri M, Bonafe M, Marfella R: **LL-paraoxonase genotype is associated with a more severe degree of homeostasis model assessment IR in healthy subjects.** *J Clin Endocrinol Metab* 2002, **87**:222-225.
30. Senti M, Tomas M, Fito M: **Antioxidant paraoxonase 1 activity in the metabolic syndrome.** *J Clin Endocrinol Metab* 2003, **88**:5422-5426.
31. Boemi M, Leviev I, Sirolla C, Pieri C, Marra M, James RW: **Serum paraoxonase is reduced in type 1 diabetic patients compared to non-diabetic, first degree relatives; influence on the ability of HDL to protect LDL from oxidation.** *Atherosclerosis* 2001, **155**:229-235.
32. Kordonouri O, James RW, Bennetts B: **Modulation by blood glucose levels of activity and concentration of paraoxonase in young patients with type 1 diabetes mellitus.** *Metab Clin Exp* 2001, **50**:657-660.
33. Letellier C, Durou MR, Jouanolle AM, Le Gall JY, Poirier JY, Rueland A: **Serum paraoxonase activity and paraoxonase gene polymorphism in type 2 diabetic patients with or without vascular complications.** *Diabetes Metab* 2002, **28**:297-304.
34. Mackness B, Durrington PN, Boulton AJ, Hine D, Mackness MI: **Serum paraoxonase activity in patients with type 1 diabetes compared to healthy controls.** *Eur J Clin Invest* 2002, **32**:259-256.
35. Agachan B, Yilmaz H, Karaali Z, Isbir T: **Paraoxonase 55 and 192 polymorphisms and relationship to serum paraoxonase activity and serum lipids in Turkish patients with non-insulin dependent diabetes mellitus.** *Cell Biochem Funct* 2004, **22**:163-168.

36. Altuner D, Suzena S, Atesa I, Kocb G, Aralb Y, Karakaya A: **Are PON1 Q/R 192 and M/L 55 polymorphisms risk factors for diabetes complications in Turkish population?** *Clin Biochem* 2012, **44**(5-6):372-376.
37. Regieli JJ, Jukema JW, Doevendans PA, Zwinderman AH, Kastelein JJ, Grobbee DE, van der Graaf Y: **Paraoxonase variants relate to 10-year risk in coronary artery disease: impact of a high-density lipoprotein-bound antioxidant in secondary prevention.** *J Am Coll Cardiol* 2009, **54**:1238-1245.
38. Mackness B, Mackness MI, Arrol S, Turkie W, Durrington PN: **Effect of the human serum paraoxonase 55 and 192 genetic polymorphisms on the protection by high density lipoprotein against low density lipoprotein oxidative modification.** *FEBS Lett* 1998, **423**:57-60.
39. Mohamed RH, Mohamed RH, Karam RA, Abd El-Aziz TA: **The relationship between paraoxonase-1-192 polymorphism and activity with coronary artery disease.** *Clin Biochem* 2010, **43**:553-558.
40. Ito T, Yasue H, Yoshimura M, Nakamura S, Nakayama M, Shimasaki Y, Harada E, Mizuno Y, Kawano H, Ogawa H: **Paraoxonase gene Gln192Arg (Q192R) polymorphism is associated with coronary artery spasm.** *Hum Genet* 2002, **110**(1):89-94.
41. Wang M, Lang X, Zou L, Huang S, Xu Z: **Four genetic polymorphisms of paraoxonase gene and risk of coronary heart disease: a meta-analysis based on 88 case-control studies.** *Atherosclerosis* 2011, **214**:377-385.
42. Rainwater DL, Rutherford S, Dyer TD, Rainwater ED, Cole SA, Vandeberg JL, Almasy L, Blangero J, Maccluer JW, Mahaney MC: **Determinants of variation in human serum paraoxonase activity.** *Heredity (Edinb)* 2009, **102**:147-154.
43. Prcourt LP, Amre D, Denis MC, Lavoie JC, Delvin E, Seidman E, Levy E: **The three-gene paraoxonase family: physiologic roles, actions and regulation.** *Atherosclerosis* 2011, **214**:20-36.
44. Luu HN, Kingah PL, North K, Boerwinkle E, Volcik KA: **Interaction of folate intake and the paraoxonase Q192R polymorphism with risk of incident coronary heart disease and ischemic stroke: the atherosclerosis risk in communities study.** *Ann Epidemiol* 2011, **21**:815-823.
45. Rodriguez-Esparragn F, Rodriguez-Przez JC, Hernandez-Trujillo Y, Macas-Reyes A, Medina A, Caballero A, Ferrario CM: **Allelic variants of the human scavenger receptor class B type 1 and paraoxonase 1 on coronary heart disease: genotype-phenotype correlations.** *Arterioscler Thromb Vasc Biol* 2005, **25**:854-860.

doi:10.1186/s40200-014-0125-y

Cite this article as: El-Lebedy *et al.*: Paraoxonase-1 gene Q192R and L55M polymorphisms and risk of cardiovascular disease in Egyptian patients with type 2 diabetes mellitus. *Journal of Diabetes & Metabolic Disorders* 2014 **13**:125.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

