

RESEARCH

Open Access



Risk factors associated with post-weaning diarrhoea in Austrian piglet-producing farms

René Renzhammer^{1*}, Sebastian Vetter², Marlies Dolezal³, Lukas Schwarz¹, Annemarie Käsbohrer² and Andrea Ladinig¹

Abstract

Post-weaning diarrhoea (PWD) is a frequent, multifactorial disease of piglets leading to increased mortality rates and high economic losses. Due to the emergence of multi-resistant *Escherichia coli* isolates and the ban of zinc oxide (ZnO) in the EU since June 2022, alternative measures to prevent PWD are urgently needed. While an abundance of feed supplements is described to prevent PWD, there are hardly any studies reflecting the current situation of PWD in the field. Thus, we aimed to identify differences in management practices, housing and feeding strategies between farms with PWD and farms without PWD. Data were personally collected using a semi-structured questionnaire in 257 Austrian piglet-producing farms. Farms with PWD in more than 10% of all weaned groups within twelve months prior to data collection were defined as case farms (n = 101), while the remaining 136 farms were defined as control farms. Data from 237 farms and 69 explanatory variables were analysed via penalized binary logistic regression using elastic-net in 100 different splits into randomly selected training and test datasets (80:20). Treatment with ZnO and/or colistin (136 farms) was negatively associated with PWD in all splits and had the biggest estimated absolute log odds ratio out of all tested variables. Implementation of an all-in/all-out system in the nursery units and administration of probiotics or horseradish also had preventive effects in most splits ($\geq 97\%$). A higher number of feeding phases for piglets within the first seven weeks of life and housing on fully slatted floors was associated negatively with the occurrence of PWD as well in $> 95\%$ of all splits. PWD was more likely to occur on farms having problems with neonatal diarrhoea or postpartum dysgalactia syndrome. While our data demonstrate that treatment with ZnO or colistin had the biggest statistical effect on PWD, we were able to identify other preventive measures like supplementation with probiotics or horseradish. Since implementation of all-in/all-out measures and fully slatted floors were also negatively associated with the occurrence of PWD on visited farms, we assume that reduction of bacterial load by the implementation of simple hygiene measures are still crucial to prevent PWD.

Keywords Post-weaning diarrhoea, PWD, Colistin, Zinc, Horseradish, Probiotics, Management, Housing, Hygiene

*Correspondence:

Renzhammer René
Rene.Renzhammer@vetmeduni.ac.at

¹Department for Farm Animals and Veterinary Public Health, University Clinic for Swine, University of Veterinary Medicine, Veterinärplatz 1, Vienna 1210, Austria

²Unit of Veterinary Public Health and Epidemiology, Institute of Food Safety, Food Technology and Veterinary Public Health, University of Veterinary Medicine, Veterinärplatz 1, Vienna 1210, Austria

³Platform for Bioinformatics and Biostatistics, Department of Biomedical Sciences, University of Veterinary Medicine, Veterinärplatz 1, Vienna 1210, Austria



© The Author(s) 2023, corrected publication 2023. **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated in a credit line to the data.

Background

Post-weaning diarrhoea (PWD) occurs within the first two weeks after weaning of piglets and can lead to high economic losses due to increased mortality rates, reduced growth rates and increased costs for treatment [1]. Moreover, PWD is also considered as one of the most frequent diseases in pig production worldwide. Although PWD is a multifactorial disease, ubiquitous enterotoxigenic *Escherichia (E.) coli* (ETEC) play a pivotal role in the pathogenesis of PWD [1, 2]. To a lesser extent other pathogens like enteropathogenic *E. coli* (EPEC) and *Salmonella* spp. are also potential infectious causatives of PWD [1, 3, 4]. Furthermore, the impact of rotavirus A and C on PWD is currently discussed controversially [5]. Nevertheless, feeding, management and housing conditions play a crucial role in the pathogenesis of PWD. In particular, the abrupt transition from highly digestible milk to less-digestible solid feed consisting of mainly plant based proteins, complex carbohydrates and several anti-nutritional factors overburdens the intestinal tract of piglets, which are usually weaned between their 21st to 28th day of life [2, 6]. Most notably, weaning is associated with anorexia of piglets within the first two days after weaning leading to villous atrophy and crypt hyperplasia in the small intestine [7–9]. Besides the reduction of nutrient absorption due to the villous atrophy, there is evidence of a decline in brush-border enzymes leading to inferior protein digestibility and malnutrition [9]. While low gastric pH is a crucial defence mechanism against ETEC, pH values in the stomach and small intestine increase significantly after weaning and consequently facilitate intestinal colonization with ETEC [10]. Higher gastric pH values after weaning are the result of a reduced HCl production due to anorexia, higher amounts of less-digestible proteins in the feed and decreased production of lactic acid produced from lactose [6, 11, 12]. Furthermore, deprivation of milk and increased pH values cause significant shifts in the intestinal microbial composition characterized by a decrease of gram-positive bacteria like *Lactobacillus* spp. and *Bifidobacterium* spp. and an increase of *Enterobacteriaceae* [13]. Besides multiple nutritional causes of PWD, the process of weaning significantly increases the level of immunosuppressive cortisol due to several stressors like separation from the sow, transfer to new pens and mixing of different litters [14].

While antimicrobials are widely applied to treat piglets with PWD, numerous reports on high antimicrobial resistance rates of porcine *E. coli* within the last few years emphasize the difficulties to choose adequate treatment of diseased piglets [15–17]. Thus, veterinarians frequently apply colistin (polymyxin E) for oral treatment of piglets with PWD, since resistance rates of porcine *E. coli* against colistin are considered to be low [18, 19]. However, since colistin is crucial to treat humans affected by

multi-resistant gram-negative bacteria, its use to treat animals is under debate [18].

As an alternative to antimicrobials, zinc oxide (ZnO) has been frequently applied in swine farms worldwide to treat piglets with PWD [20]. In general, zinc is a pivotal trace element for pigs, as it is essential for multiple biochemical processes [21]. Therefore, commercially available feed contains zinc at a concentration of 150 ppm, which is defined as the maximum nutritional concentration. First beneficial effects including a decreasing incidence of PWD can be observed, if piglets are treated with a concentration of 1,000 ppm ZnO [22]. In contrast to antimicrobial substances like colistin, there is evidence that ZnO does not act with a specific mechanism against *E. coli*, but rather improves gut health, digestion and the immune system of treated piglets in various ways. Described beneficial mechanisms include an increased expression of occludin, increased levels of superoxide dismutase and a reduction of pro-inflammatory cytokines [20, 23–25]. The combination of anti-oxidative, anti-inflammatory and immune stimulating mechanisms of ZnO might lead to lower incidences of intestinal inflammation and diarrhoea. Furthermore, treatment with ZnO (2,500 ppm) can also improve digestibility of nutrients due to increased production of amylase, carbopeptidase A, chymotrypsin, trypsin and lipase [26].

On the other hand, there is evidence that treatment of piglets with ZnO (2,500–3,000 ppm) for 21 days increases the prevalence and persistence of methicillin-resistant *Staphylococcus aureus* [27, 28] and antimicrobial resistance rates of porcine *E. coli* towards ampicillin, tetracycline and trimethoprim-sulfonamide [29–31]. In particular, environmental concerns led to the ban of ZnO in the European Union [32]. A prospective hazard analysis emphasizes the potential danger of ZnO applied in livestock for soil and ground water in the Netherlands, Flanders, north-western Germany and Denmark [33]. After the conduction of the latter study, the European Medicines Agency (EMA) concluded that disadvantages of ZnO usage outweigh its benefits leading to the EU-wide ban of all licensed medicinal products containing ZnO with a concentration beyond 150 ppm [32]. However, applicable alternatives for ZnO are yet to be identified. High antimicrobial resistance rates of porcine *E. coli* isolates, the goal of the European Commission to reduce usage of antimicrobial substances in livestock and the ban of ZnO since 26th of June 2022 emphasize the urgent need for applicable strategies to prevent PWD reliably. Although a plethora of potential alternatives to ZnO is described, ZnO has still been applied in most swine farms in Austria and other EU countries until 26th of June 2022, emphasizing the lack of a strategy to successfully prevent the occurrence of PWD.

Late weaning, reduction of crude protein and increase of crude fibre in the feed are well described methods to reduce the incidence of PWD [34–38]. Numerous feed additives have already been proposed as potential alternatives to ZnO [6, 20]. Feed additives, which potentially reduce the incidence of PWD include for example organic acids [39–41], probiotics [42–45], prebiotics [46, 47], essential oils and other plant based substances [48, 49], exogenous enzymes [50, 51], dehydrated porcine plasma [52], yeast [53] and antimicrobial peptides [54, 55]. However, while most studies on alternative strategies to reduce the usage of antimicrobial substances focus on feed additives, only a few studies report on the impact of management, housing conditions and biosecurity [56]. Due to the fact, that PWD is a multifactorial disease, it is crucial to evaluate all parameters, which potentially prevent or provoke the onset of diarrhoea. Furthermore, since most studies on prevention of PWD were conducted under experimental settings [56], they cannot reflect the situation in the field appropriately.

Thus, we aimed to reflect the frequency of PWD and the use of ZnO and/or colistin treatment in Austrian piglet-producing farms. Furthermore, we wanted to identify differences of management practices, housing conditions, feeding strategies and biosecurity measures between farms with PWD and farms without PWD.

Results

Descriptive analysis

In all 101 case farms (42.6%) first symptoms of PWD appeared between the first and 8th day post weaning (mean=3.8 days post weaning). Out of all 142 farms with regular treatment against PWD, 132 farms applied products containing ZnO, 56 of them in combination with colistin. While products containing only colistin were applied in four farms, marbofloxacin and gentamicin were used in two and four farms respectively. Means of most numerically coded variables like age at weaning, crude protein, crude fibre, feeding space per piglet, number of piglets per drinking spot, square meters per piglet and piglets per pen did not vary among case and control farms (Table 1).

Since PWD was perceived as a general herd problem, 97 farmers (40.9%) stated to have supplemented ZnO and/or colistin to the feed of at least 50% of all batches of weaned piglets and did not observe PWD within twelve months prior to data collection (Table 2). Further 45 farmers (19.0%) claimed to observe PWD regularly despite treatment with ZnO or colistin. PWD occurred on 56 other farms (23.6%), neither administering ZnO nor colistin at weaning. In 39 farms (16.5%) PWD was no problem and therefore administration of ZnO or colistin was not necessary at all. The occurrence of PWD was lower in farms supplementing probiotics (27%) compared

Table 1 Exploratory arithmetic means for numerically coded explanatory variables in the elastic-net model from 237 farms

	Age at weaning (days)	Number of feeding phases	Crude Protein g/kg DM	Crude Fiber g/kg DM	Feeding space per piglet (cm)	Number of piglets per drinking spot	Square meters per piglet	Piglets per batch
PWD	No	3.2	165.7	43.6	8.4	13.5	0.3	43.7
	Yes	2.9	164.9	45.2	8.1	15.4	0.3	44.7

PWD: post-weaning diarrhoea; DM: dry matter

to those without supplementation of probiotics (47%). Administered probiotics belonged to the genera *Pedio-coccus* (n=19), *Lactobacillus* (n=17), *Enterococcus* (n=9) and *Streptococcus* (n=3). Out of 32 farms adding horse-radish to the feed after weaning, three quarters did not have problems with PWD (Table 2). The effect of other supplementary substances based on plants could not be evaluated individually, as declaration of content was rarely provided. Thus, other feed additives based on plants were grouped together to the variable “herbs”.

Table 2 Occurrence of PWD and number of observations for categorical variables from 237 farms

		PWD				PWD	
		No	Yes			No	Yes
Treatment	No	39	56	Lactic acid	No	83	64
	Yes	97	45		Yes	53	37
Neonatal diarrhoea	No	56	24	Benzoic acid	No	85	56
	Yes	80	77		Yes	51	45
Postpartum dysgalactia syndrome	No	76	41	Sorbic acid	No	98	79
	Yes	60	60		Yes	38	22
Rearing of own gilts	No	78	55	Coke, phosphoric acid	No	110	75
	Yes	58	46		Yes	26	26
Nurse sows	No	95	62	Electrolytes, glucose	No	121	87
	Yes	41	39		Yes	15	14
Cross fostering	No	22	10	Apple pomace	No	123	92
	Yes	114	91		Yes	13	9
Weaning in farrowing units	No	98	74	Dairy products	No	122	96
	Yes	38	27		Yes	14	5
Grouping of weaned piglets	Sorted by size	63	40	Rock salt	No	122	81
	Litter wise	73	61		Yes	14	20
Automatic feeding system	No	96	80	Yeast	No	93	81
	Yes	40	21		Yes	43	20
Stimulation of feed intake	No	98	71	Tannins	No	119	92
	Yes	38	30		Yes	17	9
Weaner diet before weaning	No	33	23	Anticoccidials	No	93	66
	Yes	103	78		Yes	43	35
Main starch source	Corn	44	34	Water source	Mains water	34	26
	Barely	92	67		Well water	102	75
Wheat, rye, triticale	No	75	67	Calcareous water	No	54	46
	Yes	61	34		Yes	82	55
Main protein source	Plant	122	93	Disinfection of water pipes	No	110	85
	Animal	14	8		Yes	26	16
Main soy processing method	Heated	74	52	Looped water distribution system	No	119	95
	Extracted	62	49		Yes	17	6
Potato protein	No	43	31	Rinsing of water pipes	No	65	46
	Yes	93	70		Yes	71	55

Table 2 Occurrence of PWD and number of observations for categorical variables from 237 farms

		PWD				PWD	
		No	Yes			No	Yes
Linseed, chicory	No	80	60	Diameter of water pipes in nursery	1/2 inch	116	79
	Yes	56	41		> 1/2 inch	20	22
Oats	No	53	47	Drinkers	Only nipples	69	49
	Yes	83	54		Also bowls	67	52
Wheat bran	No	70	47	Rodent burden	Low	77	44
	Yes	66	54		High	59	57
Cellulose	No	77	59	Enrichment material	Only wood/chains	73	54
	Yes	59	42		Also straw/ropes	63	47
Fibre in creep feed	No	73	67	Microclimate zones	No	96	75
	Yes	63	34		Yes	40	26
Milk substitution	No	76	57	Heating of nursery before weaning	No	25	28
	Yes	60	44		Yes	111	73
Muesli	No	100	80	All-in/all-out in nursery units	No	73	72
	Yes	36	21		Yes	63	29
Commercial peat	No	111	92	Hospital pens	No	53	46
	Yes	25	9		Yes	83	55
Probiotics	No	101	88	Fully slatted floors	No	70	36
	Yes	35	13		Yes	66	65
Horseradish	No	112	93	Plastic or metal floors	No	92	53
	Yes	24	8		Yes	44	48
Herbs	No	119	89	Regular cleaning and disinfection	No	60	36
	Yes	17	12		Yes	76	65
Formiate, acetate, or propionate	No	41	35	Usage of detergents	No	66	48
	Yes	95	66		Yes	70	53
Citric acid	No	78	56	Equipment only for nursery units	No	81	61
	Yes	58	45		Yes	55	40
Lauric acid	No	120	90	Hygiene lock	No	105	85
	Yes	16	11		Yes	31	16

PWD: post-weaning diarrhoea

Elastic-net results

Variables associated with PWD in at least 60% of elastic-net splits (equivalent to a minimum of 60% of non-zero penalized logit estimates for the variable across 100 different splits into training and test data), were considered to have at least some impact on the occurrence of PWD. A potential influence on the output of PWD could be determined for 22 out of 69 analysed explanatory variables using elastic-net (Table 3). Eight variables (treatment, neonatal diarrhoea, all-in/all-out system, probiotics, postpartum dysgalactiae syndrome (PDS),

Table 3 Variables associated with PWD in over 60% of all training-test data splits using elastic-net (n = 22/69)

Variable	5% Percentile	Median	95% Percentile	% Non zero	Reduced chance of PWD
Treatment	-1.40	-1.07	-0.71	100	Yes
Neonatal diarrhoea	0.15	0.43	0.73	100	No
All in-all out in nursery units	-0.82	-0.51	-0.19	99	Yes
Probiotics	-0.99	-0.62	-0.18	99	Yes
PDS	0.05	0.36	0.68	98	No
Horseradish	-1.03	-0.66	-0.24	97	Yes
Number of feeding phases	-0.38	-0.20	-0.03	96	More phases
Solid floors	0.02	0.35	0.65	95	No
Commercial peat	-0.94	-0.54	0.00	94	Yes
Rodent burden	0.00	0.22	0.51	90	Low
Nurse sows	0.00	0.28	0.53	90	No
Plastic or metal floors	-0.60	-0.30	0.00	89	Yes
Dairy products	-0.74	-0.30	0.00	86	Yes
Looped water distribution system	-0.64	-0.23	0.00	82	Yes
Rock salt	-0.67	-0.21	0.00	82	Yes
Coke / phosphoric acid	0.00	0.28	0.61	82	No
Cross fostering	0.00	0.28	0.75	80	No
Regular cleaning and disinfection of nursery units	-0.44	-0.14	0.00	76	Yes
Automatic feeding	-0.55	-0.20	0.00	75	Yes
Yeast	-0.48	-0.16	0.00	73	Yes
Fibre in creep feed	-0.34	-0.07	0.00	69	Yes
Electrolytes, glucose	-0.63	-0.07	0.00	60	Yes

% Non zero: percentage of splits with log odds (ratios) unequal zero; ZnO: zinc oxide; PDS: postpartum dysgalactia syndrome; Variables for which the 5 and 95 percentile of the logit distribution across 100 elastic-net splits did not include zero are marked in bold




Fig. 1 Violin plots for area under the curve (AUC) and misclassification rates across 100 elastic-net models

horseradish, number of feeding phases and solid floors) were associated with the occurrence of PWD in ≥95% of all splits indicating a high impact on the incidence of diarrhoea (Table 3). Regular administration of ZnO and/or colistin, implementation of an all-in/all-out system in the nursery units, usage of probiotics and administration of horseradish had negative estimated log odds ratios in ≥97% of all splits demonstrating a potential preventive effect on the occurrence of PWD. The number of feeding phases for piglets from farrowing until the 21st day

after weaning and housing on fully slatted floors were also negatively associated with the occurrence of PWD in at least 95% of all splits. Log odds ratios were positive for neonatal diarrhoea and PDS in ≥98% of all splits demonstrating that farms having problems with neonatal diarrhoea or PDS were also more likely to have problems with PWD.

Out of all 69 variables tested, treatment with ZnO and/or colistin had the biggest absolute median (-1.07) of the estimated log-odds ratio distribution across 100 elastic-net splits. (Additional file 1). To assess model fit we looked at the area under the curve (AUC) which varied between 0.51 and 0.88. Misclassification rates ranged from 0.21 to 0.49 in the test data across all 100 splits (Fig. 1).

Variables excluded from statistical analysis

PWD was no problem in eight out of nine farms not offering an additional transition feed between creep feed and weaner diet. Variables on transition feed were excluded from analysis due to collinearity with the number of feeding phases. Neither problems with PWD nor treatment against PWD were reported from any of the six farms supplementing creep feed with seaweed. PWD occurred in one out of six farms using disinfection

troughs in front of the nursery units. Out of eight farmers applying a modified live vaccine against ETEC, three claimed that problems with PWD occurred despite vaccination. Supplementation of seaweed, usage of disinfection troughs and vaccination of piglets against ETEC were excluded from analysis because the number of observations for one factor level was <10 (Additional file 2).

Discussion

Administration of ZnO or colistin revealed strongest effects on the occurrence of PWD out of all 69 analysed explanatory variables. Similar to the results of an experimental study demonstrating that treatment with ZnO (2,500 ppm) reduced the incidence of diarrhoea by approximately 50% [57], in our study the percentage of farms with PWD not using ZnO was almost twice as high (58.9%) compared to farms with regular administration of ZnO (31.7%). However, 45 farmers reported to observe PWD regularly in nursery pens despite treatment of weaned piglets. Thus, ZnO does not prevent PWD completely, especially if other protective measures might be neglected. On the other hand, our data demonstrate that administration of other feed additives such as probiotics or horseradish can prevent PWD, although effects were less pronounced compared to ZnO. Therefore, we assume that a combination of different management practices along with the administration of other feed additives may compensate at least partially for the administration of ZnO to prevent PWD.

Besides treatment with ZnO and/or colistin, only neonatal diarrhoea was also associated with the occurrence of PWD in all splits, which was not surprising since ETEC expressing F4 fimbriae can cause both, neonatal diarrhoea and PWD [1]. Furthermore, piglets which have already had neonatal diarrhoea might be more susceptible to PWD due to previous damage in the intestines and worse body condition compared to piglets without diarrhoea during the suckling period. Therefore, several approaches to prevent neonatal diarrhoea like sow vaccination against ETEC, *Clostridium perfringens* or rotavirus A, as well as treatment of piglets with anticoccidials might prevent PWD indirectly. However, those variables were excluded from statistical analysis due to collinearity with the occurrence of neonatal diarrhoea. Similar to PWD, PDS is a multifactorial disease with *E. coli* playing a key role in its pathogenesis [58]. Thus, parameters increasing the risk for PDS might also increase the risk of PWD. Moreover, dysgalactia of sows leads to malnutrition and reduced growth rates of suckling piglets resulting in a higher susceptibility to disease.

Despite the fact that good feeding management and choice of beneficial ingredients in the feed are described to be very crucial to prevent PWD [20], the number of

feeding phases until 21 days after weaning was the only variable on feeding management which was associated with the occurrence of PWD in at least 95% of all splits. This could be due to the lack of information on quantities of ingredients contained in purchased commercial products. While in most Austrian farms corn, barley and wheat are produced on-site and exact quantities used in feed were available, protein sources were predominantly supplemented with commercial products. Since information on exact quantities of all ingredients contained in each commercial product (n=96) was not available, precise evaluation of feed ingredients was not feasible. Although we had to exclude most variables on creep feed from statistical analysis due to collinearity, information whether creep feed was offered or not, is still contained in the variable “number of feeding phases”. This could explain why farms with more feeding phases were less likely to have problems with PWD, since it was demonstrated before that administration of creep feed can decrease the risk of PWD [59]. In addition, creep feeding containing fibre had preventive effects in 69% of all splits. While van Hees et al. did not observe significant differences in average daily gain (ADG) between piglets getting low-fibre creep feed and piglets with high-fibre creep feed after challenge with ETEC [60], Choudhury et al. demonstrated a reduced incidence and duration of PWD in piglets fed fibrous pre-weaning diet compared to the control group receiving only milk [61].

Despite the fact that reduction of crude protein and increase of crude fibre are widely known strategies to prevent PWD [36], we did not observe any differences of both variables among case and control farms. Our results also go in line with observations of another investigation [62]. In most studies evaluating the effect of dietary protein level, incidence of diarrhoea was higher in piglets receiving feed with protein levels ranging from 20 to 26% compared to groups receiving 16–19% crude protein in feed [34–38, 63, 64]. Therefore, we assume that the level of protein did not have an influence on the occurrence of PWD in our data, since protein levels of most analysed farms were already comparably low (16–17%). Hence, reduction of crude protein and increase of crude fibre in the diet of weaned piglets alone might not be sufficient to prevent PWD under those circumstances. Albeit in our study farms with automatic feeding in the nursery units had less problems with PWD compared to farms feeding piglets manually, no differences between automatic and manual feeding were observed in a similar study conducted in Finland [65].

In contrast to feeding management and feed composition, several variables on feed additives (8/18) were associated with PWD in at least 60% of all splits. Out of all analysed feed additives, supplementation of probiotics or horseradish had the strongest effects on the occurrence

of PWD (Table 3). While strains of *Lactobacillus* spp., *Bifidobacterium* spp. and *Streptococcus* spp. are administered most frequently in pig production worldwide [66], *Bifidobacterium* spp. was not administered in a single analysed farm. Administration of probiotics might have multiple effects on the intestines, similar to treatment with ZnO, including an increase of villus heights in the jejunum [67] and decreased levels of tumour necrosis factor alpha [67, 68] and interleukin-1 beta [68]. Horseradish (*Armoracia rusticana*) is a plant commonly grown in Styria and has not been described to have preventive properties against PWD so far. Potential explanations why farms administering horseradish were less likely to have problems with PWD could be provided by confirmed antimicrobial properties of mustard essential oils, which are contained in horseradish [69, 70]. Unique effects of the administration of horseradish on the prevalence of PWD are yet to be evaluated in experimental trials or additional field studies.

Humic acids contained in peat [71] could explain why the majority of farms administering peat reported to have no problems with PWD (25/34). Administration of 2,000 ppm sodium humate to 24 piglets resulted in significantly higher ADG and decreased levels of pro-inflammatory cytokines compared to the untreated control group [72]. Since offering pig peat permits expression of natural behaviour and reduces fighting at weaning, it might also decrease stress levels of piglets and further decrease the probability of PWD [73]. Nevertheless, peat is a frequently described hazard for the introduction of pathogens [74, 75]. Therefore, offering peat to prevent PWD should be reconsidered based on biosecurity protocols. Since whey protein and casein are well digestible for three to four week old piglets, supplementation of various dairy products, which was also negatively associated with the occurrence of PWD in 86% of all splits, could be considered as a preventive measure against PWD as well [76].

There is currently no evidence that coke could also prevent PWD. In contrast, our data demonstrate that farms administering coke to piglets were more likely to have problems with PWD. Phosphoric acid, an ingredient of coke is an inorganic acid and has never been described to prevent PWD [77]. We assume that administration of coke to piglets might be linked to the fact that coke is still used as a home remedy against diarrhoea. This old wives' tale could be related to the fact that citric acid was used instead of phosphoric acids in previous formulations of coke. In comparison to phosphoric acid, it has already been demonstrated that supplementation of citric acid can prevent diarrhoea evidently [41, 77, 78].

PWD was also less likely to occur in farms administering live yeast and electrolytes, but effects were weaker compared to administration of probiotics or horseradish.

Capability of yeast (*Saccharomyces cerevisiae*) to reduce incidence of PWD and shedding of *E. coli* has already been demonstrated in experimental studies [53, 79]. Additionally, similar to ZnO it was demonstrated that administration of yeast decreased serum concentrations of pro-inflammatory cytokines [80]. Application of glucose or electrolytes could generally temper symptoms of PWD [81].

Pronounced effects of variables like all-in/all-out system in the nursery or the type of floor in most splits highlight the importance of general management measures in order to prevent PWD (Table 3). Consequent compliance of an all-in/all-out system in the nursery units can reduce the frequency of PWD, probably due to a reduction of pathogen transmission between piglets of different age groups [82]. While implementation of an all-in/all-out system proved to reduce the prevalence of various enteropathogens [83–85], it had no significant effect on the prevalence of ETEC in a Canadian study [62]. In addition to an all-in/all-out system, other measures reducing pathogen transmission between individuals might also be beneficial to prevent PWD. One example could be to avoid or reduce cross fostering, as implementation of this management practice was associated positively with the occurrence of PWD in 80% of all splits.

Despite discussions on fully slatted floors in pig production in the context of animal welfare, our data emphasize the advantages of fully slatted floors in the context of hygiene measures, since PWD was less likely to occur in farms keeping piglets on fully slatted floors. Due to easier cleaning and disinfection and the quick removal of faeces through the slats, bacterial load is considered to be lower in pens with fully slatted floors compared to solid floor pens [86, 87]. On the other hand, diarrhoea can be observed more easily in pens with solid floor. Similar to our results, Lozano et al. demonstrated that piglets housed on fully slatted plastic floors were less likely to develop PWD than piglets housed on concrete floors [88]. Berrocoso et al. demonstrated that husbandry of weaned piglets under optimal hygienic conditions including regular cleaning with detergents and disinfection resulted in significantly lower incidence of PWD and better ADG of weaned piglets [89]. Since investigated farms which regularly cleaned and disinfected nursery units were less likely to have problems with PWD, our results emphasize that hygienic measures are crucial to decrease ETEC burden and shall not be neglected in order to prevent PWD.

Besides environment, drinking water and water pipes could be another reservoir of ETEC [90]. Due to the fact that looped water distribution systems have less dead endings compared to branched water distribution systems, accumulation of sediments and biofilm formation is less likely [91]. Thus, infections with ETEC via

contaminated water coming from water pipelines with branched endings could be more likely, since ETEC are frequently recovered from biofilms in water pipes [90, 92].

In addition, rats and mice can also act as a vehicle for introduction of various *Enterobacteriaceae* like *Salmonella* spp. [93, 94]. Even though there are currently no data on the relation between rodent burden and incidence of PWD, our results emphasize that high rodent burden could presumably lead to increased transmission rates of ETEC and higher bacterial loads. Thus, strict measures against rodents might potentially decrease the probability of occurrence of PWD in swine farms.

Data collection through questionnaires heavily depends on perspectives and honesty of participants. Thus, bias due to dishonesty, subjective assessments of farmers and recall bias cannot be excluded. However, in order to reduce further bias related to subjectivity, data were always collected by the same person, who visited each farm to ensure consistent quality of collected data. Especially, recall bias for classification into case and control farms might have been high due to difficulties to estimate the number of weaned batches having problems with PWD. Additionally, farmers spending more time in the stables might be more likely to observe PWD. Furthermore, separate analysis for ZnO and colistin was not feasible, since products containing both substances were administered in numerous farms. Categorization of the variable treatment into “yes” or “no” for farms treating at least 50% of all batches of weaned piglets with antibiotics other than colistin, was done based on the results of a previous study with *E. coli* isolates recovered from Austrian swine farms [16]. However, antimicrobial resistance testing was not performed for *E. coli* from the visited farms in the current study.

Since an abundance of factors contributes to the pathogenesis of PWD, the number of variables could not be further limited without deleting pivotal parameters. Thus, elastic-net model was chosen to assess the influence of each variable for the output of PWD. While elastic-net is able to handle datasets with numerous predictors, it cannot deliver p-values. Therefore, we decided to renounce p-values. On the other hand, by using this method we were able to identify more variables with an influence on the output of PWD compared to previous similar analyses on PWD [62, 65]. McFadden pseudo R^2 accounting for 0.47 implicates that our dataset was able to partially explain the observed variance. However, since PWD is a complex, multifactorial disease the unexplained variance was no surprise. Further, certain factors influencing the output of PWD might have not been included in the questionnaire or during data processing.

Conclusions

Out of all analysed variables, treatment with ZnO and/or colistin was most preventive against PWD. Nevertheless, we were able to identify alternative preventive measures like supplementation of probiotics or horseradish, which might contribute to a lower incidence of PWD. However, their effects need to be evaluated further. We could also demonstrate that the occurrence of PDS and neonatal diarrhoea was associated positively with the occurrence of PWD on visited farms. Since implementation of all-in/all-out measures and fully slatted floors were clearly negatively associated with the probability of PWD, reduction of bacterial load by the implementation of simple hygiene measures might still be pivotal to prevent PWD.

Methods

Data collection

A semi-structured questionnaire was used in order to receive sufficient information from Austrian piglet producing farms (Additional file 3). In total, the questionnaire consisted of 156 questions concerning (a) general farm data, (b) suckling piglet period, (c) weaning management, (d) housing of piglets, (e) feed & feeding management, (f) feed additives & other preventive methods, (g) water supply, (h) hygiene and biosecurity, and (i) occurrence of PWD, including treatments with antimicrobials or ZnO (Table 4).

Study design & setting of the study

Pig farmers were first contacted with the help of their respective herd veterinarian ($n=40$) and asked to participate in the study. Farmers agreeing to participate signed a consent form and were always interviewed on the farm by the same person in a standardised way. Data were collected from 257 piglet-producing farms throughout Austria from May 2020 to March 2021 (Tables 5 and 6). The estimated time for the survey including measuring the size of two pens in the nursery units accounted for approximately 75 min.

Table 4 Number of questions in each category included in the questionnaire

Category	Number of questions
General farm data	14
Suckling piglet period	19
Weaning management	12
Housing of piglets	13
Feed & feeding management	18
Feed additives & other preventive methods	9
Post-weaning diarrhoea & treatments	28
Water supply	15
Hygiene and biosecurity	28

Table 5 Regional distribution and farming system of the 257 visited farms

	Number of visited farms	
	Federal state	Upper Austria
	Lower Austria	81
	Styria	70
	Carinthia	16
	Burgenland	5
Farming system	Conventional farming	242
	Organic farming	15

Table 6 Farm characteristics of all 257 visited farms

	Median	Minimum	Maximum
Number of sows	80	12	2300
Weaned piglets/sow/year	25.2	14.8	39.5

Data processing

All data (257 farms with 225 variables) were transferred to Microsoft® Excel and analysed anonymously. In order to represent the majority of piglet-producing farms in Austria, organic farms (n=15), farms with less than 14 sows (n=3) and farms with over 500 sows (n=2) were excluded from further analysis. All variables with missing data were excluded from statistical analysis (n=39). Those variables predominantly derived from answers of sub-questions, which were NA by default depending on the answer to the previous question (30/39). Several variables, like those on general farm data and vaccination protocols against pathogens associated with respiratory disease were excluded as well (n=46), since they were considered as irrelevant for PWD. Categorical variables were also excluded if the number of observations for one factor level was below ten (n=21). To avoid exclusion of too many variables, certain values (i.e.: metal floors & plastic floors) or variables (i.e.: coke & phosphoric acid) were combined, if the combination was biologically reasonable. The variable treatment was defined as yes, if either ZnO with a concentration beyond 1,000 ppm or colistin were supplemented at weaning for at least 50% of all batches of a farm within the year prior to data collection (n=136). Six additional farms applying marbofloxacin (n=2) or gentamicin (n=4) instead of ZnO or colistin were also included in the treatment group. For the sake of simplicity, treatment with marbofloxacin and gentamicin are not explicitly mentioned further. The variable treatment was defined as “no” for the remaining farms including farms regularly applying aminopenicillins (n=10), tetracyclines (n=3) or macrolides (n=3) after weaning, due to high resistance rates of ETEC towards those substances [16]. Nevertheless, it is likely that those antibiotics were predominantly applied to treat piglets against other diseases. Questions on PWD and treatment aimed to reflect the frequency of PWD and ZnO and colistin usage in Austrian swine farms. Thus, besides the

variable treatment, variables from that category were also excluded from further analysis (n=27). Details on complete data processing are provided in Supplementary File 2.

Definition of case and control farms

If the farmer stated to have observed at least one piglet with diarrhoea in at least 10% of all weaning groups in the first two weeks after weaning within twelve months prior to data collection, the farm was categorized as case farm (101/237).

Statistical analysis

All statistical analyses were performed in R version 4.0.2 [95]. The remaining data (237 farms and 92 explanatory variables) were imported into the statistics software R and tested for collinearity. In case of collinearity (variance inflation factor > 3) calculated in package *car* v3.0.12 [96], applying function *vif* on a linear model containing all explanatory variables, the variable with a lower expected biological relevance was excluded (n=23) [58]. In the end, 69 variables remained in the final data set for statistical analysis (Tables 1 and 2). Since application of a standard (generalized) linear model with 237 farms and 69 explanatory variables would have led to a severe overfitting problem, data were analysed via penalized binary logistic regression using elastic-net implemented in package *glmnet* v4.1.4 [97]. Elastic-net can be thought of as a combination of shrinkage (Ridge Regression) and variable selection (LASSO) methodology. The binary response was coded “0” for control farms and “1” for case farms. The function *createDataPartition* from package *caret* v6.0-92 [98] was applied to split questionnaires from 237 farms into a randomly selected independent training- and test dataset comprising of 80% and 20% of farms, respectively.

Cross validation was used to fine-tune the elastic-net parameters in the training data by applying function *cv.glmnet* with option *family*="binomial"; *standardize*=T and *maxit*=500,000. Model performance was assessed in the independent test data applying function *assess.glmnet* using option *family*="binomial" and *s*="lambda.min". “lambda.min” contains stored regularization parameter giving the minimum mean cross-validated error in the training data. Due to sample size limitations, we noticed that different seeds used for splitting the data into training and test data, resulted in somewhat different variable selections and size of estimated logits. Thus, we decided to create 100 different random splits (training data:test data=80:20) using function *set.seed(i)* in a loop over our elastic-net analysis, conducted as described above. Afterwards, model performance was assessed via the AUC metric and misclassification rates summarized and visualized as violin-plots (Fig. 1). Furthermore, we report size

of penalized estimated coefficients on a logit level and the number of times out of 100 (i.e.%) an explanatory variable was selected for the model (Supplementary File 1). Negative estimated coefficients on logit level, imply an odds ratio < 1, and the variable is negatively associated with the occurrence of PWD or has a preventive effect towards PWD. Variables with positive coefficients imply an odds ratio > 1 and are positively associated with the occurrence of PWD increasing the chance of PWD on a farm.

Abbreviations

PWD	Post-weaning diarrhoea
<i>E. coli</i>	<i>Escherichia coli</i>
ETEC	Enterotoxigenic <i>E. coli</i>
EPEC	Enteropathogenic <i>E. coli</i>
ZnO	Zinc oxide
PDS	Postpartum dysgalactia syndrome
ADG	Average daily gain

Supplementary Information

The online version contains supplementary material available at <https://doi.org/10.1186/s40813-023-00315-z>.

Additional file 1: Title: Logit distribution of all 69 tested explanatory variables across 100 elastic-net splits. Description: Results from elastic net analysis including percentiles (p5, p10, p25, p50, p75, p90, p95) and % non zero (percentage of splits with log odds (ratios) unequal zero) of estimated logits from all 69 explanatory variables across 100 elastic-net splits.

Additional file 2: Title: Data processing. Description: Complete data processing including the pathway from the original question provided in Additional file 3 to the final variables included in the statistical analysis. Additional file 2 includes English translations of all 156 original German questions. Information on variable type and reason for exclusion of the respective variable are also provided in Additional file 2.

Additional file 3: Title: Questionnaire. Description: Original questionnaire including all 156 questions in German language.

Acknowledgements

The authors thank all 257 farmers, who participated in the study and all herd veterinarians as well as the Landwirtschaftskammer Niederösterreich, who helped to get in contact with the respective farmers. The authors thank Prof. Barbara Metzler for nutritional consultation. The authors thank Dr. Heinrich Kreutzmann for critically reading the manuscript.

Author Contribution

A.L., A.K., L.S. and R.R. designed the study; R.R. was responsible for data collection on all 257 farms; R.R. performed data processing with the help of A.L., M.D. and A.K.; M.D., R.R. and S.V. conducted the statistical analysis; R.R. wrote the draft; A.L., A.K., L.S., M.D. and S.V. proofread the manuscript. All authors read and approved the final manuscript.

Funding

The study was conducted under the responsibility of the University Clinic for Swine without any funding.

Data Availability

All collected data are available as supplementary material.

Declarations

Ethics approval and consent to participate

Each participant of the questionnaire signed a consent for participation.

Consent for publication

Each participant of the questionnaire accepted by signature on a written consent that data is used for publication.

Competing interests

The authors declare that they have no competing interests.

Received: 20 September 2022 / Accepted: 19 April 2023

Published online: 11 May 2023

References

1. Fairbrother JM, Nadeau É, Gyles CL. *Escherichia coli* in postweaning diarrhea in pigs: an update on bacterial types, pathogenesis, and prevention strategies. *Anim Health Res Rev.* 2005;6(1):17–39.
2. Lallès JP, Bosi P, Smidt H, Stokes CR. Nutritional management of gut health in pigs around weaning. *Proc Nutr Soc.* 2007;66(2):260–8.
3. Janke BH, Francis DH, Collins JE, Libal MC, Zeman DH, Johnson DD. Attaching and effacing *Escherichia coli* Infections in Calves, Pigs, lambs, and Dogs. *J VET Diagn Invest.* 1989;1(1):6–11.
4. Griffith RW, Carlson SA, Krull AC. Salmonellosis. In: Zimmerman JJ, Kariker LA, Ramirez A, Schwartz KJ, Stevenson GW, Zhang J, editors. *Diseases of Swine*. Hoboken: Wiley-Blackwell; 2019. pp. 912–25.
5. Lecce JG, Balsbaugh RK, Clare DA. Rotavirus and hemolytic enteropathogenic *Escherichia coli* in Weanling Diarrhea of Pigs. *J Clin Microbiol.* 1982;16:715–23.
6. Heo JM, Opapeju FO, Pluske JR, Kim JC, Hampson DJ, Nyachoti CM. Gastro-intestinal health and function in weaned pigs: a review of feeding strategies to control post-weaning diarrhoea without using in-feed antimicrobial compounds: feeding strategies without using in-feed antibiotics. *J Anim Physiol Anim Nutr.* 2013;97(2):207–37.
7. Pluske JR, Hampson DJ, Williams IH. Factors influencing the structure and function of the small intestine in the weaned pig: a review. *Livest Prod Sci.* 1997;51(1–3):215–36.
8. van Beers-Schreurs HMG, Nabuurs MJA, Vellenga L, van der Valk HJK, Wensing T, Breukink HJ. Weaning and the Weanling Diet Influence the Villous height and crypt depth in the small intestine of Pigs and alter the concentrations of short-chain fatty acids in the large intestine and blood. *Nutr.* 1998;128(6):947–53.
9. Hedemann MS, Hojsgaard S, Jensen BB. Small intestinal morphology and activity of intestinal peptidases in piglets around weaning. *J Anim Physiol Anim Nutr.* 2003;87(1–2):32–41.
10. Prohászka L, Baron F. The Predisposing Role of High Dietary Protein Supplies in Enteropathogenic *E. coli* Infections of Weaned Pigs. *Zentralblatt Veterinarmedizin Reihe B.* 2010;27(3):222–32.
11. Manners MJ. The development of digestive function in the pig. *Proc Nutr Soc.* 1976;35(1):49–55.
12. Shimizu M, Terashima T. Appearance of Enterotoxigenic *Escherichia coli* in Piglets with Diarrhea in connection with feed changes. *Microbiol Immunol.* 1982;26(6):467–77.
13. Guevarra RB, Lee JH, Lee SH, Seok MJ, Kim DW, Kang BN, Johnson TJ, Iaconson RE, Kim HB. Piglet gut microbial shifts early in life: causes and effects. *J Anim Sci Biotechnol.* 2019;10(1):1.
14. Colson V, Martin E, Orgeur P, Prunier A. Influence of housing and social changes on growth, behaviour and cortisol in piglets at weaning. *Physiol Behav.* 2012;107(1):59–64.
15. Luppi A, Bonilauri P, Dottori M, Gherpelli Y, Biasi G, Meriardi G, Maioli G, Martelli P. Antimicrobial Resistance of F4+ *Escherichia Coli* isolated from Swine in Italy. *Transbound Emerg Dis.* 2015;62(1):67–71.
16. Renzhammer R, Loncaric I, Roch FF, Pinior B, Käsbohrer A, Spermser J, Ladinig A, Unterwiesing C. Prevalence of virulence genes and Antimicrobial Resistances in *E. coli* Associated with neonatal Diarrhea, Postweaning Diarrhea, and Edema Disease in Pigs from Austria. *Antibiotics.* 2020;9(4):208.
17. Smith MG, Jordan D, Chapman TA, Chin JJC, Barton MD, Do TN, Fahy VA, Fairbrother JM, Trott DJ. Antimicrobial resistance and virulence gene profiles in multi-drug resistant enterotoxigenic *Escherichia coli* isolated from pigs with post-weaning diarrhoea. *Vet Micro.* 2010;145(3–4):299–307.
18. Rhouma M, Beaudry F, Letellier A. Resistance to colistin: what is the fate for this antibiotic in pig production? *Int J Antimicrob Agents.* 2016;48(2):119–26.
19. Luppi A. Swine enteric colibacillosis: diagnosis, therapy and antimicrobial resistance. *Porc Health Manag.* 2017;3(1):16.

20. Bonetti A, Tugnoli B, Piva A, Grilli E. Towards zero zinc oxide: feeding strategies to Manage Post-Weaning Diarrhea in Piglets. *Animals*. 2021;11(3):642.
21. Sloup V, Jankovská I, Nechybová S, Peřínková P, Langrová I. Zinc in the animal organism: a review. *Sci Agric Bohem*. 2017;48(1):13–21.
22. Hill GM, Mahan DC, Carter SD, Cromwell GL, Ewan RC, Harrold RL, Lewis AJ, Miller PS, Shurson GC, Veum TL. Effect of pharmacological concentrations of zinc oxide with or without the inclusion of an antibacterial agent on nursery pig performance. *J Anim Sci*. 2001;79:934–41.
23. Zhang B, Guo Y. Supplemental zinc reduced intestinal permeability by enhancing occludin and zonula occludens protein-1 (ZO-1) expression in weaning piglets. *Br J Nutr*. 2009;102(5):687–93.
24. Zhu C, Lv H, Chen Z, Wang L, Wu X, Chen Z, Zhang W, Liang R, Zongyong J. Dietary zinc oxide modulates antioxidant capacity, small Intestine Development, and Jejunal Gene expression in weaned piglets. *Biol Trace Elem Res*. 2017;175(2):331–8.
25. Grilli E, Tugnoli B, Vitari F, Domeneghini C, Morlacchini M, Piva A, Prandini A. Low doses of microencapsulated zinc oxide improve performance and modulate the ileum architecture, inflammatory cytokines and tight junctions expression of weaned pigs. *Animal*. 2015;9(11):1760–8.
26. Hedemann MS, Jensen BB, Poulsen HD. Influence of dietary zinc and copper on digestive enzyme activity and intestinal morphology in weaned pigs1. *J Anim Sci*. 2006;84(12):3310–20.
27. Moodley A, Nielsen SS, Guardabassi L. Effects of tetracycline and zinc on selection of methicillin-resistant *Staphylococcus aureus* (MRSA) sequence type 398 in pigs. *Vet Microbiol*. 2011;152(3–4):420–3.
28. Slifierz MJ, Friendship R, Weese JS. Zinc oxide therapy increases prevalence and persistence of Methicillin-Resistant *Staphylococcus aureus* in Pigs: a Randomized Controlled Trial. *Zoonoses Public Health*. 2015;62(4):301–8.
29. Bednorz C, Oelgeschläger K, Kinnemann B, Hartmann S, Neumann K, Pieper R. u. a. The broader context of antibiotic resistance: zinc feed supplementation of piglets increases the proportion of multi-resistant *Escherichia coli* in vivo. *Inte J Med Microbiol*. 2013;303(6–7):396–403.
30. Vahjen W, Pietruszyńska D, Starke IC, Zentek J. High dietary zinc supplementation increases the occurrence of tetracycline and sulfonamide resistance genes in the intestine of weaned pigs. *Gut Pathog*. 2015;7(1):23.
31. Ciesinski L, Guenther S, Pieper R, Kalisch M, Bednorz C, Wieler LH. High dietary zinc feeding promotes persistence of multi-resistant *E. coli* in the swine gut. *Brockmann GA, Herausgeber. PLoS ONE*. 2018;13(1):e0191660.
32. EMA: Annex II Scientific conclusions and grounds for the refusal of the marketing authorisation and for withdrawal of the existing marketing authorisations. EMAE: Amsterdam, 2017; https://ec.europa.eu/health/documents/community-register/2017/20170626136754/anx_136754_en.pdf. Accessed 15 September 2022.
33. Monteiro SC, Lofts S, Boxall ABA. Pre-assessment of environmental impact of zinc and copper used in animal nutrition. *EFS3*. 2010;7(9). <https://doi.org/10.2903/sp.efsa.2010.EN-74>. Accessed 15 September 2022.
34. Heo JM, Kim JC, Hansen CF, Mullan BP, Hampson DJ, Pluske JR. Feeding a diet with decreased protein content reduces indices of protein fermentation and the incidence of postweaning diarrhea in weaned pigs challenged with an enterotoxigenic strain of *Escherichia coli*1. *J Anim Sci*. 2009;87(9):2833–43.
35. Hermes RG, Molist F, Ywazaki M, Nofrarias M, Gomez de Segura A, Gasa J, Pérez JF. Effect of dietary level of protein and fiber on the productive performance and health status of piglets1. *J Anim Sci*. 2009;87(11):3569–77.
36. Heo JM, Kim JC, Hansen CF, Mullan BP, Hampson DJ, Maribo H, Kjeldsen N, Pluske JR. Effects of dietary protein level and zinc oxide supplementation on the incidence of post-weaning diarrhoea in weaner pigs challenged with an enterotoxigenic strain of *Escherichia coli*. *Livest Sci*. 2010;133(1–3):210–3.
37. Nyachoti CM, Omogbenigun FO, Rademacher M, Blank G. Performance responses and indicators of gastrointestinal health in early-weaned pigs fed low-protein amino acid-supplemented diets1. *J Anim Sci*. 2006;84(1):125–34.
38. Reynoso E, Cervantes M, Figueroa JL. Respuesta productiva de lechones a dietas bajas en proteína adicionadas con aminoácidos sintéticos y cultivo de levaduras. *Rev cuba cienc agric*. 2004;38(3):9.
39. Lei XJ, Park JW, Baek DH, Kim JK, Kim IH. Feeding the blend of organic acids and medium chain fatty acids reduces the diarrhea in piglets orally challenged with enterotoxigenic *Escherichia coli* K88. *Anim Feed Sci Technol*. 2017;224:46–51.
40. Tugnoli G, Piva A, Grilli E. From Acidifiers to Intestinal Health Enhancers: how Organic acids can improve Growth Efficiency of Pigs. *Animals*. 2020;10(1):134.
41. Tsiloyiannis VK, Kyriakis SC, Vlemmas J, Sarris K. The effect of organic acids on the control of porcine post-weaning diarrhoea. *Res Vet Sci*. 2001;70(3):287–93.
42. Lu X, Zhang M, Zhao L, Ge K, Wang Z, Jun L, Ren F. Growth performance and post-weaning diarrhea in piglets Fed a Diet supplemented with Probiotic Complexes. *J Microbiol Biotechnol*. 2018;28(11):1791–9.
43. Luise D, Bertocchi M, Motta V, Salvarani C, Bosi P, Luppi A, Fanelli F, Mazzoni M, Archetti I, Maiorano G, Nielsen BKK, Trevisi P. *Bacillus* sp. probiotic supplementation diminish the *Escherichia coli* F4ac infection in susceptible weaned pigs by influencing the intestinal immune response, intestinal microbiota and blood metabolomics. *J Anim Sci Biotechnol*. 2019;10(1):74.
44. Taras D, Vahjen W, Macha M, Simon O. Performance, diarrhea incidence, and occurrence of *Escherichia coli* virulence genes during long-term administration of a probiotic *Enterococcus faecium* strain to sows and piglets1. *J Anim Sci*. 2006;84(3):608–17.
45. Nordeste R, Tessema A, Sharma S, Kovač Z, Wang C, Morales R, Griffiths MW. Molecules produced by probiotics prevent enteric colibacillosis in pigs. *BMC Vet Res*. 2017;13(1):335.
46. Silva CA, Dias CP, Callegari MA, Bridi AM, Santos RKS, Luiggi FG, Santos VL, Silva JB. Prebiotics and butyric acid can replace colistin as a growth promoter for nursery piglets. *Arq Bras Med Vet Zootec*. 2020;72(4):1449–57.
47. Mukhopadhyaya A, O'Doherty JV, Sweeney T. A combination of yeast beta-glucan and milk hydrolysate is a suitable alternative to zinc oxide in the race to alleviate post-weaning diarrhoea in piglets. *Sci Rep*. 2019;9(1):616.
48. Bakkali F, Averbeck S, Averbeck D, Idaomar M. Biological effects of essential oils – A review. *Food Chem Toxicol*. 2008;46(2):446–75.
49. Rossi B, Toschi A, Piva A, Grilli E. Single components of botanicals and nature-identical compounds as a non-antibiotic strategy to ameliorate health status and improve performance in poultry and pigs. *Nutr Res Rev*. 2020;33(2):218–34.
50. Yu J, Yu G, Yu B, Zhang Y, He J, Zheng P. u. a. dietary protease improves growth performance and nutrient digestibility in weaned piglets fed diets with different levels of soybean meal. *Livest Sci*. 2020;241:104179.
51. Vangroenweghe F, Poulsen K, Thas O. Supplementation of a β -mannanase enzyme reduces post-weaning diarrhea and antibiotic use in piglets on an alternative diet with additional soybean meal. *Porc Health Manag Dezember*. 2021;7(1):8.
52. Torralardona D. Spray dried animal plasma as an alternative to antibiotics in Weanling Pigs - A Review -. *Asian Australas J Anim Sci*. 2009;23(1):131–48.
53. Trckova M, Faldyna M, Alexa P, Zajacova ZS, Gopfert E, Kumprechtova D. u. a. The effects of live yeast *Saccharomyces cerevisiae* on postweaning diarrhea, immune response, and growth performance in weaned piglets1,2. *J Anim Sci*. 2014;92(2):767–74.
54. Feng J, Wang L, Xie Y, Chen Y, Yi H, He D. Effects of antimicrobial peptide cathelicidin-BF on diarrhea controlling, immune responses, intestinal inflammation and intestinal barrier function in piglets with postweaning diarrhea. *Int Immunopharmacol*. 2020;85:106658.
55. Zhang L, Guo T, Zhan N, Sun T, Shan A. Effects of the antimicrobial peptide WK3 on diarrhea, growth performance and intestinal health of weaned piglets challenged with enterotoxigenic *Escherichia coli* K88. *Food Nutr Res*. 2021;65:3448.
56. Wisener LV, Sargeant JM, O'Sullivan TL, O'Connor AM, McEwen SA, Reist M. u. a. non-antibiotic approaches for Disease Prevention and Control in Nursery Pigs: a scoping review. *Front Vet Sci*. 2021;8:620347.
57. Poulsen HD. Zinc oxide for Weanling Piglets. *Acta Agri Scand*. 1995;45(3):159–67.
58. Papadopoulos GA, Vanderhaeghe C, Janssens GPJ, Dewulf J, Maes DGD. Risk factors associated with postpartum dysgalactia syndrome in sows. *Veterinary J Mai*. 2010;184(2):167–71.
59. Yan L, Jang HD, Kim IH. Effects of varying Creep feed duration on pre-weaning and post-weaning performance and behavior of Piglet and sow. *Asian Australas J Anim Sci*. 2011;24(11):1601–6.
60. van Hees H, Maes D, Millet S, den Hartog L, van Kempen T, Janssens G. Fibre supplementation to pre-weaning piglet diets did not improve the resilience towards a post - weaning enterotoxigenic *E. coli* challenge. *J Anim Physiol Anim Nutr*. 2021;105(2):260–71.
61. Choudhury R, Middelkoop A, Boekhorst J, Gerrits WJJ, Kemp B, Bolhuis JE, Kleerebezem M. Early life feeding accelerates gut microbiome maturation and suppresses acute post-weaning stress in piglets. *Environ Microbiol*. 2021;23(11):7201–13.
62. Amezcua R, Friendship R, Gyles C. A case-control study investigating risk factors associated with postweaning *Escherichia coli* diarrhea in southern Ontario. *J Swine Health Prod*. 2002;10(6):5.
63. García KE, de Souza TCR, Landin GM, Barreyro AA, Santos MGB, Soto JGG. Microbial fermentation patterns, Diarrhea Incidence, and performance in

- weaned piglets Fed a low protein Diet supplemented with Probiotics. *FNS*. 2014;05(18):1776–86.
64. Yue LY, Qiao SY. Effects of low-protein diets supplemented with crystalline amino acids on performance and intestinal development in piglets over the first 2 weeks after weaning. *Livest Sci*. 2008;115(2–3):144–52.
 65. Laine TM, Lyytikäinen T, Yliahio M, Anttila M. Risk factors for post-weaning diarrhoea on piglet producing farms in Finland. *Acta Vet Scand*. 2008;50(1):21.
 66. Patil AK, Kumar S, Verma AK, Baghel RPS. Probiotics as feed additives in weaned Pigs: a review. *Int J Livest Res*. 2015;3(2):9.
 67. Sun Y, Duarte ME, Kim SW. Dietary inclusion of multispecies probiotics to reduce the severity of post-weaning diarrhea caused by *Escherichia coli* F18 + in pigs. *Anim Nutr*. 2021;7(2):326–33.
 68. Roselli M, Finamore A, Britti MS, Mengheri E. Probiotic bacteria *bifidobacterium animalis* MB5 and *Lactobacillus rhamnosus* GG protect intestinal Caco-2 cells from the inflammation-associated response induced by enterotoxigenic *Escherichia coli* K88. *Br J Nutr*. 2006;95(6):1177–84.
 69. Turgis M, Han J, Caillet S, Lacroix M. Antimicrobial activity of mustard essential oil against *Escherichia coli* O157:H7 and *Salmonella typhi*. *Food Control*. 2009;20(12):1073–9.
 70. Delaquis PJ, Sholberg PL. Antimicrobial activity of Gaseous Allylthioiocyanate. *J Food Product*. 1997;60(8):943–47.
 71. Visscher C, Hankel J, Nies A, Keller B, Galvez E, Strowig T, Keller C, Breves G, Performance. Fermentation characteristics and composition of the Microbiome in the Digest of Piglets kept on a feed with Humic Acid-Rich Peat. *Front Vet Sci*. 2019;6:29.
 72. Wang Q, Ying J, Zou P, Zhou Y, Wang B, Yu D. u. a. Effects of Dietary Supplementation of Humic Acid Sodium and Zinc Oxide on Growth Performance, Immune Status and antioxidant capacity of weaned piglets. *Animals*. 2020;10(11):2104.
 73. Vanheukelom V, Driessen B, Maenhout D, Geers R. Peat as environmental enrichment for piglets: the effect on behaviour, skin lesions and production results. *Appl Anim Behav Sci*. 2011;134(1–2):42–7.
 74. Johansen TB, Agdestein A, Lium B, Jørgensen A, Djønne B. *Mycobacterium avium* subsp. *hominissuis* infection in Swine Associated with Peat used for bedding. *Biomed Res Int*. 2014;2014:1–8.
 75. Matlova L, Dvorska L, Bartl J, Ayele WY, Bartos M, Alexa M et al. Mycobacterial contamination of environment in pig farms in the Czech Republic between 1996 and 2002. In: International Conference on the Epidemiology and Control of Biological, Chemical and Physical Hazards in Pigs and Pork [Internet]. Heraklion, Crete, Greece: Iowa State University, Digital Press; 2003; p. 226–8. <https://lib.dr.iastate.edu/safepork/2003/allpapers/87/> accessed on 15 September 2022.
 76. Li Y, Liu Y, Wu J, Chen Q, Zhou Q, Wu F. u. a. comparative effects of enzymatic soybean, fish meal and milk powder in diets on growth performance, immunological parameters, SCFAs production and gut microbiome of weaned piglets. *J Anim Sci Biotechnol*. 2021;12(1):106.
 77. Murphy J. Determination of phosphoric acid in cola beverages: a colorimetric and pH titration experiment for general chemistry. *J Chem Educ*. 1983;60(5):420.
 78. Deng Q, Shao Y, Wang Q, Li J, Li Y, Ding X, Huang P, Yin J, Yang H, Yin Y. Effects and interaction of dietary electrolyte balance and citric acid on the intestinal function of weaned piglets. *J Anim Sci*. 2020;98(5):106.
 79. Boontiam W, Wachirapakorn C, Phaengphairee P. Effects of hydrolyzed yeast supplementation on growth performance, immunity, antioxidant capacity, and microbial shedding in weaning pigs. *Vet World*. 2020;13(9):1902–9.
 80. Bautista-Marín S, Escobar-García K, Molina-Aguilar C, Mariscal-Landín G, Aguilera-Barreyro A, Díaz-Muñoz M, Reis, de Souza TC. Antibiotic-free diet supplemented with live yeasts decreases inflammatory markers in the ileum of weaned piglets. *SA J An Sci*. 2020;50(3):353–65.
 81. Fairbrother JM, Nadeau é. Colibacillosis. In: Zimmerman JJ, Karriker LA, Ramirez A, Schwartz KJ, Stevenson GW, Zhang J, editors. *Diseases of Swine*. Hoboken: Wiley-Blackwell; 2019. pp. 807–34.
 82. Jayaraman B, Nyachoti CM. Husbandry practices and gut health outcomes in weaned piglets: a review. *Anim Nutr*. 2017;3(3):205–11.
 83. Davies PR, Morrow WEM, Jones FT, Deen J, Fedorka-Cray PJ, Harris IT. Prevalence of salmonella in finishing swine raised in different production systems in North Carolina, USA. *Epidemiol Infect*. 1997;119(2):237–44.
 84. Dewey CE. Herd- and litter-level factors associated with the incidence of diarrhea morbidity and mortality in piglets 4–14 days of age. *J Swine Health Prod*. 1995;3(3):105–12.
 85. Skampardonis V, Sotiraki S, Kostoulas P, Leontides L. Factors associated with the occurrence and level of Isospora suis oocyst excretion in nursing piglets of greek farrow-to-finish herds. *BMC Vet Res*. 2012;8(1):228.
 86. Le Dividich J, Herpin P. Effects of climatic conditions on the performance, metabolism and health status of weaned piglets: a review. *Livest Prod Sci*. 1994;38(2):79–90.
 87. Rantzer D, Svendsen J. Slatted versus solid floors in the Dung Area of Farrowing Pens: Effects on Hygiene and Pig Performance, Birth to Weaning. *Acta Agric Scand Anim Sci*. 2001;51(3):167–74.
 88. Lozano AP, Poveda-Parra AR, Raelé E, Novais AK, da Silva RAM, Pacheco GD, Gavioli D, Dalto B, da Silva CA. Effects of housing conditions and glutamine levels on growthperformance of post-weaning piglets. *SCA*. 2016;37:2387.
 89. Berrococo JD, Menoyo D, Guzmán P, Saldaña B, Cámara L, Mateos GG. Effects of fiber inclusion on growth performance and nutrient digestibility of piglets reared under optimal or poor hygienic conditions. *J Anim Sci*. 2015;93(8):3919–31.
 90. Gonzales-Siles L, Sjöling Å. The different ecological niches of enterotoxigenic *Escherichia coli*: Ecology of ETEC. *Environ Microbiol*. 2016;18(3):741–51.
 91. Little S, Woodward A, Browning G, Billman-Jacobe H. Water distribution Systems in Pig Farm buildings: critical elements of design and management. *Animals*. 2021;11(11):3268.
 92. Moreira S, Brown A, Ha R, Iserhoff K, Yim M, Yang J. u. a. persistence of *Escherichia coli* in freshwater periphyton: biofilm-forming capacity as a selective advantage. *FEMS Microbiol Ecol*. 2012;79(3):608–18.
 93. Andrés-Barranco S, Vico JP, Garrido V, Samper S, Herrera-León S, de Frutos C. u. a. role of wild bird and rodents in the epidemiology of subclinical salmonellosis in Finishing Pigs. *Foodborne Pathog Dis*. 2014;11(9):689–97.
 94. Meerburg BG, Kijlstra A. Role of rodents in transmission of Salmonella and Campylobacter. *J Sci Food Agric*. 2007;87(15):2774–81.
 95. R Core Team. R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. 2020. <https://www.R-project.org/>. Accessed 10 July 2022.
 96. Fox J, Weisberg S. An {R} Companion to Applied Regression, Third Edition. Thousand Oaks CA, USA. 2019. <https://socialsciences.mcmaster.ca/fox/Books/Companion/>. Accessed 14 July 2022.
 97. Friedman J, Hastie T, Tibshirani R. Regularization Paths for Generalized Linear Models via Coordinate Descent. *J Stat Soft*. 2010. <http://www.jstatsoft.org/v33/i01/>. Accessed 14 July 2022.
 98. Kuhn MR, caret. Classification and Regression Training. R package version 6.0–92. 2022. <https://CRAN.R-project.org/package=caret>. Accessed 14 July 2022.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.