

# Determining the Safety and Efficacy of Enhanced Recovery Protocols in Major Oncologic Surgery: An Institutional NSQIP Analysis

Rebecca K. Marcus, MD<sup>1</sup>, Heather A. Lillemoe, MD<sup>1</sup>, David C. Rice, MB, BCh, BAO, FRCSI<sup>4</sup>, Gabriel Mena, MD<sup>3</sup>, Brian K. Bednarski, MD<sup>1</sup>, Barbra B. Speer, DO<sup>3</sup>, Pedro T. Ramirez, MD<sup>2</sup>, Javier D. Lasala, MD<sup>3</sup>, Neema Navai, MD<sup>5</sup>, Wendell H. Williams, MD<sup>3</sup>, Bradford J. Kim, MD, MHS<sup>1</sup>, Rachel K. Voss, MD, MHS<sup>1</sup>, Vijaya N. Gottumukkala, MD, FRCA<sup>3</sup>, and Thomas A. Aloia, MD, FACS<sup>1</sup>

<sup>1</sup>Department of Surgical Oncology, The University of Texas MD Anderson Cancer Center, Houston, TX ; <sup>2</sup>Department of Gynecologic Oncology and Reproductive Medicine, The University of Texas MD Anderson Cancer Center, Houston, TX; <sup>3</sup>Department of Anesthesiology and Perioperative Medicine, The University of Texas MD Anderson Cancer Center, Houston, TX; <sup>4</sup>Department of Thoracic and Cardiovascular Surgery, The University of Texas MD Anderson Cancer Center, Houston, TX; <sup>5</sup>Department of Urology, The University of Texas MD Anderson Cancer Center, Houston, TX

## ABSTRACT

**Background.** Enhanced-recovery (ER) protocols are increasingly being utilized in surgical practice. Outside of colorectal surgery, however, their feasibility, safety, and efficacy in major oncologic surgery have not been proven. This study compared patient outcomes before and after multispecialty implementation of ER protocols at a large, comprehensive cancer center.

**Methods.** Surgical cases performed from 2011 to 2016 and captured by an institutional NSQIP database were reviewed. Following exclusion of outpatient and emergent surgeries, 2747 cases were included in the analyses. Cases were stratified by presence or absence of ER compliance, defined by preoperative patient education and electronic medical record order set-driven opioid-sparing analgesia,

goal-directed fluid therapy, and early postoperative diet advancement and ambulation.

**Results.** Approximately half of patients were treated on ER protocols (46%) and the remaining on traditional postoperative (TP) protocols (54%). Treatment on an ER protocol was associated with decreased overall complication rates (20% vs. 33%,  $p < 0.0001$ ), severe complication rates (7.4% vs. 10%,  $p = 0.010$ ), and median hospital length of stay (4 vs. 5 days,  $p < 0.0001$ ). There was no change in readmission rates (ER vs. TP, 8.6% vs. 9.0%,  $p = 0.701$ ). Subanalyses of high magnitude cases and specialty-specific outcomes consistently demonstrated improved outcomes with ER protocol adherence, including decreased opioid use.

**Conclusions.** This assessment of a large-scale ER implementation in multispecialty major oncologic surgery indicates its feasibility, safety, and efficacy. Future efforts should be directed toward defining the long-term oncologic benefits of these protocols.

---

The study was presented as an oral abstract at Society of Surgical Oncology Annual Meeting, Chicago, IL, March 22, 2018.

---

**Electronic supplementary material** The online version of this article (<https://doi.org/10.1245/s10434-018-07150-5>) contains supplementary material, which is available to authorized users.

---

© Society of Surgical Oncology 2019

First Received: 22 September 2018;  
Published Online: 9 January 2019

T. A. Aloia, MD, FACS  
e-mail: [taaloia@mdanderson.org](mailto:taaloia@mdanderson.org)

For several decades fast-track and enhanced-recovery (ER) surgical protocols have been described, mainly by European centers.<sup>1</sup> More recently, the concept of ER has been integrated into surgical practice in North America, resulting in a paradigm shift in perioperative care. ER is a multimodality approach to patient care that is implemented by a multidisciplinary team. The driving principle behind these evidence-based care pathways is that by reducing stress associated with surgery, patients are able to recover

more rapidly and completely, thereby minimizing complications and delays in further treatment.<sup>1,2</sup> As ER has been incorporated into more surgical specialties, data are rapidly accumulating that demonstrate the benefits of ER protocols on patient outcomes. These benefits include reduced postoperative complications and inpatient length of stay (LOS).<sup>3-8</sup> As personalized care and precision medicine have become increasingly important concepts, studies examining more patient-centric, long-term outcomes reflecting functional recovery also are being reported.<sup>9-11</sup>

Despite this revolution in perioperative care, much of the ER literature focuses on the impact of ER on patients undergoing nononcologic procedures.<sup>6,8,12</sup> When oncologic patients are described, they often have early-stage disease amenable to less invasive surgical intervention.<sup>13,14</sup> Because colorectal surgeons were amongst the earliest to incorporate ER into their practice, the largest amount of oncologic data on ER exists in this patient cohort. However, the majority of oncologic specialties are underrepresented in the ER literature.<sup>15-18</sup> To address this knowledge gap, we assessed the impact of ER implementation on patient outcomes across multiple oncologic surgical specialties at a large, comprehensive cancer center.

## METHODS

### *Defining Enhanced-Recovery Protocols*

ER protocol implementation began in 2012 with a phased roll-out performed across disease sites and surgical specialties over a 3-year period. Surgical specialties included in the analyses were Colorectal, Gynecology, Hepatobiliary, Thoracic, and Urology. Each service area program was formed around a multidisciplinary team that included surgery, anesthesia, nursing, pharmacy, nutrition, rehabilitation services, and informatics support. Provider education regarding the core components of ER was supplemented by presentations at regularly scheduled CME meetings, surgical service orientations, and other didactics. For the purpose of this study, ER protocol compliance was defined as patient care that included preoperative patient education on ER and use of both pre- and postoperative ordersets in the institutional electronic medical record that incorporated the four pillars of ER: multimodal opioid-sparing analgesia, goal-directed fluid therapy, early feeding, and early ambulation.<sup>1,2,19</sup>

Patient education was performed by providers and nursing staff at preoperative clinic visits. Patients received education on both the indicated operation and on ER principles and goals, with the objectives of reducing anxiety by transparently sharing plans of care and setting patient expectations regarding their perioperative

experiences.<sup>2,19</sup> For opioid-sparing, multimodal analgesia, although each service had latitude to select the exact components, all used some combination of regional anesthetic options (e.g., epidural and transverse abdominus plane [TAP] blocks) with acetaminophen, nonsteroidal anti-inflammatories, and nonnarcotic neuromodulators. Goal-directed fluid therapy, performed both intraoperatively and postoperatively, involved titrating the amount of fluid being administered based on hemodynamic indices, low maintenance fluid rates, early discontinuance with *per oral* intake, and data-driven bolus strategies.<sup>19,20</sup> Early feeding was built into order sets, begun in most cases on postoperative day 0, and programmatically advanced to regular diet. Early ambulation was likewise hard-wired into order sets and initiated in most cases on the day of surgery.<sup>21-28</sup>

### *National Surgical Quality Improvement Program Analysis*

This study was approved by the University of Texas MD Anderson Cancer Center institutional review board and performed in compliance with the National Surgical Quality Improvement Program (NSQIP) data use agreement. For the entire duration of the study, the center participated in the Essentials data collection model, capturing approximately 14% of the surgical volume across all specialties. Patient data were obtained from the institutional American College of Surgeons (ACS)-NSQIP database from January 2011 to December 2016 for the following surgical specialties: Colorectal, Gynecology, Hepatobiliary, Thoracic, and Urology. Both open and minimally invasive surgical cases were included in the analyses, but all outpatient and emergent cases were excluded. For patients undergoing surgeries involving more than one surgical team, the operation was classified by the primary team's specialty. The date of ER protocol initiation was determined for each specialty, and the cases from each specialty were subsequently divided into two cohorts: those treated on ER versus TP protocols.

Variable collection was performed by Surgical Clinical Reviewers trained in the analysis of medical records and audited by the ACS. Demographic and clinical information, operative details, and 30-day outcomes were included in the analyses. Age, American Society of Anesthesiologists (ASA) classification, preoperative functional status (defined as independent if assistance from another person for any activities of daily living was not required vs. dependent if some or total assistance was required), and preoperative comorbidities were compared between patients treated on ER and TP protocols.<sup>29</sup> Wound class, operative time, and hospital length of stay (LOS) were assessed. Postoperative complications, as defined by the

NSQIP Participant Use Data File, were compared between ER and TP cohorts and included wound infection, pneumonia, pulmonary embolism, urinary tract infection, renal failure, myocardial infarction, cardiac arrest, stroke, sepsis, and unplanned return to the operating room. A severe complication was defined per prior publications as having one or more of the following: organ space infection, wound dehiscence, pneumonia, unplanned intubation, acute renal failure, myocardial infarction, sepsis, septic shock, or unplanned return to the operating room.<sup>10</sup> Thirty-day readmission and mortality rates also were compared. Mortalities that occurred during a continuous postoperative admission lasting longer than 30 days were included. Before implementation, throughout the study period, and after implementation, NSQIP patient outcomes reports were shared with each team and associated frontline providers. Likewise, individual teams internally reported their general and specialty-specific outcomes from existing department-level databases.

### *Analysis of High Magnitude Surgeries*

A subanalysis was performed to assess the effects of implementation of ER protocols on patients undergoing high-magnitude surgery. Case magnitude was determined by primary CPT RVU, and cases with RVU >30 were considered high magnitude. All variables included in the complete ER versus TP cohort analyses also were assessed in this subanalysis.

### *Statistics*

Statistical analyses were performed using JMP Pro software program (Version 12, SAS institute Inc., Cary, NC). Univariate analyses were performed using Chi square or Fisher's exact test for categorical variables and the Mann-Whitney *U* test for continuous variables. Continuous variables are reported as median values and interquartile ranges. A two-tailed univariate  $p < 0.05$  was considered significant.

## **RESULTS**

### *Study Population Demographics and Clinical Profile*

A total of 2747 patients entered into the institutional NSQIP database between January 2011 and December 2016 were included in the analyses (Supplementary Fig. 1). Median age at the time of surgery was 60 years (interquartile range [IQR] 51-68 years), 1,548 (56%) were female, and the mean ASA score was 3. Fourteen percent of cases were performed laparoscopically, and the

remainder were performed via an open surgical approach. Preoperative comorbidities are detailed in Table 1: 360 patients (13%) had diabetes, 1208 (44%) medication-controlled hypertension, and 82 (3%) required chronic steroids. Also, 365 patients (13%) were documented smokers within the 12 months before surgery, although the overall rates of COPD and dyspnea were low. With regard to their oncologic diagnoses, 1335 (48%) received preoperative chemotherapy within 30 days of surgery, and 495 (18%) received preoperative radiation within 90 days of surgery. At the time of surgery, 746 (27%) of patients had disseminated cancer and 72 (3%) reported loss of > 10% of their body weight during the 6 months before surgery.

A total of 1252 (46%) patients were treated on ER protocols. Compared with patients treated on TP protocols, patients on ER protocols had similar median age (60 vs. 60 years,  $p = 0.742$ ) and rates of ASA  $\geq 3$  classification (88% vs. 87%,  $p = 0.270$ ). Although patients treated using ER protocols had slightly lower rates of medication-controlled hypertension (42% vs. 46%,  $p = 0.027$ ) and disseminated cancer (24% vs. 30%,  $p = 0.001$ ), they were more likely to have undergone preoperative treatment with chemotherapy (51% vs. 47%,  $p = 0.014$ ) and radiation (20% vs. 16%,  $p = 0.004$ ).

### *Outcomes of Patients Treated using ER Versus TP Protocols*

Compared with TP protocols, treatment using ER protocols was associated with decreased perioperative complications (Table 2). Specifically, patients on ER protocols were less likely to have complications (20% vs. 33%,  $p < 0.0001$ ), including severe complications (7% vs. 10%,  $p = 0.010$ ). The median hospital LOS was lower for patients on ER protocols (4 vs. 5 days,  $p < 0.0001$ ). Despite the shorter LOS, unplanned 30-day readmission rates did not increase in the ER cohort (107 vs. 134 patients, 8.6% vs. 9.0%,  $p = 0.701$ ). Lastly, while the 30-day mortality rates were low for both cohorts, a reduced mortality rate was observed amongst patients treated on ER protocols (0.2% vs. 0.6%,  $p = 0.245$ ).

### *Individual Specialty Outcomes of Patients treated on ER Versus TP Protocols*

The previously described analyses also were performed on an individual basis for each of the surgical specialties included in the NSQIP database. These analyses demonstrated that many of the beneficial effects of ER protocol adoption observed for all cases were seen within each specialty. Although the exact factors that significantly improved with ER-protocol implementation differed for each specialty, these protocols uniformly resulted in

**TABLE 1** Clinical characteristics, based on recovery protocol

Factor	Total	Traditional	Enhanced recovery	<i>p</i> Value
No. of patients	2747	1495 (54)	1252 (46)	
Surgical subspecialty				
Colorectal	663 (24)	273 (18)	390 (31)	
Gynecology	634 (23)	401 (27)	233 (19)	
Hepatobiliary	291 (11)	230 (15)	61 (5)	
Thoracic/vascular	856 (31)	390 (26)	466 (37)	
Urology	303 (11)	201 (13)	102 (8)	
Preoperative				
Age, year, median (IQR)	60 (51–68)	60 (50–69)	60 (51–68)	0.742
Gender, female	1548 (56)	856 (57)	692 (55)	0.296
ASA score				
I/II	347 (13)	199 (13)	148 (12)	–
III/IV	2375 (87)	1287 (87)	1088 (88)	0.270
Diabetes	360 (13)	182 (12)	178 (14)	0.114
Current smoker	365 (13)	215 (14)	150 (12)	<b>0.065</b>
Dyspnea	190 (7)	98 (7)	92 (7)	0.415
COPD	128 (5)	72 (5)	56 (4)	0.671
HTN requiring medication	1208 (44)	686 (46)	522 (42)	<b>0.027</b>
Disseminated cancer	746 (27)	443 (30)	303 (24)	<b>0.001</b>
Steroid use	82 (3)	48 (3)	34 (3)	0.448
> 10% weight loss	72 (3)	46 (3)	26 (2)	0.102
Bleeding disorder	97 (4)	61 (4)	36 (3)	<b>0.088</b>
Preoperative radiation	495 (18)	240 (16)	255 (20)	<b>0.004</b>
Preoperative chemotherapy	1335 (48)	693 (47)	642 (51)	<b>0.014</b>
Perioperative				
MIS procedure	394 (14)	174 (12)	220 (18)	<b>&lt;0.0001</b>
Wound classification				
Clean	84 (3)	50 (3)	34 (3)	–
Clean/contaminated	2543 (93)	1379 (92)	1164 (93)	–
Contaminated	74 (3)	38 (3)	36 (3)	–
Dirty/infected	46 (2)	28 (2)	18 (1)	0.862
Operative time, min, median (IQR)	218 (141–316)	234 (152–335)	198 (126–294)	<b>&lt;0.0001</b>

Bold values are statistically significant ( $p < 0.05$ )

*IQR* interquartile range, *ASA* American Society of Anesthesiologists, *COPD* chronic obstructive pulmonary disease, *HTN* hypertension, *MIS* minimally invasive surgery

decreased rates of complications (Fig. 1a), including severe complications. This included decreased rates of surgical site infections following ER protocol implementation for all specialties except thoracic surgery, for which no change in the already low rates of SSIs was observed (1.0% vs. 2.8%,  $p = 0.085$ ). Patients across all specialties treated using ER protocols also had shorter median hospital LOS (Fig. 1b). Lastly, evaluation by the individual specialties of their patients' average postoperative opioid use, measured as morphine-equivalent daily doses, demonstrated a 42–85% decrease in opioid use after ER protocol implementation.<sup>30</sup> Although no data regarding patient-reported pain scores are collected within the NSQIP database,

results of previously published studies utilizing ER protocols suggest that the lower observed opioid use was associated with improved patient-reported pain scores.<sup>31–33</sup>

#### Assessment of High Magnitude Case Outcomes

To further assess the effects of ER protocol implementation on patient outcomes and to better control for case complexity, a subanalysis was performed on those patients undergoing high-magnitude surgeries (primary CPT RVU >30); 1,022 cases (37%) satisfied this definition. High-magnitude cases by surgical subspecialty are shown in Supplementary Fig. 2. Within this cohort of patients,


**TABLE 2** Postoperative complications, based on recovery protocol

Factor	Total ( <i>n</i> = 2747)	Traditional ( <i>n</i> = 1495)	Enhanced recovery ( <i>n</i> = 1252)	<i>p</i> Value
Superficial incisional SSI	95 (3)	64 (4)	31 (2)	<b>0.010</b>
Deep incisional SSI	15 (1)	13 (1)	2 (0.2)	<b>0.016</b>
Organ space infection	88 (3)	51 (3)	37 (3)	0.499
Wound disruption/dehiscence	35 (1)	26 (2)	9 (1)	<b>0.018</b>
Pneumonia	80 (3)	52 (3)	28 (2)	0.054
Urinary tract infection	81 (3)	63 (4)	18 (1)	<b>&lt;0.0001</b>
Transfusion ≤ 72 h of OR	512 (19)	344 (23)	168 (13)	<b>&lt;0.0001</b>
Unplanned return to OR	75 (3)	46 (3)	29 (2)	0.241
Any complication	747 (27)	493 (33)	254 (20)	<b>&lt;0.0001</b>
Any severe complication <sup>a</sup>	244 (9)	152 (10)	92 (7)	<b>0.010</b>
LOS, days, median (IQR)	4 (3–7)	5 (3–7)	4 (2–5)	<b>&lt;0.0001</b>
Mortality ≤ 30 days	12 (0.4)	9 (1)	3 (0.2)	0.245
Readmission ≤ 30 days	241 (9)	134 (9)	107 (9)	0.701

Bold values are statistically significant ( $p < 0.05$ )

SSI surgical site infection, OR operating room, IQR interquartile range, LOS length of stay

<sup>a</sup>Severe complications include: organ space infection, wound disruption/dehiscence, pneumonia, unplanned intubation, acute renal failure, myocardial infarction, sepsis, septic shock, unplanned return to OR



**FIG. 1** Specialty-specific outcomes based on recovery protocol. Comparison of **a** complication rates and **b** median hospital length of stay between patients managed on traditional versus enhanced recovery protocols

371 (36%) were treated using ER protocols. Compared with patients on TP protocols, ER-protocol patients were less likely to have disseminated cancer at the time of surgery (19% vs. 36%,  $p < 0.0001$ ) but were more likely to have undergone radiation (27% vs. 19%,  $p = 0.001$ ).

Median procedure duration was similar between the ER and TP cohorts (327 vs. 326 min,  $p = 0.466$ ; Table 3).

Compared with patients undergoing surgeries of any magnitude, patients undergoing high-magnitude operations had increased rates of perioperative complications (41% of cases with RVU > 30 vs. 27% of all cases,  $p < 0.0001$ ), longer median LOS (6 vs. 4 days,  $p < 0.0001$ ), and higher readmission rates (13% vs. 9%,  $p = 0.0003$ ). Despite this confirmation of increased case complexity, a beneficial effect of ER protocols was still observed, and many of the key findings from the analysis of the entire study population also held true for patients who underwent high-magnitude surgery. Specifically, patients undergoing more complex cancer surgery and on ER protocols were less likely than traditionally treated patients to have any complication (35% vs. 44%,  $p = 0.002$ ). The median LOS for the ER protocol-treated cohort also was shorter (5 vs. 6 days,  $p < 0.0001$ ), while no corresponding increase in 30-day readmissions was observed (14% vs. 12%,  $p = 0.348$ ). No significant differences in the rates of severe complications or 30-day mortality were found in this sub-analysis; however, both were trending in a confirmatory direction in the ER cohort (Table 4).

## DISCUSSION

This study found that the adoption of enhanced recovery protocols for the perioperative care of oncologic patients resulted in improved patient outcomes, including decreased rates of complications, reduced hospital LOS without a

**TABLE 3** Clinical characteristics for high-magnitude cases<sup>a</sup>, based on recovery protocol

Factor	Total	Traditional	Enhanced recovery	<i>p</i> Value
No. of patients	1022	651 (64)	371 (36)	
Surgical subspecialty				
Colorectal	262 (26)	111 (17)	151 (41)	
Gynecology	128 (13)	82 (13)	46 (12)	
Hepatobiliary	291 (28)	230 (35)	61 (16)	
Thoracic/vascular	49 (5)	32 (5)	17 (5)	
Urology	292 (29)	196 (30)	96 (26)	
Preoperative				
Age, year, median (IQR)	60 (50–69)	60 (50–60)	60 (50–68)	0.879
Gender, female	462 (45)	287 (44)	175 (47)	0.341
ASA score				
I/II	126 (12)	80 (12)	46 (13)	–
III/IV	889 (88)	569 (88)	320 (87)	0.911
Diabetes	136 (13)	82 (13)	54 (15)	0.375
Current smoker	129 (13)	92 (14)	37 (10)	0.054
Dyspnea	54 (5)	36 (6)	18 (5)	0.641
COPD	39 (4)	26 (4)	13 (4)	0.694
HTN requiring medication	448 (44)	289 (44)	159 (43)	0.634
Disseminated cancer	302 (30)	232 (36)	70 (19)	<b>&lt;0.0001</b>
Steroid use	28 (3)	21 (3)	7 (2)	0.207
> 10% weight loss	22 (2)	17 (3)	5 (1)	0.262
Bleeding disorder	40 (4)	27 (4)	13 (4)	0.610
Preoperative radiation	225 (22)	123 (19)	102 (27)	<b>0.001</b>
Preoperative chemotherapy	616 (60)	395 (61)	221 (60)	0.730
Perioperative				
Wound classification				
Clean	18 (2)	18 (3)	0 (0)	–
Clean/contaminated	952 (93)	600 (92)	352 (95)	–
Contaminated	33 (3)	19 (3)	14 (4)	–
Dirty/infected	19 (2)	14 (2)	5 (1)	<b>0.002</b>
Operative time, min, median (IQR)	327 (255–419)	326 (251–422)	327 (266–409)	0.466

Bold values are statistically significant ( $p < 0.05$ )

*IQR* interquartile range, *ASA* American Society of Anesthesiologists, *COPD* chronic obstructive pulmonary disease, *HTN* hypertension

<sup>a</sup>Defined as RVU > 30

concomitant increase in readmission rates, and lower opioid utilization. These findings of ER protocol implementation at a large comprehensive cancer center mirror those reported in the ER literature for other surgical specialties.<sup>3–8</sup> To our knowledge, this is the largest study to date looking at multispecialty ER implementation within the setting of oncologic surgery, and so provides substantial evidence that ER protocols can be successfully utilized in the treatment of this complex patient population.

Unlike other reported ER studies, this analysis looked at ER implementation across multiple surgical specialties. Overall, the findings for each specialty were similar to those for the entire cohort with regard to patient outcome improvement. Even if a significant improvement in an

individual outcome was not seen, a trend in the improved direction was observed. An exception to this was the outcomes observed for the HPB group, which demonstrated equivalent patient outcomes following adoption of ER protocols. These findings were likely the result of smaller included case volume and later provider adoption of ER protocols. It should be noted that this group has previously published on their larger experience with implementation of ER protocols in liver surgery and demonstrated that these protocols resulted in improved patient outcomes.<sup>10,31,34</sup>

Perhaps more important than the effects seen following ER implementation for all cases were those seen amongst patients undergoing high-magnitude surgery. While it is

**TABLE 4** Clinical characteristics for high-magnitude cases<sup>a</sup>, based on recovery protocol

Factor	Total <i>n</i> = 1,022	Traditional <i>n</i> = 651 (64)	Enhanced recovery <i>n</i> = 371 (36)	<i>p</i> Value
Superficial incisional SSI	51 (5)	36 (6)	15 (4)	0.294
Deep incisional SSI	10 (1)	8 (1)	2 (0.5)	0.281
Organ space infection	61 (6)	38 (6)	23 (6)	0.814
Wound disruption/dehiscence	24 (2)	16 (2)	8 (2)	0.760
Pneumonia	29 (3)	19 (3)	10 (3)	0.836
Mechanical ventilation > 48 h	8 (1)	6 (1)	2 (1)	0.718
Urinary tract infection	48 (5)	35 (5)	13 (4)	0.174
Transfusion ≤ 72 h of OR	305 (30)	214 (33)	91 (25)	<b>0.005</b>
Sepsis	24 (2)	17 (3)	7 (2)	0.462
Unplanned return to OR	45 (4)	31 (5)	24 (5)	0.459
Any complication	416 (41)	288 (44)	128 (35)	<b>0.002</b>
Any severe complication <sup>b</sup>	136 (13)	90 (14)	46 (12)	0.519
LOS, days, median (IQR)	6 (4–8)	6 (5–8)	5 (3–7)	<b>&lt;0.0001</b>
Mortality ≤ 30 days	3 (0.3)	3 (0.5)	0 (0)	0.558
Readmission ≤ 30 days	130 (13)	78 (12)	52 (14)	0.348

Bold values are statistically significant ( $p < 0.05$ )

SSI surgical site infection, OR operating room, IQR interquartile range, LOS length of stay

<sup>a</sup>Defined as RVU > 30

<sup>b</sup>Severe complications include: organ space infection, wound disruption/dehiscence, pneumonia, unplanned intubation, acute renal failure, myocardial infarction, sepsis, septic shock, unplanned return to OR

challenging to amalgamate outcomes across multiple specialties, focusing on high-magnitude cases enabled analysis of a cohort of patients undergoing operations of more uniform complexity. This is reflected by the equivalent median procedure durations for patients undergoing high-magnitude surgery and treated on ER versus TP protocols (327 vs. 326 min,  $p = 0.466$ ). That patients in the high magnitude cohort on ER protocols also were found to have improved outcomes, including decreased complications and reduced LOS, suggests the results of the analysis for all cases were not driven by a single specialty performing a large number of lower-complexity cases.

Because all surgical programs engage in continuous quality improvement at some level, one critique of this study is that it is possible that other simultaneous initiatives contributed to the observed improved patient outcomes. For example, the hepatobiliary and colorectal groups developed a quality improvement program around urinary tract infections that preceded ER implementation, and the gynecology group designed a perioperative bundle to reduce their rates of surgical site infections in parallel with their implementation. However, there were no ongoing system-wide improvements that could explain the consistency or magnitude of our cross-specialty findings, including the dramatic decrease in opioid utilization shared by all groups.

A second critique of the methodology may be the definition of ER protocol compliance. As previously discussed, the four pillars of ER protocols have been well described,

and each pillar includes more than one patient intervention. Although specific interventions were not proscribed at an institutional level, each team designed ordersets and pathways that facilitated high compliance rates with multiple accepted individual elements. Several studies have shown that ER protocol compliance of  $\geq 70\%$  results in a demonstrable effect on patient outcomes.<sup>1,35,36</sup> Regarding the adherence to guidelines in this study, preoperative education, preoperative carbohydrate loading (where indicated), and preoperative multimodal nonnarcotic oral analgesia were uniform. To facilitate consistent implementation and trainee education, electronically implemented ordersets (including the postoperative protocols and the analgesia plans) were utilized in more than 90% of ER pathway patients across all disciplines. Each of these ordersets was designed to include > 70% of recommended postoperative elements. Case audits demonstrated 60% to 100% compliance for each core component of the ER protocols (i.e., preoperative patient education and the four pillars of ER clinical care).

Lastly, the retrospective nature of this study makes it possible that other unmeasured confounding factors impacted our results. Although it is possible that confounding factors may have resulted in overestimation of the observed benefits of ER protocols, several elements of this study counter the potential for selection bias and support the conclusion that improvements in outcomes were attributable to ER protocol implementation. These include

the NSQIP methodology, random case sampling, consistency of implementation within and across specialties, correlation with high-magnitude cases, and large patient numbers.

## CONCLUSIONS

This analysis of a multispecialty enhanced recovery implementation at a large comprehensive cancer center demonstrates that not only is ER protocol implementation feasible for oncologic patients, but it is safe and effective in improving postoperative outcomes. This study provides evidence in support of ER utilization for a population of patients previously underrepresented in the ER literature. Considering the number of major oncologic operations performed annually across the world, wide-scale adoption of ER protocols is likely to improve the outcomes of many patients. As postoperative complications and delayed postoperative recovery is known to prevent return to adjuvant treatment for oncologic patients, future work should focus on defining the potential long-term oncologic benefits of ER protocols.<sup>9,16,37–39</sup>

**ACKNOWLEDGMENTS** The authors recognize the contributions of Dr. Jay Shah, who was the original surgical lead for the Urology ER program and has since moved to another institution. In addition, they thank our institution Surgical Clinical Reviewers, Annie Philip and Melony Levy, for their diligence and contributions to this study and to our broader patient quality improvement efforts.

**FUNDING** Drs. Rebecca Marcus and Heather Lillemoe are supported by National Institutes of Health grant T32CA009599 and the MD Anderson Cancer Center support grant P30CA016672.

**DISCLOSURE** None of the authors have any conflicts of interest associated with this study.

**ACS NSQIP DISCLAIMER** American College of Surgeons National Surgical Quality Improvement Program and the hospitals participating in the ACS NSQIP are the source of the data used herein. They have not verified and are not responsible for the statistical validity of the data analysis or the conclusions derived by the authors.

## REFERENCES

- Ljungqvist O, Scott M, Fearon KC. Enhanced recovery after surgery: a review. *JAMA Surg.* 2017;152:292–8.
- Manso M, Schmelz J, Aloia T. ERAS-anticipated outcomes and realistic goals. *J Surg Oncol.* 2017;116:570–7.
- Connor S, Cross A, Sakowska M, et al. Effects of introducing an enhanced recovery after surgery programme for patients undergoing open hepatic resection. *HPB (Oxford).* 2013;15:294–301.
- Coolsen MM, van Dam RM, van der Wilt AA, et al. Systematic review and meta-analysis of enhanced recovery after pancreatic surgery with particular emphasis on pancreaticoduodenectomies. *World J Surg.* 2013;37:1909–18.
- Hughes MJ, McNally S, Wigmore SJ. Enhanced recovery following liver surgery: a systematic review and meta-analysis. *HPB (Oxford).* 2014;16:699–706.
- Lau CS, Chamberlain RS. Enhanced recovery after surgery programs improve patient outcomes and recovery: a meta-analysis. *World J Surg.* 2017;41:899–913.
- Song W, Wang K, Zhang RJ, et al. The enhanced recovery after surgery (ERAS) program in liver surgery: a meta-analysis of randomized controlled trials. *SpringerPlus.* 2016;5:207.
- Visioni A, Shah R, Gabriel E, et al. Enhanced recovery after surgery for noncolorectal surgery? A systematic review and meta-analysis of major abdominal surgery. *Ann Surg.* 2018;267:57–65.
- Aloia TA, Zimmitti G, Conrad C et al. Return to intended oncologic treatment (RIOT): a novel metric for evaluating the quality of oncosurgical therapy for malignancy. *J Surg Oncol.* 2014;110:107–14.
- Day RW, Cleeland CS, Wang XS, et al. Patient-reported outcomes accurately measure the value of an enhanced recovery program in liver surgery. *J Am Coll Surg.* 2015;221:1023–30 e1021–22.
- Neville A, Lee L, Antonescu I, et al. Systematic review of outcomes used to evaluate enhanced recovery after surgery. *Br J Surg.* 2014;101:159–70.
- Takamoto T, Hashimoto T, Inoue K, et al. Applicability of enhanced recovery program for advanced liver surgery. *World J Surg.* 2014;38:2676–82.
- Alcantara-Moral M, Serra-Aracil X, Gil-Egea MJ, et al. Observational cross-sectional study of compliance with the fast track protocol in elective surgery for colon cancer in Spain. *Int J Colorectal Dis.* 2014;29:477–83.
- Schultz NA, Larsen PN, Klarskov B, et al. Evaluation of a fast-track programme for patients undergoing liver resection. *Br J Surg.* 2013;100:138–43.
- Bagnall NM, Malietzis G, Kennedy RH, et al. A systematic review of enhanced recovery care after colorectal surgery in elderly patients. *Colorectal Dis.* 2014;16:947–56.
- Gustafsson UO, Opielstrup H, Thorell A, et al. Adherence to the ERAS protocol is associated with 5-year survival after colorectal cancer surgery: a retrospective cohort study. *World J Surg.* 2016;40:1741–7.
- Li P, Fang F, Cai JX, et al. Fast-track rehabilitation vs conventional care in laparoscopic colorectal resection for colorectal malignancy: a meta-analysis. *World J Gastroenterol.* 2013;19:9119–26.
- van Vugt JL, Reisinger KW, Derikx JP, et al. Improving the outcomes in oncological colorectal surgery. *World J Gastroenterol.* 2014;20:12445–57.
- Kim BJ, Aloia TA. What is “enhanced recovery,” and how can it do it? *J Gastrointest Surg.* 2018;22:164–71.
- Patel SH, Kim BJ, Tzeng CD, et al. Reduction of cardiopulmonary/renal complications with serum bnp-guided volume status management in posthepatectomy patients. *J Gastrointest Surg.* 2018;22:467–76.
- Gustafsson UO, Scott MJ, Schwenk W, et al. Guidelines for perioperative care in elective colonic surgery: Enhanced Recovery After Surgery (ERAS(R)) Society recommendations. *Clin Nutr.* 2012;31:783–800.
- Lassen K, Coolsen MM, Slim K, et al. Guidelines for perioperative care for pancreaticoduodenectomy: Enhanced Recovery After Surgery (ERAS(R)) Society recommendations. *Clin Nutr.* 2012;31:817–30.
- Melloul E, Hubner M, Scott M, et al. Guidelines for perioperative care for liver surgery: enhanced recovery after surgery (ERAS) society recommendations. *World J Surg.* 2016;40:2425–40.
- Mortensen K, Nilsson M, Slim K, et al. Consensus guidelines for enhanced recovery after gastrectomy: Enhanced Recovery After



- Surgery (ERAS(R)) Society recommendations. *Br J Surg*. 2014;101:1209–29.
25. Nelson G, Altman AD, Nick A, et al. Guidelines for postoperative care in gynecologic/oncology surgery: Enhanced Recovery After Surgery (ERAS(R)) Society recommendations—Part II. *Gynecol Oncol*. 2016;140:323–32.
  26. Nelson G, Altman AD, Nick A, et al. Guidelines for pre- and intra-operative care in gynecologic/oncology surgery: Enhanced Recovery After Surgery (ERAS(R)) Society recommendations—Part I. *Gynecol Oncol*. 2016;140:313–22.
  27. Nygren J, Thacker J, Carli F, et al. Guidelines for perioperative care in elective rectal/pelvic surgery: Enhanced Recovery After Surgery (ERAS(R)) Society recommendations. *Clin Nutr*. 2012;31:801–16.
  28. Thorell A, MacCormick AD, Awad S, et al. Guidelines for perioperative care in bariatric surgery: Enhanced Recovery After Surgery (ERAS) Society Recommendations. *World J Surg*. 2016;40:2065–83.
  29. Scarborough JE, Bennett KM, Englum BR, et al. The impact of functional dependency on outcomes after complex general and vascular surgery. *Ann Surg*. 2015;261:432–7.
  30. Lasala JD, Heir JS, Mena GE, et al. Implementation of an Enhanced Surgical Recovery Programme (ESRP) gynaecologic oncology: how much did intra-operative opioid administration decrease in first year post implementation? *Clin Nutr ESPEN*. 2016;12:e42–3.
  31. Aloia TA, Kim BJ, Seagraves-Chun YS, et al. A randomized controlled trial of postoperative thoracic epidural analgesia versus intravenous patient-controlled analgesia after major hepatopancreatobiliary surgery. *Ann Surg*. 2017;266:545–54.
  32. Meyer LA, Lasala J, Iniesta MD, et al. Effect of an enhanced recovery after surgery program on opioid use and patient-reported outcomes. *Obstet Gynecol*. 2018; 132:281–90.
  33. Van Haren RM, Mehran RJ, Mena GE, et al. Enhanced recovery decreases pulmonary and cardiac complications after thoracotomy for lung cancer. *Ann Thorac Surg*. 2018;106:272–9.
  34. Zimmitti G, Soliz J, Aloia TA, et al. Positive impact of epidural analgesia on oncologic outcomes in patients undergoing resection of colorectal liver metastases. *Ann Surg Oncol*. 2016;23:1003–11.
  35. Gotlib Conn L, McKenzie M, Pearsall EA, McLeod RS. Successful implementation of an enhanced recovery after surgery programme for elective colorectal surgery: a process evaluation of champions' experiences. *Implement Sci*. 2015;10:99.
  36. Pedziwiatr M, Kisialeuski M, Wierdak M, et al. Early implementation of Enhanced Recovery After Surgery (ERAS(R)) protocol—Compliance improves outcomes: a prospective cohort study. *Int J Surg*. 2015;21:75–81.
  37. Kim BJ, Caudle AS, Gottumukkala V, Aloia TA. The impact of postoperative complications on a timely Return to Intended Oncologic Therapy (RIOT): the role of enhanced recovery in the cancer journey. *Int Anesthesiol Clin*. 2016;54:e33–46.
  38. Merkow RP, Bentrem DJ, Mulcahy MF, et al. Effect of postoperative complications on adjuvant chemotherapy use for stage III colon cancer. *Ann Surg*. 2013;258:847–53.
  39. Merkow RP, Bilimoria KY, Tomlinson JS, et al. Postoperative complications reduce adjuvant chemotherapy use in resectable pancreatic cancer. *Ann Surg*. 2014;260:372–7.

**Publisher's Note** Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.