

COHERENCE RELATIVE TO A WEAK TORSION CLASS

ZHANMIN ZHU, Jiaxing

Received September 22, 2016. First published February 10, 2018.

Abstract. Let R be a ring. A subclass \mathcal{T} of left R -modules is called a weak torsion class if it is closed under homomorphic images and extensions. Let \mathcal{T} be a weak torsion class of left R -modules and n a positive integer. Then a left R -module M is called \mathcal{T} -finitely generated if there exists a finitely generated submodule N such that $M/N \in \mathcal{T}$; a left R -module A is called (\mathcal{T}, n) -presented if there exists an exact sequence of left R -modules

$$0 \longrightarrow K_{n-1} \longrightarrow F_{n-1} \longrightarrow \dots \longrightarrow F_1 \longrightarrow F_0 \longrightarrow M \longrightarrow 0$$

such that F_0, \dots, F_{n-1} are finitely generated free and K_{n-1} is \mathcal{T} -finitely generated; a left R -module M is called (\mathcal{T}, n) -injective, if $\text{Ext}_R^n(A, M) = 0$ for each $(\mathcal{T}, n+1)$ -presented left R -module A ; a right R -module M is called (\mathcal{T}, n) -flat, if $\text{Tor}_n^R(M, A) = 0$ for each $(\mathcal{T}, n+1)$ -presented left R -module A . A ring R is called (\mathcal{T}, n) -coherent, if every $(\mathcal{T}, n+1)$ -presented module is $(n+1)$ -presented. Some characterizations and properties of these modules and rings are given.

Keywords: (\mathcal{T}, n) -presented module; (\mathcal{T}, n) -injective module; (\mathcal{T}, n) -flat module; (\mathcal{T}, n) -coherent ring

MSC 2010: 16D40, 16D50, 16P70

1. INTRODUCTION

Recall that a *torsion theory*, see [14], $\tau = (\mathcal{T}, \mathcal{F})$ for the category of all left R -modules consists of two subclasses \mathcal{T} and \mathcal{F} such that:

- (1) $\text{Hom}(T, F) = 0$ for all $T \in \mathcal{T}$ and $F \in \mathcal{F}$.
- (2) If $\text{Hom}(T, F) = 0$ for all $F \in \mathcal{F}$, then $T \in \mathcal{T}$.
- (3) If $\text{Hom}(T, F) = 0$ for all $T \in \mathcal{T}$, then $F \in \mathcal{F}$.

In this case, \mathcal{T} is called a torsion class.

This research was supported by the Natural Science Foundation of Zhejiang Province, China (LY18A010018).

A torsion theory $\tau = (\mathcal{T}, \mathcal{F})$ is called hereditary if \mathcal{T} is closed under submodules. By [14], page 139, Proposition 2.1, a class \mathcal{T} of left R -modules is a torsion class for some torsion theory if and only if \mathcal{T} is closed under quotient modules, direct sums and extensions. Inspired by this result, in this paper we will call a nonempty subclass \mathcal{T} of left R -modules a weak torsion class if \mathcal{T} is closed under homomorphic images and extensions.

Let $\tau = (\mathcal{T}, \mathcal{F})$ be a hereditary torsion theory for the category of all left R -modules. Then according to [8], a left R -module M is called τ -finitely generated (or τ -FG for short) if there exists a finitely generated submodule N such that $M/N \in \mathcal{T}$; a left R -module A is called τ -finitely presented (or τ -FP for short) if there exists an exact sequence of left R -modules $0 \rightarrow K \rightarrow F \rightarrow A \rightarrow 0$ with F finitely generated free and K τ -finitely generated. In Section 2, we will give the concepts of \mathcal{T} -finitely generated modules and \mathcal{T} -finitely presented modules by taking \mathcal{T} to be a weak torsion class of left R -modules, which extends the two concepts of Jones's τ -finitely generated modules and τ -finitely presented modules respectively. And then we will establish some properties of \mathcal{T} -finitely generated modules and \mathcal{T} -finitely presented modules.

Let n be a nonnegative integer. Then according to [4], a left R -module A is called n -presented in case there exists an exact sequence of left R -modules $F_n \rightarrow F_{n-1} \rightarrow \dots \rightarrow F_1 \rightarrow F_0 \rightarrow M \rightarrow 0$ in which every F_i is finitely generated free. Motivated by the concepts of n -presented modules and \mathcal{T} -finitely presented modules, in Section 3 we will define and investigate (\mathcal{T}, n) -presented modules.

Recall that a left R -module M is called *FP-injective*, see [13], or *absolutely pure*, see [11], if $\text{Ext}_R^1(A, M) = 0$ for any finitely presented left R -module A ; a right R -module M is flat if and only if $\text{Tor}_1^R(M, A) = 0$ for any finitely presented left R -module A ; a ring R is *left coherent*, see [1], if every finitely generated left ideal of R is finitely presented, or equivalently, if every finitely generated submodule of a projective left R -module is finitely presented. The FP-injective modules, flat modules, coherent rings and their generalizations have been studied extensively by many authors (see, for example, [1], [3], [4], [8], [10], [13], [18], [17]).

In 1994, Costa introduced the concept of *left n -coherent rings* in [4]. According to [4], a ring R is called left n -coherent in case every n -presented left R -module is $(n + 1)$ -presented. In 1996, Chen and Ding introduced the concepts of *n -FP-injective modules* and *n -flat modules*, see [3]. According to [3], a left R -module M is called n -FP-injective in case $\text{Ext}_R^n(A, M) = 0$ for any n -presented left R -module A , a right R -module M is called n -flat in case $\text{Tor}_n^R(M, A) = 0$ for any n -presented left R -module A . By using the concepts of n -FP-injective and n -flat modules, they characterized n -coherent rings. In 2012, Mao and Ding introduced the concepts of *τ - f -injective modules*, *τ -flat modules* and *τ -coherent rings*, see [10]. According to [10], a left R -module M is called τ - f -injective in case $\text{Ext}_R^1(R/I, M) = 0$ for any τ -finitely

presented left ideal I ; a right R -module M is called τ -flat in case $\text{Tor}_1^R(M, R/I) = 0$ for any τ -finitely presented left ideal I ; a ring R is called τ -coherent in case every τ -finitely presented left ideal is finitely presented. By using the concepts of τ -f-injective and τ -flat modules, they characterized τ -coherent rings.

Motivated by the characterization of n -coherent rings and τ -coherent rings (where τ is a hereditary torsion theory), in Section 5 we extend the concept of n -coherent rings and introduce the concept of (\mathcal{T}, n) -coherent rings (where \mathcal{T} is a weak torsion class). To characterize (\mathcal{T}, n) -coherent rings, (\mathcal{T}, n) -injective modules and (\mathcal{T}, n) -flat modules are introduced and studied in Section 4; some elementary properties of (\mathcal{T}, n) -injective modules and (\mathcal{T}, n) -flat modules are obtained in that section.

In Section 5, a series of characterizations and properties of (\mathcal{T}, n) -coherent rings are given. For instance, we prove: (1) A ring R is (\mathcal{T}, n) -coherent \Leftrightarrow any direct product of (\mathcal{T}, n) -flat right R -modules is (\mathcal{T}, n) -flat \Leftrightarrow any direct limit of (\mathcal{T}, n) -injective left R -modules is (\mathcal{T}, n) -injective \Leftrightarrow every right R -module has a (\mathcal{T}, n) -flat preenvelope \Leftrightarrow if N is a (\mathcal{T}, n) -injective left R -module, N_1 is an FP-injective submodule of N , then N/N_1 is (\mathcal{T}, n) -injective. (2) If R is a (\mathcal{T}, n) -coherent ring, then every left R -module has a (\mathcal{T}, n) -injective cover. (3) Every right R -module has a monic (\mathcal{T}, n) -flat preenvelope $\Leftrightarrow R$ is (\mathcal{T}, n) -coherent and ${}_R R$ is (\mathcal{T}, n) -injective $\Leftrightarrow R$ is (\mathcal{T}, n) -coherent and every left R -module has an epic (\mathcal{T}, n) -injective cover $\Leftrightarrow R$ is (\mathcal{T}, n) -coherent and every injective right R -module is (\mathcal{T}, n) -flat $\Leftrightarrow R$ is (\mathcal{T}, n) -coherent and every flat left R -module is (\mathcal{T}, n) -injective. As corollaries, some interesting results on n -coherent rings are obtained.

Throughout this paper, R is an associative ring with identity and all modules considered are unitary, n is a positive integer, \mathcal{T} is a weak torsion class of left R -modules. $R\text{-Mod}$ denotes the class of all left R -modules. For any R -module M , $M^+ = \text{Hom}(M, \mathbb{Q}/\mathbb{Z})$ will be the character module of M . Given a class \mathcal{L} of R -modules, we denote by $\mathcal{L}^\perp = \{M : \text{Ext}_R^1(L, M) = 0, L \in \mathcal{L}\}$ the right orthogonal class of \mathcal{L} , and by ${}^\perp\mathcal{L} = \{M : \text{Ext}_R^1(M, L) = 0, L \in \mathcal{L}\}$ the left orthogonal class of \mathcal{L} .

2. \mathcal{T} -FINITELY GENERATED AND \mathcal{T} -FINITELY PRESENTED MODULES

We begin with the following definition.

Definition 2.1. A nonempty subclass \mathcal{T} of left R -modules is called a *weak torsion class* if \mathcal{T} is closed under homomorphic images and extensions. If a class \mathcal{T} of left R -modules is a weak torsion class, then a left R -module M is called *\mathcal{T} -finitely generated* (or \mathcal{T} -FG for short) if there exists a finitely generated submodule N such that $M/N \in \mathcal{T}$. A left R -module A is called *\mathcal{T} -finitely presented* (or \mathcal{T} -FP for short)

if there exists an exact sequence of left R -modules $0 \rightarrow K \rightarrow F \rightarrow A \rightarrow 0$ with F finitely generated free and K \mathcal{T} -finitely generated.

Example 2.2.

- (1) Let R be a non-left noetherian left hereditary ring and \mathcal{T} the class of all injective left R -modules. Then by [16], Section 39.16, \mathcal{T} is a weak torsion class. But \mathcal{T} is not a torsion class.
- (2) Let \mathcal{T} be the class of all finitely generated left R -modules. Then by [16], Section 13.9 (1), \mathcal{T} is a weak torsion class. But \mathcal{T} is not a torsion class.
- (3) Let \mathcal{T} be the class of all finitely generated semisimple left R -modules. Then \mathcal{T} is a weak torsion class but not a torsion class.
- (4) Let \mathcal{T} be the class of all finitely generated left R -modules. Then a left R -module A is \mathcal{T} -finitely generated if and only if it is finitely generated.
- (5) Let $\mathcal{T} = R\text{-Mod}$. Then a left R -module A is \mathcal{T} -finitely presented if and only if it is finitely generated.
- (6) Let $\mathcal{T} = 0$. Then a left R -module A is \mathcal{T} -finitely presented if and only if it is finitely presented.

Theorem 2.3. (1) Any homomorphic image of a \mathcal{T} -FG module is \mathcal{T} -FG.

- (2) Any finite direct sum of \mathcal{T} -FG modules is \mathcal{T} -FG.
- (3) Any sum of a finite number of \mathcal{T} -FG submodules of a module M is \mathcal{T} -FG.
- (4) A direct summand of a \mathcal{T} -FP module is \mathcal{T} -FP.

Proof. (1) Let M be a \mathcal{T} -FG module and N a submodule of N . Since M is \mathcal{T} -FG, there exists a finitely generated submodule K of M such that $M/K \in \mathcal{T}$. Since \mathcal{T} is closed under homomorphic images, we have $(M/K)/[(K+N)/K] \in \mathcal{T}$, so $M/(K+N) \in \mathcal{T}$, and thus $(M/N)/(K+N)/N \in \mathcal{T}$. Observing that $(K+N)/N$ is finitely generated, we have that M/N is \mathcal{T} -FG.

(2) Let N_1, N_2 be two \mathcal{T} -FG modules. Then there exists a finitely generated submodule K_i of N_i such that $N_i/K_i \in \mathcal{T}$, $i = 1, 2$. So, $K_1 \oplus K_2$ is finitely generated and $(N_1 \oplus N_2)/(K_1 \oplus K_2) \cong N_1/K_1 \oplus N_2/K_2 \in \mathcal{T}$ because \mathcal{T} is closed under extensions. And thus $N_1 \oplus N_2$ is \mathcal{T} -FG.

(3) Let M_1, M_2 be two \mathcal{T} -FG submodules of M . Then by (2), $M_1 \oplus M_2$ is \mathcal{T} -FG. Note that $M_1 + M_2$ is a homomorphic image of $M_1 \oplus M_2$; by (1), $M_1 + M_2$ is \mathcal{T} -FG.

(4) Suppose that $M \cong F/K$ where F is finitely generated free and K is \mathcal{T} -FG. If $F/K = (A + K)/K \oplus (B + K)/K$, where A, B are finitely generated, then by (3), $B + K$ is \mathcal{T} -FG. But $(A + K)/K \cong F/(B + K)$, so $(A + K)/K$ is \mathcal{T} -FP. \square

Corollary 2.4. A direct summand of a \mathcal{T} -FG module is \mathcal{T} -FG.

Theorem 2.5. Let $0 \rightarrow A \xrightarrow{i} B \xrightarrow{p} C \rightarrow 0$ be an exact sequence of left R -modules.

- (1) If both A and C are \mathcal{T} -FG, then B is \mathcal{T} -FG.
- (2) If both A and C are \mathcal{T} -FP, then B is \mathcal{T} -FP.
- (3) If B is FG and C is \mathcal{T} -FP, then A is \mathcal{T} -FG.
- (4) If B is \mathcal{T} -FP and A is \mathcal{T} -FG, then C is \mathcal{T} -FP.

Proof. (1) Suppose that A and C are \mathcal{T} -FG. Then there exist a finitely generated submodule A' of A and a finitely generated submodule C' of C such that $A/A' \in \mathcal{T}$ and $C/C' \in \mathcal{T}$. Choose a finitely generated submodule B' of B such that $p(B') = C'$, let $A'' = A \cap (A' + B') = A' + (A \cap B')$, and define

$$\alpha: A/A'' \rightarrow B/(A' + B'); \quad a + A'' \mapsto a + (A' + B')$$

and

$$\bar{p}: B/(A' + B') \rightarrow C/C'; \quad b + (A' + B') \mapsto p(b) + C'.$$

Then we get an exact sequence $0 \rightarrow A/A'' \xrightarrow{\alpha} B/(A' + B') \xrightarrow{\bar{p}} C/C' \rightarrow 0$. Thus $A/A'' \cong (A/A')/(A''/A') \in \mathcal{T}$ and $C/C' \in \mathcal{T}$, so $B/(A' + B') \in \mathcal{T}$, and hence B is \mathcal{T} -FG.

(2) Since A and C are \mathcal{T} -FP, we have two exact sequences $0 \rightarrow K' \xrightarrow{\iota_1} F' \xrightarrow{f} A \rightarrow 0$ and $0 \rightarrow K'' \xrightarrow{\iota_2} F'' \xrightarrow{g} C \rightarrow 0$, where F', F'' are finitely generated free, K', K'' are \mathcal{T} -FG, ι_1, ι_2 are inclusion maps. Since F'' is projective, there exists a homomorphism $\sigma: F'' \rightarrow B$ such that $g = p\sigma$. And so we have the following commutative diagram with exact rows and columns:

$$\begin{array}{ccccccc}
 & & 0 & & 0 & & 0 \\
 & & \downarrow & & \downarrow & & \downarrow \\
 0 & \longrightarrow & K' & \xrightarrow{\lambda \iota_1} & \text{Ker}(h) & \xrightarrow{\pi \iota} & K'' \longrightarrow 0 \\
 & & \downarrow \iota_1 & & \downarrow \iota & & \downarrow \iota_2 \\
 0 & \longrightarrow & F' & \xrightarrow{\lambda} & F' \oplus F'' & \xrightarrow{\pi} & F'' \longrightarrow 0 \\
 & & \downarrow f & & \downarrow h & & \downarrow g \\
 0 & \longrightarrow & A & \xrightarrow{i} & B & \xrightarrow{p} & C \longrightarrow 0 \\
 & & \downarrow & & \downarrow & & \downarrow \\
 & & 0 & & 0 & & 0
 \end{array}$$

where λ is the natural injection, ι is the inclusion map, π is the natural projection, and

$$h: F' \oplus F'' \rightarrow B; \quad (x', x'') \mapsto \iota f(x') + \pi(x'').$$

By (1), $\text{Ker}(h)$ is \mathcal{T} -FG, and hence B is \mathcal{T} -FP.

(3) Suppose that B is FG and C is \mathcal{T} -FP. Let $F \xrightarrow{\varphi} B \rightarrow 0$ be exact with F FG free, let $K = \text{Ker}(p\varphi)$. Then $0 \rightarrow K \rightarrow F \rightarrow C \rightarrow 0$ is exact. Since C is \mathcal{T} -FP, there exists an exact sequence $0 \rightarrow K' \rightarrow F' \rightarrow C \rightarrow 0$ with F' FG free and K' \mathcal{T} -FG. By Schanuel's lemma, we have $K' \oplus F \cong K \oplus F'$, and thus K is \mathcal{T} -FG because a finite direct sum and a direct summand of \mathcal{T} -FG modules are \mathcal{T} -FG. Now let $\psi = \varphi|_K$. Observing that φ is epic, it is easy to see that ψ is an epimorphism from K to A . Hence, by Theorem 2.3 (1), A is \mathcal{T} -FG.

(4) Since B is \mathcal{T} -FP, there exists an exact sequence of left R -modules $0 \rightarrow K \rightarrow F \rightarrow B \rightarrow 0$ such that F is finitely generated free and K is \mathcal{T} -FG. Therefore, we can now from the pullback of $A \rightarrow B$ and $F \rightarrow B$ get the following commutative diagram:

$$\begin{array}{ccccccc}
 & & 0 & & 0 & & \\
 & & \downarrow & & \downarrow & & \\
 & & K & \xlongequal{\quad} & K & & \\
 & & \downarrow & & \downarrow & & \\
 0 & \longrightarrow & P & \longrightarrow & F & \longrightarrow & C \longrightarrow 0 \\
 & & \downarrow & & \downarrow & & \parallel \\
 0 & \longrightarrow & A & \longrightarrow & B & \longrightarrow & C \longrightarrow 0 \\
 & & \downarrow & & \downarrow & & \\
 & & 0 & & 0 & &
 \end{array}$$

with exact rows and columns. Since both K and A are \mathcal{T} -FG, by (1), P is also \mathcal{T} -FG, and so C is \mathcal{T} -FP. □

3. (\mathcal{T}, n) -PRESENTED MODULES

Definition 3.1. Let \mathcal{T} be a weak torsion class and n a positive integer. Then a left R -module A is said to be (\mathcal{T}, n) -presented if there exists an exact sequence of left R -modules

$$0 \rightarrow K_{n-1} \rightarrow F_{n-1} \rightarrow \dots \rightarrow F_1 \rightarrow F_0 \rightarrow M \rightarrow 0$$

such that F_0, \dots, F_{n-1} are finitely generated free and K_{n-1} is \mathcal{T} -finitely generated.

Clearly, a left R -module A is \mathcal{T} -finitely presented if and only if it is $(\mathcal{T}, 1)$ -presented. It is easy to see that every (\mathcal{T}, n) -presented module is $(\mathcal{T}, n-1)$ -presented. We also call \mathcal{T} -finitely generated modules $(\mathcal{T}, 0)$ -presented.

Example 3.2. (1) Let $\mathcal{T} = R\text{-Mod}$. Then a left R -module A is (\mathcal{T}, n) -presented if and only if it is $(n-1)$ -presented.

(2) Let $\mathcal{T} = 0$. Then a left R -module A is (\mathcal{T}, n) -presented if and only if it is n -presented.

Lemma 3.3. *Let A, B be two left R -modules and n a positive integer. If both A and B are (\mathcal{T}, n) -presented, then $A \oplus B$ is also (\mathcal{T}, n) -presented.*

Proof. It is a consequence of Theorem 2.3 (2). □

Proposition 3.4. *The following statements are equivalent for a left R -module A :*

- (1) A is (\mathcal{T}, n) -presented.
- (2) A is $(n-1)$ -presented, and if there exists an exact sequence of left R -modules

$$0 \longrightarrow K_{n-1} \longrightarrow F_{n-1} \longrightarrow \dots \longrightarrow F_1 \longrightarrow F_0 \longrightarrow A \longrightarrow 0$$

such that F_0, \dots, F_{n-1} are finitely generated free, then K_{n-1} is \mathcal{T} -finitely generated.

- (3) There exists an exact sequence of left R -modules

$$0 \longrightarrow K \longrightarrow F \longrightarrow A \longrightarrow 0$$

such that F is finitely generated free and K is $(\mathcal{T}, n-1)$ -presented.

If $n \geq 2$, then the above conditions are also equivalent to:

- (4) A is $(n-2)$ -presented, and if there exists an exact sequence of left R -modules

$$0 \longrightarrow K_{n-2} \longrightarrow F_{n-2} \longrightarrow \dots \longrightarrow F_1 \longrightarrow F_0 \longrightarrow A \longrightarrow 0$$

such that F_0, \dots, F_{n-2} are finitely generated free, then K_{n-2} is \mathcal{T} -finitely presented.

Proof. (1) \Rightarrow (2) Since A is (\mathcal{T}, n) -presented, there exists an exact sequence of left R -modules

$$0 \longrightarrow L_{n-1} \longrightarrow F'_{n-1} \longrightarrow \dots \longrightarrow F'_1 \longrightarrow F'_0 \longrightarrow A \longrightarrow 0$$

such that F'_0, \dots, F'_{n-1} are finitely generated free and L_{n-1} is \mathcal{T} -finitely generated, so A is $(n-1)$ -presented. Now if there exists an exact sequence of left R -modules

$$0 \longrightarrow K_{n-1} \longrightarrow F_{n-1} \longrightarrow \dots \longrightarrow F_1 \longrightarrow F_0 \longrightarrow A \longrightarrow 0$$

such that F_0, \dots, F_{n-1} are finitely generated free, then by the generalization of Schanuel's lemma [12], Exercise 3.37, and by Theorem 2.3 (2) and Corollary 2.4, K_{n-1} is \mathcal{T} -finitely generated.

(2) \Rightarrow (1); (1) \Leftrightarrow (3); and (2) \Leftrightarrow (4) are obvious. □

Proposition 3.5. *Let $0 \rightarrow A \rightarrow B \rightarrow C \rightarrow 0$ be an exact sequence of left R -modules. Then:*

- (1) *If both A and C are (\mathcal{T}, n) -presented, then so is B .*
- (2) *If B is (\mathcal{T}, n) -presented and A is $(\mathcal{T}, n-1)$ -presented, then C is (\mathcal{T}, n) -presented.*

Proof. (1) Use induction on n . If $n = 1$, then (1) holds by Theorem 2.5 (2). Suppose that (1) holds for $n - 1$. Let A and C be (\mathcal{T}, n) -presented. Then by Proposition 3.4, we have two exact sequences $0 \rightarrow K' \xrightarrow{\iota_1} F' \xrightarrow{f} A \rightarrow 0$ and $0 \rightarrow K'' \xrightarrow{\iota_2} F'' \xrightarrow{g} C \rightarrow 0$, where F', F'' are finitely generated free, K', K'' are $(\mathcal{T}, n-1)$ -presented, ι_1, ι_2 are inclusion maps. Using a method similar to the proof of Theorem 2.5 (2), by induction hypothesis and Proposition 3.4 we can get that B is also (\mathcal{T}, n) -presented.

(2) Since B is (\mathcal{T}, n) -presented, by Proposition 3.4 there exists an exact sequence of left R -modules $0 \rightarrow K \rightarrow F \rightarrow B \rightarrow 0$ such that F is finitely generated free and K is $(\mathcal{T}, n-1)$ -presented. Now, using a method similar to the proof of Theorem 2.5 (4), by (1) and Proposition 3.4, we can get that C is (\mathcal{T}, n) -presented. □

Corollary 3.6. *A direct summand of a (\mathcal{T}, n) -presented module is (\mathcal{T}, n) -presented.*

Proof. Use induction on n . If $n = 1$, then the conclusion holds by Theorem 2.3 (4). Suppose that the conclusion holds for $n - 1$. Let B be (\mathcal{T}, n) -presented and $B = A \oplus C$. Then by hypothesis, A is $(\mathcal{T}, n-1)$ -presented, and so C (\mathcal{T}, n) -presented by Proposition 3.5 (2), as required. □

Corollary 3.7. *The following statements are equivalent for a left R -module M :*

- (1) *M is (\mathcal{T}, n) -presented.*
- (2) *M is finitely generated and, if the sequence of left R -modules $0 \rightarrow K \rightarrow F \rightarrow M \rightarrow 0$ is exact with F finitely generated free, then K is $(\mathcal{T}, n-1)$ -presented.*

Proof. (1) \Rightarrow (2). Since M is (\mathcal{T}, n) -presented, by Proposition 3.4 (3) there exists an exact sequence of left R -modules $0 \rightarrow K' \rightarrow F' \rightarrow M \rightarrow 0$ such that F' is finitely generated free and K' is $(\mathcal{T}, n-1)$ -presented. So, by Schanuel's lemma, we have $K' \oplus F \cong K \oplus F'$, and thus K is $(\mathcal{T}, n-1)$ -presented because finite direct

sums and direct summands of $(\mathcal{T}, n - 1)$ -presented modules are $(\mathcal{T}, n - 1)$ -presented by Lemma 3.3 and Corollary 3.6.

(2) \Rightarrow (1). It follows from Proposition 3.4 (3). □

Corollary 3.8. *Let $n > 1$ and let $0 \rightarrow A \rightarrow B \rightarrow C \rightarrow 0$ be an exact sequence of left R -modules. If C is (\mathcal{T}, n) -presented and B is $(\mathcal{T}, n - 1)$ -presented, then A is $(\mathcal{T}, n - 1)$ -presented.*

Proof. Since $n > 1$ and B is $(\mathcal{T}, n - 1)$ -presented, we have the following commutative diagram:

$$\begin{array}{ccccccc}
 & & 0 & & 0 & & \\
 & & \downarrow & & \downarrow & & \\
 & & K & \xlongequal{\quad} & K & & \\
 & & \downarrow & & \downarrow & & \\
 0 & \longrightarrow & P & \longrightarrow & F & \longrightarrow & C \longrightarrow 0 \\
 & & \downarrow & & \downarrow & & \parallel \\
 0 & \longrightarrow & A & \longrightarrow & B & \longrightarrow & C \longrightarrow 0 \\
 & & \downarrow & & \downarrow & & \\
 & & 0 & & 0 & &
 \end{array}$$

with exact rows and columns, where F is finitely generated free. Moreover, by Corollary 3.7, K is $(\mathcal{T}, n - 2)$ -presented. Since C is (\mathcal{T}, n) -presented, by Corollary 3.7, P is $(\mathcal{T}, n - 1)$ -presented, and so A is $(\mathcal{T}, n - 1)$ -presented by Proposition 3.5 (2). □

4. (\mathcal{T}, n) -INJECTIVE AND (\mathcal{T}, n) -FLAT MODULES

Definition 4.1. A left R -module M is called (\mathcal{T}, n) -injective, if $\text{Ext}_R^n(A, M) = 0$ for each $(\mathcal{T}, n + 1)$ -presented left R -module A . A right R -module M is called (\mathcal{T}, n) -flat, if $\text{Tor}_n^R(M, A) = 0$ for each $(\mathcal{T}, n + 1)$ -presented left R -module A .

Clearly, n -FP-injective left R -modules are (\mathcal{T}, n) -injective, n -flat right R -modules are (\mathcal{T}, n) -flat. By Proposition 3.4 (3), it is easy to see that a (\mathcal{T}, n) -injective module is $(\mathcal{T}, n + 1)$ -injective, a (\mathcal{T}, n) -flat module is $(\mathcal{T}, n + 1)$ -flat. We denote by $\mathcal{T}_n\mathcal{I}$ the class of all (\mathcal{T}, n) -injective left R -modules, and denote by $\mathcal{T}_n\mathcal{F}$ the class of all (\mathcal{T}, n) -flat right R -modules. We recall that if n, d are nonnegative integers, then according to [18], a right R -module M is called (n, d) -injective if $\text{Ext}_R^{d+1}(A, M) = 0$ for every n -presented right R -module A ; a left R -module M is called (n, d) -flat if $\text{Tor}_{d+1}^R(A, M) = 0$ for every n -presented right R -module A .

Example 4.2. (1) Let $\mathcal{T} = R\text{-Mod}$. Then a left R -module M is (\mathcal{T}, n) -injective if and only if M is n -FP-injective, a right R -module M is (\mathcal{T}, n) -flat if and only if M is n -flat. In particular, a left R -module M is $(\mathcal{T}, 1)$ -injective if and only if M is FP-injective, a right R -module M is $(\mathcal{T}, 1)$ -flat if and only if M is flat.

(2) Let $\mathcal{T} = \{0\}$. Then a left R -module M is (\mathcal{T}, n) -injective if and only if M is $(n + 1, n - 1)$ -injective, a right R -module M is (\mathcal{T}, n) -flat if and only if M is $(n + 1, n - 1)$ -flat. In particular, a left R -module M is $(\mathcal{T}, 1)$ -injective if and only if M is $(2, 0)$ -injective, a right R -module M is $(\mathcal{T}, 1)$ -flat if and only if M is $(2, 0)$ -flat.

Recall that an exact sequence of left R -modules $0 \rightarrow M \rightarrow M' \rightarrow M'' \rightarrow 0$ is said to be pure if every finitely presented left R -module is projective with respect to this exact sequence.

Definition 4.3. Let $0 \rightarrow M \rightarrow M' \rightarrow M'' \rightarrow 0$ be an exact sequence of left R -modules. Then it is said to be \mathcal{T} -pure if every $(\mathcal{T}, 2)$ -presented left R -module is projective with respect to it.

Example 4.4. (1) Let $\mathcal{T} = R\text{-Mod}$. Then it is easy to see that an exact sequence of left R -modules $0 \rightarrow M \rightarrow M' \rightarrow M'' \rightarrow 0$ is pure if and only if it is \mathcal{T} -pure.

(2) Let $\mathcal{T} = \{0\}$. Then it is easy to see that an exact sequence of left R -modules $0 \rightarrow M \rightarrow M' \rightarrow M'' \rightarrow 0$ is \mathcal{T} -pure if and only if every 2-presented left R -module is projective with respect to it.

Let $\dots \rightarrow P_1 \xrightarrow{d_1} P_0 \xrightarrow{d_0} A \rightarrow 0$ be a projective resolution of a module A . As usual, we will denote $\text{Ker}(d_i)$ by K_i , and we will call K_i an i -syzygy of A . If $n \geq 2$, then it is easy to see that a left R -module A is $(\mathcal{T}, n + 1)$ -presented if and only if it is $(n - 2)$ -presented; and if the sequence of right R -modules $0 \rightarrow K_{n-2} \rightarrow F_{n-2} \rightarrow \dots \rightarrow F_1 \rightarrow F_0 \rightarrow A \rightarrow 0$ is exact, where F_0, \dots, F_{n-2} are finitely generated free, then K_{n-2} is $(\mathcal{T}, 2)$ -presented.

Theorem 4.5. Let M be a left R -module and $n \geq 2$. Then the following statements are equivalent:

- (1) M is (\mathcal{T}, n) -injective.
- (2) If the sequence $0 \rightarrow K_{n-2} \rightarrow F_{n-2} \rightarrow \dots \rightarrow F_1 \rightarrow F_0 \rightarrow A \rightarrow 0$ is exact, where F_0, \dots, F_{n-2} are finitely generated free and K_{n-2} is $(\mathcal{T}, 2)$ -presented, then $\text{Ext}_R^1(K_{n-2}, M) = 0$.
- (3) For every $(n - 1)$ -presentation $F_{n-1} \rightarrow \dots \rightarrow F_0 \rightarrow A \rightarrow 0$ of a $(\mathcal{T}, n + 1)$ -presented module A with $F_0, \dots, F_{n-2}, F_{n-1}$ finitely generated free, every homomorphism from the $(n - 1)$ -syzygy K_{n-1} to M can be extended to a homomorphism from F_{n-1} to M .
- (4) There exists a \mathcal{T} -pure exact sequence $0 \rightarrow M \rightarrow M' \rightarrow M'' \rightarrow 0$ of left R -modules with M' (\mathcal{T}, n) -injective.

Proof. (1) \Leftrightarrow (2). By the isomorphism $\text{Ext}_R^n(A, M) \cong \text{Ext}_R^1(K_{n-2}, M)$.

(2) \Leftrightarrow (3). By the exact sequence

$$\text{Hom}(F_{n-1}, M) \longrightarrow \text{Hom}(K_{n-1}, M) \longrightarrow \text{Ext}_R^1(K_{n-2}, M) \longrightarrow \text{Ext}_R^1(F_{n-1}, M) = 0.$$

(1) \Rightarrow (4). It is obvious.

(4) \Rightarrow (2). Since $0 \rightarrow M \rightarrow M' \rightarrow M'' \rightarrow 0$ is \mathcal{T} -pure and K_{n-2} is $(\mathcal{T}, 2)$ -presented, we have that the map $\text{Hom}(K_{n-2}, M') \rightarrow \text{Hom}(K_{n-2}, M'')$ is epic. So from the exact sequence

$$\text{Hom}(K_{n-2}, M') \longrightarrow \text{Hom}(K_{n-2}, M'') \longrightarrow \text{Ext}_R^1(K_{n-2}, M) \longrightarrow 0$$

we have $\text{Ext}_R^1(K_{n-2}, M) = 0$. □

Proposition 4.6. *Let $\{M_i : i \in I\}$ be a family of left R -modules. Then the following statements are equivalent:*

- (1) *Each M_i is (\mathcal{T}, n) -injective.*
- (2) $\prod_{i \in I} M_i$ *is (\mathcal{T}, n) -injective.*
- (3) $\bigoplus_{i \in I} M_i$ *is (\mathcal{T}, n) -injective.*

Proof. (1) \Leftrightarrow (2). By the isomorphism $\text{Ext}_R^n\left(A, \prod_{i \in I} M_i\right) \cong \prod_{i \in I} \text{Ext}_R^n(A, M_i)$.

(2) \Rightarrow (3). For every $(n-1)$ -presentation $F_{n-1} \rightarrow \dots \rightarrow F_0 \rightarrow A \rightarrow 0$ of a $(\mathcal{T}, n+1)$ -presented module A with $F_0, \dots, F_{n-2}, F_{n-1}$ finitely generated free, by Proposition 3.4 (4), the $(n-1)$ -syzygy K_{n-1} is \mathcal{T} -finitely presented and hence finitely generated. Let f be any homomorphism from K_{n-1} to $\bigoplus_{i \in I} M_i$. Then there exists a finite subset I_0 of I such that $\text{Im}(f) \subseteq \bigoplus_{i \in I_0} M_i$. By (2), $\bigoplus_{i \in I_0} M_i$ is (\mathcal{T}, n) -injective. So, by Theorem 4.5 (3), f can be extended to a homomorphism from F_{n-1} to $\bigoplus_{i \in I_0} M_i$, and then f can be extended to a homomorphism from F_{n-1} to $\bigoplus_{i \in I} M_i$. Therefore

$\bigoplus_{i \in I} M_i$ is (\mathcal{T}, n) -injective by Theorem 4.5 (3) again.

(3) \Rightarrow (1). It is trivial. □

Proposition 4.7. *Let $\{M_i : i \in I\}$ be a family of right R -modules. Then the following conditions are equivalent:*

- (1) *Every M_i is (\mathcal{T}, n) -flat.*
- (2) $\bigoplus_{i \in I} M_i$ *is (\mathcal{T}, n) -flat.*

Proof. By the isomorphism $\text{Tor}_n^R\left(\bigoplus_{i \in I} M_i, A\right) \cong \bigoplus_{i \in I} \text{Tor}_n^R(M_i, A)$. □

Theorem 4.8. *Let M be a right R -module. Then M is (\mathcal{T}, n) -flat if and only if M^+ is (\mathcal{T}, n) -injective.*

Proof. It follows from the isomorphism $\text{Tor}_n^R(M, A)^+ \cong \text{Ext}_R^n(A, M^+)$. \square

Proposition 4.9.

- (1) *Pure submodules of (\mathcal{T}, n) -injective modules are (\mathcal{T}, n) -injective.*
- (2) *Pure submodules of (\mathcal{T}, n) -flat modules are (\mathcal{T}, n) -flat.*

Proof. (1) Let N be a pure submodule of a (\mathcal{T}, n) -injective module M . Then N is \mathcal{T} -pure in M , and so, by Theorem 4.5 (4), N is (\mathcal{T}, n) -injective.

(2) Let M be a (\mathcal{T}, n) -flat module and N a pure submodule of M . Then the pure exact sequence $0 \rightarrow N \rightarrow M \rightarrow M/N \rightarrow 0$ induces a split exact sequence $0 \rightarrow (M/N)^+ \rightarrow M^+ \rightarrow N^+ \rightarrow 0$. By Theorem 4.8, M^+ is (\mathcal{T}, n) -injective, so N^+ is (\mathcal{T}, n) -injective by Proposition 4.6, and hence N is (\mathcal{T}, n) -flat by Theorem 4.8 again. \square

Remark 4.10. From Theorem 4.8, the (\mathcal{T}, n) -flatness of M_R can be characterized by the (\mathcal{T}, n) -injectivity of M^+ . On the other hand, by [3], Lemma 2.7 (1), the sequence $\text{Tor}_n^R(M^+, A) \rightarrow \text{Ext}_R^n(A, M^+) \rightarrow 0$ is exact for any n -presented left R -module A and any left R -module M . So, for any left R -module M , if M^+ is (\mathcal{T}, n) -flat, then M is (\mathcal{T}, n) -injective.

Let \mathcal{F} be a class of R -modules and M an R -module. Following [6], we say that a homomorphism $\varphi: M \rightarrow F$ where $F \in \mathcal{F}$ is an \mathcal{F} -preenvelope of M if for any morphism $f: M \rightarrow F'$ with $F' \in \mathcal{F}$ there is a $g: F \rightarrow F'$ such that $g\varphi = f$. An \mathcal{F} -preenvelope $\varphi: M \rightarrow F$ is said to be an \mathcal{F} -envelope if every endomorphism $g: F \rightarrow F$ such that $g\varphi = \varphi$ is an isomorphism. Dually, we have the definitions of an \mathcal{F} -precover and an \mathcal{F} -cover. The \mathcal{F} -envelopes (\mathcal{F} -covers) may not exist in general, but if they exist, they are unique up to isomorphism.

A pair $(\mathcal{A}, \mathcal{B})$ of classes of R -modules is called a cotorsion theory, see [6], if $\mathcal{A}^\perp = \mathcal{B}$ and ${}^\perp\mathcal{B} = \mathcal{A}$. A cotorsion theory $(\mathcal{A}, \mathcal{B})$ is called perfect, see [7], if every R -module has a \mathcal{B} -envelope and an \mathcal{A} -cover. A cotorsion theory $(\mathcal{A}, \mathcal{B})$ is called complete (see [6], Definition 7.1.6, and [15], Lemma 1.13) if for any R -module M there are exact sequences $0 \rightarrow M \rightarrow B \rightarrow A \rightarrow 0$ with $A \in \mathcal{A}$ and $B \in \mathcal{B}$, and $0 \rightarrow B' \rightarrow A' \rightarrow M \rightarrow 0$ with $A' \in \mathcal{A}$ and $B' \in \mathcal{B}$.

For a class \mathcal{F} of R -modules, we put $\mathcal{F}^+ = \{F^+ : F \in \mathcal{F}\}$. We recall that a left R -module M is said to be *pure injective* if it is injective with respect to all pure exact sequences of left R -modules. Following [15], we denote by \mathcal{PI} the class of pure injective left R -modules.

Theorem 4.11. *Let R be a ring. Then:*

- (1) $({}^\perp(\mathcal{T}_n\mathcal{I}), \mathcal{T}_n\mathcal{I})$ is a complete cotorsion theory.
- (2) $(\mathcal{T}_n\mathcal{F}, (\mathcal{T}_n\mathcal{F})^\perp)$ is a perfect cotorsion theory.

Proof. (1) Let X be the set of representatives of all K_{n-2} 's in Theorem 4.5 (2). Then by Theorem 4.5, $\mathcal{T}_n\mathcal{I} = X^\perp$, and so $({}^\perp(\mathcal{T}_n\mathcal{I}), \mathcal{T}_n\mathcal{I}) = ({}^\perp(X^\perp), X^\perp)$ is a complete cotorsion theory by [15], Theorem 2.2 (2).

(2) Write $\mathcal{A} = \mathcal{T}_n\mathcal{F}$ and let \mathcal{X} be the class of all K_{n-2} 's in Theorem 4.5 (2). Then by dimension shifting one shows that $A \in \mathcal{T}_n\mathcal{F}$ if and only if $\text{Tor}_1^R(A, X) = 0$ for each $X \in \mathcal{X}$. Thus, by the isomorphism $\text{Tor}_1^R(A, B)^+ \cong \text{Ext}_R^1(A, B^+)$, we have $\mathcal{A} = {}^\perp(\mathcal{X}^+)$, and so $(\mathcal{T}_n\mathcal{F}, (\mathcal{T}_n\mathcal{F})^\perp) = ({}^\perp(\mathcal{X}^+), ({}^\perp(\mathcal{X}^+))^\perp)$ is a cotorsion theory generated by \mathcal{X}^+ . Since every character module is pure injective by [6], Proposition 5.3.7, we have $\mathcal{X}^+ \subseteq \mathcal{P}\mathcal{I}$, and so it is a perfect cotorsion theory by [15], Theorem 2.8. \square

Following [6], Definition 5.3.22, a right R -module M is said to be *cotorsion* if $\text{Ext}_R^1(F, M) = 0$ for all flat right R -modules F . We call a right R -module M (\mathcal{T}, n) -cotorsion if $\text{Ext}_R^1(F, M) = 0$ for all (\mathcal{T}, n) -flat right R -modules F . By Theorem 4.11, we have the following results.

Corollary 4.12. *Let R be a ring. Then:*

- (1) Every right R -module has a (\mathcal{T}, n) -flat cover.
- (2) Every right R -module has a (\mathcal{T}, n) -cotorsion envelope.

5. (\mathcal{T}, n) -COHERENT RINGS

We begin this section with the concepts of (\mathcal{T}, n) -coherent rings and \mathcal{T} -coherent rings.

Definition 5.1. A ring R is called (\mathcal{T}, n) -coherent, if every $(\mathcal{T}, n+1)$ -presented module is $(n+1)$ -presented. A ring R is called \mathcal{T} -coherent if it is $(\mathcal{T}, 1)$ -coherent.

It is easy to see that a ring R is (\mathcal{T}, n) -coherent if and only if every (\mathcal{T}, n) -presented submodule of a finitely generated free left R -module is n -presented, and a ring R is \mathcal{T} -coherent if and only if every \mathcal{T} -finite presented submodule of a finitely generated free left R -module is finitely presented.

Example 5.2. (1) Let $\mathcal{T} = R\text{-Mod}$. Then R is (\mathcal{T}, n) -coherent if and only if R is left n -coherent. In particular, R is $(\mathcal{T}, 1)$ -coherent if and only if R is left coherent.

(2) Let $\mathcal{T} = \{0\}$. Then R is (\mathcal{T}, n) -coherent for any positive integer n .

Next we will characterize (\mathcal{T}, n) -coherent rings in terms of, among others, (\mathcal{T}, n) -injective modules and (\mathcal{T}, n) -flat modules. These results extend the theory of coherence of rings.

Theorem 5.3. *The following statements are equivalent for the ring R :*

- (1) R is (\mathcal{T}, n) -coherent.
- (2) $\varinjlim \text{Ext}_R^n(A, M_i) \cong \text{Ext}_R^n(A, \varinjlim M_i)$ for any $(\mathcal{T}, n+1)$ -presented module A and direct system $(M_i)_{i \in I}$ of left R -modules.
- (3) $\text{Tor}_n^R(\prod N_i, A) \cong \prod \text{Tor}_n^R(N_i, A)$ for any family $\{N_i\}$ of right R -modules and any $(\mathcal{T}, n+1)$ -presented module A .
- (4) Any direct product of copies of R_R is (\mathcal{T}, n) -flat.
- (5) Any direct product of (\mathcal{T}, n) -flat right R -modules is (\mathcal{T}, n) -flat.
- (6) Any direct limit of (\mathcal{T}, n) -injective left R -modules is (\mathcal{T}, n) -injective.
- (7) Any direct limit of injective left R -modules is (\mathcal{T}, n) -injective.
- (8) A left R -module M is (\mathcal{T}, n) -injective if and only if M^+ is (\mathcal{T}, n) -flat.
- (9) A left R -module M is (\mathcal{T}, n) -injective if and only if M^{++} is (\mathcal{T}, n) -injective.
- (10) A right R -module M is (\mathcal{T}, n) -flat if and only if M^{++} is (\mathcal{T}, n) -flat.
- (11) For any ring S , $\text{Tor}_n^R(\text{Hom}_S(B, E), A) \cong \text{Hom}_S(\text{Ext}_R^n(A, B), E)$ for the situation $({}_R A, {}_R B_S, E_S)$ with A $(\mathcal{T}, n+1)$ -presented and E_S injective.
- (12) Every right R -module has a (\mathcal{T}, n) -flat preenvelope.

Proof. (1) \Rightarrow (2). follows from [3], Lemma 2.9 (2).

(1) \Rightarrow (3). follows from [3], Lemma 2.10 (2).

(2) \Rightarrow (6) \Rightarrow (7) and (3) \Rightarrow (5) \Rightarrow (4) are trivial.

(7) \Rightarrow (1). Let A be $(\mathcal{T}, n+1)$ -presented with a finite n -presentation $F_n \xrightarrow{d_n} F_{n-1} \xrightarrow{d_{n-1}} \dots \rightarrow F_2 \xrightarrow{d_2} F_1 \xrightarrow{d_1} F_0 \xrightarrow{d_0} A \rightarrow 0$. Write $K_{n-1} = \text{Ker}(d_{n-1})$ and $K_{n-2} = \text{Ker}(d_{n-2})$. Then K_{n-1} is finitely generated, and we get an exact sequence of left R -modules $0 \rightarrow K_{n-1} \rightarrow F_{n-1} \rightarrow K_{n-2} \rightarrow 0$. Let $(E_i)_{i \in I}$ be any direct system of injective left R -modules (with I directed). Then $\varinjlim E_i$ is (\mathcal{T}, n) -injective by (7), so $\text{Ext}_R^n(A, \varinjlim E_i) = 0$ and then $\text{Ext}_R^1(K_{n-2}, \varinjlim E_i) = 0$. Thus, we have a commutative diagram

$$\begin{array}{ccccccc}
 \varinjlim \text{Hom}(K_{n-2}, E_i) & \longrightarrow & \varinjlim \text{Hom}(F_{n-1}, E_i) & \longrightarrow & \varinjlim \text{Hom}(K_{n-1}, E_i) & \longrightarrow & 0 \\
 \downarrow f & & \downarrow g & & \downarrow h & & \\
 \text{Hom}(K_{n-2}, \varinjlim E_i) & \longrightarrow & \text{Hom}(F_{n-1}, \varinjlim E_i) & \longrightarrow & \text{Hom}(K_{n-1}, \varinjlim E_i) & \longrightarrow & 0
 \end{array}$$

with exact rows. Since f and g are isomorphisms by [16], 25.4(d), h is an isomorphism by the Five lemma. Now, let $(M_i)_{i \in I}$ be any direct system of left R -modules (with

I directed). Then we have a commutative diagram with exact rows

$$\begin{array}{ccccc}
 0 \longrightarrow \varinjlim \operatorname{Hom}(K_{n-1}, M_i) & \longrightarrow & \varinjlim \operatorname{Hom}(K_{n-1}, E(M_i)) & \longrightarrow & \varinjlim \operatorname{Hom}(K_{n-1}, E(M_i)/M_i) \\
 & \searrow \varphi_1 & & \searrow \varphi_2 & \searrow \varphi_3 \\
 0 \longrightarrow \operatorname{Hom}(K_{n-1}, \varinjlim M_i) & \longrightarrow & \operatorname{Hom}(K_{n-1}, \varinjlim E(M_i)) & \longrightarrow & \operatorname{Hom}(K_{n-1}, \varinjlim E(M_i)/M_i)
 \end{array}$$

where $E(M_i)$ is the injective hull of M_i . Since K_{n-1} is finitely generated, by [16], Section 24.9, the maps φ_1 , φ_2 and φ_3 are monic. By the above proof, φ_2 is an isomorphism. Hence φ_1 is also an isomorphism by the Five lemma again, so K_{n-1} is finitely presented by [16], Section 25.4 (d), again, and thus A is $(n+1)$ -presented. Therefore R is (\mathcal{T}, n) -coherent.

(4) \Rightarrow (1). It follows similarly to (7) \Rightarrow (1).

(5) \Rightarrow (12). Let N be any left R -module. By [6], Lemma 5.3.12, there is a cardinal number \aleph_α dependent on $\operatorname{Card}(N)$ and $\operatorname{Card}(R)$ such that for any homomorphism $f: N \rightarrow F$ with F (\mathcal{T}, n) -flat, there is a pure submodule S of F such that $f(N) \subseteq S$ and $\operatorname{Card} S \leq \aleph_\alpha$. Thus f has a factorization $N \rightarrow S \rightarrow F$ with S (\mathcal{T}, n) -flat by Proposition 4.9 (2). Now let $(\varphi_\beta)_{\beta \in B}$ be all such homomorphisms $\varphi_\beta: N \rightarrow S_\beta$ with $\operatorname{Card} S_\beta \leq \aleph_\alpha$ and S_β (\mathcal{T}, n) -flat. Then any homomorphism $N \rightarrow F$ with F (\mathcal{T}, n) -flat has a factorization $N \rightarrow S_i \rightarrow F$ for some $i \in B$. Thus the homomorphism $N \rightarrow \prod_{\beta \in B} S_\beta$ induced by all φ_β is a (\mathcal{T}, n) -flat preenvelope since $\prod_{\beta \in B} S_\beta$ is (\mathcal{T}, n) -flat by (5).

(12) \Rightarrow (5). For any family $\{F_i\}_{i \in I}$ of (\mathcal{T}, n) -flat left R -modules, by hypothesis, $\prod_{i \in I} F_i$ has a (\mathcal{T}, n) -flat preenvelope $\varphi: \prod_{i \in I} F_i \rightarrow F$. Let $p_i: \prod_{i \in I} F_i \rightarrow F_i$ be the projection. Then there exists $f_i: F \rightarrow F_i$ such that $p_i = f_i \varphi$. Define $\psi: F \rightarrow \prod_{i \in I} F_i$ by $\psi(x) = (f_i(x))$ for every $x \in F$, then it is easy to check that $\psi \varphi = 1$. Hence $\prod_{i \in I} F_i$ is isomorphic to a direct summand of F , and so $\prod_{i \in I} F_i$ is (\mathcal{T}, n) -flat.

(1) \Rightarrow (11). For any $(\mathcal{T}, n+1)$ -presented module A , since R is (\mathcal{T}, n) -coherent, A is $(n+1)$ -presented. And so (11) follows from [3], Lemma 2.7 (2).

(11) \Rightarrow (8). Let $S = \mathbb{Z}$, $E = \mathbb{Q}/\mathbb{Z}$ and $B = M$. Then $\operatorname{Tor}_n^R(M^+, A) \cong \operatorname{Ext}_R^n(A, M)^+$ for any $(\mathcal{T}, n+1)$ -presented module A by (11), and hence (8) holds.

(8) \Rightarrow (9). Let M be a left R -module. If M is (\mathcal{T}, n) -injective, then M^+ is (\mathcal{T}, n) -flat by (8), and so M^{++} is (\mathcal{T}, n) -injective by Theorem 4.8. Conversely, if M^{++} is (\mathcal{T}, n) -injective, then M , being a pure submodule of M^{++} (see [14], Exercise 41, page 48), is (\mathcal{T}, n) -injective by Proposition 4.9 (1).

(9) \Rightarrow (10). If M is a (\mathcal{T}, n) -flat right R -module, then M^+ is a (\mathcal{T}, n) -injective left R -module by Theorem 4.8, and so M^{+++} is (\mathcal{T}, n) -injective by (9). Thus M^{++}

is (\mathcal{T}, n) -flat by Theorem 4.8 again. Conversely, if M^{++} is (\mathcal{T}, n) -flat, then M is (\mathcal{T}, n) -flat by Proposition 4.9 (2) as M is a pure submodule of M^{++} .

(10) \Rightarrow (5). Let $\{N_i\}_{i \in I}$ be a family of (\mathcal{T}, n) -flat right R -modules. Then by Proposition 4.7, $\bigoplus_{i \in I} N_i$ is (\mathcal{T}, n) -flat, and so $\left(\prod_{i \in I} N_i^+\right)^+ \cong \left(\bigoplus_{i \in I} N_i\right)^{++}$ is (\mathcal{T}, n) -flat by (10). Since $\bigoplus_{i \in I} N_i^+$ is a pure submodule of $\prod_{i \in I} N_i^+$ by [2], Lemma 1 (1), $\left(\prod_{i \in I} N_i^+\right)^+ \rightarrow \left(\bigoplus_{i \in I} N_i^+\right)^+ \rightarrow 0$ splits, and hence $\left(\bigoplus_{i \in I} N_i^+\right)^+$ is (\mathcal{T}, n) -flat. Thus $\prod_{i \in I} N_i^{++} \cong \left(\bigoplus_{i \in I} N_i^+\right)^+$ is (\mathcal{T}, n) -flat. Since $\prod_{i \in I} N_i$ is a pure submodule of $\prod_{i \in I} N_i^{++}$ by [2], Lemma 1 (2), $\prod_{i \in I} N_i$ is (\mathcal{T}, n) -flat by Proposition 4.9 (2). \square

Corollary 5.4. *The following statements are equivalent for a ring R :*

- (1) R is left n -coherent.
- (2) $\varinjlim \text{Ext}_R^n(C, M_\alpha) \cong \text{Ext}_R^n(C, \varinjlim M_\alpha)$ for any n -presented left R -module C and direct system $(M_\alpha)_{\alpha \in A}$ of left R -modules.
- (3) $\text{Tor}_n^R(\prod N_\alpha, C) \cong \prod \text{Tor}_n^R(N_\alpha, C)$ for any family $\{N_\alpha\}$ of right R -modules and any n -presented left R -module C .
- (4) Any direct product of copies of R_R is n -flat.
- (5) Any direct product of n -flat right R -modules is n -flat.
- (6) Any direct limit of n -FP-injective left R -modules is n -FP-injective.
- (7) Any direct limit of injective left R -modules is n -FP-injective.
- (8) A left R -module M is n -FP-injective if and only if M^+ is n -flat.
- (9) A left R -module M is n -FP-injective if and only if M^{++} is n -FP-injective.
- (10) A right R -module M is n -flat if and only if M^{++} is n -flat.
- (11) For any ring S , $\text{Tor}_n^R(\text{Hom}_S(B, E), C) \cong \text{Hom}_S(\text{Ext}_R^n(C, B), E)$ for the situation $({}_R C, {}_R B_S, E_S)$ with C n -presented and E_S injective.
- (12) Every right R -module has an n -flat preenvelope.

We note that the equivalences of (1)–(6), (8)–(11) in Corollary 5.4 appeared in [3], Theorem 3.1.

Lemma 5.5. *Let A be an $(n-1)$ -presented left R -module. Then A is n -presented if and only if $\text{Ext}_R^n(A, M) = 0$ for any FP-injective module M .*

Proof. Let A have a finite $(n-1)$ -presentation $F_{n-1} \xrightarrow{d_{n-1}} \dots \rightarrow F_2 \xrightarrow{d_2} F_1 \xrightarrow{d_1} F_0 \xrightarrow{\varepsilon} A \rightarrow 0$. Write $K_{n-2} = \text{Ker}(d_{n-2})$. Then K_{n-2} is finitely generated. By the isomorphism $\text{Ext}_R^n(A, M) \cong \text{Ext}_R^1(K_{n-2}, M)$, we have that $\text{Ext}_R^n(A, M) = 0$ for any FP-injective module M if and only if $\text{Ext}_R^1(K_{n-2}, M) = 0$ for any FP-injective module M . So, by [5], we have that $\text{Ext}_R^n(A, M) = 0$ for any FP-injective module M if and only if K_{n-2} is finitely presented, that is, A is n -presented. \square

Theorem 5.6. *The following statements are equivalent for a ring R .*

- (1) R is (\mathcal{T}, n) -coherent.
- (2) $\text{Ext}_R^{n+1}(A, N) = 0$ for any $(\mathcal{T}, n+1)$ -presented left R -module A and any FP-injective left R -module N .
- (3) If N is a (\mathcal{T}, n) -injective left R -module, N_1 is an FP-injective submodule of N , then N/N_1 is (\mathcal{T}, n) -injective.
- (4) For any FP-injective left R -module N , $E(N)/N$ is (\mathcal{T}, n) -injective, where $E(N)$ is the injective hull of N .

Proof. (1) \Rightarrow (2). For any $(\mathcal{T}, n+1)$ -presented left R -module A , there exists an exact sequence of left R -modules $0 \rightarrow K \rightarrow F \rightarrow A \rightarrow 0$, where F is finitely generated free and K is (\mathcal{T}, n) -presented. Since R is (\mathcal{T}, n) -coherent, K is n -presented, and so from the exact sequence

$$0 = \text{Ext}_R^n(F, N) \rightarrow \text{Ext}_R^n(K, N) \rightarrow \text{Ext}_R^{n+1}(A, N) \rightarrow \text{Ext}_R^{n+1}(F, N) = 0$$

we have $\text{Ext}_R^{n+1}(A, N) \cong \text{Ext}_R^n(K, N) = 0$ by Lemma 5.5 since N is FP-injective.

(2) \Rightarrow (3). For any $(\mathcal{T}, n+1)$ -presented left R -module A , the exact sequence $0 \rightarrow N_1 \rightarrow N \rightarrow N/N_1 \rightarrow 0$ induces the exactness of the sequence

$$0 = \text{Ext}_R^n(A, N) \rightarrow \text{Ext}_R^n(A, N/N_1) \rightarrow \text{Ext}_R^{n+1}(A, N_1) = 0.$$

Therefore $\text{Ext}_R^n(A, N/N_1) = 0$, as required.

(3) \Rightarrow (4) is obvious.

(4) \Rightarrow (1). Let A be a $(\mathcal{T}, n+1)$ -presented left R -module. Then there exists an exact sequence of left R -modules $0 \rightarrow K \rightarrow F \rightarrow A \rightarrow 0$, where F is finitely generated free and K is $(n-1)$ -presented. For any FP-injective module N , $E(N)/N$ is (\mathcal{T}, n) -injective by (4). From the exactness of the two sequences

$$0 = \text{Ext}_R^n(F, N) \rightarrow \text{Ext}_R^n(K, N) \rightarrow \text{Ext}_R^{n+1}(A, N) \rightarrow \text{Ext}_R^{n+1}(F, N) = 0$$

and

$$0 = \text{Ext}_R^n(A, E(N)) \rightarrow \text{Ext}_R^n(A, E(N)/N) \rightarrow \text{Ext}_R^{n+1}(A, N) \rightarrow \text{Ext}_R^{n+1}(A, E(N)) = 0$$

we have $\text{Ext}_R^n(K, N) \cong \text{Ext}_R^{n+1}(A, N) \cong \text{Ext}_R^n(A, E(N)/N) = 0$. Thus, K is n -presented by Lemma 5.5, and so A is $(n+1)$ -presented. Therefore, R is (\mathcal{T}, n) -coherent. \square

Corollary 5.7. *The following statements are equivalent for a ring R :*

- (1) R is left n -coherent.
- (2) $\text{Ext}_R^{n+1}(A, N) = 0$ for any n -presented left R -module A and any FP-injective left R -module N .
- (3) If N is an n -FP-injective left R -module, N_1 is an FP-injective submodule of N , then N/N_1 is n -FP-injective.
- (4) For any FP-injective left R -module N , $E(N)/N$ is n -FP-injective.

Corollary 5.8. *Let R be a (\mathcal{T}, n) -coherent ring. Then every left R -module has a (\mathcal{T}, n) -injective cover.*

Proof. Let $0 \rightarrow A \rightarrow B \rightarrow C \rightarrow 0$ be a pure exact sequence of left R -modules with B (\mathcal{T}, n) -injective. Then $0 \rightarrow C^+ \rightarrow B^+ \rightarrow A^+ \rightarrow 0$ is split exact. Since R is (\mathcal{T}, n) -coherent, B^+ is (\mathcal{T}, n) -flat by Theorem 5.3 (8), so C^+ is (\mathcal{T}, n) -flat, and hence C is (\mathcal{T}, n) -injective by Remark 4.10. Thus, the class of (\mathcal{T}, n) -injective modules is closed under pure quotients. By [9], Theorem 2.5, and Proposition 4.6, every left R -module has a (\mathcal{T}, n) -injective cover. \square

Corollary 5.9. *Let R be a left n -coherent ring. Then every left R -module has an n -FP-injective cover.*

Corollary 5.10. *The following statements are equivalent for a (\mathcal{T}, n) -coherent ring R :*

- (1) Every (\mathcal{T}, n) -flat right R -module is n -flat.
- (2) Every (\mathcal{T}, n) -injective left R -module is n -FP-injective.

In this case, R is left n -coherent.

Proof. (1) \Rightarrow (2). Let M be any (\mathcal{T}, n) -injective left R -module. Then M^+ is a (\mathcal{T}, n) -flat right R -module by Theorem 5.3 (8) since R is (\mathcal{T}, n) -coherent, and so M^+ is n -flat by (1). Thus M^{++} is n -FP-injective. Since M is a pure submodule of M^{++} , and a pure submodule of an n -FP-injective module is n -FP-injective, so M is n -FP-injective.

(2) \Rightarrow (1). Let M be any (\mathcal{T}, n) -flat right R -module. Then M^+ is a (\mathcal{T}, n) -injective left R -module by Theorem 4.8, and so M^+ is n -FP-injective by (2). Thus M is n -flat.

In this case, any direct product of n -flat right R -modules is n -flat by Theorem 5.3 (5), and so R is left n -coherent by Corollary 5.4 (5). \square

Proposition 5.11. *The following statements are equivalent for a ring R :*

- (1) Every right R -module has a monic (\mathcal{T}, n) -flat preenvelope.

- (2) R is (\mathcal{T}, n) -coherent and ${}_R R$ is (\mathcal{T}, n) -injective.
- (3) R is (\mathcal{T}, n) -coherent and every left R -module has an epic (\mathcal{T}, n) -injective cover.
- (4) R is (\mathcal{T}, n) -coherent and every injective right R -module is (\mathcal{T}, n) -flat.
- (5) R is (\mathcal{T}, n) -coherent and every flat left R -module is (\mathcal{T}, n) -injective.

Proof. (1) \Rightarrow (4). Assume (1). Then it is clear that R is a (\mathcal{T}, n) -coherent ring by Theorem 5.3 (12). Let E be any injective right R -module. E has a monic (\mathcal{T}, n) -flat preenvelope F , so E is isomorphic to a direct summand of F , and thus E is (\mathcal{T}, n) -flat.

(4) \Rightarrow (5). Let M be a flat left R -module. Then M^+ is injective, and so M^+ is (\mathcal{T}, n) -flat by (4). Hence M is (\mathcal{T}, n) -injective by Theorem 5.3 (8).

(5) \Rightarrow (2). It is obvious.

(2) \Rightarrow (1). Let M be any right R -module. Then M has a (\mathcal{T}, n) -flat preenvelope $f: M \rightarrow F$ by Theorem 5.3 (12). Since $({}_R R)^+$ is a cogenerator, there exists an exact sequence $0 \rightarrow M \xrightarrow{g} \prod ({}_R R)^+$. Since ${}_R R$ is (\mathcal{T}, n) -injective, by Theorem 5.3, $\prod ({}_R R)^+$ is (\mathcal{T}, n) -flat, and so there exists a right R -homomorphism $h: F \rightarrow \prod ({}_R R)^+$ such that $g = hf$, which shows that f is monic.

(2) \Rightarrow (3). Let M be a left R -module. Then M has a (\mathcal{T}, n) -injective cover $\varphi: C \rightarrow M$ by Corollary 5.8. On the other hand, there is an exact sequence $F \xrightarrow{\alpha} M \rightarrow 0$ with F free. Since F is (\mathcal{T}, n) -injective by (2) and Proposition 4.6, there exists a homomorphism $\beta: F \rightarrow C$ such that $\alpha = \varphi\beta$. It follows that φ is epic.

(3) \Rightarrow (2). Let $f: N \rightarrow {}_R R$ be an epic (\mathcal{T}, n) -injective cover. Then the projectivity of ${}_R R$ implies that ${}_R R$ is isomorphic to a direct summand of N , and so ${}_R R$ is (\mathcal{T}, n) -injective. \square

Corollary 5.12. *The following statements are equivalent for a ring R :*

- (1) Every right R -module has a monic n -flat preenvelope.
- (2) R is left n -coherent and ${}_R R$ is n -FP-injective.
- (3) R is left n -coherent and every left R -module has an epic n -FP-injective cover.
- (4) R is left n -coherent and every injective right R -module is n -flat.
- (5) R is left n -coherent and every flat left R -module is n -FP-injective.

Acknowledgment. The author wishes to thank the referee for careful reading of the paper and giving a detailed and helpful report.

References

- [1] *S. U. Chase*: Direct products of modules. *Trans. Am. Math. Soc.* *97* (1960), 457–473. [zbl](#) [MR](#) [doi](#)
- [2] *T. J. Cheatham, D. R. Stone*: Flat and projective character modules. *Proc. Am. Math. Soc.* *81* (1981), 175–177. [zbl](#) [MR](#) [doi](#)
- [3] *J. Chen, N. Ding*: On n -coherent rings. *Commun. Algebra* *24* (1996), 3211–3216. [zbl](#) [MR](#) [doi](#)
- [4] *D. L. Costa*: Parameterizing families of non-Noetherian rings. *Commun. Algebra* *22* (1994), 3997–4011. [zbl](#) [MR](#) [doi](#)
- [5] *E. Enochs*: A note on absolutely pure modules. *Canad. Math. Bull.* *19* (1976), 361–362. [zbl](#) [MR](#) [doi](#)
- [6] *E. E. Enochs, O. M. G. Jenda*: Relative Homological Algebra. *De Gruyter Expositions in Mathematics* *30*, Walter de Gruyter, Berlin, 2000. [zbl](#) [MR](#) [doi](#)
- [7] *E. E. Enochs, O. M. G. Jenda, J. A. Lopez-Ramos*: The existence of Gorenstein flat covers. *Math. Scand.* *94* (2004), 46–62. [zbl](#) [MR](#) [doi](#)
- [8] *M. Finkel Jones*: Coherence relative to an hereditary torsion theory. *Commun. Algebra* *10* (1982), 719–739. [zbl](#) [MR](#) [doi](#)
- [9] *H. Holm, P. Jørgensen*: Covers, precovers, and purity. *Illinois J. Math.* *52* (2008), 691–703. [zbl](#) [MR](#)
- [10] *L. Mao, N. Ding*: Relative coherence of rings. *J. Algebra Appl.* *11* (2012), 1250047, 16 pages. [zbl](#) [MR](#) [doi](#)
- [11] *C. Megibben*: Absolutely pure modules. *Proc. Am. Math. Soc.* *26* (1970), 561–566. [zbl](#) [MR](#) [doi](#)
- [12] *J. J. Rotman*: An Introduction to Homological Algebra. *Pure and Applied Mathematics* *85*, Academic Press, Harcourt Brace Jovanovich Publishers, New York-London, 1979. [zbl](#) [MR](#)
- [13] *B. Stenström*: Coherent rings and FP-injective modules. *J. Lond. Math. Soc., II. Ser.* *2* (1970), 323–329. [zbl](#) [MR](#) [doi](#)
- [14] *B. Stenström*: Rings of Quotients. *An Introduction to Methods of Ring Theory. Die Grundlehren der mathematischen Wissenschaften, Band 217*, Springer, New York, 1975. [zbl](#) [MR](#) [doi](#)
- [15] *J. Trlifaj*: Cover, Envelopes, and Cotorsion Theories. *Lecture notes for the workshop. Homological Methods in Module Theory*, Cortona, 2000.
- [16] *R. Wisbauer*: Foundations of Module and Ring Theory. *A Handbook for Study and Research. Algebra, Logic and Applications* *3*, Gordon and Breach Science Publishers, Philadelphia, 1991. [zbl](#) [MR](#)
- [17] *X. Yang, Z. Liu*: n -flat and n -FP-injective modules. *Czech. Math. J.* *61* (2011), 359–369. [zbl](#) [MR](#) [doi](#)
- [18] *D. Zhou*: On n -coherent rings and (n, d) -rings. *Commun. Algebra* *32* (2004), 2425–2441. [zbl](#) [MR](#) [doi](#)

Author's address: Zhanmin Zhu, Department of Mathematics, Jiaying University, 118 Jiahang Rd, Nanhu, 314001 Jiaying, Zhejiang, P. R. China e-mail: zhuzhanminzjxu@hotmail.com.