

A Modification-Specific Peptide-Based Immunization Approach Using CRM197 Carrier Protein: Development of a Selective Vaccine Against Pyroglutamate A β Peptides

Valérie Vingtdeux,* Haitian Zhao, Pallavi Chandakkar, Christopher M Acker, Peter Davies, and Philippe Marambaud

The Litwin-Zucker Research Center for the Study of Alzheimer's Disease, The Feinstein Institute for Medical Research, Manhasset, New York, United States of America. *Current affiliation: INSERM, Jean-Pierre Aubert Research Center, Université de Lille, Faculté de Médecine, Lille, France

Strategies aimed at reducing cerebral accumulation of the amyloid- β (A β) peptides have therapeutic potential in Alzheimer's disease (AD). A β immunization has proven to be effective at promoting A β clearance in animal models, but adverse effects have hampered its clinical evaluation. The first anti-A β immunization clinical trial, which assessed a full-length A β 1-42 vaccine, showed an increased risk of encephalitis, most likely because of autoimmune proinflammatory T helper 1 (Th1) response against all forms of A β . Immunization against less abundant but potentially more pathologically relevant A β products, such as N-terminally truncated pyroglutamate-3 A β (A β pE3), could provide efficacy and improve tolerability in A β immunotherapy. Here, we describe a selective vaccine against A β pE3 that uses the diphtheria toxin mutant CRM197 as a carrier protein for epitope presentation. CRM197 is currently used in licensed vaccines and has demonstrated excellent immunogenicity and safety in humans. In mice, our A β pE3:CRM197 vaccine triggered the production of specific anti-A β pE3 antibodies that did not cross-react with A β 1-42, non-cyclized A β E3 or N-terminally truncated pyroglutamate-11 A β (A β pE11). A β pE3:CRM197 antiserum strongly labeled A β pE3 in insoluble protein extracts and decorated cortical amyloid plaques in human AD brains. Anti-A β pE3 antibodies were almost exclusively of the IgG1 isotype, suggesting an antiinflammatory Th2 response bias to the A β pE3:CRM197 vaccine. To the best of our knowledge, this study shows for the first time that CRM197 has potential as a safe and suitable vaccine carrier for active and selective immunization against specific protein sequence modifications or conformations such as A β pE3.

Online address: <http://www.molmed.org>

doi: 10.2119/molmed.2016.00218

INTRODUCTION

Anti-amyloid- β (A β) immunotherapy is under intense investigation in Alzheimer's disease (AD) (1–3). A β is the core component of amyloid plaques (a hallmark of the AD brain) and mutations in its precursor APP or in presenilins, the catalytic components of the A β -producing enzyme γ -secretase, cause familial AD (4–6). Thus, strategies aimed

at preventing or lowering A β cerebral accumulation might interfere with AD pathogenesis (7). A β immunization has proven to be very effective at promoting A β clearance, at least in animal models. Preclinical and clinical studies, however, have been hampered by unforeseen side effects. The first clinical trial, AN-1792, which evaluated an A β 1-42/QS-21 vaccine, was halted in phase II when about

6% of the patients developed meningoencephalitis (8). Although the exact mechanism that led to acute brain inflammation in this clinical trial remains unclear, it is believed that encephalitis arose from an autoimmune reaction triggered by a vaccine directed against the abundant self-protein A β coupled to the strong adjuvant QS-21, thus favoring proinflammatory T helper 1 (Th1) immune response (9). In this context, second-generation anti-A β vaccines were designed to prevent T cell response during anti-A β immunization. These vaccines tested, for instance, N-terminal epitopes within the A β sequence and adjuvants that minimize T cell engagement and favor B cell response (2). Another approach is passive immunization, which has the advantage of bypassing T cell engagement and allowing better

Address correspondence to Philippe Marambaud, The Feinstein Institute for Medical Research, 350 Community Drive, Manhasset, NY 11030, USA. Phone: 516-562-0425; Fax: 516-562-0401; E-mail: PMaramba@northwell.edu
Submitted November 8, 2016; Accepted for Publication November 21, 2016; Published Online (www.molmed.org) November 28, 2016.

control of monoclonal antibody (mAb) dosage and epitope targeting. However, recent phase III AD trials of two anti-A β mAbs, solanezumab and bapineuzumab, failed to slow cognitive or functional decline in patients with mild to moderate AD (10). The main argument put forward to explain the lack of efficacy of these passive immunization approaches was that treatment might have started too late to reverse or delay the disease process (11). It is also possible that passive immunization might not deliver enough mAbs to promote plaque clearance. Passive immunization also raised concerns because of the practical and financial sustainability of injecting and monitoring mAb injections on a regular basis for several years (12).

In the amyloidogenic pathway, APP is sequentially endoproteolyzed by the proteases β -secretase/BACE1 and the presenilin/ γ -secretase complex to produce various A β peptides, including the most abundant isoforms, A β 1-40 and A β 1-42 (13). In addition to these major A β isoforms, N-terminally truncated A β products have been identified in the AD brain, including peptides, starting with pyroglutamate residues at positions 3 (A β pE3) and 11 (A β pE11) (14–16). N-terminal truncation was proposed to be mediated, at least in part, by aminopeptidase A (17), and cyclization of N-terminally exposed glutamates is catalyzed by the enzyme glutamyl cyclase (18). Pyroglutamate A β is a promising target because it appears to play a key role in A β oligomerization, seeding and stabilization (19–24). Furthermore, pyroglutamate A β has specific neurotoxic properties in cell cultures and leads to cerebral neuronal loss and synaptic function impairment in mice (25–28).

In this context, recent studies have proposed that immunization against less abundant but potentially more amyloidogenic and neurotoxic isoforms of A β , such as A β pE3, could improve tolerability and efficacy in A β immunotherapy (29–31). These studies, however, are limited to passive immunization using antibodies previously screened *in vitro*

for their specificity to A β pE3 and non-cross-reactivity to full-length A β . Here, we describe a novel A β pE3 vaccine using the nontoxic mutant of diphtheria toxin CRM197 (cross-reacting material 197) as a carrier protein for epitope presentation. CRM197 has been extensively used in licensed vaccines directed against capsular polysaccharides of several bacterial pathogens and has demonstrated excellent efficacy and tolerability in humans (32,33). Because recent data have shown that CRM197 is also suitable for conjugation to and presentation of peptides (34), we speculated that it could be conjugated to minimal peptide epitopes to facilitate the generation of conformation/modification-specific antibodies directed against pyroglutamate A β . We show that our vaccines in mice, composed of A β pE3-8 or A β pE11-16 peptides covalently conjugated to CRM197, triggered the production of fully specific antibodies directed against A β pE3 and A β pE11, respectively. Anti-A β pE3 antibodies stained brains from AD patients by Western Blotting (WB) and immunohistochemistry (IHC), and were almost exclusively of the IgG1 isotype, indicating the engagement of a Th2 response.

MATERIALS AND METHODS

Preparation of A β pE3-8:CRM197 and A β pE11-16:CRM197 Conjugate Vaccines

C-terminally amidated A β pE3-8 and A β pE11-16 peptides containing a C-terminal cysteine residue preceded by a two-glycine bridge (pEFRHDSGGC and pEVHHQKGGC, respectively; GenScript) were solubilized in phosphate-buffered saline (PBS) containing 2 mM ethylenediaminetetraacetic acid (EDTA) to obtain 4 mg/mL solutions. CRM197 (List Biological Labs) was reconstituted with PBS/EDTA to obtain a 2 mg/mL solution. To irreversibly crosslink the peptides to the carrier protein, CRM197 was first activated with a 20-fold molar excess of succinimidyl-4-(N-maleimidomethyl) cyclohexane-1-carboxylate (SMCC) cross-linker (Pierce, at 1.5 mg/mL in dimethyl

sulfoxide). After 30 min incubation at room temperature (RT), the SMCC/CRM197 mixture was desalted on a resin column (Zeba Desalt Spin Column, Pierce). Activated CRM197 was then combined with A β pE3-8 and A β pE11-16 peptide solutions and incubated for 30 min at RT. A ratio of 1:10 (CRM197 to peptides) was empirically chosen. Successful conjugation was confirmed by sodium dodecyl sulfate polyacrylamide gel electrophoresis (SDS-PAGE) and coomassie staining (GelCode, Pierce; see Figure 2A).

Immunization and Sample Collection

Animal experiments were performed according to procedures approved by The Feinstein Institute for Medical Research Institutional Animal Care and Use Committee. C57BL/6J male mice (~3 months old) were injected subcutaneously with 10 μ g of A β pE3-8:CRM197 and A β pE11-16:CRM197 conjugate vaccines or with 10 μ g of free A β pE3-8 and A β pE11-16 peptides used as negative controls. A prime injection (d 0) was followed by a boost injection at d 14 of the same amount of vaccines or free peptides. Blood samples were taken at d 0 and d 30.

ELISA for Measurement of Antibody Responses

Anti-A β pE3-8, anti-A β pE11-16 and anti-CRM197 antibody levels in mouse serum were determined by enzyme-linked immunosorbent assay (ELISA). Ninety-six-well plates (Maxisorp, Nunc) were coated with 100 μ L of 2 μ g/mL A β pE3-8, A β pE11-16 or CRM197 in carbonate buffer (0.05 M, pH 9.6) and incubated overnight at 4°C. Control plates were also coated with non-cyclized A β E3-8 and A β E11-16 (GenScript) to determine the levels of antibodies directed against these peptides. The following morning, plates were blocked for 1 h at RT with 5% skim milk in Tris-buffered saline (TBS) containing 0.05% Tween 20 (TBST). After washing with TBST, serial dilutions of individual mouse serum samples (diluted in PBST containing 1% skim milk) were prepared and 100 μ L/well of mouse serum was

incubated for 2 h at RT. After five more washes, 100 μ L/well horseradish peroxidase (HRP)-conjugated goat anti-mouse immunoglobins (Igs) secondary antibody (Southern Biotech, diluted 1:500 in PBST containing 1% skim milk) was incubated for 1 h at RT. TMB substrate was added after another wash and the reaction was allowed to develop for 30 min at RT. The optical density was measured at 405 nm using a Tecan GENios Pro plate reader. Antibody responses were expressed as titers. Antibody titers were determined via a linear fit for optical density values of nine dilutions and expressed in arbitrary units by calculating the reciprocal dilution that gave 50% of the maximum absorbance response. For measurement of Ig isotype specificity (IgG1, IgG2b, IgG2c, IgG3, IgA and IgM), an ELISA protocol similar to that described above was followed. Briefly, ELISA plates were coated as described above. Reference mouse IgG1, IgG2b, IgG2c, IgG3, IgA and IgM antibodies were diluted in TBST containing 1% skim milk. To detect specific binding, HRP-conjugated anti-mouse IgG1, IgG2b, IgG2c, IgG3, IgA and IgM antibodies (Southern Biotech) were used at a dilution of 1:500. TMB was then added as described above. The presence of antigen-specific isotype-specific antibodies was detected and measured as described above. Antibody levels were expressed in micrograms per milliliter.

Statistical Analysis

For comparison of results within experimental groups, a Student *t* test was performed. For all multigroup comparisons, one-way analysis of variance and post-hoc Bonferoni test for multiple comparisons were used. Statistical significance was defined as $p < .05$.

Human Cases and Brain Protein Extraction

Cases were obtained from the Albert Einstein College of Medicine human brain bank, Bronx, NY (see Table 1). Brain samples (Table 1) were processed as previously described (35). Briefly,

Table 1. Human cases analyzed in this study.

Case #	Clinical Dx	Brack	Age	Sex	Region
3665	Normal	0	85	Female	Mid-temporal cortex
4323	Normal	0	93	Female	Mid-temporal cortex
4359	Normal	0	92	Male	Mid-temporal cortex
4428	Normal	0	84	Female	Mid-temporal cortex
4860	Normal	0	92	Female	Mid-temporal cortex
4870	Normal	0	99	Female	Mid-temporal cortex
3410	AD	V-VI	81	Male	Mid-temporal cortex
4073	AD	V-VI	78	Female	Mid-temporal cortex
4137	AD	V-VI	88	Male	Mid-temporal cortex
5175	AD	V-VI	56	Female	Mid-temporal cortex
5288	AD	V-VI	81	Male	Mid-temporal cortex
5340	AD	V-VI	54	Male	Mid-temporal cortex

AD, Alzheimer's disease. Cases were obtained from the Albert Einstein College of Medicine human brain bank, Bronx, NY.

human brains were sequentially extracted to obtain a soluble SDS fraction and an insoluble formic acid (FA) fraction. Samples were first homogenized and sonicated in TBS containing 2% SDS and 1 \times complete protease inhibitor mixture (Roche Applied Science) and centrifuged at 100,000 \times g for 1 h at 4°C. The supernatant was removed and the resulting pellet was then extracted with 70% FA in water. FA extracts were dried under vacuum in a speed vacuum and then dissolved in dimethyl sulfoxide.

Western Blot

Protein extracts from SDS and FA fractions obtained from human brain samples were analyzed by SDS-PAGE using the indicated antibodies. Samples were electrophoresed on 10% Tris-HCl gels (phospho-tau and actin analysis) or on 16.5% Tris-Tricine gels (A β isoforms, BioRad) and transferred onto nitrocellulose membranes. A β was detected as previously described (36). Briefly, membranes were microwaved for 5 min in PBS. Membranes were then blocked in 5% fat-free milk in TBS, and incubated with primary antibodies overnight at 4°C. A standard enhanced chemiluminescence detection procedure was then used.

Immunohistochemistry

Five- μ m-thick sections of formalin-fixed paraffin-embedded brain tissue

were immunostained with 6E10 mAb (Covance, 1:1000 dilution) or A β pE3:CRM197 antiserum (1:2 dilution). Sections were deparaffinized by immersion in xylene and hydration through graded ethanol solutions. Denaturation for antigen recovery was performed by incubation of the slides in 70% FA for 30 min at RT. Denaturation was stopped by incubation in 100 mM Tris-HCl, pH 7.4, for 5 min. After washing once in TBS containing 0.05% Triton-X100 (TBSTx), endogenous peroxidase activity was inhibited by incubation in 5% hydrogen peroxide in TBSTx for 30 min at RT. After washing twice in TBSTx for 5 min and once in water, sections were blocked in 10% fat-free milk (6E10) or 5% normal goat serum (A β pE3:CRM197 antiserum) in TBSTx containing 1 mg/mL BSA and 1 mM NaF for 1 h at RT. Sections were then incubated in the presence of primary antibodies diluted in 10% fat-free milk (6E10) or 5% normal goat serum (A β pE3:CRM197 antiserum) in TBSTx containing 1 mg/mL BSA and 1 mM NaF overnight at 4°C in a humidified chamber. After washing, the sections were incubated with biotin-coupled anti-mouse IgG1 secondary antibodies (1:1,000 dilution in TBSTx with 20% Superblock, Pierce) before incubation with streptavidin-HRP (1:1,000 dilution in TBSTx with 20% Superblock, Southern Biotech) and visualization with diaminobenzidine tetrahydrochloride.

RESULTS

Immunogenicity of CRM197 in Mice

We first verified that unconjugated CRM197 without adjuvant was immunogenic in C57BL/6J mice. Based on previously published dose-ranging immunogenicity analyses of different CRM197-based vaccines (34,37), we chose to inject 10 μ g of CRM197. A prime injection of unconjugated CRM197 followed by a boost injection at d 14 of the same amount of carrier protein elicited robust immunogenicity against CRM197 at d 30 (Figure 1A). Saline control




Figure 1. Anti-CRM197 antibody response in mice immunized with CRM197. (A) Specific anti-CRM197 Ig titers and (B) Ig isotype concentration (IgG1, IgG2b, IgG2c, IgG3, IgA and IgM) quantified by ELISA in sera obtained from C57BL/6J mice immunized (CRM197) or not (saline) with CRM197. Sera were collected 30 d after prime injection. Data are mean \pm standard error of the mean from six mice per group; each dot represents an individual mouse serum.

injections as expected did not elicit an anti-CRM197 antibody response (Figure 1A). Ig isotype determination revealed that anti-CRM197 antibodies were exclusively of the IgG1 isotype and reached levels of \sim 35 μ g/mL at d 30 post prime injection (Figure 1B).

Immunogenicity of A β pE3:CRM197 and A β pE11:CRM197 Conjugate Vaccines

A short peptide sequence of six residues starting at pE3 (pE3-8) or pE11 (pE11-16) linked in C-terminal to a three-residue spacer (see Materials and Methods) was chosen to facilitate the production of conformation/modification-specific antibodies directed against A β . A six-residue peptide epitope has also the advantage of being shorter than conventional T cell epitopes and thus might prevent unwanted T cell activation and detrimental proinflammatory response (38). Molecular weight analysis by protein staining after SDS-PAGE revealed that pE3-8 and pE11-16 peptides could be covalently conjugated to CRM197 (Figure 2A). The generated A β pE3:CRM197 and A β pE11:CRM197 conjugates (10 μ g) elicited robust antibody responses against A β pE3-8 and A β pE11-16 peptides, respectively, when assessed by ELISA (Figures 2B and 2D). Total serum Ig titers averaged 1,000 at d 30 after one prime injection and one boost injection at d 15 (Figures 2B and 2D). Importantly, the vaccines did not generate antibodies directed against the corresponding non-cyclized A β E3-8 and A β E11-16 peptides (Figures 2C and 2E), showing that immunogenicity was specific to the pE modification. No anti-A β pE3- or anti-A β pE11-specific antibodies were detected in control mice injected with unconjugated CRM197 or saline (Figures 2B and 2D). Injection of free A β pE3-8 and A β pE11-16 peptides (unconjugated to CRM197) did not elicit immunogenic responses against these peptides (Figures 2B and 2D).

As observed for the immunogenic responses to unconjugated CRM197 (Figure 1), the anti-A β pE3 and

anti-A β pE11 antibodies were exclusively of the IgG1 isotype (Figures 3A and 3B). At d 30 post prime injection, A β pE3:CRM197 and A β pE11:CRM197 vaccines elicited the production of \sim 1.5-2 μ g/mL of serum-specific IgG1.

The A β pE3:CRM197 Vaccine Produces Antibodies that React with Amyloid Deposits in AD Brains

Antiserum produced after immunization with the A β pE3:CRM197 vaccine labeled synthetic A β pE3-42 peptide by WB (Figure 4A). Importantly, A β pE3:CRM197 antiserum did not cross-react with synthetic A β pE11-42 or synthetic A β 1-42 (Figure 4A), showing that the produced antibodies did not recognize nonspecifically the pE modification, nor did they interact with an internal epitope containing residues 3–8 of A β 1-42. A β pE3:CRM197 antiserum is thus fully specific to A β pE3. Strikingly, A β pE3:CRM197 antiserum also labeled A β pE3 in insoluble/aggregated brain protein preparations (FA fractions, see Methods) obtained from six independent well-characterized AD patients (see Table 1 and Figure 4B). Almost no immunoreactivity was observed in the corresponding soluble preparations (SDS fractions, Figure 4B), indicating that the anti-A β pE3 antibodies produced by the A β pE3:CRM197 vaccine recognized human amyloid material in the AD brain, and that A β pE3 was mostly aggregated. Of note, A β pE3 immunoreactivity was also observed in normal control brains that contained high levels of aggregated total A β (also detected with 6E10 mAb in the insoluble fractions, Figure 4B). In fact, a strong parallel was observed between the levels of aggregated total A β and aggregated A β pE3 in normal and AD brains (Figure 4B), suggesting that A β pE3 is a marker for amyloid deposition but is not specific to AD.

In addition, A β pE3:CRM197 antiserum decorated amyloid plaques in human AD brain slices by IHC. Most of the anti-A β pE3 immunoreactivity was observed on plaques present in the cortex (entorhinal and parahippocampal




Figure 2. Anti-AβpE3-8 and anti-AβpE11-16 antibody response. (A) Coomassie blue staining of CRM197 alone or conjugated to AβpE3-8 (AβpE3-8:CRM197) or AβpE11-16 (AβpE11-16:CRM197). (B) Anti-AβpE3-8, (C) anti-Aβ3-8, (D) anti-AβpE11-16 and (E) anti-Aβ11-16 Ig titers in sera obtained from mice immunized with AβpE3-8 or AβpE11-16 conjugated (AβpE3-8:CRM197, AβpE11-16:CRM197) or not (AβpE3-8, AβpE11-16) with CRM197. Data represent mean ± standard error of the mean from three to nine mice per group. MW, molecular weight markers.

cortices, Figure 4C). Although amyloid plaque deposition was very pronounced in the AD case analyzed, only a few plaques were labeled with the AβpE3:CRM197 antiserum in the hippocampal formation (the dentate gyrus, CA1 and subiculum, Figure 4C). Interestingly, in adjacent sections of the parahippocampal cortex, identical plaques stained for both total Aβ (6E10 mAb) and AβpE3 (Figure 4C, arrows). This co-staining revealed that anti-AβpE3 antiserum preferentially decorated highly dense plaques. Of note, anti-IgG1 secondary antibodies were used in combination with AβpE3:CRM197 antiserum for WB and IHC, further showing that the immunoreactive anti-AβpE3 antibodies were of the IgG1 isotype.

DISCUSSION

In this study, we show that minimal epitopes can be designed to generate vaccines that are fully specific to the pyroglutamate modifications of Aβ. Indeed, antiserum obtained after immunization with the AβpE3:CRM197 conjugate vaccine demonstrated excellent immunoreactivity against synthetic and AD brain-derived AβpE3 with no detectable cross-reactivity with Aβ1-42, non-cyclized AβE3 or AβpE11. Further analysis using AD brain samples revealed that AβpE3:CRM197 antiserum mainly decorated highly aggregated amyloid material, supporting the notion that AβpE3 is associated with the dense core of senile plaques (39).

CRM197 is routinely used as a conjugate vaccine in licensed vaccines directed against bacterial capsular polysaccharides (32,33). The use of CRM197 for peptide epitope presentation has not yet been approved for human use, but preclinical evidence has already suggested that it has therapeutic potential (34). In addition, an experimental vaccine against Aβ using CRM197 is currently being evaluated in a phase II trial (ACC-001, Elan/J&J/Wyeth). In the ACC-001 vaccine, CRM197 is conjugated to the N-terminal residues 1-7 of Aβ and thus is aimed at targeting all Aβ isoforms containing this N-end of Aβ (40). Our study strengthens the notion that




Figure 3. Isotype specificity of the Ig response elicited by A β pE3-8:CRM197 and A β pE11-16:CRM197 vaccines. (A) Anti-A β pE3-8 and (B) anti-A β pE11-16 Ig isotype (IgG1, IgG2b, IgG2c, IgG3, IgA and IgM) concentrations in sera obtained from mice immunized with A β pE3-8 or A β pE11-16 conjugated (A β pE3-8:CRM197, A β pE11-16:CRM197) or not (A β pE3-8, A β pE11-16) with CRM197. Data represents mean \pm standard error of the mean from three to seven mice per group.

CRM197 has strong potential for peptide presentation. This work further demonstrates that CRM197 might particularly be useful for the generation of conformation/modification-specific anti-peptide vaccines.

Active and passive immunization directed against A β have their respective pros and cons (see Introduction), and there is no consensus yet as to whether one approach has a stronger therapeutic potential for AD. Both approaches, targeting different A β epitopes, are therefore actively investigated at the preclinical and clinical levels. Our approach was aimed at increasing the anti-A β immunotherapy toolkit by proposing an active immunization strategy specifically targeting N-terminally truncated pyroglutamate A β . Further studies will be required to validate the utility of such a vaccine in AD mouse models of amyloid deposition. It should be noted, however, that several studies using passive immunization specifically targeting A β pE3 have already been conducted in different mouse amyloid models. These studies provide evidence that anti-A β pE3 antibodies have an overall plaque-lowering effect and can improve the associated cognitive/behavioral deficits (29,41,42). Thus, it will be important to determine whether selective and active anti-A β pE3 immunization could also reduce the pathology in these models.

In this context, new immunization formulations for the A β pE3:CRM197 vaccine will have to be tested to boost antibody production. Indeed, although the A β pE3:CRM197 antiserum was able to react with both synthetic A β pE3 and AD brain-derived amyloid material, its antibody levels in the serum, around 2 μ g/mL, are likely to be too low to allow clearance of A β pE3 and amyloid in mouse AD models. Thus, future experiments will have to determine whether immunogenicity of the A β pE3:CRM197 vaccine can be improved by repeated injections and/or addition of adjuvants. Interestingly, recent data demonstrated that combined addition of the adjuvants aluminum hydroxide and CpG can increase the antibody titer of a nicotine:CRM197 conjugate vaccine more than 100-fold




Figure 4. AβpE3-8:CRM197 antiserum immunoreactivity in AD brains. (A) WB analysis of recombinant Aβ1-42, AβpE3-42 and AβpE11-42 using 6E10 (anti-Aβ1-16) and 4G8 (anti-Aβ17-24) antibodies, or AβpE3-8:CRM197 antiserum showing the specificity of the AβpE3-8:CRM197 antiserum toward AβpE3-42. (B) Western Blot analysis of human brain homogenates obtained from normal or AD cases (Braak stage V-VI, see Table 1) using PHF1 (anti-phospho-Ser396/404 tau), actin and 6E10 antibodies or AβpE3-8:CRM197 antiserum. (C) Immunohistochemistry of serial brain sections from an AD case stained with 6E10 antibody or AβpE3-8:CRM197 antiserum.

(43), and QS-21 elicited consistently higher anti-A β IgG titers in a phase IIa trial for the A β 1-7:CRM197 ACC-001 vaccine (44).

CONCLUSION

We propose that conjugation of peptides A β pE3-8 and A β pE11-16 to CRM197 can generate fully specific vaccines directed against A β pE3 and A β pE11, respectively. The main strength of this immunization strategy is to combine the advantages of a vaccine with the high specificity of targeting particularly amyloidogenic and neurotoxic subspecies of A β , while sparing the more abundant and maybe more physiologically relevant full-length A β 40 and A β 42.

ACKNOWLEDGMENTS

This work was supported in part by National Institutes of Health grant R01AG042508 (to PM).

DISCLOSURE

The authors declare they have no competing interests as defined by *Molecular Medicine* or other interests that might be perceived to influence the results and discussion reported in this paper.

REFERENCES

- Lemere CA. (2013) Immunotherapy for Alzheimer's disease: hoops and hurdles. *Mol. Neurodegener.* 8:36.
- Wisniewski T, Goñi F. (2015) Immunotherapeutic approaches for Alzheimer's disease. *Neuron.* 85:1162–76.
- Sterner RM, Takahashi PY, Yu Ballard AC. (2016) Active Vaccines for Alzheimer Disease Treatment. *J. Am. Med. Dir. Assoc.* 17:862.e811–65.
- Selkoe DJ. (2001) Alzheimer's disease: genes, proteins, and therapy. *Physiol. Rev.* 81:741–66.
- Checler F. (1995) Processing of the beta-amyloid precursor protein and its regulation in Alzheimer's disease. *J. Neurochem.* 65:1431–44.
- Marambaud P, Robakis NK. (2005) Genetic and molecular aspects of Alzheimer's disease shed light on new mechanisms of transcriptional regulation. *Genes Brain Behav.* 4:134–46.
- Citron M. (2010) Alzheimer's disease: strategies for disease modification. *Nat. Rev. Drug. Discov.* 9:387–98.
- Holmes C, et al. (2008) Long-term effects of Abeta42 immunisation in Alzheimer's disease: follow-up of a randomised, placebo-controlled phase I trial. *Lancet.* 372:216–23.
- Tabira T. (2010) Immunization therapy for Alzheimer disease: a comprehensive review of active immunization strategies. *Tohoku J. Exp. Med.* 220:95–106.
- Hampel H, et al. (2015) Advances in the therapy of Alzheimer's disease: targeting amyloid beta and tau and perspectives for the future. *Expert Rev. Neurother.* 15:83–105.
- Karran E, Hardy J. (2014) A critique of the drug discovery and phase 3 clinical programs targeting the amyloid hypothesis for Alzheimer disease. *Ann. Neurol.* 76:185–205.
- Golde TE. (2014) Open questions for Alzheimer's disease immunotherapy. *Alzheimers Res. Ther.* 6:3.
- De Strooper B, Vassar R, Golde T. (2010) The secretases: enzymes with therapeutic potential in Alzheimer disease. *Nat. Rev. Neurol.* 6:99–107.
- Mori H, Takio K, Ogawara M, Selkoe DJ. (1992) Mass spectrometry of purified amyloid beta protein in Alzheimer's disease. *J. Biol. Chem.* 267:17082–6.
- Saido TC, et al. (1995) Dominant and differential deposition of distinct beta-amyloid peptide species, A beta N3(pE), in senile plaques. *Neuron.* 14:457–66.
- Portelius E, et al. (2010) Mass spectrometric characterization of brain amyloid beta isoform signatures in familial and sporadic Alzheimer's disease. *Acta Neuropathol.* 120:185–93.
- Sevalle J, et al. (2009) Aminopeptidase A contributes to the N-terminal truncation of amyloid beta-peptide. *J. Neurochem.* 109:248–56.
- Schilling S, et al. (2008) Glutaminyl cyclase inhibition attenuates pyroglutamate Abeta and Alzheimer's disease-like pathology. *Nat. Med.* 14:1106–11.
- Piccini A, et al. (2005) Beta-amyloid is different in normal aging and in Alzheimer disease. *J. Biol. Chem.* 280:34186–92.
- Schlenzig D, et al. (2009) Pyroglutamate formation influences solubility and amyloidogenicity of amyloid peptides. *Biochemistry.* 48:7072–8.
- Nussbaum JM, et al. (2012) Prion-like behaviour and tau-dependent cytotoxicity of pyroglutamylation of amyloid- β . *Nature.* 485:651–5.
- He W, Barrow CJ. (1999) The A beta 3-pyroglutamyl and 11-pyroglutamyl peptides found in senile plaques have greater beta-sheet forming and aggregation propensities in vitro than full-length A beta. *Biochemistry.* 38:10871–7.
- Schilling S, et al. (2006) On the seeding and oligomerization of pGlu-amyloid peptides (in vitro). *Biochemistry.* 45:12393–9.
- Dammers C, et al. (2015) Structural Analysis and Aggregation Propensity of Pyroglutamate A β (3–40) in Aqueous Trifluoroethanol. *PLoS One.* 10:e0143647.
- Russo C, et al. (2002) Pyroglutamate-modified amyloid beta-peptides—AbetaN3(pE)—strongly affect cultured neuron and astrocyte survival. *J. Neurochem.* 82:1480–9.
- Schlenzig D, et al. (2012) N-terminal pyroglutamate formation of A β 38 and A β 40 enforces oligomer formation and potency to disrupt hippocampal long-term potentiation. *J. Neurochem.* 121:774–84.
- Alexandru A, et al. (2011) Selective hippocampal neurodegeneration in transgenic mice expressing small amounts of truncated A β is induced by pyroglutamate-A β formation. *J. Neurosci.* 31:12790–801.
- Gunn AP, et al. (2016) Amyloid- β Peptide A β 3pE-42 Induces Lipid Peroxidation, Membrane Permeabilization, and Calcium Influx in Neurons. *J. Biol. Chem.* 291:6134–45.
- Frost JL, et al. (2015) An anti-pyroglutamate-3 A β vaccine reduces plaques and improves cognition in APPswe/PS1 Δ E9 mice. *Neurobiol. Aging.* 36:3187–99.
- Bayer TA, Wirths O. (2014) Focusing the amyloid cascade hypothesis on N-truncated Abeta peptides as drug targets against Alzheimer's disease. *Acta Neuropathol.* 127:787–801.
- Cynis H, Frost JL, Crehan H, Lemere CA. (2016) Immunotherapy targeting pyroglutamate-3 A β : prospects and challenges. *Mol. Neurodegener.* 11:48.
- Shinefield HR. (2010) Overview of the development and current use of CRM(197) conjugate vaccines for pediatric use. *Vaccine.* 28:4335–9.
- Bröker M, Costantino P, DeTora L, McIntosh ED, Rappuoli R. (2011) Biochemical and biological characteristics of cross-reacting material 197 CRM197, a non-toxic mutant of diphtheria toxin: use as a conjugation protein in vaccines and other potential clinical applications. *Biologicals.* 39:195–204.
- Caro-Aguilar I, et al. (2013) Immunogenicity in mice and non-human primates of the Group A Streptococcal J8 peptide vaccine candidate conjugated to CRM197. *Hum. Vaccin. Immunother.* 9:488–96.
- Vingtdeux V, Davies P, Dickson DW, Marambaud P. (2011) AMPK is abnormally activated in tangle- and pre-tangle-bearing neurons in Alzheimer's disease and other tauopathies. *Acta Neuropathol.* 121:337–49.
- Vingtdeux V, et al. (2015) CALHM1 ion channel elicits amyloid- β clearance by insulin-degrading enzyme in cell lines and in vivo in the mouse brain. *J. Cell Sci.* 128:2330–8.
- Rondini S, et al. (2011) Evaluation of the immunogenicity and biological activity of the Citrobacter freundii Vi-CRM197 conjugate as a vaccine for Salmonella enterica serovar Typhi. *Clin. Vaccine Immunol.* 18:460–8.
- Rammensee HG. (1995) Chemistry of peptides associated with MHC class I and class II molecules. *Curr. Opin. Immunol.* 7:85–96.
- Sullivan CP, et al. (2011) Pyroglutamate-A β 3 and 11 colocalize in amyloid plaques in Alzheimer's disease cerebral cortex with pyroglutamate-A β 11 forming the central core. *Neurosci. Lett.* 505:109–12.
- Winblad B, Graf A, Riviere ME, Andreasen N, Ryan JM. (2014) Active immunotherapy options for Alzheimer's disease. *Alzheimers Res. Ther.* 6:7.

41. Wirths O, *et al.* (2010) Identification of low molecular weight pyroglutamate A[β] oligomers in Alzheimer disease: a novel tool for therapy and diagnosis. *J. Biol. Chem.* 285:41517–24.
42. Demattos RB, *et al.* (2012) A plaque-specific antibody clears existing β -amyloid plaques in Alzheimer's disease mice. *Neuron.* 76:908–20.
43. McCluskie MJ, *et al.* (2015) Anti-nicotine vaccines: Comparison of adjuvanted CRM197 and Qb-VLP conjugate formulations for immunogenicity and function in non-human primates. *Int Immunopharmacol.* 29:663–71.
44. Pasquier F, *et al.* (2016) Two Phase 2 Multiple Ascending-Dose Studies of Vanutide Cridificar (ACC-001) and QS-21 Adjuvant in Mild-to-Moderate Alzheimer's Disease. *J. Alzheimers Dis.* 2016;51(4):1131–43.

Cite this article as: Vingtdeux V, *et al.* (2016) A modification-specific peptide-based immunization approach using CRM197 carrier protein: Development of a selective vaccine against pyroglutamate A β peptides. *Mol. Med.* 22:841–9.