



Effects of coenzyme Q₁₀ on the antioxidant system in SD rats exposed to lipopolysaccharide-induced toxicity

Min-Hae Song¹, Ha-Na Kim¹, Yong Lim², In-Surk Jang^{1,*}

¹Department of Animal Science and Biotechnology, and the Regional Animal Research Center, Gyeongnam National University of Science and Technology, Jinju, Korea

²Department of Clinical Laboratory Science, Dong-Eui University, Busan, Korea

The study was performed to see the effects of coenzyme Q₁₀ (CoQ₁₀) on blood biochemical components and hepatic antioxidant system in rats exposed to lipopolysaccharide (LPS)-induced toxicity. A total of 24 rats were allocated to four groups: control (CON), 100 mg/kg BW of LPS (LPS), 100 mg of CoQ₁₀/kg BW with LPS (LCQI) and 300 mg of CoQ₁₀/kg BW with LPS (LCQII). The LPS and LCQI groups showed a significant ($P<0.05$) increase in the relative spleen weight compared with the CON group without affecting body and liver weights. The blood alanine aminotransferase (ALT) level in the LPS group was significantly ($P<0.05$) greater than that in the CON group, while supplementation with 100 or 300 mg CoQ₁₀ to rats injected with LPS normalized the ALT level in the CON group. In antioxidant systems, the LPS group showed a significantly ($P<0.05$) higher mRNA and activity of superoxide dismutase (SOD) than the CON group. The supplementation with CoQ₁₀ to the LPS-treated group normalized the level of SOD, which was comparable to the level of the CON group. Both the mRNA expression and activity of glutathione peroxidase in the LCQI and LCQII groups were higher ($P<0.05$) than that of the LPS group. However, administration of LPS or CoQ₁₀ unaffected the level of catalase and total antioxidant power. The level of lipid peroxidation in the LCQII group was lower ($P<0.05$) than that in the LPS group. In conclusion, CoQ₁₀ exerted its favorable effect against liver damage by modulation of antioxidant enzymes in LPS treated rats.

Keywords: Coenzyme Q₁₀, lipopolysaccharide, antioxidant enzymes, lipid peroxidation

Received 9 January 2017; Revised version received 22 February 2017; Accepted 16 March 2017

Coenzyme Q₁₀ (CoQ₁₀) is known to be responsible for the generation of ATP via the oxidative phosphorylation by transferring electrons of the respiratory chain, which exists in the mitochondrial membrane of organisms [1]. It is also revealed as a redox-active, antioxidant lipoprotein compound that is found in the phospholipid bilayer of cell membranes of tissues [2]. The biosynthesis of CoQ₁₀ takes place in the mitochondria of the liver, heart, kidneys and muscles, where they require a greater amount of energy for their multiple biological functions [3]. Because of its essential function in cellular tissues, CoQ₁₀ deficiency is a common disorder in certain

pathological conditions due to the process of cellular aging [4]. The beneficial effects of dietary CoQ₁₀ on clinical applications have been reported to ameliorate cardiovascular disease such as congestive heart failure, cardiomyopathy and mitochondrial disorder in humans [3-5]. Thus, it has been demonstrated that the supplementation with CoQ₁₀ effectively ameliorates degenerative disease via enhancing antioxidant properties in the elderly and patients [3,5].

The free radical scavenging properties of CoQ₁₀ have been known to serve as a crucial antioxidant activity to prevent lipid and membrane oxidation against pro-

*Corresponding author: In-Surk Jang, Department of Animal Science & Biotechnology, Gyeongnam National University of Science and Technology, Chilam-Dong 150, Jinju, Gyeongnam 52725, Korea
Tel: +82-55-751-3236; Fax: + 82-55-751-3267; E-mail: isjang@gntech.ac.kr

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/3.0>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

oxidative damage in tissues [7,8]. Hence, dietary supplementation with CoQ₁₀ has an effective tool of scavenging reactive oxygen species (ROS) due to its ability to quench singlet oxygen and peroxy radicals [9]. According to the reports of several literatures [10,11], the antioxidant capacity of CoQ₁₀ was much higher than vitamin C or vitamin E in human studies.

On the contrary, it has been reported that an intracellular deficiency of CoQ₁₀ can only happen under aging and certain pathological conditions, although the major biological function of CoQ₁₀ is directly associated with antioxidant properties under normal metabolic processes [12]. Thus, some studies reported that biological functions in response to dietary supplementation with CoQ₁₀ were only effective under abnormal physiological status, since an endogenous level of CoQ₁₀ was sufficient for maintaining metabolic process under normal conditions [3,5,13]. By way of example, it was reported that hepatic damage caused by endotoxin was ameliorated by supplementation with CoQ₁₀ [14]. This result might be attributed to the high antioxidant efficacy of CoQ₁₀ to protect cell integrity against ROS and lipid peroxidation induced by toxic agents [15]. However, the effectiveness of CoQ₁₀ as a protective role against oxidative injury varies greatly according to the physiological status of animals, health and disease of animals, concentration and period of dietary CoQ₁₀ supplementation, interaction with other nutrients, etc. [3,11,16]. Thus, there is still a lack of evidence about whether the administration of CoQ₁₀ to young animals during oxidative stress can affect the antioxidant system, despite the rather well-recognized antioxidant effects of CoQ₁₀ *in vitro*.

In the above context, the aim of this study was designed to examine the effects of dietary CoQ₁₀ on blood biochemical profiles, the mRNA expression and activity of antioxidant enzymes and antioxidant status in young rats exposed to early lipopolysaccharide-induced toxicity.

Materials and Methods

Experimental animals and procedures

Male Sprague Dawley (SD) rats aged 5-weeks were obtained from Samtako (Osan, Korea). After an acclimation of 2-weeks, all rats were kept in an environmentally controlled room (22±2°C) with a 12 h light/dark cycle. A total of twenty-four rats having a similar body weight were assigned to four treatments with free access to lab

chow and sterilized water *ad libitum*. Rats in group 1 (CON) were orally administrated soybean oil daily basis with saline injection; rats in group 2 (LPS) were orally given soybean oil with the intraperitoneal (i.p.) injection of lipopolysaccharide (LPS, 1 mg/BW on day 0, 3 and 6); rats in group 3 (LCQI) and 4 (LCQII) were orally given 100 mg and 300 mg of CoQ₁₀ (kg BW) dissolved into soybean oil daily basis, respectively with i.p. injection of LPS (1 mg/BW on day 0, 3 and 6). After that, body weight was measured on a weekly basis for a 4-week experimental period. A commercial CoQ₁₀ was purchased from Inter Mongolia Kingdomway Pharmaceutical Limited (CoQ₁₀ 99.3%, Xiamen, China). The CoQ₁₀ supplement was dissolved in soybean oil carrier, and LPS (Sigma-Aldrich, St. Louis, MO, USA) was diluted in sterile saline. The animal experiment was approved by the Institutional Animal Care and Use Committee (IACUC) at the Gyeongnam National University of Science and Technology in Korea.

Tissue harvesting

At the end of 4-week treatment with CoQ₁₀, all animals were deprived of diet for 12 h and then the rats (n=6) were sacrificed with ether. Immediately after opening the abdominal cavity, blood was collected in tubes coated with sodium heparin and organs was harvested and weighed. The plasma and liver tissues were then rapidly frozen in liquid nitrogen and stored at -70°C until further assay.

Plasma biochemical composition analyses

Plasma aspartate aminotransferase (AST), alanine aminotransferase (ALT), blood urea nitrogen (BUN), glucose, and triglyceride were assayed using a clinical biochemical analyzer (Mindray, BS-120, Mindray Bio Medical Electronics co., Shenzhen, China). All analyses were conducted in duplicate.

mRNA expression of antioxidant enzymes by real time-PCR

The mRNA expression of antioxidant enzymes including superoxide dismutase (SOD), glutathione peroxidase (GPX) and catalase (CAT) was quantified by real-time PCR (Bio-rad, CA, USA). The cDNA primers used to amplify antioxidant genes are presented in Table 1. Total RNA of liver tissues was extracted using RNAzol TM B (Tel-Test Inc, TX, USA). In brief, 2 mL of RNAzol solution was added to 100 mg of hepatic tissues and then

Table 1. Primers used for the quantification of mRNA using real time-PCR

Genes	Primer sequences	Product size (bp)	Gene bank Accession No.
SOD	5'-ACT TCG AGC AGA AGG CAA GC-3' 5'-GTC TCC AAC ATG CCT CTC TTC AT-3'	194	NM017050.1
GPX	5'-CAG TTC GGA CAT CAG GAG AAT-3' 5'-AGA GCG GGT GAG CCT TCT-3'	139	NM030826.3
CAT	5'-ATT GCC GTC CGA TTC TCC-3' 5'-CCA GTT ACC ATC TTC AGT GTA G-3'	105	NM012520.2
GADPH	5'-ATG GAG AAG GCT GGG GCT CAC CT-3' 5'-AGC CCT TCC ACG ATG CCA AAG TTG-3'	199	NM017008.4

the tissues were homogenized using liquid nitrogen. After that, 100 μ L of chloroform was added to an Eppendorf-tube and incubated for 5 min on ice. Next, the aqueous phase was separated by centrifugation (15 min at 15,000 \times g), after which the isolated RNA was precipitated with the same volume of isopropanol and then centrifuged (15 min at 15,000 \times g). The isolated total RNA was then washed with 75% ethyl alcohol, dried, and diluted with diethylpyrocarbonate (DEPC)-treated H₂O, after which the optical density at 260 nm was applied to quantify the concentration of RNA (GeneQuant *pro* spectrophotometer, Amersham, Piscataway, NJ, USA). After that, 5 μ g of total RNA were incubated with 1.0 mg of oligo dT (Invitrogen Co., Carlsbad, CA, USA) at 70°C for 5 min and 4°C for 5 min to produce the first strand cDNA. The reaction mixture was incubated at 42°C for 50 min, 90°C for 10 min and 42°C for 50 min in a reaction cocktail containing 5 \times first strand buffer, 2.5 mM dNTP, 0.1 M DTT, superscript III and RT-mixture. Quantification of the antioxidant enzymes was conducted by real-time quantitative PCR with SYBR green supermix (BioRad, CA, USA) under the following conditions: 5 min at 95°C, followed by 40 cycles of denaturation at 95°C for 15 s and annealing at 60°C for 30 sec and then extension at 72°C for 30 sec. The PCR amplification cycle at which dye fluorescence passed the selected baseline (Ct) was determined by real-time monitoring. The expression of all mRNAs was calculated by the 2^[$\Delta\Delta$] method [17] to see relative changes in gene expression using GADPH as an internal control.

Antioxidant enzyme activity, lipid peroxidation and total antioxidant capacity

All of the isolation procedure was performed according to the method of Kupfer and Levin [18] to harvest cytosol and microsome fractions from the liver. Liver tissues (1 g) were homogenized with a solution containing 0.25 M sucrose, 0.05 M Tris-HCl (pH 7.4), 0.005 M

MgCl₂, 0.025 M KCl and 0.008 M CaCl₂ using a grass-grass homogenizer. The hepatic homogenate was centrifuged at 10,000 \times g for 15 min, after which time the resulting supernatant was diluted 1: 6 volume with a solution composed of 0.0125 M sucrose, 0.005 M MgCl₂, 0.025 M KCl and 0.008 M CaCl₂. And then diluted supernatant was centrifuged at 1,500 \times g for 10 min, after which time the resulting supernatant was harvested as cytosol fraction. The pellet was dispersed in 0.25 M sucrose was centrifuged at 1,500 \times g for 10 min. After that the remnant pellet was suspended in a cold 1.15% KCl solution to harvest a microsomal fraction. The harvested supernatant (cytosol) and a suspended pellet (microsomes) were frozen in liquid nitrogen and stored at -70°C until further assay. In brief, SOD activity in the cytosol fraction was measured using a commercial SOD assay kit (Sigma-Aldrich, St. Louis, MO, USA) based on an indirect assay method of xanthine oxidase as described in the manufacture's protocol. The activity of SOD is presented as units/mg of proteins, where 1 unit of activity was the amount of enzyme required to inhibit 50% of the SOD or SOD like substances. GPX was determined at 37°C in the cytosol with cumene hydroperoxide as a substrate [19]. The GPX coupled the reduction of cumene hydroperoxide to the oxidation of NADPH by glutathione reductase, and the subsequent oxidation rate was monitored with a UV-spectrophotometer based on the decrease in absorbance at 340 nm. One unit of GPX is presented as the amount of GPX required to oxidize 1 mmol of NADPH per minute. Catalase (CAT) activity was measured by the amount of hydrogen peroxide (H₂O₂), which is produced by CAT. The decomposition of the substrate was measured using a spectrophotometer at 240 nm, and one unit activity was expressed as moles of H₂O₂ consumed per minute per mg protein [20]. The lipid peroxidation concentration in the microsomes was assessed by measuring thiobarbituric acid (TBA) substances [21]. Absorbance was determined

using spectrophotometer at 532 nm, and TBA substances are expressed as nmol of malondialdehyde (MDA) per mg of protein. The plasma level of total antioxidant power was assayed using a commercially available assay kit with an ELISA reader (Oxford Biomedical Research, Inc., MI, USA). Assay procedures were carried out according to the manufacturer's protocols. Trolox was used to generate a standard curve, and data were presented as mM Trolox equivalents or in μM copper reducing equivalents. Protein concentration was analyzed by the BCA method (Pierce Assay) using ELISA reader (V_{Max} , Molecular Devices, CA, USA).

Statistical analysis

All values are expressed as means \pm standard deviation (SD). Statistical analyses were performed using Proc GLM (SAS Institute Inc.). When the treatment effect was significant at $P<0.05$, Duncan multiple range test was performed to assess significant differences among groups. A P value of <0.05 was considered statistically significant.

Results

Body and organ weights and plasma biochemical profiles

The effects of oral administration of CoQ₁₀ on body and immune organ weights and plasma biochemical

profiles in LPS-injected rats are presented in Table 2 and 3, respectively. There were no significant effects of LPS challenge and CoQ₁₀ administration on body weight, gain and the relative liver weight among groups (Table 2). However, the relative spleen weight in the LPS and LCQI groups significantly ($P<0.05$) increased compared with that in the CON group.

The plasma ALT level in the LPS group was significantly ($P<0.05$) higher than that in the CON group, while oral administration of 100 mg (LCQI) and 300 mg CoQ₁₀ (LCQII) to LPS-injected rats did not increase in ALT level compared with the CON group. The blood triglyceride level in the LPS group was significantly ($P<0.05$) lower than that in LCQI group. However, the level of other biochemical parameters including AST, BUN, creatinine and total cholesterol did not differ among treatment groups (Table 3).

mRNA expression and activity of antioxidant enzymes in the liver

The effects of orally given CoQ₁₀ on the mRNA expression and activities of SOD, GPX and CAT is shown in Table 4 and Figure 1, respectively. The LPS group showed a significantly ($P<0.05$) higher mRNA expression of SOD than the CON group. However, the oral supplementation with CoQ₁₀ to rats challenged with LPS significantly ($P<0.05$) alleviated SOD mRNA expression level, which was comparable to the CON

Table 2. Effect of coenzyme Q₁₀ on the weights of the body, liver and spleen of SD rats challenged with LPS

Item	Treatments*			
	CON	LPS	LCQI	LCQII
Initial BW (g), 7 wks	208.58 \pm 6.15	208.41 \pm 6.09	216.76 \pm 17.48	226.21 \pm 25.07
Final BW (g), 11 wks	372.84 \pm 12.26	370.46 \pm 17.26	371.54 \pm 13.64	380.00 \pm 14.67
Gain (g)	164.27 \pm 11.72	162.07 \pm 14.61	155.76 \pm 22.39	153.77 \pm 18.70
	370.46 \pm 17.26	370.46 \pm 17.26	370.46 \pm 17.26	370.46 \pm 17.26
Liver weight (g/100 g BW)	2.88 \pm 0.20	2.94 \pm 0.22	3.07 \pm 0.12	2.96 \pm 0.26
Spleen weight (g/100 g BW)	0.23 \pm 0.03 ^b	0.38 \pm 0.05 ^a	0.32 \pm 0.03 ^a	0.28 \pm 0.03 ^{ab}

*CON (saline), LPS (1 mg LPS/kg BW), LCQI (1 mg LPS/kg BW+100 mg CoQ₁₀) and LCQII (1 mg LPS/kg BW+300 mg CoQ₁₀).

Table 3. Effect of coenzyme Q₁₀ on the plasma biochemical components of SD rats challenged with LPS

Item	Treatments*			
	CON	LPS	LCQI	LCQII
AST (U/L)	110.00 \pm 14.18	110.80 \pm 9.42	106.40 \pm 5.55	95.20 \pm 18.27
ALT (U/L)	53.40 \pm 4.10 ^b	65.40 \pm 7.89 ^a	58.60 \pm 6.27 ^{ab}	58.60 \pm 6.27 ^{ab}
BUN (mg/dL)	11.66 \pm 1.46	10.26 \pm 1.17	11.76 \pm 1.42	10.72 \pm 1.77
Creatinine (mg/dL)	0.54 \pm 0.06	0.56 \pm 0.06	0.52 \pm 0.05	0.52 \pm 0.05
Triglyceride (mg/dL)	63.40 \pm 15.77 ^{ab}	50.00 \pm 10.46 ^b	89.00 \pm 34.48 ^a	83.20 \pm 33.88 ^{ab}
Total cholesterol (mg/dL)	71.90 \pm 7.50	71.28 \pm 15.52	84.70 \pm 11.07	75.58 \pm 9.95

*CON (saline), LPS (1mg LPS/kg BW), LCQI (1 mg LPS/kg BW+100 mg CoQ₁₀) and LCQII (1 mg LPS/kg BW+300 mg CoQ₁₀).

Table 4. Effect of coenzyme Q₁₀ on the mRNA expression of antioxidant enzymes (SOD, GPX and CAT) in the liver of SD rats challenged with LPS

	Treatments*							
	CON		LPS		LCQI		LCQII	
	ΔCt	$2^{-\Delta\Delta Ct}$	ΔCt	$2^{-\Delta\Delta Ct}$	ΔCt	$2^{-\Delta\Delta Ct}$	ΔCt	$2^{-\Delta\Delta Ct}$
SOD	7.90±0.95 ^a	1	6.54±0.42 ^b	2.55	7.38±0.67 ^{ab}	1.44	7.24±0.31 ^{ab}	1.58
GPX	8.90±0.36 ^{ab}	1	9.10±0.75 ^a	0.87	7.62±0.65 ^c	2.43	7.68±0.86 ^{bc}	2.43
CAT	5.20±0.36	1	4.80±0.58	1.32	5.44±0.28	0.85	5.00±0.81	1.15

*CON (saline), LPS (1 mg LPS/kg BW), LCQI (1 mg LPS/kg BW+100 mg CoQ₁₀) and LCQII (1 mg LPS/kg BW+300 mg CoQ₁₀). The values are ΔCt , which is represented as the Ct of each target gene corrected by Ct of the control gene (GADPH). The fold difference in the relative expression of the target gene was calculated as the $2^{-\Delta\Delta Ct}$. Means (Mean±SD, n=6) with different superscript differ among groups ($P<0.05$).

group (Table 4). The specific activity of SOD showed the same pattern of mRNA expression of SOD as depicted in Figure 1(A). The supplementation with CoQ₁₀ to the LPS-treated group normalized the SOD expression, showing that the LCQI and the LCQII groups showed a similar SOD expression in comparison with the CON group. The mRNA expression of GPX in the LPS group resulted in a significant ($P<0.05$) decrease compared with that in the LCQI and LCQII groups (Table 4). Similar to the mRNA expression of GPX, the specific activity of GPX in the LCQI and LCQII groups was also significantly greater ($P<0.05$) than that of the LPS group (Figure 2). However, the mRNA expression and activity of CAT were not affected by the administration of LPS or CoQ₁₀ in the liver of SD rats (Table 4, Figure 1).

Total antioxidant power and lipid peroxidation in the blood and liver

The influence of a dietary CoQ₁₀ on the level of lipid peroxidation and total antioxidant power (TAP) are presented in Figure 2. The concentration of lipid peroxidation as indicated by MDA level in the LCQII group was significantly ($P<0.05$) lower than that in the LPS group. Thus, the hepatic lipid peroxidation level was significantly ($P<0.05$) ameliorated by a dietary supplementation of 300 mg of CoQ₁₀ to the LPS-treated rats (Figure 2A), although there was no significant difference in MDA level between the LPS-treated rats and the LPS-treated rats that were given 100 mg of CoQ₁₀ supplements.

Plasma TAP in the LPS group numerically decreased compared with that in the other groups without statistical significance. Thus, the administration of CoQ₁₀ to rats challenged with LPS did not affect the level of TAS in this study (Figure 2B).




Figure 1. Specific activity of antioxidant enzymes (A: SOD, B: GPX and C: CAT) in the liver of SD rats administrated saline (CON), LPS injection (LPS), 100 mg of CoQ₁₀/kg BW with injection of LPS (LCQI) and 300 mg of CoQ₁₀/kg BW with injection of LPS (LCQII). Means (Mean±SD, n=6) with different superscript differ among groups ($P<0.05$).

Discussion

CoQ₁₀, a natural ubiquinone synthesized in the membrane




Figure 2. The levels of MDA (A) in microsomal fraction of the liver and total antioxidant power (B) in plasma of SD rats administrated saline (CON), LPS injection (LPS), 100 mg of CoQ₁₀/kg BW with injection of LPS (LCQI) and 300 mg of CoQ₁₀/kg BW with injection of LPS (LCQII). Means (Mean±SD, n=6) with different superscript differ among groups ($P<0.05$).

of the liver, heart, kidneys and muscles, is known to be a redox-active lipophilic antioxidant, scavenging ROS and inhibiting lipid peroxidation in the body [3,10]. The ability of CoQ₁₀ to scavenge ROS is much greater than vitamin E and it was also known to facilitate the absorption of lipid soluble antioxidant vitamins such as vitamin A and E [10,11].

In this study, the possible protective effect of CoQ₁₀ against hepatic oxidative damage in young rats challenged with LPS was investigated. As expected, the LPS challenge significantly increased the levels of blood ALT and spleen weight in rats. The leakage of hepatic enzymes including ALT and AST into the blood stream was directly associated with marked liver injury [22]. Our results were in agreement with earlier studies that LPS challenge resulted in a significant increase in blood AST and ALT in pigs and rodents [23,24]. Interestingly, CoQ₁₀ administration to rats challenged with LPS in this study significantly normalized LPS-induced increases in ALT activity and the relative spleen weight of rats. In accordance with this study, CoQ₁₀ significantly normalized the levels of serum AST and ALT activities in several studies, indicating that dietary CoQ₁₀ might be associated with alleviating liver damage induced by toxic agents

[25,26].

In general, it was reported that hepatic oxidative stress induced by LPS was closely associated with the generation of ROS in the liver [27]. Therefore, the mRNA expression and activity of antioxidant enzymes, lipid peroxidation and total antioxidant capacity were analyzed to investigate the effects of CoQ₁₀ against the oxidative liver damage of rats challenged with LPS in the study. From our observation, the level of hepatic SOD mRNA and activity in rats challenged with LPS resulted in a marked increase in comparison with the CON rats. However, the administration of CoQ₁₀ to LPS-treated rats showed a similar level of mRNA and activity of SOD compared with the CON rats, indicating that supplementation with CoQ₁₀ normalized an increased level of SOD expression in the LPS group. Furthermore, the administration of CoQ₁₀ to LPS-treated rats markedly increased GPX mRNA expression and activity compared with the LPS treated rats. However, the administration of CoQ₁₀ did not affect hepatic lipid peroxidation and total antioxidant capacity in LPS or CoQ₁₀ treated rats. Thus, it is postulated that CoQ₁₀ has protective or alleviative effects against LPS-induced oxidative stress in the hepatic tissue of rats by a free radical scavenging capacity through antioxidant defense mechanism.

It has been well recognized that excess ROS are eliminated by antioxidant enzymes, which are represented by SOD, GPX and CAT in the body. SOD directly converts superoxide anion to hydrogen peroxide [28], thereafter both GPX and CAT, the two main enzymes, are involved in the detoxification of H₂O₂ in a cellular antioxidant defense system [29]. Numerous studies have demonstrated that CuZn-SOD level is up-regulated in patients suffering from hepatitis, renal failure, diabetes and uremia [30,31]. LPS, a cell wall components of gram negative bacteria, seems to induce oxidative stress in the liver, which was involved in the hepatic injury of laboratory animals. A study [32] reported that LPS activated macrophages to induce numerous inflammatory ROS including superoxide anion, which was responsible for the activation of SOD expression to eliminate these radicals in the liver of rats exposed to enterotoxin [33]. In accordance with the result of our study, a study [34] demonstrated that hepatic SOD mRNA expression and activity were much greater from the LPS-treated rats than that from the control rats.

According to a recent report [8], dietary supplementation with CoQ₁₀ resulted in a significant decrease

in LPS-induced ROS production and a significant improvement of mitochondrial functions *in vitro* study. There are numerous reports that the application of CoQ₁₀ to laboratory animals, which were treated with toxic substances such as LPS, could help to maintain a normal level of antioxidant enzymes and antioxidant status in the liver [2,10,35]. In addition, CoQ₁₀ supplementation in young mice significantly increased the endogenous level of vitamin E in the plasma, heart and liver [7], suggesting that the elevated antioxidant potential of CoQ₁₀ is partially associated with increasing level of vitamin E. In partial agreement with the results of this study, alloxan-induced diabetes rats orally administered CoQ₁₀ (15 mg/kg of BW, i.p) showed significantly increased the serum level of glutathione and activities of SOD and CAT compared with diabetic rats [36]. Our findings also indicate that the administration of CoQ₁₀ at the level of above 100 mg/kg BW has a potent alleviating effect against LPS-induced oxidative stress in rats.

On the other hand, contradictory studies concerning the effects of CoQ₁₀ on antioxidant defense system have been reported [16,37]. Dietary CoQ₁₀ did not directly affect the changes in antioxidant enzymes including SOD, GPX and CAT as well as in the life span of laboratory rats [16,37]. However, the antioxidant properties of CoQ₁₀ may be attributed to potentiating electron transport chain where it plays a crucial role in electron donor and acceptor in the mitochondria of the liver [15]. This discrepancy might be due to differences in factors such as age of laboratory animals, dosage and duration of CoQ₁₀ level and dietary formulation [7,16].

In general, however, it is clear that dietary supplementation with CoQ₁₀ attenuates hepatic oxidative injury via the modulation of antioxidant enzymes and their antioxidant capacities under severe oxidative stress circumstance [8,36], although the detailed antioxidant mechanisms of CoQ₁₀ were not fully elucidated. In agreement with this hypothesis, our study also demonstrated that supplementation with CoQ₁₀ in rats challenged with LPS maintained the same level of SOD, GPX and lipid peroxidation in comparison with the control rats. Therefore, supplementation with CoQ₁₀ appeared to have potentiating effects on the antioxidant defense mechanism by directly scavenging ROS through the induction of antioxidant enzymes.

Overall, it could be proposed that CoQ₁₀ may exert its favorable effect against liver damage by the modulation of antioxidant enzymes in LPS-treated rats. Therefore,

this study suggests that dietary CoQ₁₀ has beneficial effects on antioxidant defense system in laboratory animals under severe oxidative stress conditions.

Acknowledgments

This research was supported by a grant from Gyeongnam National University of Science and Technology in 2016. The authors also thank the Regional Animal Research Center at GNTECH to use the analysis instruments.

Conflict of interests The authors declare that there is no financial conflict of interests to publish these results.

References

- Littarru GP. Energy and Defense. Facts and perspectives on coenzyme Q₁₀ in biology and medicine. Casa Editrice Scientifica Internazionale 1995. Roma.
- Quiles JL, Ochoa JJ, Battino M, Gutierrez-Rios P, Nepomuceno EA, Frías ML, Huertas JR, Mataix J. Life-long supplementation with a low dosage of coenzyme Q₁₀ in the rat: effects on antioxidant status and DNA damage. *Biofactors* 2005; 25: 73-86.
- Acosta MJ, Vazquez Fonseca L, Desbats MA, Cerqua C1, Zordan R, Trevisson E, Salviati L. Coenzyme Q biosynthesis in health and disease. *Biochim Biophys Acta* 2016; 1857(8): 1079-1085.
- Turunen M, Olsson J, Dallner G. Metabolism and function of coenzyme Q. *Biochim Biophys Acta* 2004; 1660: 171-199.
- Littarru GP, Tian L, Belardinelli R, Watts GF. Coenzyme Q(10), endothelial function, and cardiovascular disease. *Biofactors* 2011; 37(5): 366-373.
- Yang YK, Wang LP, Chen L, Yao XP, Yang KQ, Gao LG, Zhou XL. Coenzyme Q₁₀ treatment of cardiovascular disorders of ageing including heart failure, hypertension and endothelial dysfunction. *Clin Chim Acta* 2015; 450: 83-89.
- Kamzalov S, Sumien N, Forster MJ, Sohal RS. Coenzyme Q intake elevates the mitochondrial and tissue levels of Coenzyme Q and alpha-tocopherol in young mice. *J Nutr* 2003; 133(10): 3175-3180.
- Bullón P, Román-Malo L, Marín-Aguilar F, Alvarez-Suarez JM, Giampieri F, Battino M, Cordero MD. Lipophilic antioxidants prevent lipopolysaccharide-induced mitochondrial dysfunction through mitochondrial biogenesis improvement. *Pharmacol Res* 2015; 91: 1-8.
- James AM, Smith RA, Murphy MP. Antioxidant and prooxidant properties of mitochondrial Coenzyme Q. *Arch Biochem Biophys* 2004; 423(1): 47-56.
- Niklowitz P, Menke T, Andler W, Okun JG. Simultaneous analysis of coenzyme Q₁₀ in plasma, erythrocytes and platelets: comparison of the antioxidant level in blood cells and their environment in healthy children and after oral supplementation in adults. *Clin Chim Acta* 2004; 342: 219-226.
- Shekelle P, Morton S, Hardy ML. Effect of supplemental antioxidants vitamin C, vitamin E, and coenzyme Q₁₀ for the prevention and treatment of cardiovascular disease. *Evid Rep Technol Assess (Summ)* 2003; 83: 1-3.
- Rötig A, Appelkvist EL, Geromel V, Chretien D, Kadhom N, Edery P, Lebeideau M, Dallner G, Munnich A, Ernster L, Rustin P. Quinone-responsive multiple respiratory-chain dysfunction due to widespread coenzyme Q₁₀ deficiency. *Lancet* 2000; 356(9227): 391-395.
- Novoselova EG, Lunin SM, Novoselova TV, Khrenov MO,

- Glushkova OV, Avkhacheva NV, Safronova VG, Fesenko EE. Naturally occurring antioxidant nutrients reduce inflammatory response in mice. *Eur J Pharmacol* 2009; 615: 234-240.
14. Maruyama H, Furukawa K, Onda M. Effect of coenzyme Q₁₀ on endotoxin induced hepatocyte injury modulation of endotoxin-activated polymorphonuclear neutrophils. *Nihon Ika Daigaku Zasshi* 1995; 62(3): 271-282.
 15. Frei B, Kim MC, Ames BN. Ubiquinol-10 is an effective lipid-soluble antioxidant at physiological concentrations. *Proc Natl Acad Sci U S A* 1990; 87(12): 4879-4883.
 16. Sohal RS, Kamzalov S, Sumien N, Ferguson M, Rebrin I, Heinrich KR, Forster MJ. Effect of coenzyme Q₁₀ intake on endogenous coenzyme Q content, mitochondrial electron transport chain, antioxidative defenses, and life span of mice. *Free Radic Biol Med* 2006; 40(3): 480-487.
 17. Livak KJ, Schmittgen TD. Analysis of relative gene expression data using real-time quantitative PCR and the 2(-Delta Delta C(T)) Method. *Methods* 2001; 25(4): 402-408.
 18. Kupfer D, Levin E. Monooxygenase drug metabolizing activity in CaCl₂-aggregated hepatic microsomes from rat liver. *Biochem Biophys Res Commun* 1972; 47(3): 611-618.
 19. Tappel AL. Glutathione peroxidase and hydroperoxides. *Methods in Enzymol* 1978; 52: 506-513.
 20. Melega S, Canistro D, De Nicola GR, Lazzeri L, Sapone A, Paolini M. Protective effect of Tuscan black cabbage sprout extract against serum lipid increase and perturbations of liver antioxidant and detoxifying enzymes in rats fed a high-fat diet. *Br J Nutr* 2013; 110(6): 988-997.
 21. Bidlack WR, Tappel AL. Damage to microsomal membrane by lipid peroxidation. *Lipids* 1973; 8(4): 177-182.
 22. Giannini E, Botta F, Fasoli A, Ceppa P, Risso D, Lantieri PB, Celle G, Testa R. Progressive liver functional impairment is associated with an increase in AST/ALT ratio. *Dig Dis Sci* 1999; 44(6): 1249-1253.
 23. Li Q, Liu Y, Che Z, Zhu H, Meng G, Hou Y, Ding B, Yin Y, Chen F. Dietary L-arginine supplementation alleviates liver injury caused by *Escherichia coli* LPS in weaned pigs. *Innate Immun* 2012; 18(6): 804-814.
 24. El-Tanbouly DM, Abdelsalam RM, Attia AS, Abdel-Aziz MT. Pretreatment with magnesium ameliorates lipopolysaccharide-induced liver injury in mice. *Pharmacol Rep* 2015; 67(5): 914-920.
 25. Ali SA, Faddah L, Abdel-Baky A, Bayoumi A. Protective effect of L-carnitine and coenzyme Q₁₀ on CCl₄-induced liver injury in rats. *Sci Pharm* 2010; 78(4): 881-896.
 26. Fouad AA, Jresat I. Hepatoprotective effect of coenzyme Q₁₀ in rats with acetaminophen toxicity. *Environ Toxicol Pharmacol* 2012; 33(2): 158-167.
 27. Bian K, Murad F. Diversity of endotoxin-induced nitrotyrosine formation in macrophage-endothelium-rich organs. *Free Radic Biol Med* 2001; 31(4): 421-429.
 28. Fridovich I. Superoxide dismutase. *Enzymology* 1974; 41: 36-40.
 29. Matés JM. Effects of antioxidant enzymes in the molecular control of reactive oxygen species toxicology. *Toxicology* 2000; 153: 83-104.
 30. Chauhan DP, Gupta PH, Nampoothiri MR, Singhal PC, Chugh KS, Nair CR. Determination of erythrocyte superoxide dismutase, catalase, glucose-6-phosphate dehydrogenase, reduced glutathione and malonyldialdehyde in uremia. *Clin Chim Acta* 1982; 123: 153-159.
 31. Nishimura N, Ito Y, Adachi T, Hirano K, Sugiura M, Sawaki S. Enzyme immunoassay for cuprozinc-superoxide dismutase in serum and urine. *J Pharmacobiodyn* 1982; 5(6): 394-402.
 32. Rossol M, Heine H, Meusch U, Quandt D, Klein C, Sweet MJ, Hauschildt S. LPS-induced cytokine production in human monocytes and macrophages. *Crit Rev Immunol* 2011; 31(5): 379-446.
 33. Chou ST, Peng HY, Hsu JC, Lin CC, Shih Y. *Achillea millefolium* L. essential oil inhibits LPS-induced oxidative stress and nitric oxide production in RAW 264.7 Macrophages. *Int J Mol Sci* 2013; 14(7): 12978-12993.
 34. Spolarics Z. Endotoxin stimulates gene expression of ROS-eliminating pathways in rat hepatic endothelial and Kupffer cells. *Am J Physiol* 1996; 270: G660-666.
 35. Sena CM, Nunes E, Gomes A, Santos MS, Proença T, Martins MI, Seica RM. Supplementation of coenzyme Q₁₀ and alpha-tocopherol lowers glycated hemoglobin level and lipid peroxidation in pancreas of diabetic rats. *Nutr Res* 2008; 28(2): 113-121.
 36. Ahmadvand H, Tavafi M, Khosrowbeygi A. Amelioration of altered antioxidant enzymes activity and glomerulosclerosis by coenzyme Q₁₀ in alloxan-induced diabetic rats. *J Diabetes Complications* 2012; 26(6): 476-482.
 37. Lee CK, Pugh TD, Klopp RG, Edwards J, Allison DB, Weindruch R, Prolla TA. The impact of alpha-lipoic acid, coenzyme Q₁₀ and caloric restriction on life span and gene expression patterns in mice. *Free Radic Biol Med* 2004; 36(8): 1043-1057.